

3-Oxabicyclo[3,2,0]hepta-1,4-diene

By R. G. BERGMAN* and K. PETER C. VOLLHARDT*

(Gates and Crellin Laboratories of Chemistry, California Institute of Technology, Pasadena, California 91109)

Summary 3-Oxabicyclo[3,2,0]hepta-1,4-diene (**3**) has been synthesized by partial hydrogenation of 3-oxabicyclo[3,2,0]hepta-1,4,6-triene (**2**).


HETERO[5]ANNULENES fused to four-membered rings have become molecules of considerable current interest¹⁻³ since the strain in such systems is increased when compared with the corresponding benzene analogues and the chemical and physical properties of the aromatic ring are expected to be unusual.⁴ 3-Oxabicyclo[3,2,0]hepta-1,4-diene (**3**), the first oxygen analogue in this series, can also, in analogy to the corresponding sulphur compounds¹⁻³ [*e.g.*, (**4**) and (**5**)], be

used as a suitable model with which to compare the properties of furocyclobutadiene (**2**)⁵ and related systems.¹

Oxabicyclo[3,2,0]hepta-1,4,6-triene (**2**),⁵ was hydrogenated with reduced PtO₂ (-65°, 2 h) to give (**3**) (10%, g.l.c.) as an unstable colourless liquid, τ (CCl₄) 3.12br (1H, s, H_A) and 6.86br (2H, s, H_B) λ_{\max} (ether), end absorption only; *m/e* 94 (*M*⁺), 66 (*M* - CO, 25%).

Although more reactive than furan, compound (**3**) appears to be much less sensitive (*e.g.*, to oxygen and g.l.c.) than its potentially anti-aromatic precursor (**2**). The physical properties of (**3**) also differ dramatically from those of (**2**). Whereas the chemical shift [τ (CCl₄) 3.89] of H_A in (**2**) is at relatively high field, it is in (**3**) (τ 3.12) that expected for a 3,4-dialkyl-substituted furan,⁶ the chemical shift difference (0.77 p.p.m.) being unexplained by the mere removal of vinyl substitution in the 3- and 4-positions.⁷ In addition one can observe a striking change in the u.v. spectrum when going from (**2**) (highly complex spectrum with absorptions up to 387 nm) to (**3**), again not explained by the simple removal of one double bond.⁷

An attractive explanation for these differences is the assumption that (**2**) behaves as a truly anti-aromatic planar 8 π -system which manifests itself in the development of a strong paramagnetic component in the n.m.r. experiment, as previously suggested,¹ and in the relatively low-energy photoexcitation,[‡] whereas (**3**) just behaves as a perturbed furan.



† The completely hydrogenated compound 3-oxabicyclo[3,2,0]heptane could, in addition to other products, also be isolated. Prolonged hydrogenation increased its relative amount while the yield of (**3**) decreased. Details will be reported in a full paper.

‡ Anti-aromatic systems are expected to have relatively high-lying highest occupied and low-lying lowest unoccupied molecular orbitals.⁸

We thank Professor J. L. Beauchamp for the determination of the mass spectrum and the National Science Foundation, the Alfred P. Sloan Foundation, and the Camille and Henry Dreyfus Foundation for financial support.

(Received, 26th January 1973; Com. 110.)

¹ P. J. Garratt and K. P. C. Vollhardt, *Chem. Comm.*, 1970, 109; *J. Amer. Chem. Soc.*, 1972, **94**, 1022, 7087.

² P. J. Garratt, and D. N. Nicolaides, *J.C.S. Chem. Comm.*, 1972, 1014.

³ S. W. Longworth and J. F. W. McOmie, *J.C.S. Chem. Comm.*, 1972, 623.

⁴ H. Wynberg, *Accounts Chem. Res.*, 1971, **4**, 65.

⁵ R. G. Bergman and K. P. C. Vollhardt, *J. Amer. Chem. Soc.*, 1972, **94**, 8950.

⁶ L. M. Jackman and S. Sternhell, 'Applications of Nuclear Magnetic Resonance in Organic Chemistry,' Pergamon, Oxford, 1969, p. 214.

⁷ For vinyl substituted furans, see: J. A. Elix, M. V. Sargent, and F. Sondheimer, *J. Amer. Chem. Soc.*, 1970, **92**, 973; J. W. van Reijendam, G. J. Heeres, and M. J. Janssen, *Tetrahedron*, 1970, **26**, 1291.

⁸ M. J. S. Dewar, *Angew. Chem. Internat. Edn.*, 1971, **10**, 761.