



CHICAGO JOURNALS



100-year DASCH Light Curves of Kepler Planet-Candidate Host Stars

Author(s): Sumin Tang, Dimitar Sasselov, Jonathan Grindlay, Edward Los and Mathieu Servillat

Source: *Publications of the Astronomical Society of the Pacific*, Vol. 125, No. 929 (July 2013), pp. 793-797

Published by: [The University of Chicago Press](#) on behalf of the [Astronomical Society of the Pacific](#)

Stable URL: <http://www.jstor.org/stable/10.1086/671759>

Accessed: 29/08/2013 16:49

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at
<http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



The University of Chicago Press and Astronomical Society of the Pacific are collaborating with JSTOR to digitize, preserve and extend access to Publications of the Astronomical Society of the Pacific.

<http://www.jstor.org>

100-year DASCH Light Curves of Kepler Planet-Candidate Host Stars

SUMIN TANG,^{1,2,3} DIMITAR SASSELOV,¹ JONATHAN GRINDLAY,¹ EDWARD LOS,¹ AND MATHIEU SERVILLAT^{1,4}

Received 2013 April 28; accepted 2013 June 03; published 2013 June 14

ABSTRACT. We present 100 year light curves of Kepler planet-candidate host stars from the Digital Access to a Sky Century at Harvard (DASCH) project. 261 out of 997 host stars have at least 10 good measurements on DASCH scans of the Harvard plates. 109 of them have at least 100 good measurements, including 70% (73 out of 104) of all host stars with $g \leq 13$ mag, and 44% (100 out of 228) of all host stars with $g \leq 14$ mag. Our typical photometric uncertainty is $\sim 0.1\text{--}0.15$ mag. No variation is found at 3σ level for these host stars, including 21 confirmed or candidate hot Jupiter systems which might be expected to show enhanced flares from magnetic interactions between dwarf primaries and their close and relatively massive planet companions.

Online material: color figures

1. INTRODUCTION

In 2011 the Kepler Mission announced 997 stars with a total of 1,235 planetary candidates that show transit-like signatures (Borucki et al. 2011). It is estimated that more than 95% of these candidates are planets (Morton & Johnson 2011), although a much higher false positive rate of 35% for Kepler close-in giant candidates is reported by Santerne et al. (2012). Kepler has completed 3 years of observations and with a 4-year extension could accumulate 7.5 years of data for these host stars. Searching the host stars for significant very long-term photometric variations on timescales of decades and for rare flare events is important for understanding fully the environments of their planets and for constraining the habitability of exoplanets in general. For example, a recent study by Lecavelier des Etangs et al. (2012) found increased evaporating atmosphere of hot Jupiter HD 189733b during a transit, which is probably related to a X-ray flare 8 hr before the transit. Here we present the 100-year light curves of Kepler planet-candidate host stars from the Digital Access to a Sky Century at Harvard (DASCH) project (Grindlay et al. 2009, 2012).

The photometric variability of about 150,000 Kepler target stars, including the sample with planetary candidates, shows low-level variability due to modulation by spots and other manifestations of magnetic activity, e.g., white light flares (Basri et al. 2011). None of these variabilities have amplitudes that would be

detectable (>0.3 mag if only a single observation) in the DASCH data. Here we aim to complement the short-timescale and high-precision photometry by Kepler with the much longer timescales available from DASCH. The DASCH project can reveal stars with rare flare events or extremely slow changes in luminosity. It is of interest to search for both kinds of variability with DASCH. Extremely large (>1 mag) flares could drive significant mass loss from the atmospheres of hot Jupiters, and long term changes in host star luminosity could drive changes in planetary winds and spectral composition. This paper provides DASCH light curves for the initial sample of Kepler planets presented by Borucki et al. (2011).

2. DASCH LIGHT CURVES

In order to take advantage of the unprecedented Kepler data on short timescales (Borucki et al. 2010) which complements DASCH data on long timescales, we have scanned and processed 3,735 plates taken from the 1880 to 1990 in or covering part of the Kepler field. Each plate covers $5\text{--}40^\circ$ on a side, and most of them are blue-sensitive (close to Johnson B). Most plates have limiting magnitudes 12–14 mag, while $\sim 9\%$ of them (mc series) are down to ~ 17 mag. More details on the coverage and limiting magnitudes of the plates in the Kepler field are described in Tang et al. (2013).

We used the Kepler Input Catalog (KIC; Brown et al. 2011) for photometric calibration. The typical relative photometric uncertainty is $\sim 0.1\text{--}0.13$ mag, as measured by the median light curve rms of stars (Laycock et al. 2010; Tang et al. 2013; see also Table 1 and final figure in this article). Given that most stars are constant at the 0.1 mag level, their light curve rms values are dominated by photometric uncertainty, and thus the median light curve rms represents the relative photometric uncertainty. The typical absolute photometric uncertainty, as measured by the difference between our

¹Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138.

² Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106.

³ Division of Physics, Mathematics, and Astronomy, California Institute of Technology, Pasadena, CA 91125.

⁴Laboratoire AIM (CEA/DSM/IRFU/SAp, CNRS, Université Paris Diderot), CEA Saclay, Bat. 709, 91191 Gif-sur-Yvette, France.

TABLE 1
DASCH LIGHT CURVE PROPERTIES OF HOST STARS OF CONFIRMED AND CANDIDATE HOT JUPITER PLANETS IN THE KEPLER FIELD

KOI	KIC	KIC <i>g</i> (mag)	<i>N</i> _{lc} ^a	lc median ^b (mag)	lcrms ^c (mag)	Notes ^d
1	11446443	11.74	570	11.72	0.109	TrES-2; O'Donovan et al. (2006)
2	10666592	10.94	1069	10.73	0.113	HAT-P-7; Pál et al. (2008)
13	9941662	9.7	1348	9.76	0.103	Kepler-13; Shporer et al. (2011)
18	8191672	13.88	120	13.63	0.099	Kepler-5; Koch et al. (2010)
20	11804465	13.78	62	13.68	0.153	Kepler-12; Fortney et al. (2011)
97	5780885	13.26	170	13.13	0.185	Kepler-7; Latham et al. (2010)
98	10264660	12.36	404	12.3	0.128	Kepler-14; Buchhave et al. (2011)
100	4055765	12.91	243	12.85	0.115	
128	11359879	14.27	32	14.185	0.122	Kepler-15; Endl et al. (2011)
135	9818381	14.35	38	14.29	0.101	Kepler-43; Bonomo et al. (2012)
138	8506766	14.22	49	14.14	0.113	
191	5972334	15.56	54	15.635	0.066	
203	10619192	14.64	15	14.52	0.207	Kepler-17; Désert et al. (2011)
846	6061119	16.01	45	16.02	0.096	
897	7849854	15.82	13	15.86	0.143	
976	3441784	9.9	1407	9.73	0.097	
1020	2309719	13.34	84	13.14	0.114	
1452	7449844	13.84	79	13.81	0.116	
1474	12365184	13.28	242	13.255	0.158	
1540	5649956	16.17	45	16.36	0.108	
1549	8053552	15.73	10	15.725	0.077	

^a Number of measurements in DASCH light curve.

^b Median magnitude in DASCH light curve.

^c DASCH light curve rms.

^d Names of the confirmed hot Jupiter systems and references.

measurements and KIC *g* band magnitudes, is ~ 0.2 mag, and up to ~ 0.35 mag for bright stars ($g < 11$ mag). The larger absolute photometric uncertainty is caused by additional contributions from the uncertainties in KIC *g* mag (especially at the bright end), and the difference between the effective color of the plates (close to Johnson B) and *g* band. The magnitude uncertainties we provide in DASCH light curves are defined as the rms scatter of ($m_{\text{DASCH}} - g_{\text{KIC}}$) for stars in local spatial bins with similar magnitudes, and thus represent the absolute photometric uncertainty (Tang et al. 2013). The typical absolute astrometric uncertainty (per data point) is $\sim 0.8\text{--}5''$, depending on plate scale (Laycock et al. 2010; Los et al. 2011; Servillat et al. 2011).

Example DASCH light curves of 3 Kepler hot Jupiter host stars, as well as a binary are shown in Figure 1. Fainter stars have fewer DASCH measurements, due to smaller number of deeper plates available. For a given star, every measurement is typically separated by days to months, and could extend up to years. Therefore, one stellar flare, if it happened during the plate observation, is expected to appear on a single plate only. Some fields do have multiple exposures which our pipeline is able to process individually (see Los et al. 2011), and would allow stellar flares to be confirmed by successive mea-

surements; no such flares were seen. Among the 4 stars plotted in Figure 1, K10666592 (HAT-P-7; Pál et al. 2008), K10264660 (Kepler-14; Buchhave et al. 2011), and K8191672 (Kepler-5; Koch et al. 2010) are confirmed hot Jupiter host stars. K5122112 (KOI 552; Borucki et al. 2011) was listed as a hot Jupiter candidate, but later has been unveiled as a binary (Bouchy et al. 2011). All three planets have $R \geq R_{\text{Jupiter}}$, $a < 0.1$ AU, and equilibrium temperature > 1300 K. No variation is detected.

All the light curve data and plots used for this paper are available at the website.⁵ Only “good” measurements are included. We excluded blended images, measurements within 0.75 mag of the limiting magnitude which are more likely to be contaminated by noise, images within the outer border of the plates whose width is 10% of the plate’s minor-axis length (annular bin 9; see Laycock et al. 2010), and dubious points with image profiles different from neighbor stars and thus are suspected to be emulsion defects or dust. Stars with strong correlation between magnitude measurements and plate limiting magnitudes, or between

⁵ Please see <http://hea-www.harvard.edu/DASCH/papers/Kepler-planet-host-star-lightcurves> for more information.




FIG. 1.—Example DASCH light curves of 3 Kepler hot Jupiter host stars, and a binary. From *top* to *bottom*: KIC 10666592 (HAT-P-7; Pál et al. 2008), 10264660 (Kepler-14; Buchhave et al. 2011), 8191672 (Kepler-5; Koch et al. 2010), and 5122112 (KOI 552, binary; Bouchy et al. 2011). Number of measurements (n) in each light curve are also shown, which drops for fainter stars due to smaller number of deeper plates available. Kepler PDC corrected light curves are also shown in green dots for comparison; note that the variation in the Kepler light curves for these stars are at 10^{-4} to 10^{-3} level, much smaller than the plotting symbols. The DASCH results constrain any rare but bright (>0.4 mag) flares from Kepler planet host stars, which might be most likely from hot Jupiter systems. See the electronic edition of the *PASP* for a color version of this figure.

magnitude measurements and plate astrometry uncertainties, are also excluded, which are very likely to be polluted by noise or blends. More detailed descriptions can be found at Tang et al. (2013).

Note that these Kepler planet host stars have variations at the 0.01%–0.1% level in the Kepler light curves, which are much smaller than the plotting symbols in Figure 1 and our photometric accuracy. For comparison, the Kepler light curves of the 4 example Kepler planet-candidate host stars (from Q0 or Q1 to Q6) are also shown in Figure 1 as green dots. We used the PDC corrected flux, which are converted to magnitudes and shifted to the mean magnitudes of DASCH light curves (see <http://keplergo.arc.nasa.gov/CalibrationSN.shtml>).

Among 997 host stars, 261 stars have at least 10 good measurements on DASCH plates, and 109 stars have at least 100 good measurements. Distributions of g band magnitudes for all the host stars and host stars with at least 10 or 100 DASCH measurements are shown in Figure 2. We have at least 100 measurements for 70% (73 out of 104) of all host stars with $g \leq 13$ mag, and 44% (100 out of 228) of all host stars with $g \leq 14$ mag. Most stars brighter than $g = 13$ mag we lost (i.e., having less than 100 good measurements) were lost due to blending with neighbor stars, as expected in such a crowded




FIG. 2.—Distribution of g band magnitudes for 997 Kepler planet-candidate host stars from Borucki et al. 2011 (black dash-dotted line), 261 host stars with at least 10 DASCH measurements (red solid line), and 109 host stars with at least 100 DASCH measurements (blue dashed line). See the electronic edition of the *PASP* for a color version of this figure.

field, with 74% of them (23 out of 31) having bright neighbor stars with $g < 14$ mag within 1 arcmin. For comparison, for $g \leq 13$ stars with at least 100 good measurements, only 14% (10 out of 73) of them have $g < 14$ mag neighbor stars within 1 arcmin.

We have carefully examined light curves of the 261 planet candidate host stars, and none of them showed variations (flares or dips) at the $>3\sigma$ level, where σ here is our absolute photometric uncertainty with typical value of ~ 0.2 mag for $g > 13$ objects, and up to 0.35 mag for $g < 12$ objects. Nor did we see any long-term photometric trends. We also compare the light curve rms of these host stars versus other stars with similar magnitudes in the Kepler field, as shown in Figure 3. The raw light curve rms versus g band magnitude for the 261 host stars with at least 10 DASCH measurements are shown as blue dots. The black solid line shows the median light curve rms of all the stars with at least 10 DASCH measurements in the Kepler field, and the red dashed lines and green dash-dotted lines show the 1σ and 2σ distributions, respectively. The median, 1σ , and 2σ distributions of light curve rms are calculated in 0.5 mag bins, after three iterations of 4σ -clipping, to exclude variable stars and dubious light curves contaminated by blending or plate defects. Stars with bright neighbors ($\geq 30\%$ the flux of the star in KIC g band) within $30''$ are also excluded in the calculation of median, 1σ , and 2σ distributions to avoid contamination from blending. Note the drop of rms for stars fainter than 14 mag is due to the fact that in general deeper plates are of better quality. Our plates are not a homogeneous sample. Stars with different magnitudes, even in the same region of the sky, are actually covered by different plates.




FIG. 3.—DASCH light curve rms vs. g band magnitude. The 240 Kepler planet-candidate host stars with at least 10 DASCH measurements are plotted as blue dots. Light curve rms percentiles of 59,453 stars with at least 10 DASCH measurements and no bright neighbors in the Kepler field are shown for comparison; stars with bright neighbors ($\geq 30\%$ the flux of the star in KIC g band) within $30''$ are excluded to avoid contamination from blending. The 50th percentile (median) is shown in black solid line, the 16th and 84th percentiles (1σ) are shown in red dashed line, and the 2.5th and 97.5th percentiles are shown in green dash-dotted lines. The median, 1σ , and 2σ distributions of light curve rms are calculated in 0.5 mag bins, after three iterations of 4σ -clipping, to exclude variable stars and dubious light curves contaminated by blending or plate defects. See the electronic edition of the *PASP* for a color version of this figure.

For example, a 12th mag star is detected on all the plates deeper than 12th mag, while a 15th mag star is only detected on plates deeper than 15th mag. Most of the deeper plates, such as mc series, are of much better quality compared with the shallow plates, which leads to the decrease of typical (median, etc.) light curve rms of fainter stars ($g > 14$ mag).

None of the planet candidate host stars have a light curve rms more than 2σ greater than the median rms of stars of similar magnitudes. There is one planet host star, i.e., Kepler-21 (KIC 3632418, $g = 8.74$; Howell et al. 2012), with light curve rms close to the 2σ distribution. Further examination shows that some images are marginally contaminated by a $g = 11.43$ mag neighbor star located at $51''$ away from the star, and thus its rms excess is dubious.

It has been suggested that the magnetic interaction between hot Jupiters and their host stars should enhance stellar activity, and may lead to phase shifts of hot spots on the stellar chromosphere (Cuntz & Shkolnik 2002; Shkolnik et al. 2003, 2005; Kopp et al. 2011; Poppenhaeger & Schmitt 2011). Given the extremely long timescale covered by DASCH, it is interesting to examine the light curves of stars hosting hot Jupiters. Among the 261 planet candidate host stars with at least 10 DASCH measurements, 21 of them host hot Jupiter planet candidates ($R \geq R_{\text{Jupiter}}$ and $a < 0.4$ AU), and 9 of them have at least

100 DASCH measurements (4 are shown in Fig. 1 as examples). DASCH light curve properties of these Jupiter planet candidate host stars, including number of measurement, median magnitude, and rms in the light curves, are listed in Table 1. None of them showed variations at the $>3\sigma$ level.

3. CONCLUSION

The Kepler mission has now discovered more than 2,000 planetary candidates (only the first 1,235 considered here) and provided unparalleled precision light curves for their host stars. Here we complement that database with much longer timescale 100-year light curves from the DASCH project. Despite their inferior photometric accuracy, the DASCH light curves sample such an extended period of time (e.g., tens of solar-like cycles of activity), that rare or very slow phenomena can be studied. From the statistical sample of the Kepler planet host stars, limits on their long-term variations and rare flare events, such as the X-ray/EUV flare from HD189733 (Lecavelier des Etangs et al. 2012), help us understand the planetary environments around main sequence stars and the habitability of exoplanets in general. We note that the HD189733 system is not within the Kepler field and so has not yet been scanned by DASCH. However, given the luminosity of the flare (7×10^{28} erg s $^{-1}$; Lecavelier des Etangs et al. 2012), and the relatively luminous (K1.5 V) host star, such a flare is beyond the detection limit of DASCH. Relatively larger optical flares would be expected for cooler stars with hot Jupiter companions. Because Kepler is mostly targeting on Sun like stars, none of M dwarf and hot Jupiter system is in the sample of the 261 host stars we studied.

We have scanned and processed 3,735 plates taken from the 1880 to 1990 in or covering part of the Kepler field, and studied the light curves of 261 planet host stars that have at least 10 good measurements on DASCH plates. We find no photometric variations at the 3σ level. We have released all the DASCH data in the Kepler field to the public in DASCH Data Release 1 (DR1) which enable users to directly plot their own light curves for differing data selection criteria.⁶ DASCH light curves over ~ 100 year timescales will continue to provide unique constraints for planet host stars, as well as any other interesting objects in the Kepler field.

We thank the anonymous referee for suggestions that have helped improve this article. We thank Alison Doane, Jaime Pepper, David Słiski and Robert J. Simcoe at CfA for their work on DASCH, and many volunteers who have helped digitize logbooks, clean and scan plates (<http://hea-www.harvard.edu/DASCH/team.php>). This work was supported in part by NSF grants AST0407380 and AST0909073 and now also the Cornel and Cynthia K. Sarosdy Fund for DASCH.

⁶ Please see <http://dasch.rc.fas.harvard.edu> for more information.

REFERENCES

- Basri, G., Walkowicz, L. M., Batalha, N., et al. 2011, AJ, 141, 20
- Bonomo, A. S., et al. 2012, A&A, 538, 96
- Borucki, W. J., et al. 2010, Science, 327, 977
- Borucki, W. J., et al. 2011, ApJ, 736, 19
- Bouchy, E., et al. 2011, A&A, 533, 83
- Brown, T. M., Latham, D. W., Everett, M. E., & Esquerdo, G. A. 2011, AJ, 142, 112
- Buchhave, L. A., et al. 2011, ApJS, 197, 3
- Cuntz, M., & Shkolnik, E. 2002, Astron. Nachr., 323, 387
- Désert, J.-M., et al. 2011, ApJS, 197, 14
- Endl, M., et al. 2011, ApJS, 197, 13
- Fortney, J. J., et al. 2011, ApJS, 197, 9
- Grindlay, J., Tang, S., Los, E., & Servillat, M. 2012, Proc. IAUS285 (New York: Cambridge University Press), 29
- Grindlay, J., Tang, S., Simcoe, R., Laycock, S., Los, E., Mink, D., Doane, A., & Champine, G. 2009, in ASP Conf. Ser. 410, Preserving Astronomy's Photographic Legacy, ed. W. Osborn, & L. Robbins (San Francisco: ASP), 101
- Howell, S. B., et al. 2012, ApJ, 746, 123
- Koch, D. G., et al. 2010, ApJ, 713, L131
- Kopp, A., Schilp, S., & Preusse, S. 2011, ApJ, 729, 116
- Latham, D., et al. 2010, ApJ, 713, 140
- Laycock, S., Tang, S., Grindlay, J., Los, E., Simcoe, R., & Mink, D. 2010, AJ, 140, 1062
- Lecavelier des Etangs, A., et al. 2012, A&A, 543, L4
- Los, E., Grindlay, J., Tang, S., Servillat, M., & Laycock, S. 2011, in ASP Conf. Ser. 422, Astronomical Data Analysis Software Systems XX, ed. I. N. Evans, A. Accomazzi, D. J. Mink, & A. H. Rots (San Francisco: ASP), 269
- Morton, T. D., & Johnson, J. A. 2011, ApJ, 738, 170
- O'Donovan, F. T., et al. 2006, ApJ, 651, L61
- Pál, A., et al. 2008, ApJ, 680, 1450
- Poppenhaeger, K., & Schmitt, J. H. M. M. 2011, Astron. Nachr., 332, 1052
- Santerne, A., et al. 2012, A&A, 545, 76
- Servillat, M., Los, E., Grindlay, J., Tang, S., & Laycock, S. 2011, in ASP Conf. Ser. 422, Astronomical Data Analysis Software Systems XX, ed. I. N. Evans, A. Accomazzi, D. J. Mink, & A. H. Rots (San Francisco: ASP), 273
- Shkolnik, E., Walker, G. A. H., & Bohlender, D. A. 2003, ApJ, 597, 1092
- Shkolnik, E., Walker, G. A. H., Bohlender, D. A., Gu, P.-G., & Kurster, M. 2005, ApJ, 622, 1075
- Shporer, A., et al. 2011, AJ, 142, 195
- Tang, S., Grindlay, J., Los, E., & Servillat, M. 2013, PASP, in press