

Geophysical Research Letters

RESEARCH LETTER

10.1029/2019GL085809

Key Points:

- Tropical ascent area has been decreasing at a rate about 1%/decade from 1979 to 2016
- Associated with tropical ascent tightening, rain rate in the top 1% of monthly rainfall in the tropics has been increasing at 1.1%/decade
- Southeast Amazon experiences a drying trend of about 3.2%/decade related to tropical ascent tightening from 1979 to 2016

Supporting Information:

- Supporting Information S1

Correspondence to:

H. Su,
Hui.Su@jpl.nasa.gov

Citation:

Su, H., Wu, L., Zhai, C., Jiang, J. H., Neelin, J. D., & Yung, Y. L. (2020). Observed tightening of tropical ascent in recent decades and linkage to regional precipitation changes. *Geophysical Research Letters*, *47*, e2019GL085809. <https://doi.org/10.1029/2019GL085809>

Received 14 OCT 2019

Accepted 16 JAN 2020

Accepted article online 21 JAN 2020

Observed Tightening of Tropical Ascent in Recent Decades and Linkage to Regional Precipitation Changes

Hui Su¹ , Longtao Wu¹ , Chengxing Zhai¹ , Jonathan H. Jiang¹ , J. David Neelin² , and Yuk L. Yung³ 

¹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA, ²Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, Los Angeles, CA, USA, ³Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

Abstract Climate models predict that the tropical ascending region should tighten under global warming, but observational quantification of the tightening rate is limited. Here we show that the observed spatial extent of the relatively moist, rainy and cloudy regions in the tropics associated with large-scale ascent has been decreasing at a rate of $-1\%/decade$ ($-5\%/K$) from 1979 to 2016, resulting from combined effects of interdecadal variability and anthropogenic forcings, with the former contributing more than the latter. The tightening of tropical ascent is associated with an increase in the occurrence frequency of extremely strong ascent, leading to an increase in the average precipitation rate in the top 1% of monthly rainfall in the tropics. At the margins of the convective zones such as the Southeast Amazonia region, the contraction of large-scale ascent is related to a long-term drying trend about $-3.2\%/decade$ in the past 38 years.

Plain Language Summary The ascending branch of the tropical circulation is tied to the Intertropical Convergence Zone (ITCZ) and encompasses the climatologically wettest area with heavy rainfall and abundant convective clouds. Climate models suggest that the tropical ascent area should tighten under global warming; however, observational evidence of the tightening is limited. We analyze three leading reanalysis datasets and three direct observables as proxies of large-scale ascent in recent decades. We find that the area coverage of the tropical ascending region has been decreasing at a rate of $-1\%/decade$ from 1979 to 2016. The observed tightening is largely driven by natural climate variability and anthropogenically forced global warming plays a secondary role. The tightening of tropical ascent is associated with an increase in the average precipitation rate in the top 1% of monthly rainfall in the tropics. At the margins of the convective zones such as the Southeast Amazonia region, the contraction of large-scale ascent is related to a drying trend about $-3.2\%/decade$ in the past 38 years. The regional expression of the tropical ascent tightening has profound hydrological impacts in such locations.

1. Introduction

Large-scale circulation in the tropics including the Hadley Cell and Walker Cell plays an important role in transporting energy, moisture, and momentum in Earth's climate system. Changes to the large-scale circulation under global warming are closely linked to global and regional energy and water cycles. Climate models predict that the ascending branch of the Hadley Circulation tends to narrow with surface warming (Byrne & Schneider, 2016; Lau & Kim, 2015; Su et al., 2014; Su et al., 2017), while the descending branch expands poleward (Hu & Fu, 2007; Lu et al., 2007; Seidel et al., 2008). The tightening of the tropical ascent is consistent with the thermodynamic scaling argument that convective mass flux would reduce in a warmer climate (Held & Soden, 2006). It can also be explained by the “upped-ante mechanism” that postulates the margins of convective zones would experience suppressed convective initiation because increased convective threshold associated with warmer tropospheric temperature would be less frequently satisfied due to limited moisture supply (Neelin et al., 2003). Byrne and Schneider (2016) showed that the negative moist static energy advection by the zonal-mean flow acting on the strengthened moist static energy meridional gradient between the moist tropics and the dry subtropics contributes primarily to the narrowing tendency of the Intertropical Convergence Zone (ITCZ) in the models that participated in the Coupled Model Intercomparison Project Phase 5 (CMIP5). However, the magnitudes of the narrowing vary substantially across the models, which affect the model simulations of tropical

high-altitude cloud cover and thus atmospheric radiative cooling rate and global-mean precipitation sensitivity to surface warming (Su et al., 2017).

Although most climate models agree on the sign of the tropical ascent area change when climate warms, the magnitudes of the tightening differ substantially (Byrne & Schneider, 2016; Su et al., 2017; Su et al., 2019). Observational assessment of the tropical ascent area change is needed to constrain the model simulations. During the satellite era since 1979, global-mean surface temperature (T_s) has increased about 0.2 K/decade, primarily caused by anthropogenic radiative forcing associated with increasing greenhouse gases (GHG) (IPCC, 2013). Internal climate variability, such as the Pacific Decadal Oscillation (PDO) and Atlantic Multidecadal Oscillation, modulates the warming trend over certain periods (Gu et al., 2016). It is not clear whether the spatial extent of the tropical ascending region has experienced discernible changes in accordance with the long-term warming. Wodzicki and Rapp (2016) used precipitation data from the Global Precipitation Climatology Project (GPCP) and Tropical Rainfall Measuring Mission supplemented by reanalysis low-level divergence and potential temperature fields to define the location, width, and strength of the ITCZ in the tropical Pacific. They found that the ITCZ experienced narrowing and strengthening from 1979 to 2014, and the narrowing trends vary at different longitudinal locations.

In this study, we examine the long-term trends in tropical ascent area represented in three leading reanalysis data sets and three direct observables as proxies of large-scale ascent in recent decades. The associated spatial patterns of the precipitation changes are analyzed in both observations and CMIP5 model simulations, noting certain regions of particular sensitivity.

2. Data and Methods

The three reanalyses we use are the European Centre for Medium-Range Weather Forecasts (ECMWF) interim analysis (ERA-I, Dee et al., 2011), the Modern-Era Retrospective analysis for Research and Applications, version 2 (MERRA-2, Gelaro et al., 2017), and the Japanese 55-year ReAnalysis (JRA-55, Kobayashi et al., 2015). We define the tropical ascending region by vertical pressure velocity at 500 hPa less than zero ($\omega_{500} < 0$ Pa/s) (Figure S1a in the supporting information). Note that this definition of tropical ascent area is different from previous studies that focused on the meridional width of zonal-mean Hadley Circulation (e.g., Byrne & Schneider, 2016) or the commonly used ITCZ width (Wodzicki & Rapp, 2016). It represents a holistic measure of the tropical ascent area, regardless of the zonally symmetric or zonally asymmetric components. Su et al. (2019) elaborates the usefulness of this simplified definition in understanding the basic mass and energy balance of the tropics.

As ω_{500} is dynamically inferred from other observational constraints in the reanalysis models rather than a direct observation, we identify a few proxies of tropical ascent that are measured directly or are tightly constrained by instrument observables. These proxies are atmospheric column relative humidity (CRH), evaporation minus precipitation ($E - P$), and outgoing longwave radiation (OLR) because their connections with large-scale circulation are well established (Bretherton et al., 2004; Byrne & Schneider, 2016; Chen et al., 2002).

The CRH is defined as the ratio between column-integrated precipitable water (PW) and saturation PW (PWs) and the latter is calculated by the vertical integral of saturation water vapor at each pressure level as a function of tropospheric temperature and pressure from the surface to 100 hPa. Previous studies showed that precipitation has a strong relation with high CRH over tropical oceans (Bretherton et al., 2004). We use PW and PWs from the three reanalyses because both PW and tropospheric temperature are constrained in reanalysis by assimilating satellite radiances that are sensitive to atmospheric temperature and moisture content. We use $\text{CRH} \geq 60\%$ to represent the tropical ascending region (Figure S1b).

Based on atmospheric moisture budget, the regions with precipitation greater than evaporation ($E - P < 0$) require net moisture convergence and ascending motion. Using the precipitation estimates from the GPCP (Huffman et al., 2009) and evaporation from the Objectively Analyzed air-sea Fluxes (OAFlux) for the Global Oceans project (Yu & Weller, 2007), we find that the boundary of the tropical ascending zone extends wider than the contour of $E - P < 0$ because of the effect of moisture advection. Instead, $E - P \leq 1$ mm/day yields better correspondence to the ascent over the tropical oceans (Figure S1c).

Furthermore, as the tropical convergence zone is associated with abundant convective clouds and low OLR, we use $OLR \leq 250 \text{ W/m}^2$ as an additional index of tropical ascent based on the spatial distributions of OLR and ω_{500} (Figure S1d). The NOAA interpolated OLR from 1979 to 2016 (Liebmann & Smith, 1996) is used.

Table S1 lists the sources and resolutions of the observational data sets. The Hadley Centre and Climate Research Unit surface temperature 4.4.0.0 (HadCRUT4) data set (Morice et al., 2012) is used. Our trend analysis results are qualitatively insensitive to the choices of threshold values for the three proxy data sets.

Besides observational data, we employ 24 climate model simulations available at the CMIP5 archive (Taylor et al., 2012). Three types of model simulations are analyzed: the Atmospheric Model Intercomparison Project (AMIP)-type runs driven by the observed sea surface temperature (SST) and sea ice from 1979 to 2008, the atmosphere-ocean coupled historical runs driven by natural and anthropogenic forcings from 1979 to 2005, and the projected RCP4.5 runs from 2006 to 2100 (see Table S2).

All analyses are conducted over the tropics (30°S to 30°N) including both land and ocean except for $E - P$, for which only ocean data are available. The annual mean area coverage of the tropical ascent is based on the monthly mean data where the conditions for ascent are satisfied. The respective climatological annual means are removed in the yearly time series of tropical ascent area. All observational data sets are from January 1979 to December 2016, except that the MERRA-2 reanalysis starts from January 1980. The linear trends in all variables are computed using the least squares linear regression that minimizes chi-square error.

3. Results

3.1. Robust Tightening of Tropical Ascent Area in Recent Decades

Figure 1 shows the area fractions of the tropical ascending region based on the aforementioned indicators all demonstrate a decreasing trend. The tropical ascending area fractions represented by $\omega_{500} < 0 \text{ Pa/s}$ in the three reanalyses exhibit a trend around $-0.1\%/decade$ with large uncertainties (Figure 1a). The ascent area fractions indicated by ERA-I CRH $\geq 60\%$, observed $E - P \leq 1 \text{ mm/day}$ or $OLR \leq 250 \text{ W/m}^2$ display larger negative trends (Figure 1b) ranging from -0.3% to $-0.4\%/decade$. The units of $\%/decade$ refer to the percentage of the ascent area out of the entire tropics from 30°S to 30°N . Using MERRA-2 or JRA-55 CRH (Figures S2 and S3), the narrowing trend of the tropical moist area is considerably higher, $-1.2\%/decade$ or $-0.6\%/decade$, respectively, due to the differences in the assimilation modeling systems and observational data sets used. Despite the differences between the data sets, a robust decreasing trend is found in the spatial extent of the tropical ascending region for all indices. Out of the three proxy data, the ascent area fraction based on $E - P \leq 1 \text{ mm/day}$ yields the strongest negative trend with the lowest uncertainty, $-0.38\% \pm 0.15\%/decade$. Given the large uncertainties in the reanalysis vertical velocity field, the long-term trends from CRH, $E - P$, and OLR are likely more reliable. The averaged tightening trend from the three proxy data sets is $\sim -0.3\% \pm 0.2\%/decade$ (Figure 1b), which is about $-1\%/decade$ relative to climatology or $-5\%/K$ when normalized by global mean T_s trend.

The long-term trends in the tropical ascent area based on $\omega_{500} < 0 \text{ Pa/s}$ or $E - P \leq 1 \text{ mm/day}$ from all the model simulations are shown in Table S2. The simulated trends based on CRH or OLR indicators are analyzed, but they exhibit larger intermodel differences and thus are not shown. Using $E - P \leq 1 \text{ mm/day}$ to define the ascent area, we find that the multimodel mean of the AMIP runs from 1979 to 2008 is within 1 standard deviation of the observed tightening trend of $-0.38\% \pm 0.15\%/decade$ from 1979 to 2016 (Figure 2). The multimodel means of the coupled historical runs and RCP4.5 runs produce a tightening trend only 20% of the AMIP ensemble mean (Figure 2), although the three types of model simulations possess similar global-mean T_s trends with 0.16, 0.23, and 0.19 K/decade for the AMIP, historical, and RCP4.5 multimodel-means, respectively. The differences in the tightening trends are associated with a pattern of SST changes in the observation that differs substantially from the ensemble-average trends in the historical and RCP4.5 coupled runs (Figure 2 insets). The observed surface cooling in the eastern Pacific due to the negative phase of the PDO in the second half of the record, affects the SST gradient and thus tropical circulation intensity. Enhanced SST gradient in the AMIP runs leads to a strengthened tropical circulation and a greater tightening of ascent area (Su et al., 2019). The multimodel mean of the coupled historical runs minimizes the effects of the SST anomalies associated with the decadal-to-multidecadal climate variability while




Figure 1. Time evolution of the annual-mean spatial extent of tropical ascending region from 1979 to 2016 represented by (a) $\omega_{500} < 0$ Pa/s and (b) based on $CRH \geq 60\%$ (in green), $E - P \leq 1$ mm/day (in blue) and $OLR \leq 250$ W/m² (in red). The percentage is out of the entire tropics between 30°S and 30°N. The respective climatological annual means are removed from the time series. Linear regression fittings are shown; associated linear trends with 1 standard deviation are given in the legend.

approximately capturing the forced long-term warming (Figure 2 insets). Given that the AMIP simulations with observed SST approximately capture the observed tightening magnitude, we infer that the observed tropical ascent tightening in recent decades is mainly contributed by natural climate variability. The tightening trends in the coupled simulations suggest that there is indeed a tightening tendency associated with external forcing primarily driven by increasing GHG, but this is smaller, about 20% of the trend in the AMIP ensemble average. Over a longer time scale, for example, about 100 years, the effect of decadal-to-multidecadal climate variability becomes less important and the role of the GHG-forced surface warming would dominate, as in the RCP4.5 runs (Figure 2). The qualitative consistency in terms of the sign of the trends in the 24 AMIP runs (only one model produces a positive trend) corroborates the robustness of the tropical ascent tightening in recent decades. Similar results are found when $\omega_{500} < 0$ Pa/s is used to define the ascending region in the models (Table S2).

Even with the same SST patterns in all the AMIP runs, the differences in the model physics, especially the moist convective schemes (Schiro et al., 2019), can still create large intermodel spread in the tightening trend, as shown in Figure 2 and Table S2. Su et al. (2019) found that model differences in the cloud radiative effects at the top of atmosphere over deep convective regions are highly correlated with the diverse responses in the ascent intensity and area changes, indicating a strong coupling between circulation and clouds.

3.2. Linkages of Tropical Ascent Tightening to Regional Precipitation Changes

As precipitation variations are strongly influenced by changes in large-scale circulation, it is expected that the tightening of tropical ascent in recent decades would manifest in global and regional water cycle




Figure 2. CMIP5 model simulated linear trends of the spatial extent of tropical ascending region based on $E - P \leq 1$ mm/day with AMIP from 1979 to 2008 in green, historical from 1979 to 2005 in red, and the projected RCP4.5 trend from 32006 to 2100 in blue. The multimodel means are listed in the last column. The observed trend with 1 standard deviation based on observed $E - P \leq 1$ mm/day is marked in the gray-shaded area. The insets show the multimodel-mean spatial distributions of surface temperature trends in the AMIP, historical, and RCP4.5 runs.

changes. Employing the conditional sampling approach (e.g., Bony et al., 2004; Su et al., 2013) in which the tropical mean precipitation is expressed as an integral of precipitation rate in each circulation (represented by ω_{500} regime, $p(\omega)$, weighted by the probability density function of ω_{500} , $f(\omega)$, as shown in equation (1), the impacts of the tropical ascent tightening on precipitation change can be readily disclosed.

Denoting the tropical mean precipitation rate as P , we have

$$P = \int_{-\infty}^{\infty} p(\omega) f(\omega) d\omega. \quad (1)$$

Following Emori and Brown (2005), the change of precipitation with time (t) or T_s can be decomposed into a thermodynamic component (the first term on the right-hand side, r.h.s., of equation (2)), a dynamic component (the second term on the r.h.s. of equation (2)) and the covariant (the third term on the r.h.s. of equation (2)), whereas overbar denotes the climatological mean values,

$$\frac{dP}{dt} = \int_{-\infty}^{\infty} \frac{dp(\omega)}{dt} \bar{f}(\omega) d\omega + \int_{-\infty}^{\infty} p(\omega) \frac{d\bar{f}(\omega)}{dt} d\omega + \int_{-\infty}^{\infty} \frac{dp(\omega)}{dt} \frac{d\bar{f}(\omega)}{dt} d\omega. \quad (2)$$

The thermodynamic component is associated with the increase of water vapor when T_s increases assuming that the circulation is unchanged; the dynamic component is related to the change of circulation assuming that the moisture content is unchanged. The covariant term is generally smaller than the first two terms.




Figure 3. Trends in the occurrence frequency of vertical velocity at 500 hPa (ω_{500}) in 20 ω_{500} bins with 5% occurrence interval for three reanalysis data sets (ERA-I in red, MERRA-2 in green, and JRA-55 in blue). The gray bars are climatological rain rates in mm/day (axis on the right) in each ω_{500} bin using the GPCP precipitation and ERA-I ω_{500} .

Dividing the full circulation regime into 20 equal occurrence bins with each bin accounting for 5% occurrence, we examine the long-term trends of $f(\omega)$ in three reanalysis data sets in recent decades (Figure 3). The cutoff values of ω_{500} for each bin differ slightly for different reanalysis data sets. Only the ones for ERA-I are labeled in Figure 3. Climatological precipitation rate $p(\omega)$ in each ω_{500} bin is displayed in gray bars based on GPCP precipitation and ERA-I ω_{500} (Figure 3).

The distribution of $p(\omega)$ shows that precipitation decreases monotonically from strongly ascending to descending regimes and heavy precipitation is associated with strong ascent. Over the past 38 years, the three reanalysis data sets yield different trends in each ω_{500} bin; however, a common feature shared by all three data sets is that the occurrence of extremely strong ascent (the highest 5% of negative ω_{500}) has increased, while the occurrence of moderate ascent has decreased, corresponding to a strengthening and tightening of tropical ascent. Therefore, the dynamic component of precipitation change would cause a shift from moderate to extremely heavy rain rate (equation (2)). Combined with the thermodynamic increase of precipitation at a rate of about 7%/K governed by the Clausius-Clapeyron relation, we expect that the occurrence of extremely heavy precipitation in the tropics would increase markedly.

Indeed, the GPCP precipitation data confirm this expectation. Figure 4a shows that the mean precipitation rate in the highest 5% ascent bin increases at a rate of $0.37 \text{ mm}\cdot\text{day}^{-1}\cdot\text{decade}^{-1}$ (about 4%/decade or 20%/K relative to the long-term mean of 9.4 mm/day), while the mean precipitation rate in the top 1% of monthly precipitation over the tropics increases in tandem at a rate of $0.2 \text{ mm}\cdot\text{day}^{-1}\cdot\text{decade}^{-1}$ (about 1.1%/decade or 5.5%/K relative to the long-term mean of 17 mm/day) from 1979 to 2016. The correlation coefficient (R) between the two measures of tropical heavy precipitation is 0.72, statistically significant at 99% level. It suggests that the dynamically driven precipitation change due to the increase of extremely strong ascent contributes substantially to the intensifying extremely heavy rainfall. Compared to the climatological global mean precipitation rate of 2.2 mm/day, the heavy monthly rain rates of 8.5 mm/day and higher shown in Figure 4a are likely associated with frequent or extreme floods. Over land, a steady increase of extreme precipitation on the decadal time scale would be catastrophic to regional infrastructure, water resource management, economy, and human life.

Figure 4b shows CMIP5 AMIP and historical simulations of the trends in the averaged precipitation rate in the top 1% of monthly means in the tropics scattered against the modeled long-term trends of tropical ascent area. A strong negative correlation exists for all the simulations ($R = -0.61$ for AMIP and -0.74 for historical runs, statistically significant at 99% level). The observed trends are also close to the linear approximation. This suggests that the models with stronger tightening trends tend to have greater increases of extremely




Figure 4. (a) Time series of observed mean precipitation rate in the top 1% of monthly precipitation (black) and mean precipitation rate in the highest 5% ascent bin of ERA-I reanalysis (green). (b) CMIP5 long-term trends in the mean precipitation rate in the top 1% of monthly precipitation scattered against the trends in modeled tropical ascent tightening trends along with the observations in black. (c) Time series of mean precipitation rate over the Southeast Amazonia region from GPCP (black) and the percentage of the ascent area for this region based on ERA-I $\omega_{500} < 0$ (orange) during September-October-November (SON). (d) CMIP5 long-term trends in mean precipitation rate over the Southeast Amazonia region scattered against the trends in modeled tropical ascent tightening trends during SON along with the observations in black. A 3-year running averaging is applied to all the yearly time series. The dashed lines are the least squares linear fits to the data.

heavy precipitation, as a stronger tightening corresponds to a greater increase in ascent strength due to the mass balance (Su et al., 2019). The model results further demonstrates the relevance of tropical ascent area change to extreme precipitation.

On the other hand, at the edges of the tropical ascending zones, the contraction of tropical ascent would lead to a decrease in ascent occurrence in these regions, a negative $f(\omega < 0)$, which would cause a reduction in precipitation due to the dynamic component per equation (2). In the past decades, a contraction of convective boundary occurs at the eastern edge of equatorial South Pacific Convergence Zone, the northern edge of Central America, and the southeastern edge of the Amazonian convergence zone (Figure S4), approximately captured by CMIP5 models (Figures S5 and S6). We specifically examine the Southeast Amazonia region (60–30°W, 0–20°S) as precipitation is vital to biodiversity and agriculture there, and the region has been found sensitive to the low-level inflow of dry air (Lintner & Neelin, 2010) and is thus susceptible to suppression of convection due to the upped-ante mechanism (Neelin & Su, 2005). All four seasons were analyzed and the ascent area tightening over the Amazon was found more conspicuous during the wet seasons in September-October-November (SON) and December-January-February (DJF) than during the dry seasons (Figures S4 and S7). Furthermore, the CMIP5 simulations for SON are generally more consistent with the observations than other seasons (Figures S5 and S6). Hence, we focus on the results during SON to illustrate the impact of tropical ascent tightening on regional hydrological cycle.

Figure 4c shows the yearly precipitation and ascent area fraction time series over the Amazonia region during SON. There is a clear drying trend about $-3.2\%/decade$ and a decreasing trend in the ascent area fraction. The interannual variations of the regionally averaged rainfall is highly correlated with the ascent area fraction ($R = 0.71$, Figure 4c). Similar correlations are found for other seasons and in annual mean (Figure S7). To test whether the decrease in precipitation over the Amazonia is related to the tropical-wide ascent area decrease, we examine the CMIP5-simulated Amazonia precipitation trends during SON and find that their magnitudes are highly correlated with the tropical ascent area trends across the models ($R = 0.55$ for AMIP

and historical runs together, statistically significant at 99% level) (Figure 4d). The positive correlation between the Amazonia precipitation trends and tropical ascent area trends suggests that the drying in the Amazonia is likely a regional consequence of large-scale tropical circulation change. Previous studies (Fu et al., 2013; Neelin et al., 2006) indicated that the long-term droughts and increased dry-season length in Southern Amazonia may be a response to global warming. Our analysis suggests that the tightening of tropical ascending area associated with increasing GHG would cause a drying trend in the Southeast Amazonia; however, the observed drying in recent decades is likely a combined effect from both GHG forcing and natural climate variability with the latter dominating over the former. Nevertheless, as the tightening of tropical ascent is projected to occur on the centennial time scale, a persistent drying in the Southeast Amazonia would require advance planning to deal with ever decreasing local water resources.

4. Conclusion and Discussions

The tropical ascending region encompasses the climatologically wettest area with heavy rainfall and abundant convective clouds. Although the tropical ascending region accounts for only ~20% of the global area, the averaged rain rate within this region is about 2.6 times the global mean, nurturing a rich biodiversity in the land areas.

We find that the moist tropics has become tighter in the past decades. The tightening is associated with a shift from moderate ascent to extremely strong ascent in the circulation regimes, leading to an increase in extremely heavy precipitation in the top 1% in monthly precipitation pdf with rain rate > 16 mm/day. In the meantime, formerly wet regions at the boundary of the ascending zone such as the Southeast Amazonia region have experienced long-term drying as the tropical ascending area contracts. Our analysis of the CMIP5 model simulations suggests that anthropogenic forcings are partly responsible for the contraction of the ascending area, while the decadal-to-multidecadal climate variability such as the PDO accounts for about 80% of the observed tightening in recent decades. Moreover, the tightening trend will continue in concert with the increasing GHG despite interannual and decadal oscillations. In parts of the tropics where human life and ecosystems heavily depend on precipitation, the contraction of the ascending regions would have severe impacts on agriculture, forestry, biodiversity, and economy. The monotonic surface warming caused by increasing GHG would eventually surpass the internally driven variability, and the tightening of moist tropics would persist if no mitigation of the rise of GHG is materialized.

The observed tightening trends based on various indices should serve as a reference to constrain climate models simulations. Given the broad implications of the tropical ascent area changes for global and regional hydrological cycles (Su et al., 2017; Su et al., 2019), we propose that the ascent area metrics based on ω_{500} , CRH, $E - P$, and OLR, as examined in this study, be incorporated into the standard evaluation of climate model performance. The large discrepancies in model simulations of the tightening trends underscore the pressing need to improve model physics, especially those related to moist convection.

Data Statement

The original reanalysis data sets, observations, and CMIP5 model simulations used in this study are all publicly available. The websites to download these data are as follows:

ERA-I (<http://apps.ecmwf.int/datasets/data/interim-full-moda/levtype=sfc/>); MERRA-2 (https://gmao.gsfc.nasa.gov/reanalysis/MERRA-2/data_access/); JRA-55 (<https://rda.ucar.edu/datasets/ds628.1/#description>); GPCP precipitation (<https://www.esrl.noaa.gov/psd/data/gridded/data.gpcp.html>) OaFlux (<http://oafux.who.edu/data.html>) NOAA OLR (https://www.esrl.noaa.gov/psd/data/gridded/data.interp_OLR.html); HadCRU4 T_s (<http://www.metoffice.gov.uk/hadobs/hadcrut4>); CMIP5 (<http://cmip-pcmdi.llnl.gov/cmip5/>). The methods to process the data and generate the results are described in detail in the paper. Please contact the corresponding author at hui.su@jpl.nasa.gov for any questions.

Acknowledgments

This work was performed at Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA. This study is supported by NASA ACMAP-AST and MAP projects. J. D. N. is supported by the NSF AGS-1540518 Grant.

References

- Bony, S., Dufresne, J. L., Le Treut, H., Morcrette, J.-J., & Senior, C. (2004). On dynamic and thermodynamic components of cloud changes. *Climate Dynamics*, 22, 71–86. <https://doi.org/10.1007/s00382-003-0369-6>
- Bretherton, C. S., Peters, M. E., & Back, L. E. (2004). Relationships between water vapor path and precipitation over the tropical oceans. *Journal of Climate*, 17, 1517–1528.

- Byrne, M. P., & Schneider, T. (2016). Narrowing of the ITCZ in a warming climate: Physical mechanisms. *Geophysical Research Letters*, 43(11), 350–11,357. <https://doi.org/10.1002/2016GL070396>
- Chen, J., Carlson, B. E., & Del Genio, A. D. (2002). Evidence for strengthening of the tropical general circulation in the 1990s. *Science*, 295, 838–841. <https://doi.org/10.1126/science.1065835>
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., et al. (2011). The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597. <https://doi.org/10.1002/qj.828>
- Emori, S., & Brown, S. J. (2005). Dynamic and thermodynamic changes in mean and extreme precipitation under changed climate. *Geophysical Research Letters*, 32, L17706. <https://doi.org/10.1029/2005GL023272>
- Fu, R., Yin, L., Li, W., Arias, P., Dickinson, R. E., Huang, L., et al. (2013). Increased dry-season length over southern Amazonia in recent decades and its implication for future climate projection. *Proceedings of the National Academy of Sciences of the United States of America*, 110(45), 18110–18115. <https://doi.org/10.1073/pnas.1302584110>
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L. L., et al. (2017). The Modern-Era Retrospective Analysis for Research and Applications, version-2 (MERRA-2). *Journal of Climate*, 30, 5419–5454. <https://doi.org/10.1175/JCLI-D-16-0758.1>
- Gu, G., Adler, R. F., & Huffman, G. (2016). Long-term changes/trends in surface temperature and precipitation during the satellite era (1979–2012). *Climate Dynamics*, 46, 1091. <https://doi.org/10.1007/s00382-015-2634-x>
- Held, I. M., & Soden, B. J. (2006). Robust responses of the hydrological cycle to global warming. *Journal of Climate*, 19, 5686–5699.
- Hu, Y., & Fu, Q. (2007). Observed poleward expansion of the Hadley circulation since 1979. *Atmospheric Chemistry and Physics*, 7, 5229–5236.
- Huffman, G. J., Adler, R. F., Bolvin, D. T., & Gu, G. (2009). Improving the global precipitation record: GPCP Version 2.1. *Geophysical Research Letters*, 36, L17808. <https://doi.org/10.1029/2009GL040000>
- IPCC (2013). *Climate change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK and New York, NY, 1535 pp: Cambridge University Press. <http://www.climatechange2013.org/report/>
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., et al. (2015). The JRA-55 Reanalysis: General specifications and basic characteristics. *Journal of the Meteorological Society of Japan*. Ser. II, Online ISSN 2186-9057, Print ISSN 0026-1165. <https://doi.org/10.2151/jmsj.2015-001>
- Lau, W. K.-M., & Kim, K.-M. (2015). Robust Hadley Circulation changes and increasing global dryness due to CO₂ warming from CMIP5 model projections. *Proceedings of the National Academy of Sciences of the United States of America*, 112(12), 3630–3635. <https://doi.org/10.1073/pnas.1418682112>
- Liebmann, B., & Smith, C. A. (1996). Description of a complete (interpolated) outgoing longwave radiation dataset. *Bulletin of the American Meteorological Society*, 77, 1275–1277.
- Lintner, B. R., & Neelin, J. D. (2010). Tropical South America/Atlantic sector convective margins and their relationship to low-level inflow. *Journal of Climate*, 23, 2671–2685. <https://doi.org/10.1175/2009JCLI3301.1>
- Lu, J., Vecchi, G. A., & Reichler, T. (2007). Expansion of the Hadley cell under global warming. *Geophysical Research Letters*, 34, L06805. <https://doi.org/10.1029/2006GL028443>
- Morice, C. P., Kennedy, J. J., Rayner, N. A., & Jones, P. D. (2012). Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: The HadCRUT4 dataset. *Journal of Geophysical Research*, 117, D08101. <https://doi.org/10.1029/2011JD017187>
- Neelin, J. D., Chou, C., & Su, H. (2003). Tropical drought regions in global warming and El Niño teleconnections. *Geophysical Research Letters*, 30(24), 2275. <https://doi.org/10.1029/2003GLO018625>
- Neelin, J. D., Munnich, M., Su, H., Meyerson, J. E., & Holloway, C. E. (2006). Tropical drying trends in global warming models and observations. *Proceedings of the National Academy of Sciences of the United States of America*, 103(16), 6110–6115. <https://doi.org/10.1073/pnas.0601798103>
- Neelin, J. D., & Su, H. (2005). Moist teleconnection mechanisms for the tropical South American and Atlantic sector during El Niño. *Journal of Climate*, 18, 3928–3950.
- Schiro, K. A., Su, H., Wang, Y., Langenbrunner, B., Jiang, J. H., & Neelin, J. D. (2019). Relationships between tropical ascent and high cloud fraction changes with warming revealed by perturbation physics experiments in CESM. *Geophysical Research Letters*. <https://doi.org/10.1029/2019GL083026>
- Seidel, D. J., Fu, Q., Randel, W. J., & Reichler, T. J. (2008). Widening of the tropical belt in a changing climate. *Nature Geoscience*, 1, 21–24.
- Su, H., Jiang, J. H., David Neelin, J., Janice Shen, T., Zhai, C., Yue, Q., et al. (2017). Tightening of tropical ascent and high clouds key to precipitation change in a warmer climate. *Nature Communications*, 8, 15771. <https://doi.org/10.1038/ncomms15771>
- Su, H., Jiang, J. H., Zhai, C., Shen, T. J., Neelin, J. D., Stephens, G. L., & Yung, L. Y. (2014). Weakening and strengthening structures in the Hadley circulation change under global warming and implications for cloud response and climate sensitivity. *Journal of Geophysical Research: Atmospheres*, 119(10), 5787–5805. <https://doi.org/10.1002/2014JD021642>
- Su, H., Jiang, J. H., Zhai, C., Perun, V. S., Shen, J. T., Del Genio, A., et al. (2013). Diagnosis of regime-dependent cloud simulation errors in CMIP5 models using “A-Train” satellite observations and reanalysis data. *Journal of Geophysical Research*, 118(7), 2762–2780. <https://doi.org/10.1029/2012JD018575>
- Su, H., Zhai, C., Jiang, J. H., Wu, L., Neelin, J. D., & Yung, Y.-L. (2019). A dichotomy between model responses of tropical ascent and descent to surface warming. *npj Climate and Atmospheric Science*, 2(1), 1–8. <https://doi.org/10.1038/s41612-019-0066-8>
- Taylor, K. E., Stouffer, R. J., & Meehl, G. A. (2012). An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, 93, 485–498. <https://doi.org/10.1175/BAMS-D-11-00094.1>
- Wodzicki, K. R., & Rapp, A. D. (2016). Long-term characterization of the Pacific ITCZ using TRMM, GPCP, and ERA-Interim. *Journal of Geophysical Research – Atmospheres*, 121, 3153–3170. <https://doi.org/10.1002/2015JD024458>
- Yu, L., & Weller, R. A. (2007). Objectively analyzed air-sea heat fluxes for the global ice-free oceans (1981–2005). *Bulletin of the American Meteorological Society*, 88, 527–539.