



Supplementary Material for
**The solar nebula origin of (486958) Arrokoth, a primordial contact
binary in the Kuiper Belt**

W. B. McKinnon*, D. C. Richardson, J. C. Marohnic, J. T. Keane, W. M. Grundy,
D. P. Hamilton, D. Nesvorný, O. M. Umurhan, T. R. Lauer, K. N. Singer, S. A. Stern,
H. A. Weaver, J. R. Spencer, M. W. Buie, J. M. Moore, J. J. Kavelaars, C. M. Lisse,
X. Mao, A. H. Parker, S. B. Porter, M. R. Showalter, C. B. Olkin, D. P. Cruikshank,
H. A. Elliott, G. R. Gladstone, J. Wm. Parker, A. J. Verbiscer, L. A. Young,
the New Horizons Science Team†

*Corresponding author. Email: mckinnon@wustl.edu

Published 13 February 2020 as *Science* First Release
DOI: 10.1126/science.aay6620

This PDF file includes:

Team Members and Affiliations
Materials and Methods
Supplementary Text
Table S1
References

New Horizons Science Team (Kuiper Belt Extended Mission Co-Investigators)

S. Alan Stern, Southwest Research Institute, Boulder, CO, USA.

Harold A. Weaver, Johns Hopkins University Applied Physics Laboratory, Laurel, MD, USA.

Catherine B. Olkin, Southwest Research Institute, Boulder, CO, USA.

John R. Spencer, Southwest Research Institute, Boulder, CO, USA.

J. Wm. Parker, Southwest Research Institute, Boulder, CO, USA.

Anne Verbiscer, University of Virginia, Charlottesville, VA, USA.

Richard P. Binzel, Massachusetts Institute of Technology, Cambridge, MA, USA.

Daniel T. Britt, University of Central Florida, Orlando, FL, USA.

Marc W. Buie, Southwest Research Institute, Boulder, CO, USA.

Bonnie J. Buratti, Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA.

Andrew F. Cheng, Johns Hopkins University Applied Physics Laboratory, Laurel, MD, USA.

Dale P. Cruikshank, NASA Ames Research Center, Moffett Field, CA, USA.

Heather A. Elliot, Southwest Research Institute, San Antonio, TX, USA.

G. Randall Gladstone, Southwest Research Institute, San Antonio, TX, USA.

William M. Grundy, Lowell Observatory, Flagstaff, AZ, USA.

Matthew E. Hill, Johns Hopkins University Applied Physics Laboratory, Laurel, MD, USA.

Mihaly Horanyi, University of Colorado, Boulder, CO, USA.

Don E. Jennings, NASA Goddard Space Flight Center, Greenbelt, MD, USA.

J. J. Kavelaars, National Research Council of Canada, Victoria, BC, Canada.

Ivan R. Linscott, Stanford University, Stanford, CA, USA.

Jeffrey M. Moore, NASA Ames Research Center, Moffett Field, CA, USA.

David J. McComas, Department of Astrophysical Sciences, Princeton University, Princeton, NJ, USA.

William B. McKinnon, Washington University in St. Louis, St. Louis, MO, USA.

Ralph L. McNutt, Johns Hopkins University Applied Physics Laboratory, Laurel, MD, USA.

Alex H. Parker, Southwest Research Institute, Boulder, CO, USA.

Simon B. Porter, Southwest Research Institute, Boulder, CO, USA.

Silvia Protopapa, Southwest Research Institute, Boulder, CO, USA.

Dennis C. Reuter, NASA Goddard Space Flight Center, Greenbelt, MD, USA.

Paul M. Schenk, Lunar and Planetary Institute, Houston, TX, USA.

Mark R. Showalter, SETI Institute, Mountain View, CA, USA.

Kelsi N. Singer, Southwest Research Institute, Boulder, CO, USA.

Leslie A. Young, Southwest Research Institute, Boulder, CO, USA.

Materials and Methods

To model 2-body impacts and potential mergers of granular aggregates, we use PKDGRAV, an N-body code with an implementation of the soft-sphere discrete element method (SSDEM) for collisions between spherical particles. PKDGRAV uses the k -d hierarchical tree algorithm to reduce the computational cost of calculating interparticle forces and runs in parallel to reduce the time necessary to perform simulations with large numbers of particles (41). SSDEM allows particles to interpenetrate, with restoring forces implemented as springs with a user-adjustable spring constant. The implementation of SSDEM in PKDGRAV has been described in detail (42), with implementation of static, rolling, and twisting friction (101) and interparticle cohesion (102). We only applied cohesive forces between particles of the same progenitor body, i.e., any

contact between a particle from one body and a particle from another is treated as cohesionless. This choice was motivated by initial simulations of the binary merger with cohesion included. After an initial contact between bodies, the size of the neck would continue to grow as particles near the contact point stuck together and pulled others along with them. Because we wanted to use cohesion to capture the effect of material strength, we judged this behavior to be unphysical and so adjusted to the model.

We model each lobe of the present contact binary system separately, under the assumption that they formed separately and merged at some point in the past. We generate spherical “rubble piles” out of many smaller particles. We use approximately 135,000 equal-size particles to model LL (diameter 17.94 km) and 63,000 to model SL (diameter 13.64 km); the sizes of lobes were based on preliminary estimates (7). Particle radii are normally distributed with a mean radius of 136 m and a standard deviation of 27 m. Upper and lower radius cutoffs are 163 m and 109 m, respectively. After generating LL and SL we run simulations with each body separately to allow the particles to settle into an equilibrium between self-gravitation and repulsive contact forces. Particles are either frictionless or given gravel-like friction parameters—a static friction coefficient of 1.0, a rolling friction coefficient of 1.05, a twisting friction coefficient of 1.3, and a shape parameter of 0.5; the friction parameters mimic the shear strength of irregular particle shapes in contact (101). The normal and tangential coefficients of restitution are 0.2. Simulations of direct collisions (Figs. 4A,B) assumed no rotation of the individual bodies before contact, whereas that for Fig. 4C assumed a synchronous rotation rate appropriate to the density; the latter was done to better simulate the possible final merger conditions of a co-orbiting binary.

Supplementary Text

Accretion by hierarchical coagulation

The cold classical Kuiper belt population might have accreted by traditional hierarchical coagulation (27). CCKB object formation through a variant of HC has been shown to be viable even in a low-mass (or “light”) planetesimal disk (of $\sim 0.1M_{\oplus}$), but only if a pre-existing seed population of ~ 1 -km-scale planetesimals is invoked while simultaneously most of the mass is in cm-sized pebbles or smaller (28). Formation of km and sub-km scale planetesimals has not been demonstrated for traditional HC models, which was one of the motivations for the development of SI models (e.g., 57). Models (28) typically take 10s to 100s of Myr to accrete the larger bodies of the CCKB, ignore possible dynamical stirring by Neptune, and yield a CCKB that is too massive today (see below).

Characteristic planetesimal mass from the streaming instability

Numerical simulations of SI have focused on the formation of larger (~ 50 - 100 km scale) planetesimals in the Kuiper belt, i.e., those at the break in the observed KBO size-frequency distribution (e.g., 24). We focus on the implications of the SI for a lower-mass, outer protoplanetary disk, i.e., the cold classical region. The gravitational mass scale (M_G) in the streaming instability is given, in the CCKB object region, by $4\pi^5 G^2 \Sigma_p^3 / \Omega_K^4 \sim 10^{14} \text{ kg} \times (\Sigma_p / 0.016 \text{ kg m}^{-2})^3$, where Σ_p is the surface mass density in pebbles, Ω_K is the heliocentric orbital frequency, and G is the gravitational constant (26). If we adopt $\Sigma_p = 0.016 \text{ kg m}^{-2}$ from (28), who spread $0.1M_{\oplus}$ of solids between 42 and 48 au to form their “light disk,” the characteristic planetesimal mass produced by an SI-induced GI would be similar to the mass of Arrokoth ($\sim 10^{15} \text{ kg}$), given the sensitivity of M_G to Σ_p . In comparison, to produce a characteristic CCKB object diameter of

~ 100 km ($M_G \sim 3 \times 10^{17}$ kg) (103) requires an order of magnitude more solid mass in pebbles. Neither of these total masses, as planetesimals, violate the surface mass constraint (20) for halting Neptune's outward migration.

Because of the limited dynamical excitement of the cold population, (19) have argued for modest dynamical depletion of the CCKB, at most, since its formation, perhaps by no more than a factor of ~ 2 . This implies that the mass of sizeable objects in the cold classical region has always been low. Given the very low mass of CCKB objects today ($\sim 0.001 M_\oplus$ (103)), this suggests that the formation of CCKB objects via the streaming (or other) instability—from a larger reservoir of solids that would allow larger bodies than Arrokoth to accrete—must have been intermittent in space and time or otherwise inefficient during the lifetime of the protosolar nebula (104). The alternative, that the CCKB was originally more massive and lost substantial mass to collisional grinding (105), is not consistent with the lack of evidence for collisional processing of Arrokoth (7, 8) and the large fraction of loosely bound binaries among the cold classicals (e.g., 106). Sporadic or inefficient planetesimal formation could be related to a globally lower pebble/gas ratio owing to gas drag drift of pebbles (92, 94), but with local pebble concentrations due to zonal flows or other mechanisms (e.g., 107).

Alternative particle concentration mechanisms to the streaming instability

In addition to SI, nebular turbulence likely led to particle concentrations at corresponding eddy scales (21), but whether such concentrations led to GI and planetesimal formation has received less attention (108, 109). SI is a dynamic particle concentration mechanism, which is expected to occur over a range of protoplanetary disk conditions and pebble sizes, possibly in tandem with other particle concentration mechanisms (57, 110). For example, the surface mass density of gas

in the outermost protosolar nebula, and of the CCKB object formation zone in particular, plausibly should have been low enough for cosmic-ray and x-ray induced ionization and active magneto-rotational instability (MRI) (57). Levels of turbulence associated with MRI (111) may act to suppress SI, at least in its classic laminar form (112). MRI might not reach the disk midplane, however, where SI would take place (57, 113), and SI and MRI can act in tandem, with SI enhancing particle concentrations on a smaller scale (109, 114).

LL and SL as possible Roche ellipsoids

The spin and angular momentum of Arrokoth can be normalized by the critical rotation rate $\omega_c = \sqrt{\frac{4}{3}\pi\rho G}$ and $mR_e^2\omega_c$, respectively, where m and R_e are a body's total mass and equivalent spherical radius (e.g., 99). The normalized spin and angular momentum for Arrokoth are ~ 0.29 and ~ 0.36 , respectively, assuming a mass ratio of 2:1 for LL and SL and $\rho = 500 \text{ kg m}^{-3}$ for both lobes. These values resemble those for critically stable Roche ellipsoids of the same mass ratio, about 0.28 and 0.26, respectively (see 99), but the correspondence breaks down for lower densities (the normalized values scale as $\rho^{-1/2}$).

Shape of Arrokoth's individual lobes

The individual mapped units on LL may indicate the merger or assembly of discrete multi-km-scale planetesimals (7, 8). If so, to create such a lenticular or ellipsoidal body as LL requires that the mergers were themselves not very energetic or high velocity. These velocity conditions would have been met in a collapsing particle cloud (47, 58), but not during heliocentric hierarchical coagulation generally; the latter implies speeds in excess (or greatly in excess) of the escape speed from LL. Low cohesion and a near absence of internal friction would have been necessary

mechanically at the time of the LL merger collisions as well. Otherwise, the shape of LL would much more reflect the shapes of the individual subunits from which it was built (as in Fig. 4B).

Alternately, the LL and SL lobes could have accreted directly in a collapsing rotating particle cloud from myriad small pebbles (47), and acquired their lenticular shapes naturally. Arrokoth is a contact binary, and not a single, broadly ellipsoidal body, so the dynamical regime that fostered quasi-equilibrium shapes of the individual lobes must not have been applicable when the two bodies themselves finally merged. This suggests that the merger of two lobes (LL and SL) may not have occurred in the pebble cloud itself, but at some later time after the pebble cloud cleared (the latter on an $\sim 10^4$ yr time scale (58)), when the lobes may have acquired some modest measure of strength (87).

Arrokoth's neck appears somewhat bent or tilted in the direction of rotation [(8), clockwise in their figure 1A], as if this was due to a final, tangential mass displacement at the contact surface during a merger. Alternately, the bending could be due to some later mechanical failure/distortion at the neck. The edges of LL and SL observed on approach often display linear segments [(8), their figure 3], as if the portions of the lobes just out of sight had been sheared off, though this is not entirely clear in the available images. Perhaps these are the outcomes of earlier on-edge, glancing collisions between the lobes (as in Fig. 4A). Alternately, these apparent facets may have been caused by higher-velocity impacts and mass loss, such as have affected the asteroids (115), although a heliocentric impact explanation is not consistent with the dearth of large craters on the visible faces of LL and SL, save perhaps for the largest, "Maryland" (8).

Table S1.

Estimated YORP coefficients for near-Earth asteroids. From photometric measurements of asteroid rotational accelerations, an empirical, dimensionless torque coefficient Y can be estimated from the YORP torque equation $\frac{d\omega}{dt} = \left(\frac{Y}{2\pi\rho R^2}\right)\left(\frac{L_{\odot}}{4\pi c\bar{a}^2}\right)$, where ω , ρ , and R are the spin rate, density, and equivalent radius of the asteroid in question, L_{\odot} is the solar luminosity, c is the speed of light, and $\bar{a} = a_{\odot}\sqrt[4]{1 - e_{\odot}^2}$ is the solar-flux-weighted mean heliocentric distance (77). In the table P is the rotation period, $\omega/(d\omega/dt)$ is the spin-rate doubling time, and data sources are indicated. For Itokawa and Bennu the densities are known; for the others 1500 kg m⁻³ is assumed, except for (54509) YORP, (1862) Apollo, and (161989) Cacus, which are likely more monolithic and denser (116, 117, 118). Note that all these asteroids are spinning up; at present none are spinning down.

Object	$d\omega/dt$ ($\times 10^{-8}$ rad d ⁻²)	P (hr)	\bar{a} (au)	$\omega/(d\omega/dt)$ (yr)	R (km)	ρ (kg m ⁻³)	Y
54509 YORP (116)	350 ± 35	0.203	0.987	5.8×10 ⁵	0.06	2500	0.006
25143 Itokawa (117)	3.5 ± 0.4	12.132	1.297	9.7×10 ⁵	0.16	1195	0.0003
1620 Geographos (75)	1.2 ± 0.2	5.223	1.206	6.6×10 ⁶	0.98	1500	0.005
1862 Apollo (117)	5.5 ± 1.2	16.3	1.338	4.6×10 ⁵	0.75	2500	0.026
3103 Eger (118)	1.1 ± 0.5	15.3	1.358	2.5×10 ⁶	0.75	1500	0.003
161989 Cacus (118)	1.9 ± 0.3	3.755	1.110	5.8×10 ⁶	0.5	2500	0.003
101955 Bennu (82)	4.6 ± 1.8	4.296	1.115	2.1×10 ⁶	0.245	1190	0.0008

References and Notes

1. S. A. Stern, W. M. Grundy, W. B. McKinnon, H. A. Weaver, L. A. Young, The Pluto system after New Horizons. *Annu. Rev. Astron. Astrophys.* **56**, 357–392 (2018). [doi:10.1146/annurev-astro-081817-051935](https://doi.org/10.1146/annurev-astro-081817-051935)
2. S. A. Stern, H. A. Weaver, J. R. Spencer, H. A. Elliott, The New Horizons Kuiper Belt Extended Mission. *Space Sci. Rev.* **214**, 77 (2018). [doi:10.1007/s11214-018-0507-4](https://doi.org/10.1007/s11214-018-0507-4)
3. S. B. Porter, M. W. Buie, A. H. Parker, J. R. Spencer, S. Benecchi, P. Tanga, A. Verbiscer, J. J. Kavelaars, S. D. J. Gwyn, E. F. Young, H. A. Weaver, C. B. Olkin, J. W. Parker, S. A. Stern, High-precision orbit fitting and uncertainty analysis of (486958) 2014 MU69. *Astron. J.* **156**, 20 (2018). [doi:10.3847/1538-3881/aac2e1](https://doi.org/10.3847/1538-3881/aac2e1)
4. D. Nesvorný, Dynamical evolution of the early Solar System. *Annu. Rev. Astron. Astrophys.* **56**, 137–174 (2018). [doi:10.1146/annurev-astro-081817-052028](https://doi.org/10.1146/annurev-astro-081817-052028)
5. A. H. Parker, J. J. Kavelaars, Destruction of binary minor planets during Neptune scattering. *Astrophys. J.* **722**, L204–L208 (2010). [doi:10.1088/2041-8205/722/2/L204](https://doi.org/10.1088/2041-8205/722/2/L204)
6. J. M. Moore, W. B. McKinnon, D. P. Cruikshank, G. R. Gladstone, J. R. Spencer, S. A. Stern, H. A. Weaver, K. N. Singer, M. R. Showalter, W. M. Grundy, R. A. Beyer, O. L. White, R. P. Binzel, M. W. Buie, B. J. Buratti, A. F. Cheng, C. Howett, C. B. Olkin, A. H. Parker, S. B. Porter, P. M. Schenk, H. B. Throop, A. J. Verbiscer, L. A. Young, S. D. Benecchi, V. J. Bray, C. L. Chavez, R. D. Dhingra, A. D. Howard, T. R. Lauer, C. M. Lisse, S. J. Robbins, K. D. Runyon, O. M. Umurhan, Great expectations: Plans and predictions for New Horizons encounter with Kuiper Belt Object 2014 MU69 (“Ultima Thule”). *Geophys. Res. Lett.* **45**, 8111–8120 (2018). [doi:10.1029/2018GL078996](https://doi.org/10.1029/2018GL078996)
7. S. A. Stern, H. A. Weaver, J. R. Spencer, C. B. Olkin, G. R. Gladstone, W. M. Grundy, J. M. Moore, D. P. Cruikshank, H. A. Elliott, W. B. McKinnon, J. W. Parker, A. J. Verbiscer, L. A. Young, D. A. Aguilar, J. M. Albers, T. Andert, J. P. Andrews, F. Bagenal, M. E. Banks, B. A. Bauer, J. A. Bauman, K. E. Bechtold, C. B. Beddingfield, N. Behrooz, K. B. Beisser, S. D. Benecchi, E. Bernardoni, R. A. Beyer, S. Bhaskaran, C. J. Bierson, R. P. Binzel, E. M. Birath, M. K. Bird, D. R. Boone, A. F. Bowman, V. J. Bray, D. T. Britt, L. E. Brown, M. R. Buckley, M. W. Buie, B. J. Buratti, L. M. Burke, S. S. Bushman, B. Carcich, A. L. Chaikin, C. L. Chavez, A. F. Cheng, E. J. Colwell, S. J. Conard, M. P. Conner, C. A. Conrad, J. C. Cook, S. B. Cooper, O. S. Custodio, C. M. Dalle Ore, C. C. Deboy, P. Dharmavaram, R. D. Dhingra, G. F. Dunn, A. M. Earle, A. F. Egan, J. Eisig, M. R. El-Maarry, C. Engelbrecht, B. L. Enke, C. J. Ercol, E. D. Fattig, C. L. Ferrell, T. J. Finley, J. Firer, J. Fischetti, W. M. Folkner, M. N. Fosbury, G. H. Fountain, J. M. Freeze, L. Gabasova, L. S. Glaze, J. L. Green, G. A. Griffith, Y. Guo, M. Hahn, D. W. Hals, D. P. Hamilton, S. A. Hamilton, J. J. Hanley, A. Harch, K. A. Harmon, H. M. Hart, J. Hayes, C. B. Hersman, M. E. Hill, T. A. Hill, J. D. Hofgartner, M. E. Holdridge, M. Horányi, A. Hosadurga, A. D. Howard, C. J. A. Howett, S. E. Jaskulek, D. E. Jennings, J. R. Jensen, M. R. Jones, H. K. Kang, D. J. Katz, D. E. Kaufmann, J. J. Kavelaars, J. T. Keane, G. P. Keleher, M. Kinczyk, M. C. Kochte, P. Kollmann, S. M. Krimigis, G. L. Kruizinga, D. Y. Kusnierkiewicz, M. S. Lahr, T. R. Lauer, G. B. Lawrence, J. E. Lee, E. J. Lessac-Chenen, I. R. Linscott, C. M. Lisse, A. W. Lunsford, D. M. Mages, V. A. Mallder, N. P. Martin, B. H. May, D. J. McComas, R. L. McNutt Jr., D. S. Mehoke, T. S. Mehoke, D. S. Nelson,

- H. D. Nguyen, J. I. Núñez, A. C. Ocampo, W. M. Owen, G. K. Oxton, A. H. Parker, M. Pätzold, J. Y. Pelgrift, F. J. Pelletier, J. P. Pineau, M. R. Piquette, S. B. Porter, S. Protopapa, E. Quirico, J. A. Redfern, A. L. Regiec, H. J. Reitsema, D. C. Reuter, D. C. Richardson, J. E. Riedel, M. A. Ritterbush, S. J. Robbins, D. J. Rodgers, G. D. Rogers, D. M. Rose, P. E. Rosendall, K. D. Runyon, M. G. Ryschkewitsch, M. M. Saina, M. J. Salinas, P. M. Schenk, J. R. Scherrer, W. R. Schlei, B. Schmitt, D. J. Schultz, D. C. Schurr, F. Scipioni, R. L. Sepan, R. G. Shelton, M. R. Showalter, M. Simon, K. N. Singer, E. W. Stahlheber, D. R. Stanbridge, J. A. Stansberry, A. J. Steffl, D. F. Strobel, M. M. Stothoff, T. Stryk, J. R. Stuart, M. E. Summers, M. B. Tapley, A. Taylor, H. W. Taylor, R. M. Tedford, H. B. Throop, L. S. Turner, O. M. Umurhan, J. Van Eck, D. Velez, M. H. Versteeg, M. A. Vincent, R. W. Webbert, S. E. Weidner, G. E. Weigle 2nd, J. R. Wendel, O. L. White, K. E. Whittenburg, B. G. Williams, K. E. Williams, S. P. Williams, H. L. Winters, A. M. Zangari, T. H. Zurbuchen, Initial results from the New Horizons exploration of 2014 MU₆₉, a small Kuiper Belt object. *Science* **364**, eaaw9771 (2019). [doi:10.1126/science.aaw9771](https://doi.org/10.1126/science.aaw9771) [Medline](#)
8. J. R. Spencer *et al.*, The geology and geophysics of Kuiper Belt object (486958) Arrokoth. *Science* **367**, eaay3999 (2020).
 9. W. M. Grundy *et al.*, Color, composition, and thermal environment of Kuiper belt object (486958) Arrokoth. *Science* **367**, eaay3705 (2020).
 10. M. J. Mumma, S. B. Charnley, The chemical composition of comets—Emerging taxonomies and natal heritage. *Annu. Rev. Astron. Astrophys.* **49**, 471–524 (2011). [doi:10.1146/annurev-astro-081309-130811](https://doi.org/10.1146/annurev-astro-081309-130811)
 11. K. S. Noll, W. M. Grundy, E. I. Chiang, J.-L. Margot, S. D. Kern, in *The Solar System Beyond Neptune*, M. A. Barucci, H. Boehnhardt, D. P. Cruikshank, A. Morbidelli, Eds. (Univ. Arizona Press, 2008), pp. 345–364.
 12. K. S. Noll, W. M. Grundy, D. Nesvorný, A. Thirouin, in *The Trans-Neptunian Solar System*, D. Prialnic, A. Barucci, L.A. Young, Eds. (Elsevier, 2020), 205–224.
 13. A. Morbidelli, H. Rickman, Comets as collisional fragments of a primordial planetesimal disk. *Astron. Astrophys.* **583**, A43 (2016). [doi:10.1051/0004-6361/201526116](https://doi.org/10.1051/0004-6361/201526116)
 14. B. J. R. Davidsson, H. Sierks, C. Güttler, F. Marzari, M. Pajola, H. Rickman, M. F. A’Hearn, A.-T. Auger, M. R. El-Maarry, S. Fornasier, P. J. Gutiérrez, H. U. Keller, M. Massironi, C. Snodgrass, J.-B. Vincent, C. Barbieri, P. L. Lamy, R. Rodrigo, D. Koschny, M. A. Barucci, J.-L. Bertaux, I. Bertini, G. Cremonese, V. Da Deppo, S. Debei, M. De Cecco, C. Feller, M. Fulle, O. Groussin, S. F. Hviid, S. Höfner, W.-H. Ip, L. Jorda, J. Knollenberg, G. Kovacs, J.-R. Kramm, E. Kührt, M. Küppers, F. La Forgia, L. M. Lara, M. Lazzarin, J. J. Lopez Moreno, R. Moissl-Fraund, S. Mottola, G. Naletto, N. Oklay, N. Thomas, C. Tubiana, The primordial nucleus of comet 67P/Churyumov-Gerasimenko. *Astron. Astrophys.* **592**, A63 (2016). [doi:10.1051/0004-6361/201526968](https://doi.org/10.1051/0004-6361/201526968)
 15. M. Jutzi, W. Benz, Formation of bi-lobed shapes by subcatastrophic collisions. A late origin of comet 67P’s structure. *Astron. Astrophys.* **597**, A62 (2017). [doi:10.1051/0004-6361/201628964](https://doi.org/10.1051/0004-6361/201628964)
 16. M. Hirabayashi, D. J. Scheeres, S. R. Chesley, S. Marchi, J. W. McMahon, J. Steckloff, S. Mottola, S. P. Naidu, T. Bowling, Fission and reconfiguration of bilobate comets as

- revealed by 67P/Churyumov-Gerasimenko. *Nature* **534**, 352–355 (2016).
[doi:10.1038/nature17670](https://doi.org/10.1038/nature17670) [Medline](#)
17. K. Batygin, M. E. Brown, W. C. Fraser, Retention of a primordial cold classical Kuiper Belt in an instability-driven model of solar system formation. *Astrophys. J.* **738**, 13 (2011).
[doi:10.1088/0004-637X/738/1/13](https://doi.org/10.1088/0004-637X/738/1/13)
 18. R. I. Dawson, R. A. Murray-Clay, Neptune's wild days: Constraints from the eccentricity distribution of the classical Kuiper belt. *Astrophys. J.* **750**, 43 (2012). [doi:10.1088/0004-637X/750/1/43](https://doi.org/10.1088/0004-637X/750/1/43)
 19. D. Nesvorný, Jumping Neptune can explain the Kuiper Belt kernel. *Astron. J.* **150**, 68 (2015).
[doi:10.1088/0004-6256/150/3/68](https://doi.org/10.1088/0004-6256/150/3/68)
 20. R. S. Gomes, A. Morbidelli, H. F. Levison, Planetary migration in a planetesimal disk: Why did Neptune stop at 30 AU? *Icarus* **170**, 492–507 (2004).
[doi:10.1016/j.icarus.2004.03.011](https://doi.org/10.1016/j.icarus.2004.03.011)
 21. J. N. Cuzzi, R. C. Hogan, K. Shariff, Toward planetesimals: Dense chondrule clumps in the protoplanetary nebula. *Astrophys. J.* **687**, 1432–1447 (2008). [doi:10.1086/591239](https://doi.org/10.1086/591239)
 22. A. N. Youdin, J. Goodman, Streaming instabilities in protoplanetary disks. *Astrophys. J.* **620**, 459–469 (2005). [doi:10.1086/426895](https://doi.org/10.1086/426895)
 23. A. Johansen, J. S. Oishi, M.-M. Mac Low, H. Klahr, T. Henning, A. Youdin, Rapid planetesimal formation in turbulent circumstellar disks. *Nature* **448**, 1022–1025 (2007).
[doi:10.1038/nature06086](https://doi.org/10.1038/nature06086) [Medline](#)
 24. A. Johansen, M. M. Low, P. Lacerda, M. Bizzarro, Growth of asteroids, planetary embryos, and Kuiper belt objects by chondrule accretion. *Sci. Adv.* **1**, e1500109 (2015).
[doi:10.1126/sciadv.1500109](https://doi.org/10.1126/sciadv.1500109) [Medline](#)
 25. J. B. Simon, P. J. Armitage, R. Li, A. N. Youdin, The mass and size distribution of planetesimals formed by the streaming instability. I. The role of self-gravity. *Astrophys. J.* **822**, 55 (2016). [doi:10.3847/0004-637X/822/1/55](https://doi.org/10.3847/0004-637X/822/1/55)
 26. C. P. Abod, J. B. Simon, R. Li, P. J. Armitage, A. N. Youdin, K. A. Kretke, The mass and size distribution of planetesimals formed by the streaming instability. II. The effect of the radial gas pressure gradient. *Astrophys. J.* **883**, 192 (2019). [doi:10.3847/1538-4357/ab40a3](https://doi.org/10.3847/1538-4357/ab40a3)
 27. S. C. Kenyon, B. C. Bromley, Coagulation calculations of icy planet formation at 15–150 AU: A correlation between the maximum radius and the slope of the size distribution for trans-Neptunian objects. *Astron. J.* **143**, 63 (2012). [doi:10.1088/0004-6256/143/3/63](https://doi.org/10.1088/0004-6256/143/3/63)
 28. A. Shannon, Y. Wu, Y. Lithwick, Forming the cold classical Kuiper belt in a light disk. *Astrophys. J.* **818**, 175 (2016). [doi:10.3847/0004-637X/818/2/175](https://doi.org/10.3847/0004-637X/818/2/175)
 29. G. Groussin, N. Attree, Y. Brouet, V. Ciarletti, B. Davidsson, G. Filacchione, H.-H. Fischer, B. Gundlach, M. Knapmeyer, J. Knollenberg, R. Kokotanekova, E. Kührt, C. Leyrat, D. Marshall, I. Pelivan, Y. Skorov, C. Snodgrass, T. Spohn, F. Tosi, The thermal, mechanical, structural, and dielectric properties of cometary nuclei after Rosetta. *Space Sci. Rev.* **215**, 29 (2019). [doi:10.1007/s11214-019-0594-x](https://doi.org/10.1007/s11214-019-0594-x)

30. Short-period, Jupiter-family comets such as 67P are derived mostly from the scattered disk component of the Kuiper Belt (4, 14), which is a distinct, dynamically hot Kuiper Belt population not directly related to CCKB objects. The cold classical region did contribute to the original scattered disk population (17, 19), but we expect this contribution to have been minor compared with the population of the scattered disk as a whole.
31. K. A. Holsapple, K. R. Housen, A crater and its ejecta: An interpretation of Deep Impact. *Icarus* **191**, 586–597 (2007). [doi:10.1016/j.icarus.2006.08.035](https://doi.org/10.1016/j.icarus.2006.08.035)
32. W. D. MacMillan, *The Theory of the Potential* (McGraw-Hill, 1930).
33. M. Jutzi, E. Asphaug, The shape and structure of cometary nuclei as a result of low-velocity accretion. *Science* **348**, 1355–1358 (2015). [doi:10.1126/science.aaa4747](https://doi.org/10.1126/science.aaa4747) [Medline](#)
34. M. Jutzi, W. Benz, A. Toliou, A. Morbidelli, R. Brasser, How primordial is the structure of comet 67P? Combined collisional and dynamical models suggest a late formation. *Astron. Astrophys.* **597**, A61 (2017). [doi:10.1051/0004-6361/201628963](https://doi.org/10.1051/0004-6361/201628963)
35. T. W. Lambe, R. V. Whitman, *Soil Mechanics* (Wiley, ed. 1, 1969).
36. The steepest slopes are found on one of SL's shoulders, coincident with a prominent trough [figure 1 in (8)], possibly a sign of incipient slope failure.
37. C. Matonti, N. Attree, O. Groussin, L. Jorda, S. Viseur, S. F. Hviid, S. Bouley, D. Nébouy, A.-T. Auger, P. L. Lamy, H. Sierks, G. Naletto, R. Rodrigo, D. Koschny, B. Davidsson, M. A. Barucci, J.-L. Bertaux, I. Bertini, D. Bodewits, G. Cremonese, V. Da Deppo, S. Debei, M. De Cecco, J. Deller, S. Fornasier, M. Fulle, P. J. Gutiérrez, C. Güttler, W.-H. Ip, H. U. Keller, L. M. Lara, F. La Forgia, M. Lazzarin, A. Lucchetti, J. J. López-Moreno, F. Marzari, M. Massironi, S. Mottola, N. Oklay, M. Pajola, L. Penasa, F. Preusker, H. Rickman, F. Scholten, X. Shi, I. Toth, C. Tubiana, J.-B. Vincent, Bilobate comet morphology and internal structure controlled by shear deformation. *Nat. Geosci.* **12**, 157–162 (2019). [doi:10.1038/s41561-019-0307-9](https://doi.org/10.1038/s41561-019-0307-9)
38. The strength of the near-surface (<1 m deep) lunar regolith is estimated to be ~1 kPa (98), which is dynamically equivalent to several kilometers depth on Arrokoth.
39. J. E. Richardson, K. J. Graves, A. W. Harris, T. J. Bowling, Small body shapes and spins reveal a prevailing state of maximum topographic stability. *Icarus* **329**, 207–221 (2019). [doi:10.1016/j.icarus.2019.03.027](https://doi.org/10.1016/j.icarus.2019.03.027)
40. S. Greenstreet, B. Gladman, W. B. McKinnon, J. J. Kavelaars, K. N. Singer, Crater density predictions for New Horizons flyby target 2014 MU69. *Astrophys. J.* **872**, L5 (2019). [doi:10.3847/2041-8213/ab01db](https://doi.org/10.3847/2041-8213/ab01db)
41. J. G. Stadel, thesis, University of Washington, Seattle, WA (2001).
42. S. R. Schwartz, D. C. Richardson, P. Michel, An implementation of the soft-sphere discrete element method in a high-performance parallel gravity tree-code. *Granul. Matter* **14**, 363–380 (2012). [doi:10.1007/s10035-012-0346-z](https://doi.org/10.1007/s10035-012-0346-z)
43. Materials and methods are available as supplementary materials.
44. V. S. Safronov, *Evolution of the Protoplanetary Cloud and Formation of the Earth and Planets* [transl. NASA TTF-667] (Nauka, Moscow, 1972)].

45. S. R. Schwartz, P. Michel, M. Jutzi, S. Marchi, Y. Zhang, D. C. Richardson, Catastrophic disruptions as the origin of bilobate comets. *Nature Astron.* **2**, 379–382 (2018). [doi:10.1038/s41550-018-0395-2](https://doi.org/10.1038/s41550-018-0395-2)
46. J.-L. Margot, P. Pravec, P. Taylor, B. Carry, S. Jacobson, in *Asteroids IV*, P. Michel, F. E. DeMeo, W. F. Bottke, Eds. (Univ. Arizona Press, 2015), pp. 355–374.
47. W. C. Fraser, M. T. Bannister, R. E. Pike, M. Marsset, M. E. Schwamb, J. J. Kavelaars, P. Lacerda, D. Nesvorný, K. Volk, A. Delsanti, S. Benecchi, M. J. Lehner, K. Noll, B. Gladman, J.-M. Petit, S. Gwyn, Y.-T. Chen, S.-Y. Wang, M. Alexandersen, T. Burdullis, S. Sheppard, C. Trujillo, All planetesimals born near the Kuiper belt formed as binaries. *Nature Astron.* **1**, 0088 (2017). [doi:10.1038/s41550-017-0088](https://doi.org/10.1038/s41550-017-0088)
48. P. Goldreich, Y. Lithwick, R. Sari, Formation of Kuiper-belt binaries by dynamical friction and three-body encounters. *Nature* **420**, 643–646 (2002). [doi:10.1038/nature01227](https://doi.org/10.1038/nature01227)
[Medline](#)
49. S. A. Astakhov, E. A. Lee, D. Farrelly, Formation of Kuiper-belt binaries through multiple chaotic scattering encounters with low-mass intruders. *Mon. Not. R. Astron. Soc.* **360**, 401–415 (2005). [doi:10.1111/j.1365-2966.2005.09072.x](https://doi.org/10.1111/j.1365-2966.2005.09072.x)
50. S. Weidenschilling, On the origin of binary transneptunian objects. *Icarus* **160**, 212–215 (2002). [doi:10.1006/icar.2002.6952](https://doi.org/10.1006/icar.2002.6952)
51. For Arrokoth, $R_{\text{Hill}} = a_{\odot}(m/3M_{\odot})^{1/3} \sim 4 \times 10^4$ km, where a_{\odot} is Arrokoth's heliocentric distance, m is the mass of Arrokoth (1.6×10^{15} kg for $\rho = 500$ kg m⁻³), and M_{\odot} is the mass of the Sun. The Hill speed, the Keplerian orbital shear velocity at R_{Hill} , is given by $v_{\text{Hill}} = \Omega_{\text{K}} R_{\text{Hill}}$ (where Ω_{K} denotes the orbital frequency of the Keplerian orbit in question), about 2 to 3 cm s⁻¹ for Arrokoth.
52. H. E. Schlichting, R. Sari, Formation of Kuiper Belt binaries. *Astrophys. J.* **673**, 1218–1224 (2008). [doi:10.1086/524930](https://doi.org/10.1086/524930)
53. D. P. Hamilton, J. A. Burns, Orbital stability zones about asteroids. *Icarus* **92**, 118–131 (1991). [doi:10.1016/0019-1035\(91\)90039-V](https://doi.org/10.1016/0019-1035(91)90039-V)
54. H. E. Schlichting, R. Sari, The ratio of retrograde to prograde orbits: A test for Kuiper Belt binary formation theories. *Astrophys. J.* **686**, 741–747 (2008). [doi:10.1086/591073](https://doi.org/10.1086/591073)
55. W. M. Grundy, K. S. Noll, H. G. Roe, M. W. Buie, S. B. Porter, A. H. Parker, D. Nesvorný, H. F. Levison, S. D. Benecchi, D. C. Stephens, C. A. Trujillo, Mutual orbit orientations of transneptunian binaries. *Icarus* **334**, 62–78 (2019). [doi:10.1016/j.icarus.2019.03.035](https://doi.org/10.1016/j.icarus.2019.03.035)
56. S. D. Benecchi, K. S. Noll, W. M. Grundy, M. W. Buie, D. C. Stephens, H. F. Levison, The correlated colors of transneptunian binaries. *Icarus* **200**, 292–303 (2009). [doi:10.1016/j.icarus.2008.10.025](https://doi.org/10.1016/j.icarus.2008.10.025)
57. A. Johansen, J. Blum, H. Tanaka, C. Ormel, M. Bizzarro, H. Rickman, in *Protostars and Planets VI*, H. Beuther, R. S. Klessen, C. P. Dullemond, T. Henning, Eds. (Univ. Arizona Press, 2014), pp. 547–570.
58. D. Nesvorný, A. N. Youdin, D. C. Richardson, Formation of Kuiper belt binaries by gravitational collapse. *Astron. J.* **140**, 785–793 (2010). [doi:10.1088/0004-6256/140/3/785](https://doi.org/10.1088/0004-6256/140/3/785)

59. K. Noll, W. Grundy, S. Benecchi, H. Levison, The relative sizes of transneptunian binaries: Evidence for different populations from a homogeneous data set. EPSC-DPS Joint Meeting 2011, 2-7 October 2011 in Nantes, France; <https://meetingorganizer.copernicus.org/EPSC-DPS2011/EPSC-DPS2011-1029.pdf>.
60. D. Nesvorný, R. Li, A. N. Youdin, J. B. Simon, W. M. Grundy, Trans-Neptunian binaries as evidence for planetesimal formation by the streaming instability. *Nature Astron.* **3**, 808–812 (2019). [doi:10.1038/s41550-019-0806-z](https://doi.org/10.1038/s41550-019-0806-z)
61. J. B. Simon, P. J. Armitage, A. N. Youdin, R. Li, Evidence for universality in the initial planetesimal mass function. *Astrophys. J.* **847**, L12–L17 (2017). [doi:10.3847/2041-8213/aa8c79](https://doi.org/10.3847/2041-8213/aa8c79)
62. R. Li, A. N. Youdin, J. B. Simon, On the numerical robustness of the streaming instability: Particle concentration and gas dynamics in protoplanetary disks. *Astrophys. J.* **862**, 14–29 (2018). [doi:10.3847/1538-4357/aaca99](https://doi.org/10.3847/1538-4357/aaca99)
63. D. Carrera, U. Gorti, A. Johansen, M. B. Davies, Planetesimal formation by the streaming instability in a photoevaporating disk. *Astrophys. J.* **839**, 16 (2017). [doi:10.3847/1538-4357/aa6932](https://doi.org/10.3847/1538-4357/aa6932)
64. S. Chandrasekhar, *Ellipsoidal Figures of Equilibrium* (Yale Univ. Press, 1969).
65. P. Descamps, Roche figures of doubly synchronous asteroids. *Planet. Space Sci.* **56**, 1839–1846 (2008). [doi:10.1016/j.pss.2008.02.040](https://doi.org/10.1016/j.pss.2008.02.040)
66. M. Čuk, D. Nesvorný, Orbital evolution of small binary asteroids. *Icarus* **207**, 732–743 (2010). [doi:10.1016/j.icarus.2009.12.005](https://doi.org/10.1016/j.icarus.2009.12.005)
67. J. Fang, J.-L. Margot, Near-earth binaries and triples: Origin and evolution of spin-orbital properties. *Astron. J.* **143**, 24 (2012). [doi:10.1088/0004-6256/143/1/24](https://doi.org/10.1088/0004-6256/143/1/24)
68. S. Naoz, The eccentric Kozai-Lidov effect and its applications. *Annu. Rev. Astron. Astrophys.* **54**, 441–489 (2016). [doi:10.1146/annurev-astro-081915-023315](https://doi.org/10.1146/annurev-astro-081915-023315)
69. L. A. M. Benner, W. B. McKinnon, Orbital behavior of captured satellites: The effect of solar gravity on Triton’s post-capture orbit. *Icarus* **114**, 1–20 (1989). [doi:10.1006/icar.1995.1039](https://doi.org/10.1006/icar.1995.1039)
70. H. B. Perets, S. Naoz, Kozai cycles, tidal friction, and the dynamical evolution of binary minor planets. *Astrophys. J.* **699**, L17–L21 (2009). [doi:10.1088/0004-637X/699/1/L17](https://doi.org/10.1088/0004-637X/699/1/L17)
71. S. B. Porter, W. M. Grundy, KCTF evolution of trans-neptunian binaries: Connecting formation to observation. *Icarus* **220**, 947–957 (2012). [doi:10.1016/j.icarus.2012.06.034](https://doi.org/10.1016/j.icarus.2012.06.034)
72. We estimate a gravitational quadrupole $J_2 \sim 0.14$ for LL from its shape (8), so the semimajor axis where the dynamics transition from shape-driven precession to solar (Kozai-Lidov) is $\left(2J_2 M_{\text{LL}} R_{\text{LL}}^2 a_{\odot}^3 / M_{\odot}\right)^{1/5}$, where M_{LL} and R_{LL} are the mass and spherical equivalent radius of the large lobe, respectively (67).
73. S. D. Benecchi, K. S. Noll, W. M. Grundy, H. F. Levison, (47171) 1999 TC₃₆, A transneptunian triple. *Icarus* **207**, 978–991 (2010). [doi:10.1016/j.icarus.2009.12.017](https://doi.org/10.1016/j.icarus.2009.12.017)

74. K. J. Walsh, S. A. Jacobson, in *Asteroids IV*, P. Michel, F. E. DeMeo, W. F. Bottke, Eds. (Univ. Arizona Press, 2015), pp. 375–393.
75. D. Vokrouhlický, W. F. Bottke, S. R. Chesley, D. J. Scheeres, T. S. Statler, in *Asteroids IV*, P. Michel, F. E. DeMeo, W. F. Bottke, Eds. (Univ. Arizona Press, 2015), pp. 509–531.
76. M. Čuk, J. A. Burns, Effects of thermal radiation on the dynamics of binary NEAs. *Icarus* **176**, 418–431 (2005). [doi:10.1016/j.icarus.2005.02.001](https://doi.org/10.1016/j.icarus.2005.02.001)
77. S. A. Jacobson, F. Marzari, A. Rossi, D. J. Scheeres, Matching asteroid population characteristics with a model constructed from the YORP-induced rotational fission hypothesis. *Icarus* **277**, 381–394 (2016). [doi:10.1016/j.icarus.2016.05.032](https://doi.org/10.1016/j.icarus.2016.05.032)
78. BYORP tends to reset outwardly migrating satellites so that they turn and migrate inward (76), making an ultimate merger of the two bodies the most likely outcome, as long as the secondary can reestablish its synchronous spin. This more complicated evolution depends on increasing orbital eccentricity as the binary orbit expands, which has been questioned (67).
79. O. Golubov, Yu. N. Krugly, Tangential component of the YORP effect. *Astrophys. J.* **752**, L11 (2012). [doi:10.1088/2041-8205/752/1/L11](https://doi.org/10.1088/2041-8205/752/1/L11)
80. P. Ševeček, M. Brož, D. Čapek, J. Ďurech, The thermal emission from boulders on (25143) Itokawa and general implications for the YORP effect. *Mon. Not. R. Astron. Soc.* **450**, 2104–2115 (2015). [doi:10.1093/mnras/stv738](https://doi.org/10.1093/mnras/stv738)
81. D. P. Rubincam, Radiative spin-up and spin-down of small asteroids. *Icarus* **148**, 2–11 (2000). [doi:10.1006/icar.2000.6485](https://doi.org/10.1006/icar.2000.6485)
82. M. C. Nolan, E. S. Howell, D. J. Scheeres, J. W. McMahon, O. Golubov, C. W. Hergenrother, J. P. Emery, K. S. Noll, S. R. Chesley, D. S. Lauretta, Detection of rotational acceleration of Bennu using HST light curve observations. *Geophys. Res. Lett.* **46**, 1956–1962 (2019). [doi:10.1029/2018GL080658](https://doi.org/10.1029/2018GL080658)
83. C. D. Murray, S. F. Dermott, *Solar System Dynamics* (Cambridge Univ. Press, 2000).
84. P. Goldreich, R. Sari, Tidal evolution of rubble piles. *Astrophys. J.* **691**, 54–60 (2009). [doi:10.1088/0004-637X/691/1/54](https://doi.org/10.1088/0004-637X/691/1/54)
85. The critical rotation rate (or surface orbital frequency) of a rigid sphere is given by $\sqrt{GM/R^3}$, where M and R are its mass and radius, respectively. This approximately corresponds to the maximum specific angular momentum of a rotating, strengthless body (a Jacobi ellipsoid) (99).
86. The second-degree Love number k_2 for a porous, granular SL is given by $\sim 10^{-5} \times \text{SL's effective radius in kilometers}$ (84), and the tidal dissipation factor Q for SL is taken to be 10 to 100.
87. D. Nesvorný, J. Parker, D. Vokrouhlický, Bi-lobed shape of comet 67P from a collapsed binary. *Astron. J.* **155**, 246 (2018). [doi:10.3847/1538-3881/aac01f](https://doi.org/10.3847/1538-3881/aac01f)
88. H. B. Perets, R. A. Murray-Clay, Wind-shearing in gaseous protoplanetary disks and the evolution of binary planetesimals. *Astrophys. J.* **733**, 56 (2011). [doi:10.1088/0004-637X/733/1/56](https://doi.org/10.1088/0004-637X/733/1/56)

89. F. L. Whipple, in *From Plasma to Planet, Proc. Twenty-First Nobel Symposium*, A. Evlius, Ed. (Wiley Interscience, 1972), pp. 211–232.
90. S. J. Desch, Mass distribution and planet formation in the solar nebula. *Astrophys. J.* **671**, 878–893 (2007). [doi:10.1086/522825](https://doi.org/10.1086/522825)
91. H. Wang, B. P. Weiss, X.-N. Bai, B. G. Downey, J. Wang, J. Wang, C. Suavet, R. R. Fu, M. E. Zucolotto, Lifetime of the solar nebula constrained by meteorite paleomagnetism. *Science* **355**, 623–627 (2017). [doi:10.1126/science.aaf5043](https://doi.org/10.1126/science.aaf5043) [Medline](#)
92. S. J. Weidenschilling, Aerodynamics of solid bodies in the solar nebula. *Mon. Not. R. Astron. Soc.* **180**, 57–70 (1977). [doi:10.1093/mnras/180.2.57](https://doi.org/10.1093/mnras/180.2.57)
93. The mean free path in the nebular gas $\lambda = 1/n\sigma_H$, where σ_H is the collisional cross section of H_2 ($2 \times 10^{-19} \text{ m}^2$) and the number density $n \equiv \rho_g/\mu m_H$ (with m_H the mass of a hydrogen atom and $\mu = 2.3$ for solar composition gas). For the midplane density quoted in the text, we found $\lambda \sim 0.17 \text{ km}$, which puts Arrokoth's gas drag interactions into the fluid (Stokes-like) regime. The kinematic viscosity is then $\lambda \times$ sound speed, which for cold, 30 K nebular gas (90, 100) is $7 \times 10^4 \text{ m}^2 \text{ s}^{-1}$, independent of ρ_g .
94. P. R. Estrada, J. N. Cuzzi, D. Morgan, Global modeling of nebulae with particle growth, drift, and evaporation fronts. I. Methodology and typical results. *Astrophys. J.* **818**, 200 (2016). [doi:10.3847/0004-637X/818/2/200](https://doi.org/10.3847/0004-637X/818/2/200)
95. F. C. Adams, D. Hollenbach, G. Laughlin, U. Gorti, Photoevaporation of circumstellar disks due to external far-ultraviolet radiation in stellar aggregates. *Astrophys. J.* **611**, 360–379 (2004). [doi:10.1086/421989](https://doi.org/10.1086/421989)
96. Total, upper limit merger times, if Arrokoth's original orbit extended to its outer Hill sphere ($\sim 10^4 \text{ km}$), would have been a few times longer.
97. C. Hayashi, Structure of the solar nebula, growth and decay of magnetic fields and effects of magnetic and turbulent viscosities on the nebula. *Prog. Theor. Phys. Suppl.* **70**, 35–53 (1981). [doi:10.1143/PTPS.70.35](https://doi.org/10.1143/PTPS.70.35)
98. J. K. Mitchell, L. G. Bromwell, W. D. Carrier III, N. C. Costes, R. F. Scott, Soil mechanical properties at the Apollo 14 site. *J. Geophys. Res.* **77**, 5641–5664 (1972). [doi:10.1029/JB077i029p05641](https://doi.org/10.1029/JB077i029p05641)
99. P. Descamps, F. Marchis, Angular momentum of binary asteroids: Implications for their possible origin. *Icarus* **193**, 74–84 (2008). [doi:10.1016/j.icarus.2007.07.024](https://doi.org/10.1016/j.icarus.2007.07.024)
100. E. I. Chiang, P. Goldreich, Spectral energy distributions of T Tauri stars with passive circumstellar disks. *Astrophys. J.* **490**, 368–376 (1997). [doi:10.1086/304869](https://doi.org/10.1086/304869)
101. Y. Zhang, D. C. Richardson, O. S. Barnouin, C. Maurel, P. Michel, S. R. Schwartz, R.-L. Ballouz, L. A. M. Benner, S. P. Naidu, J. Li, Creep stability of the proposed AIDA mission target 65803 Didymos: I. Discrete cohesionless granular physics model. *Icarus* **294**, 98–123 (2017). [doi:10.1016/j.icarus.2017.04.027](https://doi.org/10.1016/j.icarus.2017.04.027)
102. Y. Zhang, D. C. Richardson, O. S. Barnouin, P. Michel, S. R. Schwartz, R.-L. Ballouz, Rotational failure of rubble-pile bodies: Influences of shear and cohesive strengths. *Astrophys. J.* **857**, 15 (2018). [doi:10.3847/1538-4357/aab5b2](https://doi.org/10.3847/1538-4357/aab5b2)

103. W. C. Fraser, M. E. Brown, A. Morbidelli, A. Parker, K. Batygin, The absolute magnitude distribution of Kuiper Belt objects. *Astrophys. J.* **782**, 100 (2014). [doi:10.1088/0004-637X/782/2/100](https://doi.org/10.1088/0004-637X/782/2/100)
104. A. Morbidelli, D. Nesvorný, in *The Transneptunian Solar System*, D. Prialnik, M. A. Barucci, L. A. Young, Eds. (Elsevier, 2020), 25–59.
105. M. Pan, R. Sari, Shaping the Kuiper belt size distribution by shattering large but strengthless bodies. *Icarus* **173**, 342–348 (2005). [doi:10.1016/j.icarus.2004.09.004](https://doi.org/10.1016/j.icarus.2004.09.004)
106. D. Nesvorný, D. Vokrouhlický, W. F. Bottke, K. Noll, H. F. Levison, Observed binary fraction sets limits on the extent of collisional grinding in the Kuiper belt. *Astron. J.* **141**, 159 (2011). [doi:10.1088/0004-6256/141/5/159](https://doi.org/10.1088/0004-6256/141/5/159)
107. H. Klahr, T. Pfeil, A. Schreiber, Instabilities and flow structures in protoplanetary disks: Setting the stage for planetesimal formation, in *Handbook of Exoplanets*, H. Deeg, J. Belmonte J., Eds. (Springer, 2018), pp. 2251–2286.
108. J. N. Cuzzi, R. C. Hogan, W. F. Bottke, Towards initial mass functions for asteroids and Kuiper belt objects. *Icarus* **208**, 518–538 (2010). [doi:10.1016/j.icarus.2010.03.005](https://doi.org/10.1016/j.icarus.2010.03.005)
109. C. T. Lenz, H. Klahr, T. Birnstiel, Planetesimal population synthesis: Pebble flux-regulated planetesimal formation. *Astrophys. J.* **874**, 36 (2019). [doi:10.3847/1538-4357/ab05d9](https://doi.org/10.3847/1538-4357/ab05d9)
110. C.-C. Yang, A. Johansen, D. Carrera, Concentrating small particles in protoplanetary disks through the streaming instability. *Astron. Astrophys.* **606**, A80 (2017). [doi:10.1051/0004-6361/201630106](https://doi.org/10.1051/0004-6361/201630106)
111. J. B. Simon, X.-N. Bai, K. M. Flaherty, A. M. Hughes, Origin of weak turbulence in the outer regions of protoplanetary disks. *Astrophys. J.* **865**, 10 (2018). [doi:10.3847/1538-4357/aad86d](https://doi.org/10.3847/1538-4357/aad86d)
112. O. M. Umurhan, P. R. Estrada, J. N. Cuzzi, Streaming instability: Saturation in turbulent protoplanetary disks. [arXiv:1906.05371](https://arxiv.org/abs/1906.05371) [astro-ph.EP] (2019).
113. X.-N. Bai, J. M. Stone, Wind-driven accretion in protoplanetary disks. I. Suppression of the magnetorotational instability and launching of the magnetocentrifugal wind. *Astrophys. J.* **769**, 76 (2013). [doi:10.1088/0004-637X/769/1/76](https://doi.org/10.1088/0004-637X/769/1/76)
114. A. Johansen, H. Klahr, Th. Henning, High-resolution simulations of planetesimal formation in turbulent protoplanetary discs. *Astron. Astrophys.* **529**, A62 (2011). [doi:10.1051/0004-6361/201015979](https://doi.org/10.1051/0004-6361/201015979)
115. S. Marchi, C. R. Chapman, O. S. Barnouin, J. E. Richardson, J.-B. Vincent, in *Asteroids IV*, P. Michel, F. E. DeMeo, W. F. Bottke, Eds. (Univ. Arizona Press, 2015), pp. 725–744.
116. P. A. Taylor, J.-L. Margot, D. Vokrouhlický, D. J. Scheeres, P. Pravec, S. C. Lowry, A. Fitzsimmons, M. C. Nolan, S. J. Ostro, L. A. M. Benner, J. D. Giorgini, C. Magri, Spin rate of asteroid (54509) 2000 PH5 increasing due to the YORP effect. *Science* **316**, 274–277 (2007). [doi:10.1126/science.1139038](https://doi.org/10.1126/science.1139038) [Medline](#)
117. J. Ďurech, D. Vokrouhlický, M. Kaasalainen, P. Weissman, S. C. Lowry, E. Beshore, D. Higgins, Y. N. Krugly, V. G. Shevchenko, N. M. Gaftonyuk, Y.-J. Choi, R. A. Kowalski, S. Larson, B. D. Warner, A. L. Marshalkina, M. A. Ibrahimov, I. E. Molotov, T.

Michałowski, K. Kitazato, New photometric observations of asteroids (1862) Apollo and (25143) Itokawa—an analysis of YORP effect. *Astron. Astrophys.* **488**, 345–350 (2008). [doi:10.1051/0004-6361/200809663](https://doi.org/10.1051/0004-6361/200809663)

118. J. Ďurech, D. Vokrouhlický, P. Pravec, J. Hanuš, D. Farnocchia, Y. N. Krugly, R. Y. Inasaridze, V. R. Ayvazian, P. Fatka, V. G. Chiorny, N. Gaftonyuk, A. Galád, R. Groom, K. Hornoch, H. Kučáková, P. Kušnirák, M. Lehký, O. I. Kvaratskhelia, G. Masi, I. E. Molotov, J. Oey, J. T. Pollock, V. G. Shevchenko, J. Vraštil, B. D. Warner, YORP and Yarkovsky effects in asteroids (1685) Toro, (2100) Ra-Shalom, (3103) Eger, and (161989) Cacus. *Astron. Astrophys.* **609**, A86 (2018). [doi:10.1051/0004-6361/201731465](https://doi.org/10.1051/0004-6361/201731465)