

## Supplementary Online Content

Nho K, Kueider-Paisley A, Ahmad S, et al; Alzheimer's Disease Neuroimaging Initiative and the Alzheimer Disease Metabolomics Consortium. Association of altered liver enzymes with Alzheimer disease diagnosis, cognition, neuroimaging measures, and cerebrospinal fluid biomarkers. *JAMA Netw Open*. 2019;2(7):e197978.

doi:10.1001/jamanetworkopen.2019.7978

**eTable 1.** Sensitivity Analysis for Alzheimer Disease Diagnosis (CN vs AD) Group Differences in Liver Function Biomarkers

**eTable 2.** Demographic Information of ADNI Participants

**eTable 3.** Diagnostic Group Differences in Liver Function Biomarkers

**eTable 4.** Diagnostic Group Differences of Liver Function Biomarkers With Alzheimer Disease Diagnosis (CV vs AD) Adjusted for Gamma-Glutamyltransferase (GGT) and Statin Use

**eTable 5.** Association of Liver Function Biomarkers With Cognition Adjusted for Gamma-Glutamyltransferase (GGT) and Statin Use

**eFigure.** Liver Function Biomarkers and Their Association With A/T/N Biomarkers for Alzheimer Disease Adjusted for Gamma-Glutamyltransferase (GGT) and Statin Use

This supplementary material has been provided by the authors to give readers additional information about their work.

**eTable 1.** Sensitivity Analysis for Alzheimer Disease Diagnosis (CN vs AD)  
Group Differences in Liver Function Biomarkers

Liver function marker	excluding outliers			keeping outliers		
	Odds Ratio	95% CI	Corrected P Value	Odds Ratio	95% CI	Corrected P Value
<b>Albumin</b>	5.789	0.040,843.993	$4.90 \times 10^{-1}$	5.789	0.040,843.993	$4.90 \times 10^{-1}$
<b>Alkaline Phosphatase</b>	3.620	0.844,15.529	$1.25 \times 10^{-1}$	3.620	0.844,15.529	$1.25 \times 10^{-1}$
<b>ALT</b>	0.133	0.042,0.422	$3.63 \times 10^{-3}$	0.120	0.039,0.374	$2.55 \times 10^{-3}$
<b>AST</b>	0.229	0.045,1.175	$1.25 \times 10^{-1}$	0.203	0.041,1.021	$1.06 \times 10^{-1}$
<b>AST:ALT</b>	7.932	1.673,37.617	$2.73 \times 10^{-2}$	9.815	2.156,44.670	$9.41 \times 10^{-3}$
<b>Total Bilirubin</b>	1.405	0.585,3.377	$4.90 \times 10^{-1}$	1.405	0.585,3.377	$4.90 \times 10^{-1}$

*Abbreviations:* AST: Aspartate aminotransferase; ALT: alanine aminotransferase. Age, sex, body mass index, and *APOE*  $\epsilon$ 4 status included as covariates

**eTable 2.** Demographic Information of ADNI Participants

Variable	CN (N=407)	SMC (N=20)	EMCI (N=298)	LMCI (N=544)	AD (N=312)	P Value
Age (years)	74.5 (5.7)	71.6 (5.1)	70.9 (7.4)	73.5 (7.6)	74.6 (7.7)	1.94x10 <sup>-11</sup>
Sex: (Male/Female)	206/201	10/10	166/132	331/213	171/141	3.35x10 <sup>-2</sup>
Education (years)	16.3 (2.7)	16.8 (2.2)	16.0 (2.6)	15.9 (3.0)	15.1 (3.0)	1.59x10 <sup>-6</sup>
BMI (Kg/M <sup>2</sup> )	27.4 (4.5)	29.3 (4.6)	28.4 (5.3)	26.9 (4.3)	26.1 (4.4)	6.94x10 <sup>-9</sup>
APOE ε4 status (0/1)	294/113	15/5	169/129	249/295	98/214	<2.20x10 <sup>-16</sup>
Albumin (g/dL)	4.09 (0.31)	3.94 (0.25)	4.10 (0.29)	4.14 (0.32)	4.11 (0.32)	1.29x10 <sup>-2</sup>
Alkaline phosphatase (U/L)	75.3 (20.7)	74.4 (14.1)	71.1 (17.9)	74.9 (22.4)	76.2 (19.2)	2.09x10 <sup>-2</sup>
ALT (U/L)	21.5 (8.4)	24.2 (6.6)	22.2 (9.3)	20.7 (8.0)	19.5 (7.0)	5.44x10 <sup>-5</sup>
AST (U/L)	24.3 (6.2)	25.2 (8.0)	24.7 (7.4)	24.1 (6.1)	23.7 (6.3)	4.66x10 <sup>-1</sup>
AST:ALT	1.22 (0.32)	1.06 (0.22)	1.18 (0.31)	1.25 (0.31)	1.29 (0.35)	1.22x10 <sup>-5</sup>
Total bilirubin (mg/dL)	0.52 (0.26)	0.44 (0.14)	0.54 (0.28)	0.53 (0.24)	0.56 (0.30)	3.02x10 <sup>-1</sup>
Memory composite score	0.94 (0.53)	0.99 (0.57)	0.53 (0.49)	-0.04 (0.57)	-0.77 (0.54)	<2.20x10 <sup>-16</sup>
Executive function composite score	0.75 (0.71)	0.82 (0.64)	0.47 (0.76)	0.01 (0.80)	-0.88 (0.82)	<2.20x10 <sup>-16</sup>

Abbreviations: AD: Alzheimer's Disease; ALT: alanine aminotransferase; AST: Aspartate aminotransferase; BMI: Body Mass Index; CN: Cognitively Normal; EMCI: early mild cognitive impairment; LMCI: late MCI; SMC: subjective memory concerns

**eTable 3.** Diagnostic Group Differences in Liver Function Biomarkers

Liver function marker	CN vs EMCI		CN vs LMCI		CN vs AD	
	Odds Ratio	95% CI	Odds Ratio	95% CI	Odds Ratio	95% CI
Albumin (g/dL)	0.811	0.006,117.8	19.623	0.333,1156.4	5.789	0.040,844.0
Alkaline Phosphatase (U/L)	0.144	0.035,0.588	0.883	0.290,2.694	3.620	0.844,15.53
ALT (U/L)	0.561	0.196,1.604	0.291	0.112,0.755	0.133	0.042,0.422
AST (U/L)	0.742	0.170,3.230	0.393	0.102,1.506	0.229	0.045,1.175
AST:ALT	1.240	0.278,5.540	3.783	0.995,14.38	7.932	1.673,37.62
Total Bilirubin (mg/dL)	1.319	0.556,3.133	0.887	0.417,1.886	1.405	0.585,3.377

Abbreviations: AD: Alzheimer's Disease; ALT: alanine aminotransferase; AST: Aspartate aminotransferase; BMI: Body Mass Index; CN: Cognitively Normal; EMCI: early mild cognitive impairment; LMCI: late MCI. Age, sex, body mass index, and APOE  $\epsilon$ 4 status were included as covariates.

**eTable 4.** Diagnostic Group Differences of Liver Function Biomarkers With Alzheimer Disease Diagnosis (CV vs AD) Adjusted for Gamma-Glutamyltransferase (GGT) and Statin Use

Liver function marker			
	Odds Ratio	95% CI	Corrected <i>P</i> Value
Albumin (g/dL)	5.595	0.037, 840.4	5.01 x 10 <sup>-1</sup>
Alkaline Phosphatase (U/L)	4.296	0.982, 18.79	1.06 x 10 <sup>-1</sup>
ALT (U/L)	0.112	0.032, 0.386	3.16 x 10 <sup>-3</sup>
AST (U/L)	0.246	0.045, 1.348	1.59 x 10 <sup>-1</sup>
AST:ALT	8.580	1.713, 42.98	2.68 x 10 <sup>-2</sup>
Total Bilirubin (mg/dL)	1.461	0.603, 3.539	4.82 x 10 <sup>-1</sup>


*Abbreviations:* AST: Aspartate aminotransferase; ALT: alanine aminotransferase

**eTable 5.** Association of Liver Function Biomarkers With Cognition Adjusted for Gamma-Glutamyltransferase (GGT) and Statin Use

Liver function markers	Memory			Executive function		
	$\beta$	SE	Corrected <i>P</i> Value	$\beta$	SE	Corrected <i>P</i> Value
Albumin (g/dL)	-0.843	0.578	$1.93 \times 10^{-1}$	-0.238	0.691	$7.31 \times 10^{-1}$
Alkaline Phosphatase (U/L)	-0.427	0.163	$1.51 \times 10^{-2}$	-0.607	0.194	$5.52 \times 10^{-3}$
ALT (U/L)	0.476	0.139	$3.62 \times 10^{-3}$	0.733	0.165	$1.18 \times 10^{-4}$
AST (U/L)	0.366	0.187	$7.62 \times 10^{-2}$	0.631	0.224	$1.16 \times 10^{-2}$
AST:ALT	-0.511	0.188	$1.33 \times 10^{-2}$	-0.709	0.224	$5.52 \times 10^{-3}$
Total Bilirubin (mg/dL)	-0.066	0.103	$6.22 \times 10^{-1}$	-0.070	0.123	$6.22 \times 10^{-1}$

*Abbreviations:* AST: Aspartate aminotransferase; ALT: alanine aminotransferase; GGT: gamma-glutamyl transferase. Covariates included age, sex, education, body mass index, *APOE*  $\epsilon$ 4 status, statins use, and gamma-glutamyl transferase.

**eFigure.** Liver Function Biomarkers and Their Association With A/T/N Biomarkers for Alzheimer Disease Adjusted for Gamma-Glutamyltransferase (GGT) and Statin Use



**Supplementary Figure 1: Liver function biomarkers and their association with ATN biomarkers for Alzheimer’s disease adjusted for statin use and gamma-glutamyltransferase.** Heat map of q-values of association between liver function biomarkers and the “A/T/N” biomarkers for Alzheimer’s disease. P-values estimated from linear regression analyses were corrected for multiple testing using FDR (q-value). Color code: white indicates q-value>0.05, red indicates significant positive associations, and green indicates significant negative associations.