

# Nuclear Disintegration Energies. II.\*†

D. M. VAN PATTER, *Bartol Research Foundation, Swarthmore, Pennsylvania*

AND

WARD WHALING, *California Institute of Technology, Pasadena, California*

## I. INTRODUCTION

THIS supplement is intended to augment the compilation, "Nuclear Disintegration Energies,"<sup>1</sup> which included measurements of reaction energies received up to May, 1954. The data contained in the original compilation, together with other data pertaining to mass differences<sup>2</sup> have been of use in the preparation of tables of masses<sup>3-5</sup> and mass links,<sup>6</sup> and for the comparison of masses or mass differences obtained from nuclear reaction energies and mass-spectroscopic measurements.<sup>3,7</sup>

Since publication of the original compilation, there has been a considerable number of new measurements of nuclear reaction energies, e.g., measurements of some eighty previously undetermined reaction energies have been reported for  $A < 50$ . The systematic program of precise  $Q$ -value determinations by the group at Massachusetts Institute of Technology headed by W. W. Buechner has been extended up to the mass region  $A = 32-60$ . Together with data concerning beta-decay energies, these measurements have made possible the calculation of masses in the region from S<sup>33</sup> to Ti<sup>46</sup> from nuclear reaction energies only.<sup>8</sup>

In his systematic examination of masses and mass differences for  $A < 202$ , Wapstra<sup>9</sup> pointed out several instances of  $Q$  values listed in our original compilation<sup>1</sup>

\* The contribution of D. M. Van Patter was supported in part by the U. S. Air Force, through the Office of Scientific Research of the Air Research and Development Command.

† This paper, and the three which follow, by H. E. Duckworth, L. J. Lidofsky, and by F. Asaro and I. Perlman, were prepared at the suggestion of the Subcommittee on Nuclear Constants of the Committee on Nuclear Science of the National Research Council. This subcommittee consists of Ward Whaling, *chairman*, and F. A. Selove, G. A. Bartholomew, H. E. Duckworth, I. Perlman, W. H. Sullivan, and D. M. Van Patter, together with L. Lidofsky as consultant.

Reprints of these four papers as a group may be obtained from the Publications Office, National Research Council, 2101 Constitution Avenue, Washington 25, D. C.

<sup>1</sup> D. M. Van Patter and W. Whaling, *Rev. Modern Phys.* **26**, 402 (1954).

<sup>2</sup> R. W. King, *Rev. Modern Phys.* **26**, 327 (1954); Geschwind, Gunther-Mohr, and Townes, *Rev. Modern Phys.* **26**, 444 (1954); F. Asaro and I. Perlman, *Rev. Modern Phys.* **26**, 456 (1954); Duckworth, Hogg, and Pennington, *Rev. Modern Phys.* **26**, 463 (1954).

<sup>3</sup> A. H. Wapstra, *Physica* **21**, 367 (1955); *ibid.* **21**, 385 (1955).

<sup>4</sup> J. R. Huizenga, *Physica* **21**, 410 (1955).

<sup>5</sup> Mattauch, Waldmann, Bieri, and Everling, *Z. Naturforsch.* **11A**, 525 (1956).

<sup>6</sup> Way, King, McGinnis, and van Lieshout, "Nuclear Level Schemes," TID-5300 (USAEC September 1955).

<sup>7</sup> M. E. Kettner, *Phys. Rev.* **102**, 1065 (1956); Scolman, Quisenberry, and Nier, *Phys. Rev.* **102**, 1076 (1956).

<sup>8</sup> Endt, Buechner, Braams, Paris, and Sperduto, *Phys. Rev.* **105**, 1002 (1957).

which should be assigned to transitions to excited states, rather than ground-state transitions. He also discovered discrepancies between nuclear reaction and mass-spectroscopic data, particularly in the region of the nickel isotopes. On the basis of his analysis, he also reassigned some of the gamma rays observed from neutron capture by Kinsey and Bartholomew to other isotopes. A similar detailed analysis of  $(n,\gamma)$  and  $(\gamma,n)$  transitions has been made by Way *et al.*<sup>6,9</sup> in order to determine neutron binding energies in the region of  $A = 40-92$ . More recently, Quisenberry, Scolman, and Nier,<sup>10</sup> on the basis of their new mass measurements in the region Fe to Zn, have reassigned several neutron capture gamma rays to new isotopes, particularly for the nickel and zinc isotopes. It can be seen from the above brief summary that a considerable alteration of the nuclear reaction energies contained in the original compilation is now necessary.

This supplement contains new measurements of nuclear reaction energies available to the authors up to February 1, 1957. In addition, any measured values listed in the original compilation which have been revised or reassigned since its publication have been included.

## II. ARRANGEMENT OF TABLE I

The arrangement of Table I of this supplement is the same as in the original compilation.<sup>1</sup> In Columns 1 and 2, the reactions and energy determinations are listed. In the case of a reaction shown as  $(n,\gamma,\gamma)$ , the ground-state transition was not observed, and the reaction energy shown was determined from the summing of two gamma-ray energies. Some of the experimental values, designated by the superscript "a," do not appear explicitly in the reference cited, and have been calculated by the present authors from the pertinent experimental data. If it was necessary to use masses in such a calculation, those given by Wapstra<sup>9</sup> have been used. When there is doubt that a reported  $Q$  value represents the ground-state transition, it has been enclosed in parentheses.

In Columns 3 and 4, the method and energy standards used are listed, if stated by the authors. In Column 5, the most recent reference for each measured value is

<sup>9</sup> The authors wish to acknowledge the valuable contributions of the Nuclear Data Group under the leadership of K. Way. Their compilation "Nuclear Level Schemes," for  $40 \leq A \leq 92$ , and their tables of ground-state  $Q$  values in *Nuclear Science Abstracts* have aided considerably the preparation of this present supplement.

<sup>10</sup> Quisenberry, Scolman, and Nier, *Phys. Rev.* **104**, 461 (1956).

TABLE I. Nuclear reaction energies.

Reaction	Measured <i>Q</i> value (Mev)	Method	Calibration Energy standard	Reference	Average <i>Q</i> value (Mev)
D <sup>2</sup> ( <i>d,p</i> )T <sup>3</sup>	4.044±0.005	el spec	Li <sup>7</sup> ( <i>p,n</i> )Be <sup>7</sup>	Do 56a	4.038±0.005
D <sup>2</sup> ( <i>d,n</i> )He <sup>3</sup>	3.276±0.024	ph pl	Ro 51c	Su 55	3.267±0.007
	3.271±0.011	el spec	Li <sup>7</sup> ( <i>p,n</i> )Be <sup>7</sup>	Do 56a	
He <sup>4</sup> ( <i>d,p</i> )He <sup>6</sup>	-3.3±0.1	ph pl		Fr 54a	-3.3±0.1
Li <sup>6</sup> ( <i>n,γ</i> )Li <sup>7</sup>	7.26±0.05	pr spec		Ba 56	
Li <sup>6</sup> ( <i>p,d</i> )Li <sup>5</sup>	-3.0±0.15	scint spec		Li 55	preliminary
Li <sup>6</sup> ( <i>p,γ</i> )Be <sup>7</sup>	5.66±0.03	scint spec	F <sup>19</sup> ( <i>p,αγ</i> )O <sup>16</sup>	Wa 56	
Li <sup>6</sup> ( <i>d,He<sup>3</sup></i> )He <sup>6</sup>	0.91±0.09	mag spec	Po α	Le 55	
Li <sup>6</sup> ( <i>t,α</i> )He <sup>5</sup>	15.15±0.04	pulse ht	Th C' α	Cr 56a	
Li <sup>6</sup> ( <i>t,d</i> )Li <sup>7</sup>	0.986±0.007	mag spec	Li <sup>6</sup> ( <i>p,α</i> )He <sup>3</sup>	Pe 52, Al 54*	0.986±0.007 <sup>+</sup>
Li <sup>6</sup> ( <i>t,p</i> )Li <sup>8</sup>	0.790±0.011	mag spec	Li <sup>6</sup> ( <i>p,α</i> ), Li <sup>6</sup> ( <i>t,d</i> )Li <sup>7*</sup>	Pe 52, Al 54*	
Li <sup>6</sup> (He <sup>3</sup> , <i>n</i> )B <sup>8</sup>	-1.976±0.006	threshold		Du 56	
Li <sup>7</sup> ( <i>p,n</i> )Be <sup>7</sup>	-1.645 <sub>2</sub> ±0.001	threshold	Mg <sup>24</sup> ( <i>p,p'</i> ), Na <sup>24</sup> γ, Au <sup>198</sup> γ	Jo 54	-1.644 <sub>3</sub> ±0.000 <sub>4</sub>
	-1.643 <sub>7</sub> ±0.001	threshold	Mg <sup>24</sup> ( <i>p,p'</i> ), Na <sup>24</sup> γ, Au <sup>198</sup> γ	Jo 54	
Li <sup>7</sup> ( <i>d,α</i> )He <sup>5</sup>	13.719	mag spec	Li <sup>6</sup> ( <i>d,p</i> )Li <sup>7</sup>	Kh 55	
	14.26±0.09	mag spec	Po α	Le 55	
Li <sup>7</sup> ( <i>d,p</i> )Li <sup>8</sup>	-0.183	mag spec	Li <sup>6</sup> ( <i>d,p</i> )Li <sup>7</sup>	Kh 55	-0.192±0.001
Li <sup>7</sup> ( <i>α,n</i> )B <sup>10</sup>	-2.82±0.10	pulse ht	D <sup>2</sup> ( <i>d,n</i> ), B <sup>10</sup> ( <i>α,p</i> )	Ro 56	
Li <sup>7</sup> (Li <sup>7</sup> , <i>p</i> )B <sup>13</sup>	5.97±0.05	range	B <sup>10</sup> ( <i>d,p</i> )B <sup>11*</sup>	No 57	
Be <sup>8</sup> →2He <sup>4</sup>	0.090±0.005	ang corr		Tr 55	
	0.093 <sub>9</sub> ±0.000 <sub>8</sub>	mag spec		Fo 56	0.094 <sub>1</sub> ±0.000 <sub>7</sub>
Be <sup>9</sup> ( <i>γ,n</i> )Be <sup>8</sup>	-1.664±0.004	threshold		Co 56	-1.665±0.001 <sub>4</sub>
Be <sup>9</sup> ( <i>n,γ</i> )Be <sup>10</sup>	6.80	compt spec		Gr 55b	6.816±0.006
Be <sup>9</sup> ( <i>p,pn</i> )Be <sup>8</sup>	-1.664±0.005	mag spec	Po α	Bo 56	
Be <sup>9</sup> ( <i>p,n</i> )B <sup>9</sup>	-1.853±0.003	threshold	Li <sup>7</sup> ( <i>p,n</i> )Be <sup>7</sup>	Ma 55b	-1.852±0.002
Be <sup>9</sup> ( <i>d,p</i> )Be <sup>10</sup>	4.586±0.009	mag spec	Po α	Ju 54	4.587±0.005
Be <sup>9</sup> ( <i>d,n</i> )B <sup>10</sup>	4.43±0.08	ph pl	La 47	Pr 52, Pr 53*	4.35±0.02
	4.28±0.10	ph pl		Re 54	
	4.54±0.06 <sup>c</sup>	ph pl		Gr 55	
Be <sup>9</sup> (Li <sup>7</sup> , <i>p</i> )C <sup>15</sup>	9.05±0.05	range	Li <sup>7</sup> (Li <sup>7</sup> , <i>p</i> )B <sup>12</sup>	No 57	
B <sup>10</sup> ( <i>n,t</i> )Be <sup>8</sup>	(0.35±0.20)	pulse ht	D <sup>2</sup> ( <i>d,n</i> ), B <sup>10</sup> ( <i>n,α</i> )Li <sup>7*</sup>	Ja 55	
B <sup>10</sup> ( <i>n,γ</i> )B <sup>11</sup>	11.43±0.04	pr spec	absolute	Ba 56	
B <sup>10</sup> ( <i>p,n</i> )C <sup>10</sup>	-4.37±0.05	threshold		Co 55	-4.37±0.05
B <sup>10</sup> ( <i>p,γ</i> )C <sup>11</sup>	8.81±0.10	scint spec	F <sup>19</sup> ( <i>p,αγ</i> ), C <sup>13</sup> ( <i>p,γ</i> )	Ch 56	8.81±0.10
B <sup>10</sup> ( <i>d,α</i> )Be <sup>8</sup>	17.829±0.010	mag spec	absolute	El 54	17.829±0.010
B <sup>10</sup> ( <i>d,t</i> )B <sup>9</sup>	-2.187±0.010	mag spec	Po α	Bo 56	
B <sup>10</sup> ( <i>d,p</i> )B <sup>11</sup>	9.227±0.006	mag spec	absolute	El 54	9.229±0.005
B <sup>10</sup> (He <sup>2</sup> , <i>n</i> )N <sup>12</sup>	1.46±0.06	ph pl		Aj 57	preliminary
B <sup>10</sup> ( <i>α,d</i> )C <sup>12</sup>	1.36±0.09	scint spec	Al <sup>27</sup> ( <i>α,p</i> )Si <sup>30</sup>	Pi 56	1.341±0.002
	1.341±0.002	el spec	Li <sup>7</sup> ( <i>p,n</i> )Be <sup>7</sup>	Do 56a	
B <sup>10</sup> ( <i>α,p</i> )C <sup>13</sup>	4.064±0.012	mag spec	Po α	Fa 55	4.064±0.012
	4.08±0.03	scint spec	Al <sup>27</sup> ( <i>α,p</i> )Si <sup>30</sup>	Pi 56	
	4.10±0.03	ph pl	Ro 51c	Pa 56	
B <sup>11</sup> ( <i>p,n</i> )C <sup>11</sup>	-2.83±0.08	ph pl	Ro 51c	Aj 56	-2.762±0.003
	-0.05				
B <sup>11</sup> ( <i>d,α</i> )Be <sup>9</sup>	8.029±0.005	mag spec	absolute	El 54	8.024±0.004
	8.015±0.010	mag spec	Po α	Bo 56	
B <sup>11</sup> ( <i>d,p</i> )B <sup>12</sup>	1.110	spec		Kh 54	
B <sup>11</sup> ( <i>d,n</i> )C <sup>12</sup>	13.81	ph pl		Ih 55	13.8±0.1
B <sup>11</sup> ( <i>α,p</i> )C <sup>14</sup>	0.788±0.017	mag spec	Po α	Fa 55	0.788±0.017
B <sup>11</sup> ( <i>α,n</i> )N <sup>14</sup>	0.0±0.3	pulse ht	D <sup>2</sup> ( <i>d,n</i> )He <sup>3</sup> , B <sup>10</sup> ( <i>α,p</i> )C <sup>13</sup>	Qu 56	0.27±0.06 <sup>+</sup>
C <sup>12</sup> ( <i>p,α</i> )B <sup>9</sup>	-7.58±0.10	range	Be 50d, Bi 54a	Re 55	
C <sup>12</sup> ( <i>d,α</i> )B <sup>10</sup>	-1.39±0.02	range	El 51c	El 57	
C <sup>12</sup> ( <i>d,p</i> )C <sup>13</sup>	2.717±0.010	mag spec	Po α	Sp 54	2.721±0.002 <sup>+</sup>
	2.720±0.003	mag spec	absolute	El 54	

TABLE I.—Continued.

Reaction	Measured Q value (Mev)	Method	Calibration energy standard	Reference	Average Q value (Mev)
C <sup>12</sup> (He <sup>3</sup> ,n)O <sup>14</sup>	-1.147±0.002 <sup>a</sup> -1.158 <sub>b</sub> ±0.003	threshold threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup> , Li <sup>7</sup> (α,γ)B <sup>11</sup>	Bu 56a Br 56a	-1.152±0.005
C <sup>13</sup> (p,d)C <sup>12</sup>	-2.718	mag spec	Po α	Sp 56a	-2.721±0.002 <sup>+</sup>
C <sup>13</sup> (d,p)C <sup>14</sup>	5.942±0.011 5.953±0.010	mag spec mag spec	Po α ThC α	Sp 54 Ah 54d	5.943±0.003
C <sup>13</sup> (d,n)N <sup>14</sup>	5.40±0.10 5.41±0.06 5.325±0.04	ph pl ph pl ph pl	Ri 51	Be 52 Gr 55 Bi 55	5.35±0.03
C <sup>14</sup> (p,n)N <sup>14</sup>	-0.626 <sub>b</sub> ±0.000 <sub>b</sub>	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup>	Sa 56	-0.626 <sub>b</sub> ±0.000 <sub>b</sub> <sup>+</sup>
C <sup>14</sup> (d,α)B <sup>12</sup>	0.362±0.001 <sub>b</sub>	el spec	Li <sup>7</sup> (p,n)Be <sup>7</sup>	Do 56a	
C <sup>14</sup> (d,p)C <sup>15</sup>	-1.007±0.001	el spec	Li <sup>7</sup> (p,n)Be <sup>7</sup>	Do 56a	
C <sup>14</sup> (α,n)O <sup>17</sup>	-1.820±0.002	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup>	Sa 56	
N <sup>14</sup> (n,p)C <sup>14</sup>	0.609±0.005 <sup>c</sup>	pulse ht		Is 50	0.626 <sub>b</sub> ±0.000 <sub>b</sub> <sup>+</sup>
N <sup>14</sup> (d,n)O <sup>15</sup>	5.21±0.07	ph pl		No 56	5.14±0.03
N <sup>14</sup> (α,p)O <sup>17</sup>	-1.16	ph pl		Hj 53, Hj 53a	-1.16±0.04
N <sup>14</sup> (α,n)F <sup>17</sup>	-4.76±0.07	threshold, pulse ht	Be 49	Do 56	-4.76±0.07
N <sup>15</sup> (p,n)O <sup>15</sup>	-3.539±0.008 -3.5432±0.0015	threshold threshold	(p,n) threshs, (p,γ) resonances Li <sup>7</sup> (p,n)Be <sup>7</sup> , E <sub>T</sub> =1.8811	Ki 55 Li 57	-3.5432±0.0015 preliminary
N <sup>15</sup> (d,p)N <sup>16</sup>	0.286	mag spec		Wh 55	
O <sup>16</sup> (d,p)O <sup>17</sup>	1.915±0.010	mag spec	Po α	Sp 54	1.918±0.004
O <sup>16</sup> (d,n)F <sup>17</sup>	-1.622±0.004 -1.626±0.004	threshold threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup> (p,n) thresholds	Bo 51, Ma 55a* Ma 55a	-1.623±0.003
O <sup>17</sup> (d,α)N <sup>15</sup>	9.807±0.012	mag spec	F <sup>19</sup> (d,α)O <sup>17</sup> , N <sup>14</sup> (d,α)C <sup>12</sup>	Pa 54a	
O <sup>17</sup> (d,p)O <sup>18</sup>	5.821±0.010	mag spec	ThC α	Ah 54c, Ah 54d	
O <sup>18</sup> (p,α)N <sup>15</sup>	3.967±0.009	mag spec	F <sup>19</sup> (p,αγ)O <sup>16</sup> , N <sup>15</sup> (p,α)C <sup>12</sup>	Mi 54, Ah 54d*	3.967±0.009
O <sup>18</sup> (p,n)F <sup>18</sup>	-2.447±0.010	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup>	Ma 56a	-2.452±0.004
O <sup>18</sup> (d,α)N <sup>16</sup>	4.237±0.009	mag spec	C <sup>12</sup> (d,p)C <sup>13</sup> , N <sup>14</sup> (d,p)N <sup>15</sup> *	Pa 55a	
O <sup>18</sup> (d,p)O <sup>19</sup>	1.730±0.008 1.732±0.008 1.735±0.008	mag spec mag spec mag spec	B <sup>10</sup> (d,p)B <sup>11</sup> *, F <sup>19</sup> (p,αγ)O <sup>16</sup>	Ah 54b, Mi 54b Th 54 Ho 55	1.732±0.005
F <sup>19</sup> (n,γ)F <sup>20</sup>	6.599±0.011	pr spec	absolute	Ca 56	6.599±0.011
F <sup>19</sup> (p,α)O <sup>16</sup>	8.110±0.010	mag spec	Po α	Sq 56	8.114±0.007
F <sup>19</sup> (p,n)Ne <sup>19</sup>	-4.029±0.008 -4.027±0.008 -4.022±0.005	threshold threshold threshold	F <sup>19</sup> (p,αγ)O <sup>16</sup> (p,n) threshs, (p,γ) resonances Li <sup>7</sup> (p,n)Be <sup>7</sup>	Wi 52, Ki 55* Ki 55 Ma 55b	-4.025±0.004
F <sup>19</sup> (d,p)F <sup>20</sup>	4.38±0.03	range	El 51c	El 56	4.373±0.007
F <sup>19</sup> (d,t)F <sup>18</sup>	-4.17±0.02	range	El 51c	El 57	
F <sup>19</sup> (t,p)F <sup>21</sup>	6.200±0.025 6.03±0.1 <sup>a</sup>	mag spec ph pl	O <sup>16</sup> (t,α)N <sup>15</sup>	Ja 56 Bi 55a	6.200±0.025 preliminary
F <sup>19</sup> (α,p)Ne <sup>22</sup>	1.673±0.011	mag spec	Po α	Fa 55	1.673±0.011
F <sup>19</sup> (α,n)Na <sup>22</sup>	-2.0±0.2	pulse ht	D <sup>2</sup> (d,n)He <sup>3</sup> , B <sup>10</sup> (α,p)C <sup>13</sup>	Qu 56	-2.0±0.2
Ne <sup>20</sup> (n,α)O <sup>17</sup>	-0.70±0.02	pulse ht		Fl 53	-0.71±0.02
Ne <sup>20</sup> (d,α)F <sup>18</sup>	2.810±0.009	mag spec	F <sup>19</sup> (p,αγ)O <sup>16</sup> , O <sup>16</sup> (d,p)O <sup>17</sup>	Mi 54a, Mi 56*	2.810±0.009
Ne <sup>22</sup> (d,p)Ne <sup>23</sup>	2.968±0.008	mag spec	C <sup>12</sup> (d,p)C <sup>13</sup>	Ah 54a	2.966±0.005
Na <sup>23</sup> (n,γ,γ)Na <sup>24</sup>	6.96±0.03	compt spec		Gr 55b	
Na <sup>23</sup> (p,α)Ne <sup>20</sup>	2.370±0.008	mag spec	Po α	Bu 56c	2.377±0.003
Na <sup>23</sup> (p,n)Mg <sup>23</sup>	-4.849±0.010 -4.841±0.010	threshold threshold	F <sup>19</sup> (p,αγ)O <sup>16</sup> (p,n) threshs, (p,γ) resonances	Wi 52a, Ki 55* Ki 55	-4.845±0.007
Na <sup>23</sup> (α,n)Al <sup>26</sup>	-2.9±0.2 ≥ -2.970±0.004	pulse ht threshold	Be 49 Li <sup>7</sup> (p,n)Be <sup>7</sup>	Do 56 Bu 56b	preliminary
Mg <sup>24</sup> (p,γ)Al <sup>25</sup>	2.26±0.03 <sup>a</sup> 2.29±0.02 <sup>a</sup> 2.29±0.02 <sup>a</sup>	scint spec scint spec scint spec	F <sup>19</sup> (p,αγ)O <sup>16</sup> , (p,γ) reactions Pr <sup>144</sup> γ, Cs <sup>137</sup> γ, Na <sup>22</sup> γ Zn <sup>65</sup> γ, Co <sup>60</sup> γ, ThC''γ	Gr 55a Cr 56 Ag 56	2.28±0.01 <sub>b</sub>
Mg <sup>24</sup> (d,α)Na <sup>22</sup>	1.953±0.012	mag spec	Po α	Br 55, Br 55a*	preliminary
Mg <sup>24</sup> (d,p)Mg <sup>25</sup>	5.02±0.02 <sup>c</sup>	spec		Kh 53	5.097±0.007

TABLE I.—Continued.

Reaction	Measured Q value (Mev)	Method	Calibration energy standard	Reference	Average Q value (Mev)
Mg <sup>25</sup> (γ, p)Na <sup>24</sup>	-12.1	threshold		Ka 54	
Mg <sup>25</sup> (γ, n)Mg <sup>24</sup>	-7.2	threshold		Ka 54	
Mg <sup>25</sup> (n, γ)Mg <sup>26</sup>	11.086±0.025	pr spec	absolute	Ca 56	
Mg <sup>25</sup> (p, α)Na <sup>22</sup>	-3.15	mag spec	Po α	Br 55a	preliminary
Mg <sup>25</sup> (p, n)Al <sup>25</sup>	-5.084±0.024	threshold	(p, n) threshs, (p, γ) resonances	Ki 55	
Mg <sup>25</sup> (p, γ)Al <sup>26</sup>	6.35±0.08	scint spec	F <sup>19</sup> (p, αγ), Be <sup>9</sup> (α, nγ), C <sup>13</sup> (p, γ)	Kl 54	
Mg <sup>25</sup> (α, p)Al <sup>28</sup>	-1.29±0.04	pulse ht	Al <sup>27</sup> (p, α)Mg <sup>24</sup>	Gr 57	
Mg <sup>26</sup> (γ, p)Na <sup>25</sup>	-14.3	threshold		Ka 54	
Mg <sup>26</sup> (γ, n)Mg <sup>24</sup>	-11.1	threshold		Ka 54	
Mg <sup>26</sup> (p, n)Al <sup>26</sup>	-4.83±0.1 <sup>a</sup> -4.778±0.015 <sup>a</sup>	threshold	(p, n) threshs, (p, γ) resonances	Sc 54, En 54, Ka 55	-4.778±0.015
Mg <sup>26</sup> (α, p)Al <sup>29</sup>	-2.90±0.04	pulse ht	Al <sup>27</sup> (p, α)Mg <sup>24</sup>	Gr 57	
Al <sup>27</sup> (γ, n)Al <sup>26</sup>	-13.4±0.2	threshold	Cu <sup>63</sup> (γ, n)Cu <sup>62</sup>	Ha 54	-13.1±0.3
Al <sup>27</sup> (p, α)Mg <sup>24</sup>	1.61±0.04 1.596±0.006	ph pl mag spec	Li <sup>7</sup> (p, n)Be <sup>7</sup> , Po α	Gr 54 Va 57	1.595±0.002 <sup>+</sup>
Al <sup>27</sup> (p, n)Si <sup>27</sup>	-5.581±0.010 -5.584±0.010 -5.607±0.008	threshold	F <sup>19</sup> (p, αγ)O <sup>16</sup> (p, n) threshs, (p, γ) resonances	Ki 53d, Ki 55* Ki 55 Ma 55b	-5.593±0.009
Al <sup>27</sup> (d, p)Al <sup>28</sup>	5.475 5.502±0.010	spec mag spec	Po α	Kh 54 Bu 56	5.498±0.007
Al <sup>27</sup> (α, p)Si <sup>30</sup>	2.38±0.03	scint spec pulse ht ph pl		Ha 56	2.38±0.03
Al <sup>27</sup> (α, n)P <sup>30</sup>	≥ -2.662±0.004	threshold	Li <sup>7</sup> (p, n)Be <sup>7</sup>	Bu 56b	preliminary
Si <sup>28</sup> (n, γ)Si <sup>29</sup>	8.482±0.015	compt spec		Ad 56a	8.471±0.007
Si <sup>28</sup> (d, α)Al <sup>26</sup>	1.416±0.008	mag spec	Po α	Br 54c	
Si <sup>29</sup> (γ, p)Al <sup>28</sup>	-12.3	threshold		Ka 54	
Si <sup>29</sup> (γ, n)Si <sup>28</sup>	-8.5	threshold		Ka 54	
Si <sup>29</sup> (n, γ)Si <sup>30</sup>	10.59±0.03	compt spec		Ad 56a	10.600±0.010
Si <sup>29</sup> (p, γ)P <sup>30</sup>	5.55±0.06	scint spec	Be <sup>9</sup> (α, nγ)C <sup>12</sup> , F <sup>19</sup> (p, αγ)O <sup>16</sup>	En 54a	
Si <sup>30</sup> (γ, p)Al <sup>29</sup>	-12.9	threshold		Ka 54	
Si <sup>30</sup> (γ, n)Si <sup>29</sup>	-10.6	threshold		Ka 54	
P <sup>31</sup> (γ, n)P <sup>30</sup>	-12.33±0.05	threshold	(γ, n) thresholds	Ba 55a	-12.32±0.05
P <sup>31</sup> (p, α)Si <sup>28</sup>	1.911±0.005 1.909±0.010	mag spec mag spec	Li <sup>7</sup> (p, n), Al <sup>27</sup> (p, α), Po α	Va 56 En 57	1.910±0.004
P <sup>31</sup> (p, n)S <sup>31</sup>	-6.06±0.2	ph pl	Gi 54	Ru 56	
P <sup>31</sup> (d, n)S <sup>32</sup>	6.63±0.08	ph pl	El 51d	El 52, El 55*	6.63±0.08
P <sup>31</sup> (α, p)S <sup>34</sup>	0.7±0.1	scint spec	Al <sup>27</sup> (α, p)Si <sup>30</sup>	St 56	0.7±0.1
P <sup>31</sup> (α, n)Cl <sup>34</sup>	-5.7±0.2	threshold	D <sup>2</sup> (d, n)He <sup>3</sup> , B <sup>10</sup> (α, p)C <sup>13</sup>	Qu 56	-5.7±0.2
S <sup>32</sup> (n, γ)S <sup>33</sup>	8.63±0.04	compt spec		Gr 55b	8.64±0.02
S <sup>32</sup> (p, γ)Cl <sup>33</sup>	2.285±0.012	scint spec	Na <sup>24</sup> γ	Va 56a	
S <sup>32</sup> (d, α)P <sup>30</sup>	4.831±0.013	mag spec		Le 56	
S <sup>32</sup> (d, p)S <sup>33</sup>	6.408±0.020	mag spec		Le 56	6.419±0.010
S <sup>32</sup> (α, p)Cl <sup>35</sup>	-2.3	scint spec	Al <sup>27</sup> (α, p)Si <sup>30</sup>	Pi 55	-1.861±0.004 <sup>+</sup>
S <sup>34</sup> (p, n)Cl <sup>34</sup>	(-6.1)	ph pl		Aj 55	preliminary
S <sup>34</sup> (d, α)P <sup>32</sup>	5.04±0.02	mag spec		Le 56	
Cl <sup>35</sup> (γ, n)Cl <sup>34</sup>	-12.35±0.035	threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> (γ, n) threshs	De 55	
Cl <sup>35</sup> (n, γ)Cl <sup>36</sup>	8.55±0.04	compt spec		Gr 55b	8.56±0.02
Cl <sup>35</sup> (p, α)S <sup>32</sup>	1.865±0.015 1.863±0.008 1.860±0.005	mag spec mag spec mag spec	Po α Li <sup>7</sup> (p, n)Be <sup>7</sup> , Po α	Al 55 En 56 Va 56, Va 57*	1.861±0.004 <sup>+</sup>
Cl <sup>35</sup> (d, α)S <sup>33</sup>	8.277±0.010	mag spec	Po α	Pa 55	8.277±0.010
Cl <sup>35</sup> (d, p)Cl <sup>36</sup>	6.354±0.008	mag spec	Po α	Pa 55	6.354±0.008
Cl <sup>37</sup> (p, α)S <sup>34</sup>	3.015±0.015 3.026±0.008 3.028±0.006	mag spec mag spec mag spec	Po α Li <sup>7</sup> (p, n)Be <sup>7</sup> , Po α	Al 55 En 56 Va 56, Va 57*	3.026±0.005

TABLE I.—Continued.

Reaction	Measured Q value (Mev)	Method	Calibration energy standard	Reference	Average Q value (Mev)
Cl <sup>37</sup> (d, $\alpha$ )S <sup>35</sup>	7.783±0.012	mag spec	Po $\alpha$	Pa 55	
Cl <sup>37</sup> (d, $p$ )Cl <sup>38</sup>	3.877±0.008	mag spec	Po $\alpha$	Pa 55	3.877±0.008
A <sup>36</sup> ( $\alpha$ , $p$ )K <sup>39</sup>	-1.28±0.03	ph pl	N <sup>14</sup> ( $\alpha$ , $p$ )O <sup>17</sup>	Sc 56a	-1.283±0.008+
A <sup>40</sup> ( $\gamma$ , $n$ )A <sup>39</sup>	-9.85±0.15	threshold	C <sup>12</sup> , Mn <sup>55</sup> , Bi <sup>209</sup> ( $\gamma$ , $n$ ) threshs	Ha 54a	
A <sup>40</sup> (n, $\alpha$ )S <sup>37</sup>	-2.5±0.1	pulse ht	Po $\alpha$	Be 55	
A <sup>40</sup> ( $\alpha$ , $p$ )K <sup>43</sup>	-3.36±0.03	ph pl	N <sup>13</sup> ( $\alpha$ , $p$ )O <sup>17</sup>	Sc 56a	preliminary
K <sup>39</sup> ( $\gamma$ , $n$ )K <sup>38</sup>	-13.00	threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> ( $\gamma$ , $n$ ) threshs	De 55	
K <sup>39</sup> (n, $\gamma$ , $\gamma$ )K <sup>40</sup>	7.795±0.010	compt spec		Ad 56, Bu 53	7.791±0.006
K <sup>39</sup> (n, $\alpha$ )Cl <sup>36</sup>	1.25±0.20	scint spec	Po $\alpha$ , K <sup>39</sup> (n, $p$ )A <sup>39</sup>	Sc 56	
K <sup>39</sup> ( $p$ , $\alpha$ )A <sup>36</sup>	1.267±0.020	mag spec		Al 55	1.283±0.008+
	1.286±0.008	mag spec	Po $\alpha$	Sp 55, Sp 56a*	
K <sup>39</sup> ( $\alpha$ , $p$ )Ca <sup>42</sup>	-0.19±0.07	range		Sc 55	-0.118±0.007+
K <sup>41</sup> ( $p$ , $\alpha$ )A <sup>38</sup>	4.002±0.015	mag spec	mag spec	Al 55	
K <sup>41</sup> ( $\alpha$ , $p$ )Ca <sup>44</sup>	0.98±0.10	range		Sc 55	1.057±0.010+
Ca <sup>40</sup> (n, $\gamma$ , $\gamma$ )Ca <sup>41</sup>	8.37±0.03	pr spec	absolute	Ki 52, Br 56b	8.355±0.014
	8.350±0.017	compt spec		Ad 56	
Ca <sup>40</sup> (d, $\alpha$ )K <sup>38</sup>	4.650±0.010	mag spec	Po $\alpha$	Br 56	
Ca <sup>40</sup> (d, $p$ )Ca <sup>41</sup>	6.140±0.009	mag spec	Po $\alpha$	Br 54, Br 56b*	6.140±0.009
Ca <sup>40</sup> (d, $n$ )Sc <sup>41</sup>	-0.60±0.05	range		Pl 55	preliminary
Ca <sup>42</sup> ( $p$ , $\alpha$ )K <sup>39</sup>	0.118±0.007	mag spec	Po $\alpha$	Br 56	0.118±0.007+
Ca <sup>42</sup> (d, $p$ )Ca <sup>43</sup>	5.711±0.010	mag spec	Po $\alpha$	Br 54b, Br 56*	
Ca <sup>43</sup> ( $p$ , $\alpha$ )K <sup>40</sup>	-0.014±0.008	mag spec	Po $\alpha$	Br 56	
Ca <sup>43</sup> (d, $p$ )Ca <sup>44</sup>	9.07±0.07	range		Sc 55	8.913±0.014
	8.913±0.014	mag spec	Po $\alpha$	Br 56	
Ca <sup>44</sup> ( $p$ , $\alpha$ )K <sup>41</sup>	-1.057±0.010	mag spec	Po $\alpha$	Br 56	-1.057±0.010+
Ca <sup>44</sup> (d, $p$ )Ca <sup>45</sup>	5.188±0.010	mag spec	Po $\alpha$	Br 54b, Br 56*	
Ca <sup>48</sup> (d, $p$ )Ca <sup>49</sup>	2.916±0.006	mag spec	Po $\alpha$	Br 56	2.916±0.006
Sc <sup>45</sup> (p, $n$ )Ti <sup>45</sup>	-2.844±0.004	threshold	Li <sup>7</sup> (p, $n$ )Be <sup>7</sup> , E <sub>T</sub> =1.8811	Br 55b	
Ti <sup>47</sup> (n, $\gamma$ , $\gamma$ )Ti <sup>48</sup>	(11.609±0.02)			Ba 56a	
	[10.619 $\gamma$ transition (Ba 56a) to 0.990±0.015 state (He 55)]			Ki 53, Wa 55	
	(11.51±0.05)			Pi 52a, Wa 55	9.13±0.05
Ti <sup>47</sup> (d, $p$ , $\gamma$ )Ti <sup>48</sup>	9.13±0.05			Ba 56a	8.145±0.006
Ti <sup>48</sup> (n, $\gamma$ , $\gamma$ )Ti <sup>49</sup>	8.141±0.008				
	[6.756±0.006 $\gamma$ transition (Ki 53) to 1.385±0.005 state (Mo 54)]			Ki 53, Wa 55	
	8.14±0.02			Ad 56a	
	8.153±0.010			Ki 53, Wa 55	
Ti <sup>49</sup> (n, $\gamma$ , $\gamma$ )Ti <sup>50</sup>	10.97±0.07			Bu 55	
V <sup>51</sup> (p, $\alpha$ )Ti <sup>48</sup>	1.161±0.010	mag spec	Po $\alpha$	Gi 55	-1.535±0.001
V <sup>51</sup> (p, $n$ )Cr <sup>51</sup>	-1.535±0.001	threshold	Li <sup>7</sup> (p, $n$ )Be <sup>7</sup>	Ma 56	
	-1.536	threshold	Li <sup>7</sup> (p, $n$ )Be <sup>7</sup> , E <sub>T</sub> =1.8811		
Cr <sup>50</sup> (n, $\gamma$ , $\gamma$ )Cr <sup>51</sup>	9.25±0.01			Ki 53, Wa 55	
	[8.499±0.007 $\gamma$ transition (Ki 53) to 0.750±0.011 state (He 55)]			El 56a	preliminary
Cr <sup>52</sup> (d, $p$ )Cr <sup>53</sup>	5.74	range Al		El 56a	preliminary
Cr <sup>53</sup> (d, $p$ )Cr <sup>54</sup>	7.55	range Al		El 56a	preliminary
Cr <sup>54</sup> (p, $n$ )Mn <sup>54</sup>	-2.162±0.005	threshold	Li <sup>7</sup> (p, $n$ )Be <sup>7</sup>	Lo 52	
Mn <sup>55</sup> (p, $n$ )Fe <sup>55</sup>	-1.015±0.003	threshold		Jo 56	-1.014±0.003
Fe <sup>64</sup> ( $\gamma$ , $n$ )Fe <sup>63</sup>	(-13.7±0.2)	threshold	C <sup>12</sup> , O <sup>16</sup> , Cu <sup>63</sup> , Ag <sup>109</sup> ( $\gamma$ , $n$ ) thresholds	Ka 51	
	(-13.65±0.05)	threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> ( $\gamma$ , $n$ ) thresholds	Ba 55	
	-11.90±0.07	threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> ( $\gamma$ , $n$ ) thresholds	De 55	
Fe <sup>64</sup> (n, $\gamma$ )Fe <sup>65</sup>	9.295±0.015	compt spec		Ad 56a	9.297±0.006
Fe <sup>64</sup> (d, $p$ )Fe <sup>65</sup>	7.073	mag spec	Po $\alpha$	Sp 56	
Fe <sup>66</sup> (n, $\gamma$ )Fe <sup>67</sup>	7.636±0.010	compt spec		Ad 56a	7.638±0.004
Fe <sup>66</sup> (d, $p$ )Fe <sup>67</sup>	5.418	mag spec	Po $\alpha$	Sp 56	
	5.53	range		Mc 55	preliminary
Fe <sup>67</sup> (d, $p$ )Fe <sup>68</sup>	7.808	mag spec	Po $\alpha$	Sp 56	
	7.89	range		Mc 55	preliminary

TABLE I.—Continued.

Reaction	Measured Q value (Mev)	Method	Calibration energy standard	Reference	Average Q value (Mev)
Fe <sup>58</sup> (d,p)Fe <sup>59</sup>	4.350	mag spec	Po $\alpha$	Sp 56	
Co <sup>59</sup> (p,n)Ni <sup>59</sup>	-1.84	range ph pl		St 52	
	-1.858±0.004	threshold		Bu 56d	-1.858±0.002
	-1.862±0.005 <sup>b</sup>	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup> , $E_T=1.8811$	Ch 57	preliminary
Co <sup>59</sup> (d,p)Co <sup>60</sup>	5.283±0.008	mag spec	Po $\alpha$	Fo 54	5.283±0.008
Ni <sup>58</sup> ( $\gamma$ ,n)Ni <sup>57</sup>	-12.0	threshold		Ka 51	
Ni <sup>58</sup> (n, $\gamma$ )Ni <sup>59</sup>	8.996±0.010	compt spec		Ad 56	8.997±0.004
Ni <sup>58</sup> (p, $\gamma$ )Cu <sup>59</sup>	3.42±0.02	scint spec		Go 57a	
Ni <sup>58</sup> (d,p)Ni <sup>59</sup>	6.70±0.1	range ph pl		Pr 54	6.74±0.07
Ni <sup>60</sup> (n, $\gamma$ )Ni <sup>61</sup>	(7.817±0.008)	pr spec	absolute	Ki 53, Qu 56a	tgt isotope uncertain
	(7.825±0.020)	compt spec		Ad 56, Qu 56a	tgt isotope uncertain
Ni <sup>60</sup> (p, $\gamma$ )Cu <sup>61</sup>	4.81±0.03	scint spec		Go 57a	preliminary
Ni <sup>60</sup> (p,n)Cu <sup>60</sup>	-6.6±0.4	threshold		Co 54a	-6.6±0.4
Ni <sup>60</sup> (d,p)Ni <sup>61</sup>	(5.55±0.1)	range ph pl		Pr 54	
Ni <sup>61</sup> (p, $\gamma$ )Cu <sup>62</sup>	6.03±0.06	scint spec		Go 57a	preliminary
Ni <sup>62</sup> (p, $\gamma$ )Cu <sup>63</sup>	6.13±0.03	scint spec		Go 57a	
Ni <sup>64</sup> (p, $\gamma$ )Cu <sup>65</sup>	7.42±0.03	scint spec		Go 57a	
Cu <sup>63</sup> ( $\gamma$ ,n)Cu <sup>62</sup>	-10.78±0.05	threshold	D <sup>2</sup> , F <sup>19</sup> , O <sup>16</sup> ( $\gamma$ ,n) thresholds	Be 56	-10.65±0.06
	-10.73±0.05	threshold	N <sup>14</sup> , F <sup>19</sup> ( $\gamma$ ,n) thresholds	Ro 55	
	-10.54±0.04	threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> ( $\gamma$ ,n) thresholds	De 55	
Cu <sup>63</sup> ( $\gamma$ ,2n)Cu <sup>61</sup>	-20.0±0.5	threshold		Be 54	
Cu <sup>63</sup> (p, $\gamma$ )Zn <sup>64</sup>	7.69±0.04	scint spec		Go 57	preliminary
Cu <sup>63</sup> (p,n)Zn <sup>63</sup>	-4.147±0.008	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup>	Ki 55	-4.149±0.004
	-4.149±0.004	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup> , $E_T=1.8811$	Br 55b	
Cu <sup>65</sup> ( $\gamma$ ,n)Cu <sup>64</sup>	-9.94±0.08	threshold	D <sup>2</sup> , O <sup>16</sup> , F <sup>19</sup> ( $\gamma$ ,n) thresholds	Be 56	-9.95±0.08
Cu <sup>65</sup> (n, $\gamma$ )Cu <sup>66</sup>	(7.01±0.02)	pr spec	absolute	Ba 53, Qu 56a	(7.01±0.02) tgt isotope uncertain
Cu <sup>66</sup> (p, $\gamma$ )Zn <sup>66</sup>	8.85±0.04	scint spec		Go 57	
Cu <sup>66</sup> (p,n)Zn <sup>65</sup>	-2.137±0.005	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup>	Ki 55	-2.1327±0.0013
	-2.136±0.004	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup> , $E_T=1.8811$	Br 55b	
	-2.131±0.005	resonance	Li <sup>7</sup> (p,n)Be <sup>7</sup>	Ma 56	
	-2.132±0.001 <sub>b</sub>	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup> , $E_T=1.8811$	Ma 56b	
Zn <sup>64</sup> ( $\gamma$ ,2n)Zn <sup>62</sup>	-20.35±0.35	threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> ( $\gamma$ ,n) thresholds	De 55	
Zn <sup>64</sup> ( $\gamma$ ,n)Zn <sup>63</sup>	-11.6	threshold		Ka 51	-11.60±0.06
	-11.58±0.06	threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> ( $\gamma$ ,n) thresholds	De 55	
Zn <sup>64</sup> (n, $\gamma$ , $\gamma$ )Zn <sup>65</sup>	(7.928±0.007)			Ki 53, Wa 55	tgt isotope uncertain
	(7.990±0.008)			Ki 53, Qu 56a	
Zn <sup>66</sup> (n, $\gamma$ , $\gamma$ )Zn <sup>67</sup>	(7.03±0.02)			Ki 53, Qu 56a	tgt isotope uncertain
Zn <sup>67</sup> (n, $\gamma$ , $\gamma$ )Zn <sup>68</sup>	(10.22±0.01)			Wa 55, Qu 56a	tgt isotope uncertain
Zn <sup>67</sup> (p,n)Ga <sup>67</sup>	-1.777±0.005 <sup>b</sup>	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup> , $E_T=1.8811$	Ch 57	-1.781±0.003
Zn <sup>68</sup> (n, $\gamma$ )Zn <sup>69</sup>	(6.49±0.02)			Ki 53, Qu 56a	tgt isotope uncertain
Zn <sup>68</sup> (p,n)Ga <sup>68</sup>	-3.694±0.006	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup> , $E_T=1.8811$	Br 55b	-3.700±0.005
	-3.704±0.005 <sup>b</sup>	threshold	Li <sup>7</sup> (p,n)Be <sup>7</sup> , $E_T=1.8811$	Ch 57	
Ga <sup>69</sup> (n, $\gamma$ )Ga <sup>70</sup>	7.733±0.020	pr spec	absolute	Ba 56a	
Ge <sup>70</sup> ( $\gamma$ ,n)Ge <sup>69</sup>	-12.1±0.2	threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> ( $\gamma$ ,n) thresholds	De 55	
Ge <sup>76</sup> ( $\gamma$ ,n)Ge <sup>75</sup>	-9.3	threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> ( $\gamma$ ,n) thresholds	De 55	
Kr <sup>78</sup> (d,p)Kr <sup>79</sup>	5.98±0.05	range ph pl	N <sup>14</sup> (d,p)N <sup>15</sup>	Bi 56	
Kr <sup>80</sup> (d,p)Kr <sup>81</sup>	5.63±0.10	range ph pl	N <sup>14</sup> (d,p)N <sup>15</sup>	Bi 56	
Rb <sup>85</sup> ( $\gamma$ ,n)Rb <sup>84</sup>	-9.26±0.15	threshold	O <sup>16</sup> , F <sup>19</sup> ( $\gamma$ ,n) thresholds	To 56	
Rb <sup>87</sup> ( $\gamma$ ,n)Rb <sup>86</sup>	-10.14±0.15	threshold	O <sup>16</sup> , F <sup>19</sup> ( $\gamma$ ,n) thresholds	To 56	
Sr <sup>86</sup> ( $\gamma$ ,n)Sr <sup>85</sup>	-11.5	threshold		Ye 55	
Sr <sup>86</sup> (n, $\gamma$ , $\gamma$ )Sr <sup>87</sup>	8.417±0.018			Wa 55	8.417±0.018+
Sr <sup>87</sup> (n, $\gamma$ , $\gamma$ )Sr <sup>88</sup>	11.14±0.05			Wa 55	11.14±0.05+
Y <sup>89</sup> (d,p)Y <sup>90</sup>	4.41±0.05	scint spec		Wa 54	
Zr <sup>90</sup> ( $\gamma$ ,n)Zr <sup>89</sup>	-11.78±0.09		Cu <sup>63</sup> ( $\gamma$ ,n)Cu <sup>62</sup> , $E_T=10.73$	Ax 56	-11.78±0.09

The Q values for this reaction in reference (1) are now attributed to excitation of the 0.588 level in Zr<sup>89</sup> (Ax 56).

TABLE I.—Continued.

Reaction	Measured Q value (Mev)	Method	Calibration energy standard	Reference	Average Q value (Mev)
Nb <sup>93</sup> (p,n)Mo <sup>93</sup>	-1.27±0.04	ph pl		Pa 54	-1.27±0.04
	Value quoted in reference 1 is not ground-state transition.				
Mo <sup>92</sup> (γ,n)Mo <sup>91</sup>	(-13.5±0.4) (-13.28±0.15) -13.1±0.1	threshold threshold threshold		Ba 45 Ha 49a Ka 53, Ka 53a	-13.1±0.1
Mo <sup>92</sup> (n,2n)Mo <sup>91</sup>	-12.34	threshold		Br 53	
Rh <sup>103</sup> (p,n)Pd <sup>103</sup>	(-1.53)	ph pl		Pa 54	
Ag <sup>107</sup> (γ,n)Ag <sup>106</sup>	-9.45±0.05 -9.57±0.06	threshold threshold	D <sup>2</sup> , F <sup>19</sup> , O <sup>16</sup> (γ,n) thresholds C <sup>12</sup> , O <sup>16</sup> , Cu <sup>63</sup> , Ag <sup>109</sup> (γ,n)-thresholds	Be 56	-9.50±0.06
Ag <sup>109</sup> (γ,n)Ag <sup>108</sup>	-9.17±0.06 -8.78±0.04	threshold threshold	D <sup>2</sup> , F <sup>19</sup> , O <sup>16</sup> (γ,n) thresholds	Be 56 De 55	-8.90±0.18
Cd <sup>111</sup> (p,n)In <sup>111</sup>	-2.37±0.20	threshold		Mc 51c	
Cd <sup>113</sup> (n,γ)Cd <sup>114</sup>	9.04±0.03	compt spec		Ad 55	9.046±0.008
Ba <sup>138</sup> (d,p)Ba <sup>139</sup>	2.493±0.010			Pa 55	2.493±0.010
Ba <sup>138</sup> (n,γ)Ba <sup>139</sup>	(4.70±0.03)	pr spec	absolute	Ki 53c, Pa 55	tgt isotope uncertain
Pr <sup>141</sup> (d,p)Pr <sup>142</sup>	3.42±0.30	scint spec		Wa 54	
Sm <sup>144</sup> (γ,n)Sm <sup>143</sup>	-9.60±0.05 -9.6	threshold threshold	C <sup>12</sup> , N <sup>14</sup> , O <sup>16</sup> (γ,n) thresholds	De 55 Si 56	
Sm <sup>149</sup> (n,γ,γ)Sm <sup>150</sup>	8.00±0.03			Ad 55	8.00±0.03
Gd <sup>156</sup> (n,γ)Gd <sup>156</sup>	(7.78±0.05)	pr spec	absolute	Ki 53c	tgt isotope uncertain
Gd <sup>157</sup> (n,γ)Gd <sup>158</sup>	(7.36±0.05)	pr spec	absolute	Kr 53c	tgt isotope uncertain
Hf <sup>177</sup> (γ,n)Hf <sup>176</sup>	-6.70±0.09	threshold	O <sup>16</sup> , F <sup>19</sup> (γ,n) thresholds	To 56	
Hf <sup>179</sup> (γ,n)Hf <sup>178</sup>	-6.52±0.12	threshold	O <sup>16</sup> , F <sup>19</sup> (γ,n) thresholds	To 56	
Pt <sup>194</sup> (n,γ)Pt <sup>195</sup>	(6.07±0.04)	pr spec	absolute	Ki 53c	tgt isotope uncertain
Hg <sup>199</sup> (n,γ)Hg <sup>200</sup>	8.03±0.03			Ad 55	8.03±0.03
Bi <sup>209</sup> (p,2n)Po <sup>208</sup>	-9.65±0.08	threshold	range Al, Sm 47	An 56	

\* This Q value has been calculated specifically for this compilation from the experimental data, using accurate masses.

† This Q value has been corrected for the Li<sup>7</sup>(p,n)Be<sup>7</sup> threshold energy of 1.8811 Mev.

‡ This Q value has been omitted from the weighted average.

\* This average contains a later correction to the value originally reported.

+ This average value was calculated by including the measured Q value for the inverse reaction.

#### List of abbreviations used for experimental methods.

ang corr	angular correlation	ph pl	range in photographic plates
compt spec	Compton electron spectrometer	pr spec	pair spectrometer
el spec	electrostatic spectrometer	pulse ht	pulse height
E <sub>T</sub>	threshold energy	range Al	range in aluminum
mag spec	magnetic spectrometer	scint spec	scintillation spectrometer

given. When a correction to the original value has been reported by another author, a second reference with an asterisk has been added. For some measurements, such as a (n,γ,γ) reaction, two references are given. The first reference contains the measurements of the neutron capture gamma rays, while the second contains the assignment of the gamma rays to transitions for specific isotopes, and, also, other determinations of level energies if necessary for the calculation of the energy of the ground-state transition (e.g., see references 6 and 10).

In Column 6 are listed the weighted average values, calculated in the same manner as before. Measured values listed in the original compilation are included in the calculations of the weighted average values given in this supplement. Following the bibliography, a few additional corrections to the original compilation are listed, if not already included in this supplement. No attempt has been made to list revised references for those given previously if no alteration in the Q values reported were made in the later reference.

However, all references listed in Table I of this supplement are included in the bibliography for the sake of completeness.

#### BIBLIOGRAPHY

- Ad 55 Adyasevich, Groshev, and Demidov, Conf. Acad. Sci. U.S.S.R. *Peaceful Uses of Atomic Energy*, Moscow, 1955, Session Div. Phys. Math. Sci. English translation U.S.A.E.C. (1956), p. 195.
- Ad 56 Adyasevich, Groshev, Demidov, and Lutsenko, Soviet J. Atomic Energy 1, 171 (1956); J. Nuclear Energy II, 3, 325 (1956).
- Ad 56a Adyasevich, Groshev, and Demidov, Soviet J. of Atomic Energy 1, 183 (1956); J. Nuclear Energy II, 3, 258 (1956).
- Ag 56 Ager-Hanssen, Lonsjo, and Nordhagen, Phys. Rev. 101, 1779 (1956).
- Ag 54a K. Ahnlund, Arkiv Fysik 7, 459 (1954).
- Ag 54b K. Ahnlund and C. Mileikowsky, Arkiv Fysik 8, 161 (1954).
- Ah 54c K. Ahnlund, Phys. Rev. 96, 999 (1954).
- Ah 54d Ahnlund, Thulin, and Pauli, Arkiv Fysik 8, 489 (1954).

- Aj 55 Ajzenberg, Rubin, and Likely, Phys. Rev. **99**, 654(A) (1955) and verbal report.
- Aj 56 Ajzenberg-Selove, Johnson, Rubin, and Mazari, Phys. Rev. **103**, 356 (1956).
- Aj 57 Ajzenberg-Selove, Bullock, and Almqvist, Bull. Am. Phys. Soc. Ser. II, **2**, 28 (1956).
- Al 54 Allen, Almqvist, Dewan, and Pepper, Phys. Rev. **96**, 684 (1954).
- Al 55 Almqvist, Clarke, and Paul, Phys. Rev. **100**, 1265(A) (1955).
- An 56 Andre, Huizenga, Mech, Ramler, Rauh, and Rocklin, Phys. Rev. **101**, 645 (1956).
- Ax 56 P. Axel and J. D. Fox, Phys. Rev. **102**, 400 (1956).
- Ba 45 G. C. Baldwin and H. W. Koch, Phys. Rev. **67**, 1 (1945).
- Ba 53 G. A. Bartholomew and B. B. Kinsey, Phys. Rev. **89**, 386 (1953).
- Ba 53c S. J. Barne, Jr., Phys. Rev. **92**, 1096 (1953).
- Ba 55 R. Basile and C. Schuhl, J. phys. radium **16**, 372 (1955).
- Ba 55a R. Basile and C. Schuhl, Compt. rend. **240**, 2399 (1955).
- Ba 56 G. A. Bartholomew and P. J. Campion, Bull. Am. Phys. Soc. Ser. II, **1**, 324 (1956).
- Ba 56a G. A. Bartholomew (private communication).
- Be 49 H. A. Bethe, AECU-346 (1949).
- Be 50d H. Bethe, Revs. Modern Phys. **22**, 213 (1950).
- Be 52 R. E. Beneson, Phys. Rev. **90**, 420 (1952).
- Be 54 A. I. Berman and K. L. Brown, Phys. Rev. **96**, 83 (1954).
- Be 55 E. H. Bellamy and F. C. Flack, Phil. Mag. **46**, 341 (1955).
- Be 56 Bendel, McElhinney, and Tobin, Bull. Am. Phys. Soc. Ser. II, **1**, 192 (1956).
- Bi 54a H. Bichsel and R. F. Mosley, Phys. Rev. **94**, 764(A) (1954).
- Bi 55 J. R. Bird, Australian J. Phys. **8**, 314 (1955).
- Bi 55a Bigham, Allen, and Almqvist, Phys. Rev. **99**, 631(A) (1955).
- Bl 56 N. C. Blais and W. W. Watson, Phys. Rev. **104**, 202 (1956).
- Bo 51 T. W. Bonner and J. W. Butler, Phys. Rev. **83**, 1091 (1951).
- Bo 56 Bockelman, Leveque, and Buechner, Phys. Rev. **104**, 456 (1956).
- Br 53 J. E. Brolley, Phys. Rev. **89**, 877 (1953).
- Br 54 C. M. Braams, Phys. Rev. **94**, 763(A) (1954).
- Br 54b C. M. Braams, Phys. Rev. **95**, 650 (1954).
- Br 54c C. P. Browne, Phys. Rev. **95**, 860(L) (1954).
- Br 55 C. P. Browne and W. C. Cobb, Phys. Rev. **99**, 644(A) (1955).
- Br 55a C. P. Browne, Mass. Inst. Technol. Annual Progr. Rept. p. 117 (1955).
- Br 55b Brugger, Bonner, and Marion, Phys. Rev. **100**, 84 (1955).
- Br 56 C. M. Braams, Ph.D. thesis, University of Utrecht (1956).
- Br 56a Bromley, Almqvist, Gove, Litherland, Paul, and Ferguson, PD-264, Atomic Energy of Canada, Ltd. (October, 1956).
- Br 56b C. M. Braams, Phys. Rev. **103**, 1310 (1956).
- Bu 53 Buechner, Sperduto, Browne, and Bockelman, Phys. Rev. **91**, 1502 (1953).
- Bu 55 Buechner, Braams, and Sperduto, Phys. Rev. **100**, 1387 (1955).
- Bu 56 Buechner, Mazari, and Sperduto, Phys. Rev. **101**, 188 (1956).
- Bu 56a J. W. Butler, Bull. Am. Phys. Soc. Ser. II, **1**, 94 (1956) and private communication.
- Bu 56b B. S. Burton and R. M. Williamson, Bull. Am. Phys. Soc. Ser. II, **1**, 264 (1956) and private communication from R. M. Williamson.
- Bu 56c W. W. Buechner and A. Sperduto, Mass. Inst. Technol. Annual Progr. Rept., p. 107 (1956).
- Bu 56d Butler, Dunning, and Bondelid, Bull. Am. Phys. Soc. Ser. II, **1**, 327 (1956).
- Ca 56 P. J. Campion and G. A. Bartholomew, Bull. Am. Phys. Soc. Ser. II, **1**, 28 (1956).
- Ch 56 Chadwick, Alexander, and Warren, Can. J. Phys. **34**, 381 (1956).
- Ch 57 R. A. Chapman and J. C. Slattery, Phys. Rev. **105**, 633 (1957).
- Co 54a Cohen, Newman, Charpie, and Handley, Phys. Rev. **94**, 620 (1954).
- Co 55 Cook, Marion, and Bonner [private communication to F. Ajzenberg and T. Lauritsen, Revs. Modern Phys. **27**, 106 (1955)].
- Co 56 D. R. Connors and W. C. Miller, Bull. Am. Phys. Soc. Ser. II, **1**, 340 (1956).
- Cr 56 D. S. Craig, Phys. Rev. **101**, 1479 (1956).
- Cr 56a Craig, Cross, and Jarvis, Phys. Rev. **103**, 1427 (1956).
- De 55 de Sousa Santos, Goldemberg, Pieroni, Silva, Borello, Villaca, and Lopes, *Proceedings of International Conference on Peaceful Uses of Atomic Energy, Geneva, 1955* (United Nations, New York, 1956), Vol. II, p. 169.
- Do 56 W. T. Doyle and A. B. Robbins, Phys. Rev. **101**, 1056 (1956).
- Do 56a Douglas, Broer, Chiba, Herring, and Silverstein, Phys. Rev. **104**, 1059 (1956).
- Du 56 Dunning, Butler, and Bondelid, Bull. Am. Phys. Soc. Ser. II, **1**, 328 (1956).
- El 51c F. A. El-Bedewi, Proc. Phys. Soc. (London) **A64**, 1079 (1951).
- El 52 El-Bedewi, Middleton, and Tai, Nature **169**, 235 (1952).
- El 54 R. B. Elliot and D. J. Livesay, Proc. Roy. Soc. (London) **A224**, 129 (1954).
- El 55 F. A. El-Bedewi and M. A. El Wahab, Proc. Phys. Soc. (London) **A68**, 754 (1955).
- El 56 F. A. El-Bedewi, Proc. Phys. Soc. (London) **A69**, 221 (1956).
- El 56a A. J. Elwyn and F. B. Shull, Bull. Am. Phys. Soc. Ser. II, **1**, 281 (1956).
- El 57 F. A. El-Bedewi and I. Hussein, Proc. Phys. Soc. (London) **A70**, 233 (1957).
- En 54 Endt, Kluyver, and Van der Leun, Physica **20**, 1299 (1954).
- En 54a Endt, Kluyver, and Van der Leun, Phys. Rev. **95**, 580(L) (1954).
- En 56 Endt, Paris, Sperduto, and Buechner, Phys. Rev. **103**, 961 (1956).
- En 57 P. M. Endt and C. H. Paris, Phys. Rev. **106**, 764 (1957).
- Fa 55 W. J. Fader, Mass. Inst. Technol. Annual Progr. Rept., p. 123 (1955).
- Fl 53 F. C. Flack and J. B. Warren, Proc. Roy. Soc. (Canada) **47**, 131(A) (1953).
- Fo 54 G. M. Foglesong and D. G. Foxwell, Phys. Rev. **96**, 1001 (1954).
- Fo 56 Fowler, Lauritsen, Lauritsen, and Cook (private communication).
- Fr 54a Freemantle, Grotdal, Gibson, McKeague, Prowse, and Rotblat, Phil. Mag. **45**, 1090 (1954); erratum **45** (November, 1954).
- Gi 54 Gibson, Prowse, and Rotblat, Nature **173**, 1180 (1954).
- Gi 55 Gibbons, Macklin, and Schmitt, Phys. Rev. **100**, 167 (1955).
- Go 57 C. R. Gossett, Bull. Am. Phys. Soc. Ser. II, **2**, 69 (1957).
- Go 57a C. R. Gossett and J. W. Butler (private communication).
- Gr 54 G. W. Greenlees, Proc. Phys. Soc. (London) **67A**, 1107 (1954).
- Gr 55 Green, Scanlon, and Willmott, Proc. Phys. Soc. (London) **68A**, 386 (1955).
- Gr 55a Green, Singh, and Willmott, Phil. Mag. **46**, 982 (1955).

- Gr 55b Groshev, Adyasevich, and Demidov, *Proceedings of International Conference on Peaceful Uses of Atomic Energy, Geneva, 1955* (United Nations, New York, 1956), Vol. II, p. 39.
- Gr 55c G. M. Griffiths and T. B. Warren, Proc. Phys. Soc. (London) **68A**, 781 (1955).
- Gr 57 H. G. Graetzer and A. B. Robbins, Phys. Rev. **105**, 1570 (1957).
- Ha 49a Hanson, Duffield, Knight, Diven, and Palevsky, Phys. Rev. **76**, 578 (1949).
- Ha 54 Haslam, Roberts, and Robb, Can. J. Phys. **32**, 361 (1954).
- Ha 54a Halpern, Nathan, and Yergin, Phys. Rev. **95**, 1529 (1954).
- Ha 56 Hassler, Steigert, and Pieper, Bull. Am. Phys. Soc. Ser. II, **1**, 280 (1956).
- He 55 N. P. Heydenburg and G. M. Temmer, Phys. Rev. **99**, 617(A) (1955).
- Hj 53 E. Hjalmar and H. Slatis, Arkiv Fysik **6**, 451 (1953).
- Hj 53a E. Hjalmar and H. Slatis, Phys. Rev. **89**, 1151 (1953).
- Ho 55 Holmgren, Hanscome, and Willet, Phys. Rev. **98**, 241(A) (1955).
- Ih 55 M. A. Ihsan, Proc. Phys. Soc. (London) **A68**, 393 (1955).
- Is 50 R. Ishiwari and K. Yuasa, Mem. Coll. Sci. Univ. Kyoto **A26**, 151 (1950).
- Ja 55 James, Kubelka, Heiberg, and Warren, Can. J. Phys. **33**, 219 (1955).
- Ja 56 N. Jarmie, Phys. Rev. **104**, 1683 (1956).
- Jo 54 Jones, McEllistrem, Douglas, and Richards, Phys. Rev. **94**, 947 (1954).
- Jo 56 C. H. Johnson [private communication to R. A. Chapman and J. C. Slattery, Phys. Rev. **105**, 633 (1957)].
- Ju 54 J. J. Jung and C. K. Bockelman, Phys. Rev. **96**, 1353 (1954).
- Ka 51 Katz, Johns, Baker, Haslam, and Douglas, Phys. Rev. **82**, 271 (1951).
- Ka 53 Katz, Baker, and Montalbatti, Can. J. Phys. **31**, 250 (1953).
- Ka 53a L. Katz [private communication to J. E. Brolley, Phys. Rev. **89**, 877 (1953)].
- Ka 54 Katz, Haslam, Goldemberg, and Taylor, Can. J. Phys. **32**, 580 (1954).
- Ka 55 Kavanagh, Mills, and Sherr, Phys. Rev. **97**, 248(L) (1955).
- Kh 53 L. M. Khromchenko, Doklady Akad. Nauk. SSSR **93**, 451 (1953).
- Kh 54 L. M. Khromchenko, Doklady Akad. Nauk. SSSR **94**, 1037 (1954).
- Kh 55 L. M. Khromchenko and V. A. Blinov, Soviet Physics JETP **1**, 596 (1955).
- Ki 53 B. B. Kinsey and G. A. Bartholomew, Phys. Rev. **89**, 375 (1953).
- Ki 53c B. B. Kinsey and G. A. Bartholmew, Can. J. Phys. **31**, 1051 (1953).
- Ki 53d Kington, Bair, Carlson, and Willard, Phys. Rev. **89**, 530 (1953).
- Ki 55 Kington, Bair, Cohn, and Willard, Phys. Rev. **99**, 1393 (1955).
- Kl 54 Kluyver, Van der Leun, and Endt, Phys. Rev. **94**, 1795(L) (1955); Physica **20**, 1287 (1955).
- La 47 Lattes, Fowler, and Cuer, Proc. Phys. Soc. (London) **A59**, 883 (1947).
- Le 55 Levine, Bender, and McGruer, Phys. Rev. **97**, 1249 (1955).
- Le 56 L. L. Lee, Jr., and F. P. Mooring, Phys. Rev. **104**, 1342 (1956).
- Li 37 M. S. Livingston and H. A. Bethe, Revs. Modern Phys. **9**, 245 (1937).
- Li 55 J. G. Likely, Phys. Rev. **98**, 1538(A) (1955).
- Li 57 Lidofsky, Weil, and Jones, Bull. Am. Phys. Soc. Ser. II, **2**, 182 (1957).
- Lo 52 Lovington, McCue, and Preston, Phys. Rev. **85**, 585 (1952).
- Ma 55a Marion, Brugger, and Bonner, Phys. Rev. **100**, 46 (1955).
- Ma 55b Marion, Bonner, and Cook, Phys. Rev. **100**, 91 (1955).
- Ma 56 J. M. Marion and R. A. Chapman, Phys. Rev. **101**, 283 (1956).
- Ma 56a H. Mark and C. Goodman, Phys. Rev. **101**, 768 (1956).
- Ma 56b J. B. Marion and R. W. Kavanagh, Phys. Rev. **104**, 107 (1956).
- Mc 51c C. L. McGinnis, Phys. Rev. **81**, 734 (1951).
- Mc 55 McFarland, Shull, Elwyn, and Zeidman, Phys. Rev. **99**, 655(A) (1955).
- Mi 54 C. Mileikowsky, Arkiv Fysik **7**, 89 (1954).
- Mi 54a C. Mileikowsky, Arkiv Fysik **7**, 117 (1954).
- Mi 54b C. Mileikowsky and K. Ahnlund, Phys. Rev. **96**, 996 (1954).
- Mo 54 H. T. Motz, Phys. Rev. **93**, 925(A) (1954), probable error is our estimate.
- No 56 Nonaka, Morita, Kawai, Ishimatsu, Suematsu, Takeshita, Nakajima, and Wakuda, J. Phys. Soc. Japan **11**, 1 (1956).
- No 57 E. Norbeck, Jr., Phys. Rev. **105**, 204 (1957).
- Og 50a Ogle, Brown, and Carson, Phys. Rev. **78**, 63 (1950).
- Pa 54 R. Patterson, Phys. Rev. **95**, 303(A) (1954).
- Pa 54a Pauli, Ahnlund, and Mileikowsky, Arkiv Fysik **8**, 213 (1954).
- Pa 55 Paris, Buechner, and Endt, Phys. Rev. **100**, 1317 (1955).
- Pa 55a R. T. Pauli, Arkiv Fysik **9**, 571 (1955).
- Pa 56 A. Papkov, Z. Naturforsch. **11**, 776 (1956).
- Pe 52 Pepper, Allen, Almqvist, and Dewan, Phys. Rev. **85**, 155 (1952).
- Pi 52a G. F. Pieper, Phys. Rev. **88**, 1299 (1952).
- Pi 55 Pieper, Stanford, and von Herrmann, Phys. Rev. **98**, 1185(A) (1955).
- Pi 56 G. F. Pieper and G. S. Stanford, Phys. Rev. **101**, 672 (1956).
- Pl 55 H. S. Plendl and F. E. Steigert, Phys. Rev. **98**, 1538(A) (1955).
- Pr 52 Pruitt, Hanna, and Swartz, Phys. Rev. **87**, 534 (1952).
- Pr 53 Pruitt, Swartz, and Hanna, Phys. Rev. **92**, 1456 (1953).
- Pr 54 W. W. Pratt, Phys. Rev. **95**, 1517 (1954).
- Qu 56 A. R. Quinton and W. T. Doyle, Phys. Rev. **101**, 669 (1956).
- Qu 56a Quisenberry, Scolman, and Nier, Phys. Rev. **104**, 461 (1956).
- Re 54 G. C. Reid, Proc. Phys. Soc. (London) **A67**, 466 (1954).
- Re 55 J. B. Reynolds, Phys. Rev. **98**, 1289 (1955).
- Ri 51 Richards, Johnson, Ajzenberg, and Laubenstein, Phys. Rev. **83**, 994 (1951).
- Ro 51c J. Rotblat, Nature **167**, 550 (1951).
- Ro 55 Robinson, McPherson, Greenberg, Katz, and Haslam (unpublished), quoted in A. S. Penfold and B. M. Spicer, Phys. Rev. **100**, 1377 (1955).
- Ro 56 A. B. Robbins, Phys. Rev. **101**, 1373 (1956).
- Ru 56 Rubin, Johnson, and Reynolds, Phys. Rev. **104**, 1444 (1956).
- Sa 56 R. W. Sanders, Phys. Rev. **104**, 1434 (1956).
- Sc 54 Schnelder, Martin, Sempert, and Sutter, Helv. Phys. Acta **27**, 127A (1954).
- Sc 55 J. P. Schiffer, Phys. Rev. **97**, 428 (1955).
- Sc 56 M. J. Scott and R. E. Segel, Phys. Rev. **102**, 1557 (1956).
- Sc 56a Schwartz, Corbett, and Watson, Phys. Rev. **101**, 1370 (1956).
- Sh 51 Sher, Halpern, and Mann, Phys. Rev. **84**, 387 (1951).
- Sh 51a Sher, Halpern, and Stephens, Phys. Rev. **81**, 154 (1951).
- Si 56 E. Silva and J. Goldemberg, Nuovo cimento, **3**, 12 (1956).

- Sm 47 J. H. Smith, Phys. Rev. **71**, 32 (1947).  
 Sp 54 Sperduto, Buechner, Bockelman, and Browne, Phys. Rev. **96**, 1316 (1954).  
 Sp 55 A. Sperduto and W. W. Buechner, Phys. Rev. **100**, 961(A) (1955); Mass. Inst. Technol. Progr. Rept., p. 56 (November, 1955).  
 Sp 56 A. Sperduto and W. W. Buechner, Bull. Am. Phys. Soc. Ser. II, **1**, 223 (1956).  
 Sp 56a A. Sperduto and W. W. Buechner, Mass. Inst. Technol. Annual Progr. Rept., p. 111 (1956).  
 Sq 56 Squires, Bockelman, and Bueckner, Phys. Rev. **104**, 413 (1956).  
 St 52 P. H. Stelson and W. M. Preston, Phys. Rev. **86**, 807 (1952).  
 St 56 G. S. Stanford and G. F. Pieper, Phys. Rev. **103**, 637 (1956).  
 Su 55 S. Subotic and B. Maglic, Phil. Mag. **46**, 805(L) (1955).  
 Th 54 Thirion, Cohen, and Whaling, Phys. Rev. **96**, 850(A) (1954).  
 To 56 R. Tobin and J. McElhinney, Bull. Am. Phys. Soc. Ser. II, **1**, 340 (1956).  
 Tr 55 P. B. Treacy, Proc. Phys. Soc. (London) **68A**, 204 (1955).  
 Va 56 Van Patter, Swann, Porter, and Mandeville, Phys. Rev. **103**, 656 (1956).  
 Va 56a C. Van der Leun and P. M. Endt, Physica **22**, 1234 (1956).  
 Va 57 Van Patter, Porter, and Rothman, Phys. Rev. **106**, 1016 (1957).  
 Wa 54 N. S. Wall, Phys. Rev. **96**, 664 (1954).
- Wa 55 Way, King, McGinnis, and van Lieshout, "Nuclear Level Schemes," U.S. Atomic Energy Commission TID-5300 (1955).  
 Wa 56 Warren, Alexander, and Chadwick, Phys. Rev. **101**, 242 (1956).  
 Wh 55 W. Whaling [private communication to R. T. Pauli, Arkiv Fysik **9**, 571 (1955)].  
 Wi 52 Willard, Bair, Kington, Hahn, and Green, Phys. Rev. **85**, 849 (1952).  
 Wi 52a Willard, Kington, and Bair, Phys. Rev. **86**, 259 (1952).  
 Ye 55 P. F. Yergin and B. P. Fabricand, Phys. Rev. **100**, 1269(A) (1955).

**Corrections to "Nuclear Disintegration Energies"**  
**[Revs. Modern Phys. 26, 402 (1954)].**

- Li<sup>7</sup>( $p,\alpha$ )He<sup>4</sup> The average  $Q$  value should read  $17.345 \pm 0.006$  Mev.  
 Si<sup>28</sup>( $p,n$ )P<sup>28</sup> The  $Q$  value measured by Br 54a should be  $-14.9 \pm 0.6$  Mev.  
     The average  $Q$  value should read  $-15.0 \pm 0.4$  Mev.  
 Cr<sup>54</sup>( $p,n$ )Mn<sup>54</sup> The  $Q$  value measured by Lo 52 should read  $-2.162 \pm 0.005$  Mev.  
 Cu<sup>65</sup>( $n,\gamma$ )Cu<sup>66</sup> Assignment of 7.634 Mev  $\gamma$  to this reaction is now thought to be incorrect (Qu 56a, Wa 55).  
 Be 50d H. A. Bethe, Revs. Modern Phys. **22**, 213 (1950).  
 El 51d F. A. El-Bedewi, Proc. Phys. Soc. (London) **A64**, 584 (1951).