

Quantum Corrections to the Pekar Asymptotics of a Strongly Coupled Polaron

RUPERT L. FRANK

*Ludwig-Maximilians Universität München
California Institute of Technology*

ROBERT SEIRINGER

IST Austria (Institute of Science and Technology Austria)

Abstract

We consider the Fröhlich polaron model in the strong coupling limit. It is well-known that to leading order the ground state energy is given by the (classical) Pekar energy. In this work, we establish the subleading correction, describing quantum fluctuation about the classical limit. Our proof applies to a model of a confined polaron, where both the electron and the polarization field are restricted to a set of finite volume, with linear size determined by the natural length scale of the Pekar problem. © 2020 The Authors. *Communications on Pure and Applied Mathematics* published by Wiley Periodicals LLC

Contents

1. Introduction	1
2. Model and Main Results	2
3. The Pekar Functional	6
4. Proof of Theorem 2.6: Upper Bound	11
5. Multiple Lieb-Yamazaki Bound	15
6. Gross Transformation	19
7. Proof of Theorem 2.6: Lower Bound	23
Appendix A. Equivalent Formulation of Assumption 2.4	29
Appendix B. Bounds on Solutions of Poisson's Equation	31
Appendix C. Bounds on the Kernel of Functions of the Dirichlet Laplacian	39
Bibliography	44

1 Introduction

The polaron model was introduced by Fröhlich [10] as a model of an electron interacting with the quantized optical modes of a polar crystal. It represents a

Communications on Pure and Applied Mathematics, 0001–0045 (PREPRINT)

© 2020 The Authors. *Communications on Pure and Applied Mathematics* published by Wiley Periodicals LLC

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

simple and well-studied model of nonrelativistic quantum field theory, and we refer to [1, 8, 11, 23, 29] for properties, results, and further references.

In the strong coupling limit $\alpha \rightarrow \infty$, the model allows for an exact solution, in the sense that the ground state energy asymptotically equals the one given by the Pekar approximation [27], which amounts to a classical approximation to the quantum field theory. This was first shown by Donsker and Varadhan [5] using a path integral formulation of the problem. (See also [24, 25] for recent work on the construction of the Pekar process [29].) Later the result was improved by Lieb and Thomas [20, 21], who provided a quantitative bound on the difference.

We are interested here in the subleading correction to the classical (Pekar) approximation. It was predicted in the physics literature (see [2, 3, 15, 30] and references there) that this correction results from quantum fluctuations about the classical limit, and is $O(\alpha^{-2})$ smaller than the main term. It can be calculated by evaluating the ground state energy of a system of (infinitely many) harmonic oscillators with frequencies determined by the Hessian of the Pekar functional. This result is verified rigorously in this paper, by giving upper and lower bounds on the ground state energy of the Fröhlich polaron model that establish this subleading correction. Our analysis applies to a model of a confined polaron, where both the electron and the polarization field are restricted to a finite volume (with linear size of the natural length scale set by the Pekar problem).

The confinement breaks translation invariance, which removes zero modes otherwise present in the Hessian of the Pekar functional, and avoids having to localize the electron on the Pekar scale, which simplifies the problem. The singular ultraviolet behavior is unaffected by the confinement, however, and represents one of the main technical challenges. A key ingredient in our analysis is a multiple use of the commutator method of Lieb and Yamazaki [22], combined with Nelson's Gross transformation [14, 26].

2 Model and Main Results

2.1 The Model

For $\Omega \subset \mathbb{R}^3$ open, let Δ_Ω denote the Dirichlet Laplacian, and let v_x be the function $v_x(\cdot) = (-\Delta_\Omega)^{-1/2}(x, \cdot)$. The model we consider is defined by the Hamiltonian

$$(2.1) \quad \mathbb{H} := -\Delta_\Omega - a(v_x) - a^\dagger(v_x) + \mathbb{N}$$

in $L^2(\Omega) \otimes \mathcal{F}$, where \mathcal{F} is the bosonic Fock space over $L^2(\Omega)$. The creation and annihilation operators satisfy the commutation relation

$$[a(f), a^\dagger(g)] = \alpha^{-2} \langle f | g \rangle \quad \text{for } f, g \in L^2(\Omega)$$

with a parameter $\alpha > 0$. The field energy is given by the number operator $\mathbb{N} = \sum_j a^\dagger(\varphi_j) a(\varphi_j)$ for some orthonormal basis $\{\varphi_j\}$ in $L^2(\Omega)$ with spectrum $\sigma(\mathbb{N}) = \alpha^{-2} \{0, 1, 2, \dots\}$. We are interested in the ground state energy of \mathbb{H} as $\alpha \rightarrow \infty$.

We note that the expression (2.1) is somewhat formal, since $v_x \notin L^2(\Omega)$ and hence $a^\dagger(v_x)$ is not densely defined. The operator \mathbb{H} can be defined with the aid of its corresponding quadratic form, however. It is in fact well-known that \mathbb{H} defines a self-adjoint operator on a suitable domain; see [13] or Section 6 below.

Remark 2.1. By rescaling all lengths by α , \mathbb{H} is unitarily equivalent to the operator $\alpha^{-2}\tilde{\mathbb{H}}$ with

$$\tilde{\mathbb{H}} = -\Delta_{\Omega/\alpha} - \sqrt{\alpha}(\tilde{a}(\tilde{v}_x) - \tilde{a}^\dagger(\tilde{v}_x)) + \tilde{\mathbb{N}}$$

where $\tilde{v}_x(\cdot) = (-\Delta_{\Omega/\alpha})^{-1/2}(x, \cdot)$, $\tilde{\mathbb{N}} = \sum_j \tilde{a}^\dagger(\varphi_j)\tilde{a}(\varphi_j)$, and the \tilde{a} and \tilde{a}^\dagger operators satisfy $[\tilde{a}(f), \tilde{a}^\dagger(g)] = \langle f|g \rangle$ (and are thus independent of α). Large α hence corresponds to the strong coupling limit of a polaron confined to a region of linear size α^{-1} . We find it more convenient to work in the variables defined in (2.1), however.

Remark 2.2. Typically the polaron model is considered without confinement, i.e., for $\Omega = \mathbb{R}^3$, in which case the electron-phonon coupling function is taken to be $(-\Delta_{\mathbb{R}^3})^{-1/2}(x, y) = (2\pi)^{-3} \int_{\mathbb{R}^3} e^{ik \cdot (x-y)} |k|^{-1} dk = (2\pi^2)^{-1} |x-y|^{-2}$. For the proof of our main theorem the compactness of $(-\Delta_\Omega)^{-1}$ will be important; hence we need to consider bounded sets Ω here.

2.2 Pekar Functional(s)

We introduce the classical energy functional corresponding to (2.1) as

$$(2.2) \quad \mathcal{E}(\psi, \varphi) = \int_{\Omega} |\nabla \psi(x)|^2 dx + \int_{\Omega} \varphi(x)^2 dx - 2 \iint_{\Omega \times \Omega} \varphi(x) (-\Delta_\Omega)^{-1/2}(x, y) |\psi(y)|^2 dx dy$$

where $\psi \in H_0^1(\Omega)$, $\|\psi\|_2 = 1$, and $\varphi \in L^2_{\mathbb{R}}(\Omega)$, the real-valued functions in $L^2(\Omega)$. Formally, it can be obtained from (2.1) by replacing the field operators $a(f)$ and $a^\dagger(f)$ by $\int \varphi(x) f(x) dx$, and taking an expectation value with the electron wave function ψ . The Pekar energy is

$$(2.3) \quad e^P = \min_{\psi, \varphi} \mathcal{E}(\psi, \varphi).$$

For $\Omega = \mathbb{R}^3$ it was shown in [5, 20, 21] that $\inf \text{spec } \mathbb{H} \rightarrow e^P$ as $\alpha \rightarrow \infty$. The result can be shown to hold also for general Ω . Our goal here is to compute the subleading correction in this asymptotics.

We will work under the following:

ASSUMPTION 2.3. The functional \mathcal{E} in (2.2) has a unique minimizer ψ^P, φ^P (up to a trivial constant phase factor for ψ^P).

Our proof works under the more general assumption that the set of minimizers of \mathcal{E} is discrete (up to the phase degeneracy). The case where minimizers form a continuous manifold requires additional ideas, however.

Since $\mathcal{E}(|\psi|, \varphi) \leq \mathcal{E}(\psi, \varphi)$ we assume from now on that ψ^P is nonnegative. For given ψ , the choice of the minimizing φ is clearly unique, and vice versa. In particular, our Assumption 2.3 concerns uniqueness of the minimizer of the corresponding Pekar functional

$$\begin{aligned} \mathcal{E}^P(\psi) &= \min_{\varphi} \mathcal{E}(\psi, \varphi) \\ &= \int_{\Omega} |\nabla \psi(x)|^2 dx - \iint_{\Omega \times \Omega} |\psi(x)|^2 (-\Delta_{\Omega})^{-1}(x, y) |\psi(y)|^2 dx dy. \end{aligned}$$

Recall that, for $\Omega = \mathbb{R}^3$, uniqueness of minimizers of \mathcal{E}^P (up to translations and phase factor) is known [18] (see also [31]). We expect Assumption 2.3 to hold if Ω is convex, for instance. The proof in [18] can be adapted to show uniqueness in case Ω is a ball [6].

ASSUMPTION 2.4. There exists a $\kappa > 0$ such that

$$(2.4) \quad \mathcal{E}^P(\psi) \geq \mathcal{E}^P(\psi^P) + \kappa \int_{\Omega} |\nabla(\psi - \psi^P)|^2 \quad \forall \psi \in H_0^1(\Omega), \psi \geq 0, \|\psi\|_2 = 1.$$

The bound (2.4) follows from an a priori weaker spectral assumption on the absence of nontrivial zero modes of the Hessian of \mathcal{E}^P at its minimizer ψ^P by a simple compactness argument. For completeness, we spell out the details of this argument in Appendix A. The analogue of this spectral assumption in the case $\Omega = \mathbb{R}^3$ is known (up to zero modes resulting from the translation invariance) [17, 32]. Using the method in [17], one can prove Assumption 2.4 in case Ω is a ball [6].

If one minimizes $\mathcal{E}(\psi, \varphi)$ over ψ for given φ , one obtains the functional

$$(2.5) \quad \mathcal{F}^P(\varphi) = \min_{\psi} \mathcal{E}(\psi, \varphi) = \|\varphi\|_2^2 + \inf \text{spec}(-\Delta_{\Omega} + V_{\varphi}(x))$$

where $V_{\varphi} = -2(-\Delta_{\Omega})^{-1/2}\varphi$. Let H^P denote its Hessian at the unique minimizer φ^P , i.e.,

$$(2.6) \quad \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2} (\mathcal{F}^P(\varphi^P + \varepsilon\varphi) - e^P) = \langle \varphi | H^P | \varphi \rangle \quad \forall \varphi \in L_{\mathbb{R}}^2(\Omega).$$

An explicit computation gives

$$(2.7) \quad H^P = \mathbb{1} - 4(-\Delta_{\Omega})^{-1/2} \psi^P \frac{Q^P}{-\Delta_{\Omega} + V_{\varphi^P} - \mu^P} \psi^P (-\Delta_{\Omega})^{-1/2}$$

where ψ^P acts as a multiplication operator, $\mu^P = \inf \text{spec}(-\Delta_{\Omega} + V_{\varphi^P}) = e^P - \|\varphi^P\|_2^2$, and Q^P is the projection orthogonal to ψ^P , i.e., orthogonal to the kernel of $-\Delta_{\Omega} + V_{\varphi^P} - \mu^P$. It is not difficult to see that Assumption 2.4 implies that H^P is nondegenerate, i.e., strictly positive (compare with Proposition 3.4 in Section 3.2 below).

Finally, we need a regularity assumption on the domain Ω .

ASSUMPTION 2.5. The domain Ω is bounded and has a $C^{3,\delta}$ boundary for some $0 < \delta < 1$.

For a proper definition of the meaning of $C^{3,\delta}$ boundary, see Appendix B. Assumption 2.5 allows us to estimate derivatives of the integral kernel of certain functions of the Dirichlet Laplacian (see Appendix C). The required estimates certainly hold under less restrictive assumptions on Ω , and we expect our main result to hold also in case Ω is a cube, for instance. We shall not try to investigate the minimal regularity assumptions, however, and shall henceforth work with Assumption 2.5.

2.3 Main Result

Recall the definition (2.3) for the Pekar energy e^P , as well as (2.7) for the Hessian H^P of \mathcal{F}^P in (2.5) at the unique minimizer φ^P . Our main result is as follows.

THEOREM 2.6. *Under Assumptions 2.3–2.5 one has, as $\alpha \rightarrow \infty$,*

$$(2.8) \quad \inf \text{spec } \mathbb{H} = e^P - \frac{1}{2\alpha^2} \text{Tr}(\mathbb{1} - \sqrt{H^P}) + o(\alpha^{-2}).$$

More precisely, the bounds

$$(2.9) \quad \begin{aligned} -C\alpha^{-1/7} (\ln \alpha)^{5/14} &\leq \alpha^2 \inf \text{spec } \mathbb{H} - \alpha^2 e^P + \frac{1}{2} \text{Tr}(\mathbb{1} - \sqrt{H^P}) \\ &\leq C\alpha^{-2/11} \end{aligned}$$

hold for some constant $C > 0$ and α large enough.

The trace in (2.8) and (2.9) is over $L^2(\Omega)$. We shall see below that $\mathbb{1} - \sqrt{H^P}$ is actually trace class. Note also that $H^P < \mathbb{1}$; hence the coefficient of α^{-2} in (2.8) is strictly negative.

In the case $\Omega = \mathbb{R}^3$, the correctness of the leading term e^P was shown in [5, 20, 21]. The proof in [20, 21] gives an error bound of the order $\alpha^{-1/5}$. In the confined case considered here, we improve this error bound to $O(\alpha^{-2})$ and actually compute the next order correction. We conjecture that the formula (2.8) also holds true in case $\Omega = \mathbb{R}^3$, as predicted in the physics literature [2, 3, 15, 30]. Our upper bound, in fact, can easily be generalized to this case. While our methods are not strong enough to prove the corresponding lower bound, parts of our proof are applicable also to the $\Omega = \mathbb{R}^3$ case and yield an improved error bound compared to the one given in [20, 21].

The α^{-2} correction to the ground state energy in (2.8) can be interpreted as arising from quantum fluctuations around the classical limit described by the Pekar functional. The trace originates from the ground state energy of a Hamiltonian describing a system of (infinitely) many harmonic oscillators.

The remainder of the paper is devoted to the proof of Theorem 2.6. We start with a brief outline to guide the reader.

2.4 Outline of the Proof

In Section 3 we study the Pekar functional (2.5). We shall compute its Hessian at the unique minimizer φ^{P} and use it to estimate the functional in a small neighborhood of its minimizer. We shall also derive a useful quadratic lower bound that is valid globally, i.e., not just close to the minimizer.

In Section 4 we shall derive an upper bound on the ground state energy of \mathbb{H} that has the desired asymptotic form as $\alpha \rightarrow \infty$. We shall construct an appropriate trial state and utilize the estimate of the Pekar functional close to its minimizer from the previous section.

Sections 5 and 6 contain auxiliary results that are essential for the lower bound, in particular to allow for an ultraviolet regularization of the problem. In Section 5 the commutator method of Lieb and Yamazaki [22] is applied three times in order to estimate the effect of an ultraviolet cutoff in the coupling function v_x in terms of the number operator \mathbb{N} and the electron kinetic energy $-\Delta_\Omega$. The relevant operator that needs to be bounded is $\mathbb{N}^{1/2}(-\Delta_\Omega)^{3/2}$, which cannot be controlled in terms of \mathbb{H}^2 , however. The necessary bound does hold after a unitary Gross transformation, which shall be explained in Section 6. This will be sufficient for our purpose.

In Section 7 we shall give a lower bound on the ground state energy of \mathbb{H} of the desired asymptotic form. We shall use the results of Sections 5 and 6 to implement an ultraviolet cutoff, which effectively reduces the problem to finitely many modes. We shall then use an IMS localization in Fock space and the bounds in Section 3 to conclude the desired lower bound.

In Appendix A we shall give an equivalent formulation of Assumption 2.4 in terms of spectral properties of the Hessian of \mathcal{E}^{P} . In the appendices we shall derive bounds on derivatives of the integral kernel of certain functions of the Dirichlet Laplacian Δ_Ω that we need in our proof. These bounds are derived in Appendix C utilizing a theorem in Appendix B on bounds on solutions of Poisson's equation.

Throughout the proof, we shall use the symbol $a \lesssim b$ if $a \leq Cb$ for some constant $C > 0$.

3 The Pekar Functional

3.1 Hessian of the Pekar Functional

We consider the Pekar functional (2.5) and write it as

$$\mathcal{F}^{\text{P}}(\varphi) = e(\varphi) + \|\varphi\|_2^2$$

with

$$e(\varphi) = \inf \text{spec } H_\varphi \quad \text{and} \quad H_\varphi = -\Delta_\Omega + V_\varphi(x).$$

Recall that for $\varphi \in L^2_{\mathbb{R}}(\mathbb{R}^3)$ we set $V_\varphi = -2(-\Delta_\Omega)^{-1/2}\varphi$. In this section we work under Assumption 2.3, which states that $\mathcal{F}^{\text{P}}(\varphi)$ has a unique minimizer φ^{P} . We have $e(\varphi) + \|\varphi\|_2^2 \geq e(\varphi^{\text{P}}) + \|\varphi^{\text{P}}\|_2^2$, and our goal in this section is to obtain upper and lower bounds on the difference.

Recall that ψ^P denotes the unique nonnegative minimizer of \mathcal{E}^P , which is the ground state of H_{φ^P} . We have

$$(3.1) \quad \varphi^P = (-\Delta_\Omega)^{-1/2} |\psi^P|^2.$$

For later use, we record that ψ^P is a bounded function.

LEMMA 3.1. $\psi^P \in L^\infty(\Omega)$

PROOF. The Euler-Lagrange equation for the Pekar minimizer ψ^P reads

$$-\Delta_\Omega \psi^P - 2((-\Delta_\Omega)^{-1} |\psi^P|^2) \psi^P = \mu \psi^P$$

for some $\mu \in \mathbb{R}$, which we rewrite as

$$\psi^P = (-\Delta_\Omega)^{-1} ((\mu + 2((-\Delta_\Omega)^{-1} |\psi^P|^2)) \psi^P).$$

From (C.2) we deduce that $(-\Delta_\Omega)^{-1}(x, y) \leq (-\Delta_{\mathbb{R}^3})^{-1}(x, y) = (4\pi |x - y|)^{-1}$. By Sobolev's inequality $|\psi^P|^2 \in L^3(\Omega)$, and hence $(-\Delta_\Omega)^{-1} |\psi^P|^2 \in L^\infty(\Omega)$ by Hölder's inequality. Thus, $f = (\mu + 2((-\Delta_\Omega)^{-1} |\psi^P|^2)) \psi^P \in L^2(\Omega)$, and once again by Hölder's inequality, $\psi^P = (-\Delta_\Omega)^{-1} f \in L^\infty(\Omega)$, as claimed. \square

Let $P = |\psi^P\rangle\langle\psi^P|$ and $Q = \mathbb{1} - P$. We introduce the following nonnegative operators

$$(3.2) \quad K = 4(-\Delta_\Omega)^{-1/2} \psi^P \frac{Q}{H_{\varphi^P} - e(\varphi^P)} \psi^P (-\Delta_\Omega)^{-1/2}$$

and

$$L = 4(-\Delta_\Omega)^{-1/2} \psi^P (-\Delta_\Omega)^{-1} \psi^P (-\Delta_\Omega)^{-1/2},$$

where ψ^P acts as a multiplication operator. We shall see that $K = \mathbb{1} - H^P$, where H^P denotes the Hessian of $\mathcal{F}^P(\varphi)$ at $\varphi = \varphi^P$, introduced in (2.6) above.

It is easy to see that L is trace class, since $(-\Delta_\Omega)^{-1/2} \psi^P (-\Delta_\Omega)^{-1/2}$ is Hilbert-Schmidt for any multiplication operator $\psi \in L^2(\Omega)$. In fact, since $(-\Delta_\Omega)^{-1/2} \leq \sqrt{2}(-\Delta_\Omega + e_1)^{-1/2}$ (with $e_1 = \inf \text{spec}(-\Delta_\Omega) > 0$) and $(-\Delta_\Omega + e_1)^{-1/2}(x, y) \leq (-\Delta_{\mathbb{R}^3} + e_1)^{-1/2}(x, y)$ for any $x, y \in \mathbb{R}^3$ by (C.2), the Cauchy-Schwarz inequality implies that

$$(3.3) \quad \begin{aligned} & \text{Tr} [(-\Delta_\Omega)^{-1/2} \psi^P (-\Delta_\Omega)^{-1/2}]^2 \\ & \leq \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} \left(\frac{2}{k^2 + e_1} \right)^2 dk \int_\Omega |\psi(x)|^2 dx. \end{aligned}$$

To show that also K is trace class, we shall first prove the following lemma, which implies, in particular, that V_φ is operator-bounded relative to $-\Delta_\Omega$ if $\varphi \in L^2(\Omega)$.

LEMMA 3.2. *With $V_\varphi(x) = -2(-\Delta_\Omega)^{-1/2} \varphi(x)$, we have*

$$\|V_\varphi (-\Delta_\Omega)^{-1}\|^2 \lesssim \langle \varphi | (-\Delta_\Omega)^{-1} | \varphi \rangle.$$

PROOF. Note that the right side is simply the square of the L^2 -norm of V_φ . By arguing as in (3.3), one readily checks that the desired bound even holds with the Hilbert–Schmidt norm on the left side. \square

A straightforward modification of the proof shows that V_φ is actually infinitesimally operator-bounded relative to Δ_Ω , i.e., $\lim_{\kappa \rightarrow \infty} \|V_\varphi(-\Delta_\Omega + \kappa)^{-1}\| = 0$. This readily implies that

$$(-\Delta_\Omega)^{1/2} \frac{Q}{H_{\varphi^P} - e(\varphi^P)} (-\Delta_\Omega)^{1/2}$$

is bounded; hence the trace class property of K follows from the one of L .

Our main result in this section is the following.

PROPOSITION 3.3. *Assume that $\varphi \in L^2_{\mathbb{R}}(\Omega)$ is such that*

$$(3.4) \quad \|(-\Delta_\Omega)^{-1/2}(\varphi - \varphi^P)\|_2 \leq \varepsilon$$

for $\varepsilon > 0$ small enough. Then

$$(3.5) \quad |\mathcal{F}^P(\varphi) - \mathcal{F}^P(\varphi^P) - \langle \varphi - \varphi^P | \mathbb{1} - K | \varphi - \varphi^P \rangle| \lesssim \varepsilon \langle \varphi - \varphi^P | L | \varphi - \varphi^P \rangle.$$

This result implies, in particular, that $0 \leq K \leq \mathbb{1}$. It identifies $H^P = \mathbb{1} - K$ as the Hessian of $\mathcal{F}^P(\varphi) = e(\varphi) + \|\varphi\|_2^2$ at the minimizer φ^P . Our assumption on the strict positivity of the Hessian thus translates, in view of the compactness of K , to the statement $\|K\| < 1$.

PROOF. By choosing $\varepsilon > 0$ small enough and arguing as in the proof of Lemma 3.2, we can ensure that the family of operators $-\Delta_\Omega + V_\varphi(x)$ has a unique eigenvalue close to $e(\varphi^P)$, and this eigenvalue is $e(\varphi)$. The rest of the spectrum of H_φ is uniformly bounded away from $e(\varphi^P)$. Hence we can write

$$(3.6) \quad e(\varphi) = \text{Tr} \int_C \frac{z}{z - H_\varphi} \frac{dz}{2\pi i}$$

for a fixed (i.e., φ -independent) contour C that encircles $e(\varphi^P)$.

We claim that the operator $\Delta_\Omega(z - H_{\varphi^P})^{-1}$ is uniformly bounded for $z \in C$. This follows from the fact that the multiplication operator V_{φ^P} is infinitesimally operator-bounded relative to $-\Delta_\Omega$, as already argued after the proof of Lemma 3.2 above. Consequently,

$$(3.7) \quad \sup_{z \in C} \|V_{\varphi - \varphi^P}(z - H_{\varphi^P})^{-1}\| < 1$$

for small ε , by Lemma 3.2 and our assumption (3.4). We can thus use the resolvent identity in the form

$$\begin{aligned} & \frac{1}{z - H_\varphi} \\ &= \left(\mathbb{1} - \frac{Q}{z - H_{\varphi^P}} V_{\varphi - \varphi^P} \right)^{-1} \frac{Q}{z - H_{\varphi^P}} \\ & \quad + \left(\mathbb{1} - \frac{Q}{z - H_{\varphi^P}} V_{\varphi - \varphi^P} \right)^{-1} \frac{P}{z - e(\varphi^P)} \left(\mathbb{1} - V_{\varphi - \varphi^P} \frac{1}{z - H_{\varphi^P}} \right)^{-1}. \end{aligned}$$

The first term on the right side is analytic in z for all z inside the contour C , and hence gives 0 after integration when inserted in (3.6). The second term is rank 1, and Fubini's theorem implies that we can interchange the trace and the integral after inserting this term in (3.6). We thus obtain

$$(3.8) \quad \begin{aligned} e(\varphi) &= \int_C \frac{z}{z - e(\varphi^P)} \left\langle \psi^P \left| \left(\mathbb{1} - V_{\varphi - \varphi^P} \frac{1}{z - H_{\varphi^P}} \right)^{-1} \right. \right. \\ & \quad \left. \left. \times \left(\mathbb{1} - \frac{Q}{z - H_{\varphi^P}} V_{\varphi - \varphi^P} \right)^{-1} \right| \psi^P \right\rangle \frac{dz}{2\pi i}. \end{aligned}$$

For simplicity, let us introduce the notation

$$A = V_{\varphi - \varphi^P} \frac{1}{z - H_{\varphi^P}}, \quad B = \frac{Q}{z - H_{\varphi^P}} V_{\varphi - \varphi^P}.$$

Because of (3.7) these operators are smaller than 1 in norm, uniformly in $z \in C$. We shall use the identity

$$(3.9) \quad \begin{aligned} \frac{1}{\mathbb{1} - A} \frac{1}{\mathbb{1} - B} &= \mathbb{1} + A + A(A + B) + \frac{B}{\mathbb{1} - B} \\ & \quad + \frac{A^3}{\mathbb{1} - A} + \frac{A^2}{\mathbb{1} - A} B + \frac{A}{\mathbb{1} - A} \frac{B^2}{\mathbb{1} - B}. \end{aligned}$$

We insert the various terms into (3.8) and do the contour integration. The term $\mathbb{1}$ then yields $e(\varphi^P)$. The term A yields

$$\langle \psi^P | V_{\varphi - \varphi^P} | \psi^P \rangle = 2 \int_{\Omega} \varphi^P(x) (\varphi^P(x) - \varphi(x)) dx$$

using (3.1). A standard calculation shows that the term $A(A + B)$ leads to

$$\left\langle \psi^P \left| V_{\varphi - \varphi^P} \frac{Q}{e(\varphi^P) - H_{\varphi^P}} V_{\varphi - \varphi^P} \right| \psi^P \right\rangle = -\langle \varphi - \varphi^P | K | \varphi - \varphi^P \rangle.$$

Furthermore, since $Q | \psi^P \rangle = 0$, the term $B(\mathbb{1} - B)^{-1}$ yields 0. We conclude that

$$(3.10) \quad \begin{aligned} & \mathcal{F}^P(\varphi) - \mathcal{F}^P(\varphi^P) - \langle \varphi - \varphi^P | \mathbb{1} - K | \varphi - \varphi^P \rangle \\ &= \int_C \frac{z}{z - e(\varphi^P)} \left\langle \psi^P \left| \frac{A^3}{\mathbb{1} - A} + A \left(\frac{A}{\mathbb{1} - A} + \frac{1}{\mathbb{1} - A} \frac{B}{\mathbb{1} - B} \right) B \right| \psi^P \right\rangle \frac{dz}{2\pi i}. \end{aligned}$$

To bound the first term on the right side of (3.10), note that

$$(3.11) \quad \left\langle \psi^{\text{P}} \left| \frac{A^3}{\mathbb{1} - A} \right| \psi^{\text{P}} \right\rangle = \frac{1}{z - e(\varphi^{\text{P}})} \left\langle \psi^{\text{P}} \left| V_{\varphi - \varphi^{\text{P}}} \frac{1}{z - H_{\varphi^{\text{P}}} \mathbb{1} - A} V_{\varphi - \varphi^{\text{P}}} \right| \psi^{\text{P}} \right\rangle.$$

We claim that

$$(3.12) \quad \sup_{z \in \mathbb{C}} \left\| (-\Delta_{\Omega})^{1/2} \frac{1}{z - H_{\varphi^{\text{P}}} \mathbb{1} - A} (-\Delta_{\Omega})^{1/2} \right\| \lesssim \varepsilon,$$

which implies that (3.11) is bounded, in absolute value, as

$$(3.13) \quad |(3.11)| \lesssim \varepsilon \langle \psi^{\text{P}} | V_{\varphi - \varphi^{\text{P}}} (-\Delta_{\Omega})^{-1} V_{\varphi - \varphi^{\text{P}}} | \psi^{\text{P}} \rangle = \varepsilon \langle \varphi - \varphi^{\text{P}} | L | \varphi - \varphi^{\text{P}} \rangle,$$

as desired. To prove (3.12) we use the fact that $\|(-\Delta_{\Omega})^{1/2} (z - H_{\varphi^{\text{P}}})^{-1} (-\Delta_{\Omega})^{1/2}\|$ is uniformly bounded in order to reduce the problem to showing

$$\|(-\Delta_{\Omega})^{-1/2} A (\mathbb{1} - A)^{-1} (-\Delta_{\Omega})^{1/2}\| \lesssim \varepsilon.$$

Since $S^{-1} A (\mathbb{1} - A)^{-1} S = S^{-1} A S (\mathbb{1} - S^{-1} A S)^{-1}$ with $S = (-\Delta_{\Omega})^{1/2}$, it suffices to show that $\|(-\Delta_{\Omega})^{-1/2} A (-\Delta_{\Omega})^{1/2}\| \lesssim \varepsilon$, which follows from

$$\|(-\Delta_{\Omega})^{-1/2} V_{\varphi} (-\Delta_{\Omega})^{-1/2}\| \leq \|V_{\varphi} (-\Delta_{\Omega})^{-1}\|$$

and Lemma 3.2.

For the last term in (3.10), we simply bound

$$(3.14) \quad \left| \left\langle \psi^{\text{P}} \left| A \left(\frac{A}{\mathbb{1} - A} + \frac{1}{\mathbb{1} - A} \frac{B}{\mathbb{1} - B} \right) B \right| \psi^{\text{P}} \right\rangle \right| \\ \leq \left\| \frac{A}{\mathbb{1} - A} + \frac{1}{\mathbb{1} - A} \frac{B}{\mathbb{1} - B} \right\| \left\| \langle \psi^{\text{P}} | A A^{\dagger} | \psi^{\text{P}} \rangle^{1/2} \langle \psi^{\text{P}} | B^{\dagger} B | \psi^{\text{P}} \rangle^{1/2} \right\|.$$

The same bounds as above easily lead to the conclusion that also this term is bounded by the right side of (3.13). This concludes the proof of Proposition 3.3. \square

3.2 A Uniform Quadratic Lower Bound

Inequality (3.5) gives a bound on \mathcal{F}^{P} for φ near the minimizer φ^{P} . We shall also need the following rougher global bound.

PROPOSITION 3.4. *There is a constant $\kappa' > 0$ such that for all $\varphi \in L^2_{\mathbb{R}}(\Omega)$,*

$$(3.15) \quad \mathcal{F}^{\text{P}}(\varphi) \geq e^{\text{P}} + \langle \varphi - \varphi^{\text{P}} | \mathbb{1} - (\mathbb{1} + \kappa' (-\Delta_{\Omega})^{1/2})^{-1} | \varphi - \varphi^{\text{P}} \rangle.$$

We start with the following lemma.

LEMMA 3.5. *For $\psi \in H^1_0(\Omega)$ with $\|\psi\|_2 = 1$,*

$$\langle |\psi|^2 - |\psi^{\text{P}}|^2 | (-\Delta_{\Omega})^{-1/2} | \psi|^2 - |\psi^{\text{P}}|^2 \rangle \leq \frac{8}{\pi^2} \int_{\Omega} |\nabla(|\psi| - |\psi^{\text{P}}|)|^2.$$

PROOF. Given that $f(x) = |\psi(x)| + |\psi^P(x)|$ and $g(x) = |\psi(x)| - |\psi^P(x)|$, the Schwarz inequality and the symmetry and positivity of the integral kernel of $(-\Delta_\Omega)^{-1/2}$ imply that

$$\begin{aligned} & (|\psi|^2 - |\psi^P|^2)(-\Delta_\Omega)^{-1/2}(|\psi|^2 - |\psi^P|^2) \\ &= \int_\Omega \int_\Omega f(x)g(x)(-\Delta_\Omega)^{-1/2}(x,y)f(y)g(y)dx dy \\ &\leq \int_\Omega \int_\Omega f(x)^2(-\Delta_\Omega)^{-1/2}(x,y)g(y)^2 dx dy. \end{aligned}$$

For fixed x , we can use the Hardy inequality and the fact that $(-\Delta_\Omega)^{-1/2}(x,y) \leq (-\Delta_{\mathbb{R}^3})^{-1/2}(x,y) = (2\pi^2)^{-1}|x-y|^{-2}$ from (C.2) to obtain the bound

$$\int_\Omega (-\Delta_\Omega)^{-1/2}(x,y)g(y)^2 dy \leq \frac{2}{\pi^2} \int_\Omega |\nabla g|^2.$$

Since $\int_\Omega f^2 \leq 4$, the result follows. \square

PROOF OF PROPOSITION 3.4. From our assumption (2.4) on the Hessian of the Pekar functional \mathcal{E}^P and Lemma 3.5, it follows that

$$\begin{aligned} \mathcal{E}^P(\psi) &\geq \mathcal{E}^P(|\psi|) \\ &\geq \mathcal{E}^P(\psi^P) + \kappa'(|\psi|^2 - |\psi^P|^2)(-\Delta_\Omega)^{-1/2}(|\psi|^2 - |\psi^P|^2) \end{aligned}$$

for $\kappa' = \kappa\pi^2/8$. In particular,

$$\begin{aligned} \mathcal{E}(\psi, \varphi) &= \mathcal{E}^P(\psi) + \|\varphi - (-\Delta_\Omega)^{-1/2}|\psi|^2\|_2^2 \\ &\geq e^P + \kappa'(|\psi|^2 - |\psi^P|^2)(-\Delta_\Omega)^{-1/2}(|\psi|^2 - |\psi^P|^2) \\ &\quad + \|\varphi - (-\Delta_\Omega)^{-1/2}|\psi|^2\|_2^2. \end{aligned}$$

Minimizing with respect to ψ and using (3.1) leads to the desired lower bound. \square

4 Proof of Theorem 2.6: Upper Bound

In this section we construct a trial state to derive an upper bound on the polaron ground state energy. Our trial state Ψ will depend only on finitely many phonon variables. More precisely, for Π a finite rank projection on $L^2_{\mathbb{R}}(\Omega)$, we can write the Fock space $\mathcal{F}(L^2(\Omega))$ as a tensor product $\mathcal{F}(\Pi L^2(\Omega)) \otimes \mathcal{F}((\mathbb{1} - \Pi)L^2(\Omega))$, and our trial state corresponds to the vacuum vector in the second factor $\mathcal{F}((\mathbb{1} - \Pi)L^2(\Omega))$. The first factor $\mathcal{F}(\Pi L^2(\Omega))$ can naturally be identified with $L^2(\mathbb{R}^N)$ corresponding to N simple harmonic oscillators, where $N = \dim \text{ran } \Pi$.

We find it convenient to identify a point $(\lambda_1, \dots, \lambda_N) \in \mathbb{R}^N$ with a function $\varphi \in \text{ran } \Pi$ via the identification

$$(4.1) \quad \varphi = \Pi\varphi = \sum_{n=1}^N \lambda_n \varphi_n$$

for some orthonormal basis $\{\varphi_n\}$ of $\text{ran } \Pi$. With this identification, we can think of a wave function $\Psi \in L^2(\Omega) \otimes L^2(\mathbb{R}^N)$ as a function $\Psi(x, \varphi)$ with $x \in \Omega, \varphi \in \text{ran } \Pi$.

The function we choose is as follows:

$$(4.2) \quad \begin{aligned} \Psi(x, \varphi) &= e^{-\alpha^2 \langle \varphi - \varphi^P | (\mathbb{1} - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle} \\ &\quad \times \chi(\varepsilon^{-1} \|(-\Delta_\Omega)^{-1/2}(\varphi - \varphi^P)\|_2) \psi_\varphi(x) \end{aligned}$$

where

- $\varepsilon > 0$ is a small parameter that will be chosen to go to 0 as $\alpha \rightarrow \infty$.
- $0 \leq \chi \leq 1$ is a smooth cutoff function with $\chi(t) = 1$ for $t \leq 1/2$ and $\chi(t) = 0$ for $t \geq 1$.
- Π is a finite rank projection on $L^2_{\mathbb{R}}(\Omega)$ with range containing φ^P .
- ψ_φ is the unique nonnegative, normalized ground state of $H_\varphi = -\Delta_\Omega + V_\varphi$.
- $K = \mathbb{1} - H^P$, explicitly given in (3.2).

On states of the type described above (corresponding to the vacuum for all modes outside the range of Π), the Hamiltonian (2.1) simply acts as $H_\varphi + \mathbb{N}$, with

$$\mathbb{N} = \sum_{n=1}^N \left(-\frac{1}{4\alpha^4} \partial_{\lambda_n}^2 + \lambda_n^2 - \frac{1}{2\alpha^2} \right).$$

Using the eigenvalue equation $H_\varphi \psi_\varphi = e(\varphi) \psi_\varphi$, the energy of our trial state Ψ is thus given as

$$(4.3) \quad \langle \Psi | \mathbb{H} | \Psi \rangle = \langle \Psi | e(\varphi) + \mathbb{N} | \Psi \rangle.$$

Since Ψ is supported on the set $\{\|(-\Delta_\Omega)^{-1/2}(\varphi - \varphi^P)\|_2 \leq \varepsilon\}$, we can use Proposition 3.3 for an upper bound on $e(\varphi)$. This leads to

$$(4.4) \quad \begin{aligned} \langle \Psi | \mathbb{H} | \Psi \rangle &\leq e^P \langle \Psi | \Psi \rangle \\ &\quad + \langle \Psi | \mathbb{N} - \|\varphi\|_2^2 + \langle \varphi - \varphi^P | \mathbb{1} - K + \varepsilon C L | \varphi - \varphi^P \rangle | \Psi \rangle \end{aligned}$$

for a suitable constant $C > 0$.

Utilizing the fact that the Gaussian factor in Ψ satisfies

$$\begin{aligned} &\left(-\frac{1}{4\alpha^4} \sum_{n=1}^N \partial_{\lambda_n}^2 + \langle \varphi - \varphi^P | \mathbb{1} - K | \varphi - \varphi^P \rangle \right) \\ &\quad \times e^{-\alpha^2 \langle \varphi - \varphi^P | (\mathbb{1} - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle} \\ &= \frac{1}{2\alpha^2} \text{Tr}(\mathbb{1} - \Pi K \Pi)^{1/2} e^{-\alpha^2 \langle \varphi - \varphi^P | (\mathbb{1} - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle}, \end{aligned}$$

we can integrate by parts and rewrite the right side of (4.4) as

$$\left(e^P - \frac{1}{2\alpha^2} \text{Tr}[\mathbb{1} - (\mathbb{1} - \Pi K \Pi)^{1/2}] \right) \langle \Psi | \Psi \rangle + A + B$$

with

$$A = \varepsilon C \langle \Psi | \langle \varphi - \varphi^P | L | \varphi - \varphi^P \rangle | \Psi \rangle$$

and

$$B = \frac{1}{4\alpha^4} \sum_{n=1}^N \int_{\Omega} dx \int_{\mathbb{R}^N} \prod_{m=1}^N d\lambda_m e^{-2\alpha^2 \langle \varphi - \varphi^P | (1 - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle} \\ \times \left| \partial_{\lambda_n} \left(\chi(\varepsilon^{-1} \|(-\Delta_{\Omega})^{-1/2}(\varphi - \varphi^P)\|_2) \psi_{\varphi}(x) \right) \right|^2.$$

We claim that L is bounded by $(-\Delta_{\Omega})^{-1}$. This follows immediately from the boundedness of ψ^P shown in Lemma 3.1. Alternatively, one can use that $\psi(-\Delta_{\Omega})^{-1}\psi$ is a bounded operator for $\psi \in L^3(\Omega)$ by Sobolev's inequality. Hence we can use the rough bound

$$A \lesssim \varepsilon^3 \langle \Psi | \Psi \rangle.$$

Moreover, by a simple Cauchy-Schwarz inequality, $B \leq 2(B_1 + B_2)$ with

$$B_1 = \frac{1}{4\alpha^4} \int_{\mathbb{R}^N} \prod_{m=1}^N d\lambda_m e^{-2\alpha^2 \langle \varphi - \varphi^P | (1 - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle} \\ \times \chi(\varepsilon^{-1} \|(-\Delta_{\Omega})^{-1/2}(\varphi - \varphi^P)\|_2)^2 \sum_{n=1}^N \|\partial_{\lambda_n} \psi_{\varphi}\|_2^2$$

and

$$B_2 = \frac{1}{4\alpha^4} \int_{\mathbb{R}^N} \prod_{m=1}^N d\lambda_m e^{-2\alpha^2 \langle \varphi - \varphi^P | (1 - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle} \\ \times \sum_{n=1}^N \left| \partial_{\lambda_n} \chi(\varepsilon^{-1} \|(-\Delta_{\Omega})^{-1/2}(\varphi - \varphi^P)\|_2) \right|^2.$$

To bound B_1 , we use standard first-order perturbation theory for eigenvectors to compute

$$\partial_{\lambda_n} \psi_{\Pi\varphi} = -\frac{Q_{\varphi}}{H_{\varphi} - e(\varphi)} V_{\varphi_n} \psi_{\varphi}$$

where $Q_{\varphi} = 1 - |\psi_{\varphi}\rangle\langle\psi_{\varphi}|$. In particular,

$$(4.5) \quad \sum_{n=1}^N \|\partial_{\lambda_n} \psi_{\varphi}\|_2^2 \\ = 4 \operatorname{Tr} \Pi (-\Delta_{\Omega})^{-1/2} \psi_{\varphi} \left(\frac{Q_{\varphi}}{H_{\varphi} - e(\varphi)} \right)^2 \psi_{\varphi} (-\Delta_{\Omega})^{-1/2} \Pi$$

where we again interpret ψ_{φ} as a multiplication operator on the right side. It is not difficult to see that

$$(-\Delta_{\Omega})^{1/2} \frac{Q_{\varphi}}{H_{\varphi} - e(\varphi)} (-\Delta_{\Omega})^{1/2}$$

is uniformly bounded on the support of χ (compare with the proof of Proposition 3.3). Using this fact and (3.3), we see that (4.5) is uniformly bounded, independently of N . Hence $B_1 \lesssim \alpha^{-4} \langle \Psi | \Psi \rangle$.

For B_2 , we have

$$\begin{aligned} B_2 &\lesssim \frac{1}{\alpha^4 \varepsilon^2} \int_{\mathbb{R}^N} \prod_{m=1}^N d\lambda_m e^{-2\alpha^2 \langle \varphi - \varphi^P | (\mathbb{1} - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle} \\ &= \frac{1}{\alpha^4 \varepsilon^2} (\alpha \sqrt{2/\pi})^{-N} \det(\mathbb{1} - \Pi K \Pi)^{-1/4} \end{aligned}$$

where we have used the fact that φ^P is in the range of Π . We have to compare this with the norm of Ψ , which is bounded from below by

$$\begin{aligned} \langle \Psi | \Psi \rangle &\geq \int_{\mathbb{R}^N \setminus S_\varepsilon} \prod_{m=1}^N d\lambda_m e^{-\alpha^2 \langle \varphi - \varphi^P | (\mathbb{1} - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle} \\ &= (\alpha \sqrt{2/\pi})^{-N} \det(\mathbb{1} - \Pi K \Pi)^{-1/4} \\ &\quad - \int_{S_\varepsilon} \prod_{m=1}^N d\lambda_m e^{-2\alpha^2 \langle \varphi - \varphi^P | (\mathbb{1} - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle} \end{aligned}$$

where

$$S_\varepsilon = \{\vec{\lambda} \in \mathbb{R}^N : \|(-\Delta_\Omega)^{-1/2}(\varphi - \varphi^P)\|_2 \geq \varepsilon/2\}.$$

Since $\|K\| < 1$ by assumption, $(-\Delta_\Omega)^{-1} \leq \nu(1 - \Pi K \Pi)^{1/2}$ for some constant $\nu > 0$ independent of N . Hence we can bound the characteristic function of S_ε from above by

$$\exp\left(-\frac{1}{4\nu} \alpha^2 \varepsilon^2\right) \times \exp(\alpha^2 \langle \varphi - \varphi^P | (\mathbb{1} - \Pi K \Pi)^{1/2} | \varphi - \varphi^P \rangle).$$

Therefore,

$$\langle \Psi | \Psi \rangle \geq (\alpha \sqrt{2/\pi})^{-N} \det(\mathbb{1} - \Pi K \Pi)^{-1/4} (1 - 2^{N/2} e^{-\frac{1}{4\nu} \alpha^2 \varepsilon^2}).$$

In particular, as long as $\alpha \varepsilon \geq \text{const} \sqrt{N}$ with a sufficiently large constant, we have

$$\langle \Psi | \Psi \rangle \gtrsim (\alpha \sqrt{2/\pi})^{-N} \det(\mathbb{1} - \Pi K \Pi)^{-1/4},$$

and hence

$$B_2 \lesssim \alpha^{-4} \varepsilon^{-2} \langle \Psi | \Psi \rangle.$$

In summary, we have shown that

$$\frac{\langle \Psi | \mathbb{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} \leq e^P - \frac{1}{2\alpha^2} \text{Tr}[\mathbb{1} - (\mathbb{1} - \Pi K \Pi)^{1/2}] + \text{const}(\varepsilon^3 + \alpha^{-4} \varepsilon^{-2})$$

as long as $\alpha \varepsilon \geq \text{const} \sqrt{N}$ and ε is small enough. We shall choose Π to be the projection onto the span of $g_1, \dots, g_{N-1}, \varphi^P$, where we denote by $\{g_j\}_j$ an orthonormal basis of eigenfunctions of K , ordered in a way that the corresponding

eigenvalues $k_j = \langle g_j | K g_j \rangle$ form a decreasing sequence.¹ Then

$$\mathrm{Tr}[\mathbb{1} - (\mathbb{1} - \Pi K \Pi)^{1/2}] \geq \sum_{j=1}^{N-1} (1 - (1 - k_j)^{1/2})$$

and hence

$$\mathrm{Tr}[\mathbb{1} - (\mathbb{1} - \Pi K \Pi)^{1/2}] \geq \mathrm{Tr}[\mathbb{1} - (\mathbb{1} - K)^{1/2}] - \sum_{j=N}^{\infty} (1 - (1 - k_j)^{1/2}).$$

Since $K \lesssim [(-\Delta_{\Omega})^{-1/2} \psi^{\mathrm{P}} (-\Delta_{\Omega})^{-1/2}]^2$ and ψ^{P} is bounded by Lemma 3.1, we have $k_j \leq \mathrm{const} e_j^{-2}$, where e_j denotes the (ordered) eigenvalues of $-\Delta_{\Omega}$. Since Ω is assumed to be a smooth and bounded domain, we have the Weyl asymptotics $e_j \sim j^{2/3}$ for $j \gg 1$ (see, e.g., [28, sec. XIII.15]), which implies that

$$\sum_{j=N}^{\infty} (1 - (1 - k_j)^{1/2}) \lesssim N^{-1/3}.$$

In order to minimize the error term, we shall choose $\varepsilon \sim \alpha^{-8/11}$ and $N \sim \alpha^2 \varepsilon^2 \sim \alpha^{6/11}$, which leads to the bound

$$(4.6) \quad \frac{\langle \Psi | \mathbb{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} \leq e^{\mathrm{P}} - \frac{1}{2\alpha^2} \mathrm{Tr}[\mathbb{1} - (\mathbb{1} - K)^{1/2}] + \mathrm{const} \alpha^{-24/11}$$

for large enough α . This concludes the proof of the upper bound in Theorem 2.6. \square

5 Multiple Lieb-Yamazaki Bound

In [22] Lieb and Yamazaki used the fact that the interaction between the particle and the field can be written as a commutator, together with a Cauchy-Schwarz inequality, to get a uniform lower bound on the ground state energy of \mathbb{H} (for $\Omega = \mathbb{R}^3$) for large α . Their method shows that the introduction of an ultraviolet cutoff Λ in the interaction affects the ground state energy at most by $O(\Lambda^{-1/2})$. We shall apply their method three times, which will allow us to conclude that the effect of the cutoff is at most $O(\Lambda^{-5/2})$ (up to logarithmic corrections). It will be essential to use the Gross transformation explained in the next section, however, since we need relative operator boundedness of the kinetic energy with respect to the full Hamiltonian, which only holds for the transformed kinetic energy, as we shall see.

Before stating the main result of this section, we shall prove the following useful lemma. Its proof proceeds similarly to the one of lemma 10 in [9]. For its

¹ In case φ^{P} is in the span of $\{g_j\}_{j=1}^{N-1}$, we take Π to be the projection onto the span of $\{g_j\}_{j=1}^N$ instead.

statement, we introduce the Coulomb norm,

$$(5.1) \quad \|f\|_{\mathbb{C}} = \left(\frac{1}{4\pi} \int_{\mathbb{R}^6} \frac{\overline{f(x)}f(y)}{|x-y|} dx dy \right)^{1/2}.$$

By the Hardy-Littlewood-Sobolev inequality (see, e.g., [19, theorem 4.3]), this norm is dominated by the $L^{6/5}(\mathbb{R}^3)$ -norm.

Let us introduce the notation $p = -i\nabla_x = (p_1, p_2, p_3)$ for the momentum operator. We shall also use p^2 for the Dirichlet Laplacian $-\Delta_\Omega$ on Ω .

LEMMA 5.1. *Consider a function $h_x(\cdot)$ such that $k(x) = \sup_{y \in \mathbb{R}^3} |h_{x+y}(y)|$ has finite Coulomb norm. Then*

$$(5.2) \quad a^\dagger(h_x)a(h_x) \leq \|k\|_{\mathbb{C}}^2 p^2 \mathbb{N}$$

holds on $L^2(\Omega) \otimes \mathcal{F}$.

Note that the bound holds trivially with the right side replaced by $\|h_x\|_2^2 \mathbb{N}$. The point of Lemma 5.1 is that functions that are more singular (in the $x - y$ variable) can be handled, at the expense of the kinetic energy term p^2 .

PROOF. For convenience of notation, let Ψ be a one-phonon vector; the general case works in the same way. We need to bound

$$(5.3) \quad \int_{\Omega} \left| \int_{\Omega} \Psi(x, y) h_x(y) dy \right|^2 dx \leq \int_{\Omega} \left| \int_{\Omega} |\Psi(x, y)| k(x-y) dy \right|^2 dx.$$

With $\Phi(p, q)$ denoting the Fourier transform of $|\Psi(x, y)|$ (regarded as a function on $\mathbb{R}^3 \times \mathbb{R}^3$), we have

$$\begin{aligned} & \int_{\Omega} \left| \int_{\Omega} |\Psi(x, y)| k(x-y) dy \right|^2 dx \\ &= \int_{\mathbb{R}^3} \left| \int_{\mathbb{R}^3} \Phi(p-q, q) \widehat{k}(q) dq \right|^2 dp \\ &\leq \int_{\mathbb{R}^3} \left(\int_{\mathbb{R}^3} |\Phi(p-q, q)|^2 (p-q)^2 dq \right) \left(\int_{\mathbb{R}^3} |\widehat{k}(q)|^2 |p-q|^{-2} dq \right) dp \\ &\leq \sup_p \left(\int_{\mathbb{R}^3} |\widehat{k}(q)|^2 |p-q|^{-2} dq \right) \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |\Phi(p, q)|^2 p^2 dq dp. \end{aligned}$$

The last factor is smaller than $\|\sqrt{\mathbb{N}}\sqrt{p^2}\Psi\|^2$ (by the diamagnetic inequality). By writing the integral in x -space, one easily checks that

$$(5.4) \quad \sup_p \int_{\mathbb{R}^3} |\widehat{k}(q)|^2 |p-q|^{-2} dq \leq \|k\|_{\mathbb{C}}^2;$$

hence our claim (5.2) is proven. \square

The main result of this section is the following:

LEMMA 5.2. *Assume that $w_x(\cdot)$ is such that*

$$(5.5) \quad A_1 := \max_{j,k,l} \sup_{x \in \Omega} \|p_j p_k p_l |p|^{-6} w_x\|_2 < \infty,$$

$$(5.6) \quad A_2 := \max_{j,k} \sup_{x \in \Omega} \|p_j p_k |p|^{-4} w_x\|_2 < \infty,$$

and

$$(5.7) \quad A_3 := \max_{j,k} \|u_{jk}\|_C < \infty,$$

where $u_{jk}(x) = \sup_{y \in \mathbb{R}^3} |p_j p_k |p|^{-4} w_{x+y}(y)|$. Then

$$\begin{aligned} a(w_x) + a^\dagger(w_x) &\leq 12A_1(|p|^4 + 3p^2(\mathbb{N} + 1/(2\alpha^2))) \\ &\quad + 6(\alpha^{-1}A_2 + A_3)\left(|p|^4 + p^2\mathbb{N} + \frac{1}{2}\right) \end{aligned}$$

holds on $L^2(\Omega) \otimes \mathcal{F}$.

PROOF. For any w_x , we have

$$(5.8) \quad \sum_j [p_j, a(p_j |p|^{-2} w_x)] = -a(w_x).$$

Applying this three times, we also get

$$(5.9) \quad \sum_{j,k,l} [p_j, [p_k, [p_l, a(p_j p_k p_l |p|^{-6} w_x)]]] = -a(w_x).$$

In particular, we conclude that

$$(5.10) \quad \begin{aligned} &a(w_x) + a^\dagger(w_x) \\ &= \sum_{j,k,l} [p_j, [p_k, [p_l, a^\dagger(p_j p_k p_l |p|^{-6} w_x) - a(p_j p_k p_l |p|^{-6} w_x)]]]. \end{aligned}$$

We introduce the notation $B_{jkl} = a^\dagger(p_j p_k p_l |p|^{-6} w_x) - a(p_j p_k p_l |p|^{-6} w_x)$, and we rewrite the triple commutator as

$$(5.11) \quad \begin{aligned} \sum_{j,k,l} [p_j, [p_k, [p_l, B_{jkl}]]] &= \sum_{j,k,l} (p_j p_k [p_l, B_{jkl}] + [B_{jkl}^\dagger, p_l] p_j p_k) \\ &\quad - 2 \sum_{j,k,l} (p_j p_k B_{jkl} p_l + p_l B_{jkl}^\dagger p_j p_k) \end{aligned}$$

using the invariance of B_{jkl} under exchange of indices.

The Cauchy-Schwarz inequality implies that

$$(5.12) \quad -p_j p_k B_{jkl} p_l - p_l B_{jkl}^\dagger p_j p_k \leq \lambda p_j^2 p_k^2 + \lambda^{-1} p_l B_{jkl}^\dagger B_{jkl} p_l$$

for any $\lambda > 0$. Moreover,

$$(5.13) \quad B_{jkl}^\dagger B_{jkl} \leq (4\mathbb{N} + 2\alpha^{-2}) \|p_j p_k p_l |p|^{-6} w_x\|_2^2 \leq A_1^2 (4\mathbb{N} + 2\alpha^{-2}).$$

In particular, by choosing $\lambda = 2A_1$ and summing over j, k, l , we obtain the bound

$$(5.14) \quad \begin{aligned} & -2 \sum_{j,k,l} (p_j p_k B_{jkl} p_l + p_l B_{jkl}^\dagger p_j p_k) \\ & \leq 12A_1 (|p|^4 + 3p^2(\mathbb{N} + 1/(2\alpha^2))). \end{aligned}$$

We also have

$$(5.15) \quad C_{jk} = \sum_l [p_l, B_{jkl}] = a^\dagger (p_j p_k |p|^{-4} w_x) + a (p_j p_k |p|^{-4} w_x)$$

and

$$(5.16) \quad p_j p_k C_{jk} + C_{jk} p_j p_k \leq \lambda p_j^2 p_k^2 + \lambda^{-1} C_{jk}^2$$

for any $\lambda > 0$. Furthermore, we can bound

$$(5.17) \quad C_{jk}^2 \leq 4a^\dagger (p_j p_k |p|^{-4} w_x) a (p_j p_k |p|^{-4} w_x) + \frac{2}{\alpha^2} \|p_j p_k |p|^{-4} w_x\|_2^2.$$

By Lemma 5.1, the first term on the right side is bounded by $4\|u_{jk}\|_{\mathbb{C}}^2 p^2 \mathbb{N}$, and hence $C_{jk}^2 \leq 4A_3^2 p^2 \mathbb{N} + 2A_2^2 \alpha^{-2}$. The choice $\lambda = 6(A_3 + \alpha^{-1} A_2)$ then leads to the bound

$$(5.18) \quad \sum_{j,k} (p_j p_k C_{jk} + C_{jk} p_j p_k) \leq 6(A_3 + \alpha^{-1} A_2) \left(|p|^4 + p^2 \mathbb{N} + \frac{1}{2} \right).$$

In combination with (5.10), (5.11), and (5.14), this concludes the proof of the lemma. \square

In the following, we shall apply this bound to the large momentum part of the interaction in order to quantify the effect of an ultraviolet cutoff on the ground state energy. Because the Coulomb norm in (5.7) estimates the off-diagonal decay, we cannot use a sharp cutoff, however, and need to work with a smooth one instead. In fact, we shall apply Lemma 5.2 with

$$(5.19) \quad w_x(y) = z(-\Delta_\Omega)(x, y) \quad \text{for } z(t) = t^{-1/2} (1 - e^{-t/\Lambda^2})^2$$

for some $\Lambda > 0$. The function z is nonnegative and behaves like $t^{3/2} \Lambda^{-4}$ for $t \ll \Lambda^2$. Moreover, $z(t) - t^{-1/2}$ falls off like $t^{-1/2} e^{-t/\Lambda^2}$ for $t \gg \Lambda^2$.

We shall show in Appendix C that the various norms appearing in (5.5)–(5.7) can be bounded, up to a multiplicative constant, by the equivalent expressions for $\Omega = \mathbb{R}^3$, which can easily be estimated using Fourier transforms. We have

$$(5.20) \quad \|p_j p_k |p|^{-4} w_x\|_2^2 = \sum_n e_n^{-5} (1 - e^{-e_n/\Lambda^2})^4 |\partial_j \partial_k \varphi_n(x)|^2$$

where e_n and φ_n denote the eigenvalues and eigenfunctions of $-\Delta_\Omega$. In particular, from (C.10) we deduce that

$$(5.21) \quad \sup_{x \in \Omega} \max_{j,k} \|p_j p_k |p|^{-4} w_x\|_2 \lesssim \left(\int_{\mathbb{R}^3} |k|^{-6} (1 - e^{-k^2/\Lambda^2})^4 dk \right)^{1/2} \\ = \text{const } \Lambda^{-3/2}.$$

In the same way, we obtain the bound

$$(5.22) \quad \sup_{x \in \Omega} \max_{j,k,l} \|p_j p_k p_l |p|^{-6} w_x\|_2 \lesssim \Lambda^{-5/2}.$$

Moreover, in Section C.3 we shall show that

$$(5.23) \quad \max_{j,k} \|u_{jk}\|_C \lesssim \Lambda^{-5/2}.$$

We collect these results in the following corollary.

COROLLARY 5.3. *For $\Lambda > 0$ let $w_x(\cdot)$ be the function defined in (5.19). Then*

$$(5.24) \quad a(w_x) + a^\dagger(w_x) \lesssim (p^2 + \mathbb{N} + 1)^2 (\Lambda^{-5/2} + \alpha^{-1} \Lambda^{-3/2})$$

for $\alpha \gtrsim 1$.

6 Gross Transformation

In this section we shall investigate the effect of a unitary Gross transformation [14, 26] on the Hamiltonian (2.1). Let $\{f_x\}_{x \in \Omega} \subset L^2(\Omega)$ be a family of functions, parametrized by $x \in \Omega$, such that $\nabla_x f_x \in L^2(\Omega)$ for all $x \in \Omega$. We consider a unitary transformation in $L^2(\Omega) \otimes \mathcal{F}$ of the form

$$(6.1) \quad U = e^{a(\alpha^2 f_x) - a^\dagger(\alpha^2 f_x)}.$$

(This operator acts by ‘‘multiplication’’ with respect to the x -variable.) For $g \in L^2(\Omega)$ we have

$$(6.2) \quad Ua(g)U^\dagger = a(g) + \langle g | f_x \rangle \quad \text{and} \quad Ua^\dagger(g)U^\dagger = a^\dagger(g) + \langle f_x | g \rangle$$

and hence

$$(6.3) \quad U\mathbb{N}U^\dagger = \mathbb{N} + a^\dagger(f_x) + a(f_x) + \|f_x\|_2^2.$$

Moreover, for $p = -i\nabla_x$,

$$(6.4) \quad UpU^\dagger = p + \alpha^2(a^\dagger(pf_x) + a(pf_x) + \text{Re}\langle f_x | pf_x \rangle).$$

We shall choose f_x real-valued, hence the last term vanishes. Then

$$Up^2U^\dagger = p^2 + \alpha^4(a^\dagger(pf_x) + a(pf_x))^2 + 2\alpha^2 p \cdot a(pf_x) \\ + 2\alpha^2 a^\dagger(pf_x) \cdot p + \alpha^2 a(p^2 f_x) + \alpha^2 a^\dagger(p^2 f_x).$$

For the Hamiltonian (2.1), we thus have

$$(6.5) \quad \begin{aligned} U \mathbb{H} U^\dagger &= p^2 + \alpha^4 (a^\dagger(p f_x) + a(p f_x))^2 + 2\alpha^2 p \cdot a(p f_x) \\ &+ 2\alpha^2 a^\dagger(p f_x) \cdot p + \mathbb{N} + a(\alpha^2 p^2 f_x + f_x - v_x) \\ &+ a^\dagger(\alpha^2 p^2 f_x + f_x - v_x) + \|f_x\|_2^2 - 2 \operatorname{Re}\langle v_x | f_x \rangle. \end{aligned}$$

We shall choose f_x such that $\alpha^2 p^2 f_x + f_x - v_x = g_x$, i.e.,

$$(6.6) \quad f \cdot (y) = (-\alpha^2 \Delta_\Omega + 1)^{-1} (g \cdot (y) + v \cdot (y)) \quad \forall y \in \Omega$$

for some $g_x \in L^2(\Omega)$ with $\sup_{x \in \Omega} \|g_x\|_2 < \infty$. The choice $g_x \equiv 0$ would be possible, but it will be more convenient to choose

$$(6.7) \quad g_x(y) = \xi(-\Delta_\Omega)(x, y) \quad \text{for } \xi(t) = -t^{-1/2} \theta(K^2 - t)$$

for some $K > 0$, where

$$(6.8) \quad \theta(t) = \begin{cases} 0 & \text{for } t < 0, \\ 1/2 & \text{for } t = 0, \\ 1 & \text{for } t > 0. \end{cases}$$

Then

$$(6.9) \quad \|g_x\|_2^2 = \xi^2(-\Delta_\Omega)(x, x)$$

and, since $\xi(t)^2 \leq t^{-1} e^{1-t/K^2}$, the fact that the heat kernel of Δ_Ω is dominated by the one of $\Delta_{\mathbb{R}^3}$ implies as in (C.2) that

$$(6.10) \quad \sup_x \|g_x\|_2^2 \leq \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} \frac{e^{1-k^2/K^2}}{k^2} dk = \frac{e}{4\pi^{3/2}} K.$$

For the corresponding f_x , we have

$$(6.11) \quad f_x(y) = \eta(-\Delta_\Omega)(x, y) \quad \text{for } \eta(t) = -t^{-1/2} \frac{\theta(t - K^2)}{\alpha^2 t + 1}.$$

Using the fact that

$$(6.12) \quad \eta(t)^2 \leq \alpha^{-4} t^{-3} \theta(t - K^2) \leq \alpha^{-4} \left(\frac{2}{t + K^2} \right)^3,$$

one obtains in a similar way as above

$$(6.13) \quad \begin{aligned} \sup_x \|f_x\|_2^2 &= \sup_x \eta^2(-\Delta_\Omega)(x, x) \\ &\leq \frac{1}{\alpha^4 (2\pi)^3} \int_{\mathbb{R}^3} \left(\frac{2}{k^2 + K^2} \right)^3 dk = \frac{1}{4\pi} \alpha^{-4} K^{-3}. \end{aligned}$$

Moreover,

$$(6.14) \quad \sup_x |\langle v_x | f_x \rangle| \leq \frac{1}{\alpha^2 (2\pi)^3} \int_{\mathbb{R}^3} \left(\frac{2}{k^2 + K^2} \right)^2 dk = \frac{1}{2\pi} \alpha^{-2} K^{-1}$$

and, using (6.12) and (C.10),

$$(6.15) \quad \sup_x \|pf_x\|_2^2 \lesssim \frac{1}{\alpha^4} \int_{\mathbb{R}^3} k^2 \left(\frac{2}{k^2 + K^2} \right)^3 dk = 6\pi^2 \alpha^{-4} K^{-1}.$$

With the above choice of the function f_x (depending on α and the parameter K) we denote U by $U_{K,\alpha}$ from now on. With the aid of the previous estimates, we can now prove the following proposition. Its proof follows along similar lines as the corresponding argument for $\Omega = \mathbb{R}^3$ in [13].

PROPOSITION 6.1. *For any $\varepsilon > 0$ there are $K > 0$ and $C > 0$ such that for all $\alpha \gtrsim 1$ and any $\Psi \in L^2(\Omega) \otimes \mathcal{F}$ in the domain of $p^2 + \mathbb{N}$*

$$(6.16) \quad \begin{aligned} (1 + \varepsilon) \|(p^2 + \mathbb{N})\Psi\| + C \|\Psi\| &\geq \|U_{K,\alpha} \mathbb{H} U_{K,\alpha}^\dagger \Psi\| \\ &\geq (1 - \varepsilon) \|(p^2 + \mathbb{N})\Psi\| - C \|\Psi\|. \end{aligned}$$

We remark that due to the singular nature of v_x in the interaction term, it is essential to apply the unitary transformation $U_{K,\alpha}$. In its absence, the bound (6.16) fails to hold. In other words, the domain of \mathbb{H} does not coincide with the domain of $p^2 + \mathbb{N}$, but the one of $U_{K,\alpha} \mathbb{H} U_{K,\alpha}^\dagger$ does for a suitable choice of K .

PROOF. From (6.5) we see that the terms to estimate are the following:

$$(6.17) \quad \begin{aligned} \alpha^4 \|(a^\dagger(pf_x) + a(pf_x))^2 \Psi\| &\lesssim \alpha^4 \sup_x \|pf_x\|_2^2 \|(\mathbb{N} + \alpha^{-2})\Psi\| \\ &\lesssim K^{-1} \|(\mathbb{N} + \alpha^{-2})\Psi\| \end{aligned}$$

where we used (6.15),

$$(6.18) \quad \begin{aligned} \|(a(g_x) + a^\dagger(g_x))\Psi\| &\lesssim \sup_x \|g_x\|_2 \|(\sqrt{\mathbb{N} + \alpha^{-2}})\Psi\| \\ &\lesssim \delta \|(\mathbb{N} + \alpha^{-2})\Psi\| + \delta^{-1} K \|\Psi\| \end{aligned}$$

for any $\delta > 0$, using (6.10),

$$(6.19) \quad \begin{aligned} \alpha^2 \|a^\dagger(pf_x) \cdot p \Psi\| &\leq \alpha^2 \sup_x \|pf_x\|_2 \|\sqrt{\mathbb{N} + \alpha^{-2}} \sqrt{p^2} \Psi\| \\ &\lesssim K^{-1/2} \|(p^2 + \mathbb{N} + \alpha^{-2})\Psi\| \end{aligned}$$

and finally, the term

$$(6.20) \quad \alpha^2 p \cdot a(pf_x) = \alpha^2 a(pf_x) \cdot p + a(\alpha^2 p^2 f_x).$$

The first term on the right side of (6.20) can be estimated as in (6.19) above. For the second term, we write

$$(6.21) \quad \alpha^2 (p^2 f_x)(y) = h_x^{(1)}(y) + h_x^{(2)}(y)$$

where

$$(6.22) \quad h_x^{(1)}(y) = g_x(y) - f_x(y) + [(-\Delta_\Omega)^{-1/2} - (K^2 - \Delta_\Omega)^{-1/2}](x, y)$$

and

$$(6.23) \quad h_x^{(2)}(y) = (K^2 - \Delta_\Omega)^{-1/2}(x, y).$$

The L^2 -norms of g_x and f_x have already been bounded above, in (6.10) and (6.13), respectively. To bound the third function in $h_x^{(1)}$, we use

$$0 \leq t^{-1/2} - (K^2 + t)^{-1/2} \leq Kt^{-1/2}(K^2 + t)^{-1/2},$$

and find that the square of its L^2 -norm is bounded by

$$\frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} \frac{K^2}{k^2(k^2 + K^2)} dk = \frac{1}{4\pi} K.$$

By using the Schwarz inequality we conclude that

$$(6.24) \quad \|a(h_x^{(1)})\Psi\| \lesssim \delta \|\mathbb{N}\Psi\| + \delta^{-1} K(1 + (K\alpha)^{-4}) \|\Psi\|$$

for any $\delta > 0$.

The last term to bound is $a(h_x^{(2)})\Psi$. Since $|h_x^{(2)}(y)| \leq (K^2 - \Delta_{\mathbb{R}^3})^{-1/2}(x, y)$, Lemma 5.1 implies that

$$(6.25) \quad \|a(h_x^{(2)})\Psi\| \leq (2\pi)^{-3/2} \left(\int_{\mathbb{R}^3} (K^2 + q^2)^{-1} |q|^{-2} dq \right)^{1/2} \|\sqrt{\mathbb{N}}\sqrt{p^2}\Psi\|.$$

The prefactor on the right side is equal to a constant times $K^{-1/2}$. Moreover, we can bound $\|\sqrt{\mathbb{N}}\sqrt{p^2}\Psi\| \leq \frac{1}{2}\|(p^2 + \mathbb{N})\Psi\|$. In combination with (6.13) and (6.14), we hence arrive at the desired result, with $K \sim \varepsilon^{-2}$ and $C \sim \varepsilon^{-3}$. \square

From Proposition 6.1 we draw two important conclusions. First, the ground state energy of \mathbb{H} is uniformly bounded in α for large α . Second, in any state of bounded energy, in the sense that $\|\mathbb{H}\Psi\| \leq \text{const}$, both

$$\|U_{K,\alpha}^\dagger p^2 U_{K,\alpha} \Psi\| \quad \text{and} \quad \|U_{K,\alpha}^\dagger \mathbb{N} U_{K,\alpha} \Psi\|$$

are uniformly bounded (for suitable K independent of α). In particular, we conclude that in order to compute the ground state energy, it suffices to consider wave functions Ψ having this property.

We have, by a similar computation as in (6.4),

$$(6.26) \quad U_{K,\alpha}^\dagger p^2 U_{K,\alpha} = (p - A_{K,\alpha})^2 \quad \text{with} \quad A_{K,\alpha} = \alpha^2 (a^\dagger(p f_x) + a(p f_x))$$

and

$$(6.27) \quad U_{K,\alpha}^\dagger \mathbb{N} U_{K,\alpha} = \mathbb{N} - a(f_x) - a^\dagger(f_x) + \|f_x\|_2^2.$$

Since $\|f_x\|_2$ is uniformly bounded, as shown in (6.13) above, it easily follows that uniform boundedness of $\|U_{K,\alpha}^\dagger \mathbb{N} U_{K,\alpha} \Psi\|$ is equivalent to the one of $\|\mathbb{N}\Psi\|$.

7 Proof of Theorem 2.6: Lower Bound

7.1 Ultraviolet Cutoff

The first step in the lower bound is to introduce an ultraviolet cutoff in the interaction. Corollary 5.3 together with Proposition 6.1 will allow us to quantify its effect on the ground state energy.

PROPOSITION 7.1. *For $\Lambda > 0$, let*

$$(7.1) \quad \mathbb{H}^\Lambda = -\Delta_\Omega - a(v_x^\Lambda) - a^\dagger(v_x^\Lambda) + \mathbb{N}$$

where

$$(7.2) \quad v_x^\Lambda(y) = \frac{\theta(\Lambda^2 + \Delta_\Omega)}{(-\Delta_\Omega)^{1/2}}(x, y).$$

Then

$$(7.3) \quad \begin{aligned} & \inf \text{spec } \mathbb{H} - \inf \text{spec } \mathbb{H}^\Lambda \\ & \gtrsim -\Lambda^{-5/2}(\ln \Lambda)^{5/4} - \alpha^{-1} \Lambda^{-3/2}(\ln \Lambda)^{3/4} - \alpha^{-2} \Lambda^{-1}(\ln \Lambda)^{1/2} \end{aligned}$$

for $\alpha \gtrsim 1$ and $\Lambda \gtrsim 1$.

In order for the error introduced in (7.3) to be negligible compared to α^{-2} , it is sufficient to choose $\Lambda \sim \alpha^\kappa$ with $\kappa > 4/5$.

PROOF.

Step 1. Recall that $v_x(y) = (-\Delta_\Omega)^{-1/2}(x, y)$. We pick some $0 < \Lambda' < \Lambda$ and decompose v_x as $v_x(y) = u_x^{\Lambda'}(y) + w_x(y)$ where w_x is defined as in (5.19) above, but with Λ replaced by Λ' , i.e.,

$$(7.4) \quad w_x(y) = z(-\Delta_\Omega)(x, y) \quad \text{for } z(t) = t^{-1/2}(1 - e^{-t/\Lambda'^2}).$$

Corollary 5.3 states that

$$(7.5) \quad a(w_x) + a^\dagger(w_x) \lesssim (p^2 + \mathbb{N} + 1)^2 (\Lambda'^{-5/2} + \alpha^{-1} \Lambda'^{-3/2})$$

for $\alpha \gtrsim 1$. We now apply the unitary Gross transformation (6.1), with f_x given in (6.11), and K chosen such that Proposition 6.1 holds for some fixed $0 < \varepsilon < 1$, say $\varepsilon = 1/2$. We have

$$(7.6) \quad U_{K,\alpha}^\dagger a(w_x) U_{K,\alpha} = a(w_x) + \langle w_x | f_x \rangle,$$

and

$$(7.7) \quad \sup_{x \in \Omega} |\langle w_x | f_x \rangle| \lesssim \alpha^{-2} \Lambda'^{-1},$$

which can easily be seen by noting that $\langle w_x | f_x \rangle = (z\eta)(-\Delta_\Omega)(x, x)$ (with z and η defined in (7.4) and (6.11), respectively) and using that

$$|z(t)\eta(t)| \lesssim \alpha^{-2}(t + \Lambda^2)^{-2},$$

proceeding as in (C.2) to bound the expression in terms of the one for $\Omega = \mathbb{R}^3$. Proposition 6.1 thus implies that

$$(7.8) \quad a(w_x) + a^\dagger(w_x) \lesssim (\mathbb{H} + C)^2 (\Lambda'^{-5/2} + \alpha^{-1} \Lambda'^{-3/2} + \alpha^{-2} \Lambda'^{-1})$$

for a suitable constant $C > 0$ (independent of α for $\alpha \gtrsim 1$).

For computing the ground state energy, it is clearly sufficient to consider wave functions in the spectral subspace of \mathbb{H} corresponding to $|\mathbb{H}| \leq C$ for a suitable constant C . We thus conclude that

$$(7.9) \quad \inf \text{spec } \mathbb{H} \geq \inf \text{spec } \widetilde{\mathbb{H}}^{\Lambda'} - \text{const}(\Lambda'^{-5/2} + \alpha^{-1} \Lambda'^{-3/2} + \alpha^{-2} \Lambda'^{-1})$$

where $\widetilde{\mathbb{H}}^{\Lambda'}$ is obtained from \mathbb{H} by replacing v_x with $u_x^{\Lambda'} = v_x - w_x$, i.e.,

$$(7.10) \quad u_x^{\Lambda'}(y) = (-\Delta_\Omega)^{-1/2} (1 - (1 - e^{\Delta_\Omega/\Lambda'^2})^2)(x, y).$$

Step 2. We shall now further truncate $u_x^{\Lambda'}$ and replace it by

$$(7.11) \quad \widetilde{v}_x^\Lambda(y) = \frac{\theta(\Lambda^2 + \Delta_\Omega)}{(-\Delta_\Omega)^{1/2}} (1 - (1 - e^{\Delta_\Omega/\Lambda'^2})^2)(x, y).$$

With the aid of (C.3), one checks that

$$(7.12) \quad \sup_{x \in \Omega} \|u_x^{\Lambda'} - \widetilde{v}_x^\Lambda\|_2^2 \lesssim \Lambda e^{-(\Lambda/\Lambda')^2},$$

and hence, using the fact that $\sqrt{\mathbb{N}}$ is uniformly bounded for states with bounded energy, the error for introducing this additional cutoff is at most of the order $\Lambda^{1/2} e^{-(\Lambda/\Lambda')^2/2}$.

Step 3. Finally, we want to further simplify \widetilde{v}_x^Λ and replace it by v_x^Λ in (7.2). We claim that the ground state energy can only decrease under this replacement. This is the content of the following lemma.

LEMMA 7.2. *Let $\{\varphi_j\}_{j=1}^N$ be a set of orthonormal functions in $L^2(\Omega)$, and let*

$$(7.13) \quad u_x(y) = \sum_{j=1}^N \lambda_j \overline{\varphi_j(x)} \varphi_j(y) \quad \text{for } \lambda_j \geq 0, 1 \leq j \leq N.$$

Then

$$(7.14) \quad e(\lambda_1, \dots, \lambda_N) = \inf \text{spec}[-\Delta_\Omega - a(u_x) - a^\dagger(u_x) + \mathbb{N}]$$

is decreasing in each λ_j .

PROOF. We shall use a Perron-Frobenius-type argument. Let $\Psi \in L^2(\Omega) \otimes \mathcal{F}$ be given by $\{\psi_0(x), \psi_1(x, y_1), \psi_2(x, y_1, y_2), \dots\}$. We extend $\{\varphi_j\}_{j=1}^N$ to an

orthonormal basis $\{\varphi_j\}_{j \in \mathbb{N}}$ of $L^2(\Omega)$, and define $a_{i_1, \dots, i_n}^n(x)$ by the expansion $\psi_n(x, y_1, \dots, y_n) = \sum_{i_1, \dots, i_n} a_{i_1, \dots, i_n}^n(x) \varphi_{i_1}(y_1) \cdots \varphi_{i_n}(y_n)$. Then

$$\begin{aligned} & \langle \Psi | -\Delta_\Omega + \mathbb{N} | \Psi \rangle \\ &= \sum_{n \geq 0} \sum_{i_1, \dots, i_n} \left(\int_\Omega |\nabla_x a_{i_1, \dots, i_n}^n(x)|^2 dx + n \int_\Omega |a_{i_1, \dots, i_n}^n(x)|^2 dx \right) \end{aligned}$$

and

$$\begin{aligned} & \langle \Psi | a(u_x) + a^\dagger(u_x) | \Psi \rangle \\ &= 2 \sum_{j=1}^N \lambda_j \sum_{n \geq 0} \sqrt{n+1} \sum_{i_1, \dots, i_n} \Re \int_\Omega \overline{a_{i_1, \dots, i_n}^n(x)} a_{i_1, \dots, i_n, j}^{n+1}(x) \varphi_j(x) dx. \end{aligned}$$

By multiplying the functions a_{i_1, \dots, i_n}^n with an appropriate phase factor, we can make sure that

$$(7.15) \quad \int_\Omega \overline{a_{i_1, \dots, i_n}^n(x)} a_{i_1, \dots, i_n, j}^{n+1}(x) \varphi_j(x) dx \geq 0$$

for all $n \geq 0$, $1 \leq j \leq N$, and all i_1, \dots, i_n , and this can clearly only decrease the energy. When computing the ground state energy, it suffices to consider Ψ 's with such property, in which case the energy is clearly monotone decreasing in all the λ_j . \square

As a consequence, the ground state energy with interaction \tilde{v}_x^Λ is bounded below by the one with interaction v_x^Λ . In particular, we have thus shown that

$$(7.16) \quad \begin{aligned} \inf \text{spec } \mathbb{H} &\geq \inf \text{spec } \mathbb{H}^\Lambda - \text{const}(\Lambda'^{-5/2} + \alpha^{-1} \Lambda'^{-3/2} \\ &\quad + \alpha^{-2} \Lambda'^{-1} + \Lambda^{1/2} e^{-(\Lambda/\Lambda')^2/2}) \end{aligned}$$

and this holds for all $\alpha \gtrsim 1$ and $\Lambda' \gtrsim 1$. The choice $\Lambda' = \Lambda(6 \ln \Lambda)^{-1/2}$ yields (7.3). \square

7.2 Final Lower Bound

The starting point of the proof of the lower bound is Proposition 7.1, which quantifies the error in replacing \mathbb{H} by \mathbb{H}^Λ in (7.1) for computing the ground state energy. We are thus left with giving a lower bound on $\inf \text{spec } \mathbb{H}^\Lambda$.

We choose, for simplicity, Λ in such a way that Λ^2 is not an eigenvalue of $-\Delta_\Omega$. Let Π denote the projection

$$(7.17) \quad \Pi = \theta(\Lambda^2 + \Delta_\Omega) \quad \text{and} \quad N = \dim \text{ran } \Pi.$$

For later purposes we note that one has the Weyl asymptotics

$$(7.18) \quad N \sim (2\pi)^{-3} |\Omega| \Lambda^3 \quad \text{as } \Lambda \rightarrow \infty$$

(see, e.g., [28, sec. XIII.15]). If e_n and φ_n , respectively, denote the eigenvalues and (real-valued) eigenfunctions of $-\Delta_\Omega$, then

$$(7.19) \quad v_x^\Lambda(y) = \sum_{n=1}^N \frac{1}{\sqrt{e_n}} \varphi_n(x) \varphi_n(y)$$

has finite rank. The Fock space $\mathcal{F}(L^2(\Omega))$ naturally factors into a tensor product $\mathcal{F}(\Pi L^2(\Omega)) \otimes \mathcal{F}((1-\Pi)L^2(\Omega))$, and \mathbb{H}^Λ is of the form $\mathbb{A} \otimes \mathbb{1} + \mathbb{1} \otimes \mathbb{N}^>$, where \mathbb{A} acts on $L^2(\Omega) \otimes \mathcal{F}(\Pi L^2(\Omega))$ and $\mathbb{N}^> = \sum_{n>N} a^\dagger(\varphi_n) a(\varphi_n)$ is the number operator on $\mathcal{F}((1-\Pi)L^2(\Omega))$. In particular, $\inf \text{spec } \mathbb{H}^\Lambda = \inf \text{spec } \mathbb{A}$.

As in Section 4 (where a different basis was used, however), we identify the spaces $\mathcal{F}(\Pi L^2(\Omega))$ and $L^2(\mathbb{R}^N)$ via the representation

$$(7.20) \quad \varphi = \Pi \varphi = \sum_{n=1}^N \lambda_n \varphi_n,$$

thus identifying a function $\varphi \in \text{ran } \Pi$ with a point $(\lambda_1, \dots, \lambda_N) \in \mathbb{R}^N$. In this representation, we have

$$(7.21) \quad \mathbb{A} = -\Delta_\Omega + V_\varphi(x) + \sum_{n=1}^N \left(-\frac{1}{4\alpha^4} \partial_{\lambda_n}^2 + \lambda_n^2 - \frac{1}{2\alpha^2} \right)$$

on $L^2(\Omega) \otimes L^2(\mathbb{R}^N)$. For a lower bound, we can replace $-\Delta_\Omega + V_\varphi(x)$ by the infimum of its spectrum, for any fixed $\varphi \in \text{ran } \Pi$. In particular, we have

$$(7.22) \quad \inf \text{spec } \mathbb{H}^\Lambda \geq \inf \text{spec } \mathbb{K}$$

where \mathbb{K} is the operator on $L^2(\mathbb{R}^N)$

$$(7.23) \quad \mathbb{K} = -\frac{1}{4\alpha^4} \sum_{n=1}^N \partial_{\lambda_n}^2 - \frac{N}{2\alpha^2} + \mathcal{F}^P(\varphi)$$

with \mathcal{F}^P defined in (2.5). Here $\mathcal{F}^P(\varphi)$ is a function of $(\lambda_1, \dots, \lambda_N)$ via the identification (7.20).

We now introduce an IMS-type localization. Let $\chi : \mathbb{R}_+ \rightarrow [0, 1]$ be a smooth function with $\chi(t) = 1$ for $t \leq 1/2$, $\chi(t) = 0$ for $t \geq 1$. Let $\varepsilon > 0$, and let j_1 and j_2 denote the multiplication operators in $L^2(\mathbb{R}^N)$

$$(7.24) \quad \begin{aligned} j_1 &= \chi(\varepsilon^{-1} \|(-\Delta_\Omega)^{-1/2}(\varphi - \varphi^P)\|_2), \\ j_2 &= \sqrt{1 - \chi(\varepsilon^{-1} \|(-\Delta_\Omega)^{-1/2}(\varphi - \varphi^P)\|_2)^2}. \end{aligned}$$

Then clearly $j_1^2 + j_2^2 = 1$ and

$$(7.25) \quad \mathbb{K} = j_1 \mathbb{K} j_1 + j_2 \mathbb{K} j_2 - \mathbb{E}$$

where \mathbb{E} is the IMS localization error

$$(7.26) \quad \mathbb{E} = \frac{1}{4\alpha^4} \sum_{n=1}^N (|\partial_{\lambda_n} j_1|^2 + |\partial_{\lambda_n} j_2|^2).$$

It is easy to see that that $\mathbb{E} \lesssim \alpha^{-4} \varepsilon^{-2}$, independently of N . In particular, the localization error is negligible if $\varepsilon \gg \alpha^{-1}$.

On the support of j_1 , we can use the bound (3.5) on \mathcal{F}^P . This gives

$$(7.27) \quad j_1 \mathbb{K} j_1 \geq j_1^2 \inf \text{spec} \left(e^P - \frac{1}{4\alpha^4} \sum_{n=1}^N \partial_{\lambda_n}^2 - \frac{N}{2\alpha^2} + \langle \varphi - \varphi^P | \mathbb{1} - K - \varepsilon CL | \varphi - \varphi^P \rangle \right)$$

for C a positive constant. Now φ^P will not necessarily be in the range of Π . However, since $\mathbb{1} - K - \varepsilon CL$ is positive for ε small enough, we can replace φ^P by its closest point (in the norm defined via $\mathbb{1} - K - \varepsilon CL$) in the range of Π for a lower bound. That is,

$$(7.28) \quad \langle \varphi - \varphi^P | \mathbb{1} - K - \varepsilon CL | \varphi - \varphi^P \rangle \geq \langle \varphi - y | \Pi(\mathbb{1} - K - \varepsilon CL) \Pi | \varphi - y \rangle$$

where $y = (\Pi(\mathbb{1} - K - \varepsilon CL) \Pi)^{-1} \Pi(\mathbb{1} - K - \varepsilon CL) \varphi^P$. The shift by y can be removed by a unitary transformation, without affecting the ground state energy. Hence

$$\begin{aligned} j_1 \mathbb{K} j_1 &\geq j_1^2 \inf \text{spec} \left(e^P - \frac{1}{4\alpha^4} \sum_{n=1}^N \partial_{\lambda_n}^2 - \frac{N}{2\alpha^2} + \langle \varphi | \Pi(\mathbb{1} - K - \varepsilon CL) \Pi | \varphi \rangle \right) \\ &= j_1^2 \left[e^P - \frac{1}{2\alpha^2} \text{Tr}(\mathbb{1} - \sqrt{\mathbb{1} - \Pi(K + \varepsilon CL) \Pi}) \right]. \end{aligned}$$

This is of the correct form if $N \rightarrow \infty$ and $\varepsilon \rightarrow 0$ as $\alpha \rightarrow \infty$.

On the support of j_2 , we use the bound (3.15) instead. We have, for any $\eta \geq 0$,

$$\begin{aligned} j_2 \mathbb{K} j_2 &\geq j_2^2 \inf \text{spec} \left(e^P - \frac{1}{4\alpha^4} \sum_{n=1}^N \partial_{\lambda_n}^2 - \frac{N}{2\alpha^2} + \frac{\eta}{4} \varepsilon^2 \right. \\ &\quad \left. + \langle \varphi - \varphi^P | \mathbb{1} - (\mathbb{1} + \kappa'(-\Delta_\Omega)^{1/2})^{-1} - \eta(-\Delta_\Omega)^{-1} | \varphi - \varphi^P \rangle \right) \end{aligned}$$

where we have used the fact that $\|(-\Delta_\Omega)^{-1/2}(\varphi - \varphi^P)\|_2 \geq \varepsilon/2$ on the support of j_2 . We choose η independent of α (and hence also independent of Λ and ε) and small enough such that the operator in the last line is positive. Proceeding as in the case of j_1 above, we obtain

$$(7.29) \quad j_2 \mathbb{K} j_2 \geq j_2^2 \left(e^P + \frac{\eta}{4} \varepsilon^2 - \frac{1}{2\alpha^2} \text{Tr} \Pi \left[\mathbb{1} - \sqrt{\mathbb{1} - (\mathbb{1} + \kappa'(-\Delta_\Omega)^{1/2})^{-1} - \eta(-\Delta_\Omega)^{-1}} \right] \right).$$

From the Weyl asymptotics (7.18) one checks that the trace diverges like $N^{2/3} \sim \Lambda^2$ for large Λ . Hence if we choose $\Lambda\alpha^{-1} \leq \text{const } \varepsilon$ with a sufficiently small constant, the term in parenthesis in (7.29) is actually larger than e^P . Since we will choose $\Lambda \sim \alpha^\kappa$ with $\kappa > 4/5$, this is compatible with the condition $\varepsilon \ll 1$ as long as $\kappa < 1$.

We thus conclude that if $\Lambda\alpha^{-1} \leq \text{const } \varepsilon$ and ε is small enough, we have the bound

$$(7.30) \quad \inf \text{spec } \mathbb{K} \geq e^P - \frac{1}{2\alpha^2} \text{Tr}(\mathbb{1} - \sqrt{\mathbb{1} - \Pi(K + \varepsilon CL)\Pi}) - \text{const } \alpha^{-4} \varepsilon^{-2}.$$

For a lower bound, we can further drop the Π 's in the second term on the right side, and replace them by $\mathbb{1}$. Note that $\|K + \varepsilon CL\| \leq \nu < 1$ for small enough ε , and the function $f(t) = 1 - \sqrt{1-t}$ is Lipschitz-continuous and convex on $[0, \nu]$. We utilize the following simple lemma.

LEMMA 7.3. *For $\nu > 0$, let $f : [0, \nu] \rightarrow \mathbb{R}$ be a Lipschitz-continuous and convex function with $f(0) = 0$, and let A, B be nonnegative trace class operators with $A + B \leq \nu$. Then*

$$(7.31) \quad \text{Tr } f(A + B) \leq \text{Tr } f(A) + C_f \text{Tr } B$$

where C_f denotes the Lipschitz constant of f .

PROOF. With $\{g_j\}$ a basis of eigenvectors of $A + B$, we have

$$\begin{aligned} \text{Tr } f(A + B) &= \sum_j f(\langle g_j | A + B | g_j \rangle) \\ &\leq \sum_j f(\langle g_j | A | g_j \rangle) + C_f \sum_j \langle g_j | B | g_j \rangle. \end{aligned}$$

The convexity of f implies that $f(\langle g_j | A | g_j \rangle) \leq \langle g_j | f(A) | g_j \rangle$, which yields the desired result. \square

Lemma 7.3 readily implies that

$$(7.32) \quad \text{Tr}(\mathbb{1} - \sqrt{\mathbb{1} - K - \varepsilon CL}) \leq \text{Tr}(\mathbb{1} - \sqrt{\mathbb{1} - K}) + \text{const } \varepsilon \text{Tr } L.$$

We thus have

$$(7.33) \quad \inf \text{spec } \mathbb{K} \geq e^P - \frac{1}{2\alpha^2} \text{Tr}(\mathbb{1} - \sqrt{\mathbb{1} - K}) - \text{const}(\alpha^{-4} \varepsilon^{-2} + \alpha^{-2} \varepsilon).$$

In combination with (7.3) and (7.22), this is our final lower bound.

In order to minimize the error terms in (7.3) and (7.33), we shall choose $\varepsilon \sim \alpha^{-1/7} (\ln \alpha)^{5/14}$ and $\Lambda \sim \alpha^{6/7} (\ln \alpha)^{5/14}$. This yields

$$(7.34) \quad \inf \text{spec } \mathbb{H} \geq e^P - \frac{1}{2\alpha^2} \text{Tr}(\mathbb{1} - \sqrt{\mathbb{1} - K}) - \text{const } \alpha^{-15/7} (\ln \alpha)^{5/14}$$

for $\alpha \gtrsim 1$, and thus completes the proof of the lower bound in Theorem 2.6. \square

Appendix A Equivalent Formulation of Assumption 2.4

In this appendix we shall explain how Assumption 2.4 can be verified via a spectral analysis of the Hessian of \mathcal{E}^P at its minimizer $\psi^P \geq 0$, which is assumed to be unique. We partly follow ideas in [7, sec. 2].

The Euler-Lagrange equation for the minimizer is

$$(A.1) \quad -\Delta_\Omega \psi^P - 2((-\Delta_\Omega)^{-1} |\psi^P|^2) \psi^P = \mu \psi^P.$$

The relevant Hessian Z^P is defined via

$$\langle \psi | Z^P | \psi \rangle = \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2} \left(\mathcal{E}^P \left(\frac{\psi^P + \varepsilon \psi}{\|\psi^P + \varepsilon \psi\|_2} \right) - e^P \right)$$

for real-valued $\psi \in H_0^1(\Omega)$, and equals

$$\begin{aligned} Z^P &= -\Delta_\Omega - 2(-\Delta_\Omega)^{-1} |\psi^P|^2 - 4X - \mu \\ &\quad - 4 \iint_{\Omega \times \Omega} |\psi^P(x)|^2 (-\Delta_\Omega)^{-1}(x, y) |\psi^P(y)|^2 dx dy |\psi^P\rangle \langle \psi^P| \\ &\quad + 4(|\psi^P\rangle \langle (-\Delta_\Omega)^{-1} |\psi^P|^2 \rangle \psi^P | + \text{h.c.}) \end{aligned}$$

where X is the operator with integral kernel

$$X(x, y) = \psi^P(x) (-\Delta_\Omega)^{-1}(x, y) \psi^P(y).$$

There is also another Hessian defined for purely imaginary perturbations of ψ^P , but it is trivially given by the linear operator defined by the equation (A.1) and plays no role here.

Note that $Z^P \psi^P = 0$. We now show that if ψ^P spans the kernel of Z^P , then Assumption 2.4 holds.

LEMMA A.1. *If $\ker Z^P = \text{span}\{\psi^P\}$, then there exists a $\kappa > 0$ such that for all $0 \leq \psi \in H_0^1(\Omega)$ with $\|\psi\|_2 = 1$ we have*

$$(A.2) \quad \mathcal{E}^P(\psi) \geq \mathcal{E}^P(\psi^P) + \kappa \|\psi - \psi^P\|_{H^1(\Omega)}^2.$$

PROOF.

Step 1. We first show that there are $c > 0$ and $\kappa > 0$ such that (A.2) holds for all $0 \leq \psi \in H_0^1(\Omega)$ with $\|\psi\|_2 = 1$ and $\|\psi - \psi^P\|_{H^1(\Omega)} \leq c$. We set $\delta = \psi - \psi^P$ and expand

$$\begin{aligned} \mathcal{E}^P(\psi^P + \delta) &= \mathcal{E}^P(\psi^P) + 2\mu \int_{\Omega} \psi^P(x) \delta(x) dx \\ &\quad + \int_{\Omega} |\nabla \delta(x)|^2 dx - 2 \iint_{\Omega \times \Omega} \psi^P(x)^2 (-\Delta_\Omega)^{-1}(x, y) \delta(y)^2 dx dy \\ &\quad - 4\langle \delta | X | \delta \rangle + O(\|\delta\|_{H^1}^3). \end{aligned}$$

The assumption $\|\psi\|_2 = 1$ implies that

$$(A.3) \quad 2 \int_{\Omega} \psi^P(x) \delta(x) dx = -\|\delta\|_2^2,$$

and therefore, using this identity multiple times,

$$(A.4) \quad \mathcal{E}^P(\psi^P + \delta) = \mathcal{E}^P(\psi^P) + \langle \delta | Z^P | \delta \rangle + O(\|\delta\|_{H^1}^3).$$

The operator Z^P has discrete spectrum, and hence our assumption on the simplicity of the kernel implies that for some $\kappa > 0$

$$\begin{aligned} \langle \delta | Z^P | \delta \rangle &\geq \kappa \|\delta - \langle \psi^P | \delta \rangle \psi^P\|_2^2 \\ &= \kappa \left(\|\delta\|_2^2 - \left(\int_{\Omega} \psi^P \delta \right)^2 \right) = \kappa \|\delta\|_2^2 (1 - 4^{-1} \|\delta\|_2^2). \end{aligned}$$

On the other hand, it is easy to see that for some $C > 0$

$$(A.5) \quad Z^P \geq -(1/2)\Delta_{\Omega} - C.$$

Taking a mean of the previous two inequalities we obtain for any $0 \leq \theta \leq 1$,

$$(A.6) \quad \langle \delta | Z^P | \delta \rangle \geq (\theta/2) \|\nabla \delta\|_2^2 + ((1-\theta)\kappa - C\theta) \|\delta\|_2^2 - 4^{-1} \kappa (1-\theta) \|\delta\|_2^4.$$

In particular, for $\theta = \kappa / (C + \kappa + 1/2)$ we have

$$(A.7) \quad \langle \delta | Z^P | \delta \rangle \geq \frac{\kappa}{2C + 2\kappa + 1} \|\delta\|_{H^1(\Omega)}^2 - 4^{-1} \kappa \frac{2C + 1}{2C + 2\kappa + 1} \|\delta\|_2^4.$$

Inserting this into the above inequality, we obtain

$$(A.8) \quad \mathcal{E}^P(\psi^P + \delta) \geq \mathcal{E}^P(\psi^P) + \frac{\kappa}{2C + 2\kappa + 1} \|\delta\|_{H^1(\Omega)}^2 + O(\|\delta\|_{H^1(\Omega)}^3),$$

which clearly implies the assertion in Step 1.

Step 2. We now prove the full statement of the lemma. We argue by contradiction. If there were no such κ , we could find a sequence $0 \leq \psi_n \in H_0^1(\Omega)$ with $\|\psi_n\|_2 = 1$ such that

$$(A.9) \quad \mathcal{E}^P(\psi_n) < \mathcal{E}^P(\psi^P) + n^{-1} \|\psi_n - \psi^P\|_{H^1(\Omega)}^2.$$

Using (C.2), Hardy-Littlewood-Sobolev, Hölder, and Sobolev we bound

$$\begin{aligned} &\iint_{\Omega \times \Omega} \psi(x)^2 (-\Delta_{\Omega})^{-1}(x, y) \psi(y)^2 dx dy \\ &\leq \frac{1}{4\pi} \iint_{\Omega \times \Omega} \frac{\psi(x)^2 \psi(y)^2}{|x - y|} dx dy \\ &\lesssim \|\psi^2\|_{6/5}^2 \leq \|\psi\|_6 \|\psi\|_2^3 \lesssim \|\nabla \psi\|_2 \|\psi\|_2^3. \end{aligned}$$

This implies $\mathcal{E}^P(\psi) \geq (1/2) \|\nabla \psi\|_2^2 - C \|\psi\|_2^6$ for all $\psi \in H_0^1(\Omega)$. Combining this inequality with the upper bound (A.9) on $\mathcal{E}^P(\psi_n)$ we easily infer that (ψ_n) is

bounded in $H_0^1(\Omega)$ and hence that $\|\psi_n - \psi^P\|_{H^1(\Omega)}$ is bounded. Thus, (A.9) implies that (ψ_n) is a minimizing sequence for \mathcal{E}^P . Therefore, by a simple compactness argument, after passing to a subsequence, ψ_n converges in H^1 to a minimizer. Since $\psi_n \geq 0$, our assumed uniqueness of the minimizer implies that $\psi_n \rightarrow \psi^P$. Thus, for all sufficiently large n , $\|\psi_n - \psi^P\|_{H^1(\Omega)} \leq c$, where c is the constant from Step 1. Therefore the inequality from Step 1 is applicable, but this bound contradicts (A.9) for large n . This completes the proof. \square

Appendix B Bounds on Solutions of Poisson's Equation

We consider solutions u of the equation $-\Delta u = f$ in an open set $\Omega \subset \mathbb{R}^d$ with boundary conditions $u = 0$ on $\partial\Omega$. We are interested in bounds on derivatives of u in terms of derivatives of f , uniformly on small balls, possibly intersecting the boundary of Ω . While we use these bounds only for $d = 3$, it requires no extra effort to prove them in arbitrary dimension $d \geq 2$.

B.1 Statement of the Inequality

Let $k \in \mathbb{N}$ and $\delta \in (0, 1)$. We say that an open set $\Omega \subset \mathbb{R}^d$ is a $C^{k,\delta}$ set if there are constants $r_0 > 0$ and $M < \infty$ such that for any $x \in \partial\Omega$ there is a function $\Gamma : \{y' \in \mathbb{R}^{d-1} : |y'| < r_0\} \rightarrow \mathbb{R}$ satisfying $\Gamma(0) = 0$, $\nabla\Gamma(0) = 0$, and

$$(B.1) \quad \sum_{j=0}^k r_0^{j-1} \sup_{|y'| < r_0} |\partial^j \Gamma(y')| + r_0^{k-1+\delta} \sup_{|y'|, |z'| < r_0} \frac{|\partial^k \Gamma(y') - \partial^k \Gamma(z')|}{|y' - z'|^\delta} \leq M$$

such that, after a translation and a rotation (which maps x to 0 and the exterior unit normal at x to $(0, \dots, 0, -1)$, and is denoted by \mathcal{T}_x),

$$(B.2) \quad \mathcal{T}_x(\Omega \cap B_{r_0}(x)) = \{(y', y_d) \in \mathbb{R}^{d-1} \times \mathbb{R} : |y'| < r_0, y_d > \Gamma(y')\} \cap B_{r_0}(0).$$

Here and below we use the notation $|\partial^k f(x)| = (\sum_{|\beta|=k} |\partial^\beta f(x)|^2)^{1/2}$ and similarly $|\partial^k f(x) - \partial^k f(y)| = (\sum_{|\beta|=k} |\partial^\beta f(x) - \partial^\beta f(y)|^2)^{1/2}$, with $\partial^\beta = \partial_1^{\beta_1} \dots \partial_d^{\beta_d}$ for $\beta \in \mathbb{N}_0^d$, and $|\beta| = \sum_{j=1}^d \beta_j$. The above definition of a $C^{k,\delta}$ set is standard (see, e.g., [12, sec. 6.2]), except possibly for the choice of the r_0 dependence in (B.1). Our choice ensures scale invariance in the sense that if Ω is scaled by a factor λ , r_0 gets multiplied by λ while M stays the same.

THEOREM B.1. *Let $k \in \mathbb{N}$, $0 < \delta < 1$, $R_0 > 0$, and $\Omega \subset \mathbb{R}^d$ be an open $C^{k,\delta}$ set. Then we have, for all $a \in \Omega$ and all $R \leq R_0$, if $k = 1$*

$$\begin{aligned} & \sum_{j=0}^1 R^j \sup_{B_R(a) \cap \Omega} |\partial^j u| + R^{1+\delta} \sup_{x, y \in B_R(a) \cap \Omega} \frac{|\partial u(x) - \partial u(y)|}{|x - y|^\delta} \\ & \lesssim \sup_{B_{2R}(a) \cap \Omega} |u| + R^2 \sup_{B_{2R}(a) \cap \Omega} |f| \end{aligned}$$

and if $k \geq 2$

$$\begin{aligned} & \sum_{j=0}^k R^j \sup_{B_R(a) \cap \Omega} |\partial^j u| + R^{k+\delta} \sup_{x,y \in B_R(a) \cap \Omega} \frac{|\partial^k u(x) - \partial^k u(y)|}{|x-y|^\delta} \\ & \lesssim \sup_{B_{2R}(a) \cap \Omega} |u| + \sum_{j=0}^{k-2} R^{j+2} \sup_{B_{2R}(a) \cap \Omega} |\partial^j f| \\ & \quad + R^{k+\delta} \sup_{x,y \in B_{2R}(a) \cap \Omega} \frac{|\partial^{k-2} f(x) - \partial^{k-2} f(y)|}{|x-y|^\delta}. \end{aligned}$$

The constants in these bounds depend only on d, k, δ, M , and R_0/r_0 .

Dropping the Hölder seminorm on the left side and estimating it on the right side in terms of one higher derivative, we obtain:

COROLLARY B.2. *Let $k \in \mathbb{N}$, $0 < \delta < 1$, $R_0 > 0$, and $\Omega \subset \mathbb{R}^d$ be an open $C^{k,\delta}$ set. Then we have for all $a \in \Omega$ and all $R \leq R_0$,*

$$(B.3) \quad \sum_{j=0}^k R^j \sup_{B_R(a) \cap \Omega} |\partial^j u| \lesssim \sup_{B_{2R}(a) \cap \Omega} |u| + \sum_{j=0}^{k-1} R^{j+2} \sup_{B_{2R}(a) \cap \Omega} |\partial^j f|.$$

The constants in these bounds depend only on d, k, δ, M , and R_0/r_0 .

B.2 Local Estimates

The more difficult assertion in Theorem B.1 is for balls such that $B_{2R}(a) \cap \partial\Omega \neq \emptyset$. The strategy in this case will be to flatten the boundary, but this results in a second-order elliptic equation with variable coefficients. In this subsection we state and prove bounds on solutions of such equations for domains with a flat boundary portion.

Let $\Omega \subset \mathbb{R}_+^d := \mathbb{R}^{d-1} \times (0, \infty)$ be an open set with an open boundary portion T on $\partial\mathbb{R}_+^d$. We emphasize explicitly that the case $T = \emptyset$ is allowed. For $x, y \in \Omega$ we write, following [12, sec. 4.4],

$$\bar{d}_x := \text{dist}(x, \partial\Omega \setminus T), \quad \bar{d}_{x,y} := \min\{\bar{d}_x, \bar{d}_y\},$$

and introduce the norms

$$|u|_{k,\Omega \cup T}^{(\sigma)} := \sum_{j=0}^k \sup_{x \in \Omega} \bar{d}_x^{j+\sigma} |\partial^j u(x)|$$

and

$$(B.4) \quad \begin{aligned} |u|_{k,\delta,\Omega \cup T}^{(\sigma)} &:= \sum_{j=0}^k \sup_{x \in \Omega} \bar{d}_x^{j+\sigma} |\partial^j u(x)| \\ &+ \sup_{x,y \in \Omega} \bar{d}_{x,y}^{k+\delta+\sigma} \frac{|\partial^k u(x) - \partial^k u(y)|}{|x-y|^\delta}. \end{aligned}$$

One readily checks that these norms satisfy $|fg|_{k,\delta,\Omega \cup T}^{(\sigma_1+\sigma_2)} \lesssim |f|_{k,\delta,\Omega \cup T}^{(\sigma_1)} |g|_{k,\delta,\Omega \cup T}^{(\sigma_2)}$ as well as $|\partial f|_{k,\delta,\Omega \cup T}^{(\sigma)} \lesssim |f|_{k+1,\delta,\Omega \cup T}^{(\sigma-1)}$ and $|f|_{k,\delta,\Omega \cup T}^{(\sigma)} \lesssim |f|_{k+1,\delta,\Omega \cup T}^{(\sigma)}$ with implicit constants depending only on d, k, δ , and σ .

The following two lemmas are the main technical ingredients in the proof of Theorem B.1.

LEMMA B.3. *Let $0 < \delta < 1$ and $\Omega \subset \mathbb{R}_+^d$ be an open set with a boundary portion T on $\partial\mathbb{R}_+^d$. Let*

$$(B.5) \quad Lu = f + \nabla \cdot g \text{ in } \Omega \quad \text{and} \quad u = 0 \text{ on } T,$$

where

$$(B.6) \quad L = - \sum_{r,s=1}^d \partial_r a_{r,s} \partial_s.$$

Then

$$(B.7) \quad |u|_{1,\delta,\Omega \cup T}^{(0)} \lesssim |u|_{0,\Omega \cup T}^{(0)} + |f|_{0,\Omega \cup T}^{(2)} + |g|_{0,\delta,\Omega \cup T}^{(1)},$$

with the implicit constant depending only on d, δ, λ , and Λ , where

$$(B.8) \quad \sum_{r,s=1}^d |a_{r,s}|_{0,\delta,\Omega \cup T}^{(0)} \leq \Lambda$$

and $\lambda > 0$ is a uniform lower bound on the lowest eigenvalue of the symmetric matrix defined by $a_{r,s}$.

For us the bound with $g = 0$ suffices, but g appears naturally in the proof.

PROOF. A similar, but less precise bound appears in [12, cor. 8.36]. Since its proof is sketched only very briefly, we provide some more details. The starting point is [12, (4.46)], which proves the lemma in the case $L = -\Delta$ and $\Omega = B_R(x_0) \cap \mathbb{R}_+^d$ with $x_0 \in \mathbb{R}_+^d$. By the same argument as in the proof of [12, theorem 4.12] (which is not given, but which is similar to the proof of [12, theorem 4.8]), this bound leads to Lemma B.3 for $L = -\Delta$, but for general Ω . Using a simple change of variables as in the proof of [12, lemma 6.1], we obtain the lemma for $L = -\nabla \cdot A \nabla$ with a constant matrix A again for a general Ω . Finally, using the perturbation argument as in the proof of [12, lemma 6.4] (which again is not given, but which is similar to the proof of [12, theorem 6.2]), we obtain the lemma. \square

LEMMA B.4. *Let $k \geq 2, 0 < \delta < 1$, and $\Omega \subset \mathbb{R}_+^d$ be an open set with a boundary portion T on $\partial\mathbb{R}_+^d$. Let*

$$(B.9) \quad Lu = f \text{ in } \Omega \quad \text{and} \quad u = 0 \text{ on } T,$$

where

$$(B.10) \quad L = - \sum_{r,s=1}^d a_{r,s} \partial_r \partial_s + \sum_{r=1}^d b_r \partial_r.$$

Then

$$(B.11) \quad |u|_{k,\delta,\Omega \cup T}^{(0)} \lesssim |u|_{0,\Omega \cup T}^{(0)} + |f|_{k-2,\delta,\Omega \cup T}^{(2)}$$

with the implicit constant depending only on d , k , δ , λ , and Λ , where

$$(B.12) \quad \sum_{r,s=1}^d |a_{r,s}|_{k-2,\delta,\Omega \cup T}^{(0)} + \sum_{r=1}^d |b_r|_{k-2,\delta,\Omega \cup T}^{(1)} \leq \Lambda$$

and $\lambda > 0$ is a uniform lower bound on the lowest eigenvalue of the symmetric matrix defined by $a_{r,s}$.

PROOF. Lemma B.4 with $k = 2$ coincides with [12, lemma 6.4]. Estimates similar to, but less precise than our statement for $k \geq 3$, are stated as [12, problem 6.2], but without any details.

We shall show that for any integer $k \geq 2$ and any $\sigma \geq 0$,

$$(B.13) \quad |u|_{k,\delta,\Omega \cup T}^{(\sigma)} \lesssim |u|_{0,\Omega \cup T}^{(\sigma)} + |f|_{k-2,\delta,\Omega \cup T}^{(\sigma+2)}$$

where the implicit constant depends only on d , k , δ , σ , λ , and Λ . We will prove this by induction on k .

First, let $k = 2$. For $\sigma = 0$ the claimed inequality is [12, lemma 6.4] (whose proof is not given, but which is similar to the proof of [12, theorem 6.2]). The proof for $\sigma > 0$ follows by the same argument.

Now let $k \geq 3$ and $\sigma \geq 0$. We assume the inequality has already been shown for all smaller values of k and for all values of σ . For $1 \leq j \leq d - 1$ the function $v = \partial_j u$ satisfies

$$(B.14) \quad Lv = \tilde{f} \text{ in } \Omega \quad \text{and} \quad v = 0 \text{ on } T,$$

where

$$(B.15) \quad \tilde{f} = \partial_j f + \sum_{r,s=1}^d (\partial_j a_{r,s}) \partial_r \partial_s u - \sum_{r=1}^d (\partial_j b_r) \partial_r u.$$

Therefore, by the induction assumption (B.13) with σ replaced by $\sigma + 1$,

$$\begin{aligned}
& |v|_{k-1,\delta,\Omega\cup T}^{(\sigma+1)} \\
& \lesssim |v|_{0,\Omega\cup T}^{(\sigma+1)} + |\tilde{f}|_{k-3,\delta,\Omega\cup T}^{(\sigma+3)} \\
& \leq |\partial_j u|_{0,\Omega\cup T}^{(\sigma+1)} + |\partial_j f|_{k-3,\delta,\Omega\cup T}^{(\sigma+3)} + \sum_{r,s} |(\partial_j a_{r,s}) \partial_r \partial_s u|_{k-3,\delta,\Omega\cup T}^{(\sigma+3)} \\
& \quad + \sum_r |(\partial_j b_r) \partial_r u|_{k-3,\delta,\Omega\cup T}^{(\sigma+3)} \\
& \lesssim |\partial_j u|_{0,\Omega\cup T}^{(\sigma+1)} + |\partial_j f|_{k-3,\delta,\Omega\cup T}^{(\sigma+3)} + \sum_{r,s} |\partial_j a_{r,s}|_{k-3,\delta,\Omega\cup T}^{(1)} |\partial_r \partial_s u|_{k-3,\delta,\Omega\cup T}^{(\sigma+2)} \\
& \quad + \sum_r |\partial_j b_r|_{k-3,\delta,\Omega\cup T}^{(2)} |\partial_r u|_{k-3,\delta,\Omega\cup T}^{(\sigma+1)} \\
& \lesssim |u|_{1,\Omega\cup T}^{(\sigma)} + |f|_{k-2,\delta,\Omega\cup T}^{(\sigma+2)} + \sum_{r,s} |a_{r,s}|_{k-2,\delta,\Omega\cup T}^{(0)} |u|_{k-1,\delta,\Omega\cup T}^{(\sigma)} \\
& \quad + \sum_r |b_r|_{k-2,\delta,\Omega\cup T}^{(1)} |u|_{k-2,\delta,\Omega\cup T}^{(\sigma)} \\
& \lesssim |f|_{k-2,\delta,\Omega\cup T}^{(\sigma+2)} + |u|_{k-1,\delta,\Omega\cup T}^{(\sigma)},
\end{aligned}$$

where we have used the properties of the norms discussed after equation (B.4). Bounding the last term on the right side using the induction assumption with σ , we finally obtain

$$(B.16) \quad |\partial_j u|_{k-1,\delta,\Omega\cup T}^{(\sigma+1)} \lesssim |u|_{0,\Omega\cup T}^{(\sigma)} + |f|_{k-2,\delta,\Omega\cup T}^{(\sigma+2)} \quad \text{if } j = 1, \dots, d-1.$$

On the other hand, we have

$$\partial_d^2 u = \frac{1}{a_{dd}} \left(- \sum_{(r,s) \neq (d,d)} a_{rs} \partial_r \partial_s u + \sum_r b_r \partial_r u - f \right)$$

and therefore

$$\begin{aligned}
& |\partial_d^2 u|_{k-2,\delta,\Omega\cup T}^{(\sigma+2)} \\
& \leq |a_{dd}^{-1}|_{k-2,\delta,\Omega\cup T}^{(0)} \left(\sum_{(r,s) \neq (d,d)} |a_{rs}|_{k-2,\delta,\Omega\cup T}^{(0)} |\partial_r \partial_s u|_{k-2,\delta,\Omega\cup T}^{(\sigma+2)} \right. \\
& \quad \left. + \sum_r |b_r|_{k-2,\delta,\Omega\cup T}^{(1)} |\partial_r u|_{k-2,\delta,\Omega\cup T}^{(\sigma+1)} + |f|_{k-2,\delta,\Omega\cup T}^{(\sigma+2)} \right).
\end{aligned}$$

Our assumptions imply that $|a_{dd}^{-1}|_{k-2,\delta,\Omega\cup T}^{(0)}$ is bounded in terms of Λ and λ . Moreover, $|\partial_r u|_{k-2,\delta,\Omega\cup T}^{(\sigma+1)}$ is bounded above for any $1 \leq r \leq d$ by $|u|_{k-1,\delta,\Omega\cup T}^{(\sigma)}$,

which by the induction hypothesis (B.13) is bounded by $|u|_{0,\Omega \cup T}^{(\sigma)} + |f|_{k-3,\delta,\Omega \cup T}^{(\sigma+2)}$. We thus conclude that

$$\begin{aligned} & |\partial_d^2 u|_{k-2,\delta,\Omega \cup T}^{(\sigma+2)} \\ & \lesssim \sum_{(r,s) \neq (d,d)} |\partial_r \partial_s u|_{k-2,\delta,\Omega \cup T}^{(\sigma+2)} + \sum_r |\partial_r u|_{k-2,\delta,\Omega \cup T}^{(\sigma+1)} + |f|_{k-2,\delta,\Omega \cup T}^{(\sigma+2)} \\ & \lesssim \sum_{j=1}^{d-1} |\partial_j u|_{k-1,\delta,\Omega \cup T}^{(\sigma+1)} + |f|_{k-2,\delta,\Omega \cup T}^{(\sigma+2)}. \end{aligned}$$

Combining this with (B.16) we obtain the claimed estimate on $|u|_{k,\delta,\Omega \cup T}^{(\sigma)}$. This completes the proof of Lemma B.4. \square

B.3 Proof of Theorem B.1

We first assume that $\text{dist}(a, \partial\Omega) \geq 2R$. In this case $B_{2R}(a) \subset \Omega$, and we can apply Lemmas B.3 and B.4 with $L = -\Delta$, $T = \emptyset$, and $B_{2R}(a)$ playing the role of Ω . Since

$$\begin{aligned} |u|_{k,\delta,\Omega}^{(\sigma)} & \leq \sum_{j=0}^k \sup_{x \in B_{2R}(a)} (2R)^{j+\sigma} |\partial^j u(x)| \\ & \quad + \sup_{x,y \in B_{2R}(a)} (2R)^{k+\delta+\sigma} \frac{|\partial^k u(x) - \partial^k u(y)|}{|x-y|^\delta} \end{aligned}$$

(and similarly with u replaced by f) and

$$\begin{aligned} |u|_{k,\delta,\Omega}^{(\sigma)} & \geq \sum_{j=0}^k \sup_{x \in B_R(a)} R^{j+\sigma} |\partial^j u(x)| \\ & \quad + \sup_{x,y \in B_R(a)} R^{k+\delta+\sigma} \frac{|\partial^k u(x) - \partial^k u(y)|}{|x-y|^\delta}, \end{aligned}$$

we immediately obtain the bound in this case. (Of course, in order to prove the bounds, much simpler versions of Lemmas B.3 and B.4 would suffice.)

Now assume that $\text{dist}(a, \partial\Omega) < 2R$. We set

$$(B.17) \quad r_1 = \begin{cases} (2M)^{-1/\delta} r_0 & \text{if } k = 1, \\ (2M)^{-1} r_0 & \text{if } k \geq 2. \end{cases}$$

Without loss of generality we assume $M \geq \frac{1}{2}$; hence $r_1 \leq r_0$. We will first assume that $R \leq r_1/4$, which implies that if $p \in \partial\Omega$ is chosen with $|p-a| = \text{dist}(a, \partial\Omega)$; then

$$(B.18) \quad B_{2R}(a) \cap \Omega \subset B_{r_1}(p) \cap \Omega.$$

(Indeed, if $|y - a| < 2R$, then $|y - p| \leq |y - a| + |a - p| < 2R + \text{dist}(a, \partial\Omega) \leq 4R \leq r_1$.) Therefore, we can work in the boundary coordinates from the definition of a $C^{k,\delta}$ domain centered at the point p . After a translation and a rotation we may assume that $p = 0$ and that there is a function $\Gamma : \{y' \in \mathbb{R}^{d-1} : |y'| < r_0\} \rightarrow \mathbb{R}$ with $\Gamma(0) = 0$, $\nabla\Gamma(0) = 0$, and

$$(B.19) \quad \Omega \cap B_{r_0}(0) = \{(y', y_d) \in \mathbb{R}^{d-1} \times \mathbb{R} : |y'| < r_0, y_d > \Gamma(y')\} \cap B_{r_0}(0).$$

We introduce the change of variables $\Phi : \Omega \cap B_{r_0}(0) \rightarrow \mathbb{R}_+^d$,

$$(B.20) \quad \Phi_m(y) = y_m \quad \text{if } 1 \leq m \leq d-1, \quad \Phi_d(y) = y_d - \Gamma(y').$$

The following lemma shows that decreasing r_0 to r_1 ensures that Φ is bi-Lipschitz.

LEMMA B.5. *For $x, y \in \Omega \cap B_{r_1}(0)$, we have*

$$(B.21) \quad \frac{1}{2}|x - y| \leq |\Phi(x) - \Phi(y)| \leq \frac{3}{2}|x - y|.$$

PROOF. For $x, y \in \Omega \cap B_{r_0}(0)$ we have by the triangle inequality

$$(B.22) \quad \left| |\Phi(x) - \Phi(y)| - |x - y| \right| \leq |\Gamma(x') - \Gamma(y')|.$$

In order to further bound this, we write, using $\nabla\Gamma(0) = 0$,

$$(B.23) \quad \Gamma(x') - \Gamma(y') = \int_0^1 (x' - y') \cdot (\nabla\Gamma(y' + t(x' - y')) - \nabla\Gamma(0)) dt.$$

When $k = 1$, we obtain

$$(B.24) \quad \begin{aligned} |\Gamma(x') - \Gamma(y')| &\leq M r_0^{-\delta} \int_0^1 |x' - y'| |y' + t(x' - y')|^\delta dt \\ &\leq M r_0^{-\delta} \max\{|x'|, |y'|\}^\delta |x' - y'|. \end{aligned}$$

For $|x'|, |y'| \leq r_1 = (2M)^{-1/\delta} r_0$, this is $\leq |x' - y'|/2$. The argument for $k \geq 2$ is similar. \square

Let $\tilde{\Omega} = \Phi(B_{2R}(a) \cap \Omega)$. This is an open set in \mathbb{R}_+^d with a boundary portion $T = \Phi(B_{2R}(a) \cap \partial\Omega)$ on $\partial\mathbb{R}_+^d$. For a function g on $B_{2R}(a) \cap \Omega$ we define a function \tilde{g} on $\tilde{\Omega}$ by

$$(B.25) \quad \tilde{g}(x) = g(\Phi^{-1}(x)).$$

We claim that

$$(B.26) \quad \begin{aligned} |\tilde{g}|_{k,\delta,\tilde{\Omega} \cup T}^{(\sigma)} &\lesssim \sum_{j=0}^k R^{j+\sigma} \sup_{B_{2R}(a) \cap \Omega} |\partial^j g| \\ &\quad + R^{k+\delta+\sigma} \sup_{x,y \in B_{2R}(a) \cap \Omega} \frac{|\partial^k g(x) - \partial^k g(y)|}{|x - y|^\delta} \end{aligned}$$

with an implicit constant depending only on d, k, δ , and M . Indeed, by Lemma B.5, for $x \in B_{2R}(a) \cap \Omega$,

$$(B.27) \quad \text{dist}(\Phi(x), \partial\tilde{\Omega} \setminus T) \leq \frac{3}{2} \text{dist}(x, \partial(B_{2R}(a) \cap \Omega) \setminus (B_{2R}(a) \cap \partial\Omega)) \leq 3R.$$

Moreover, for $j \leq d-1$, we have $\partial_j \tilde{g} = \partial_j g + \partial_d g \partial_j \Gamma$ and $\partial_d \tilde{g} = \partial_d g$. Since $|\partial_j \Gamma| \leq M$, we see that $|\partial \tilde{g}| \lesssim |\partial g|$. When computing a second derivative, also a term like $\partial_d g \partial_j \partial_k \Gamma$ appears. Bounding $|\partial_j \partial_k \Gamma| \leq M r_0^{-1}$ and $R \lesssim r_0$, we obtain $|\partial^2 \tilde{g}| \lesssim |\partial^2 g| + R^{-1} |\partial g|$. The arguments for higher derivatives and for the Hölder term are similar.

After these preliminaries we now return to our differential equation. We have $-\Delta u = f$ in $\Omega \cap B_{2R}(a)$ and $u = 0$ on $\partial\Omega \cap B_{2R}(a)$. Therefore the functions

$$(B.28) \quad \tilde{u}(x) = u(\Phi^{-1}(x)), \quad \tilde{f}(x) = f(\Phi^{-1}(x))$$

satisfy

$$(B.29) \quad L\tilde{u} = \tilde{f} \text{ in } \tilde{\Omega} \quad \text{and} \quad \tilde{u} = 0 \text{ on } T$$

with the operator

$$(B.30) \quad L = - \sum_{r,s=1}^d \partial_r a_{r,s} \partial_s,$$

where

$$(B.31) \quad a_{r,s} = \begin{cases} \delta_{r,s} & \text{if } r, s \leq d-1, \\ 1 + (\nabla\Gamma)^2 & \text{if } r = s = d, \\ -\partial_r \Gamma & \text{if } r < d = s, \\ -\partial_s \Gamma & \text{if } s < d = r. \end{cases}$$

A straightforward computation shows that the smallest eigenvalue of the matrix defined by $a_{r,s}$ is given by $1 + \frac{1}{2}((\nabla\Gamma)^2 - \sqrt{(\nabla\Gamma)^4 + 4(\nabla\Gamma)^2})$. The function $t \mapsto 1 + \frac{1}{2}(t - \sqrt{t^2 + 4t})$ is positive for $t \geq 0$ and strictly decreasing to 0 as $t \rightarrow \infty$. Therefore, since $|\nabla\Gamma| \leq M$ by our definition of $C^{k,\delta}$ smoothness, we see that the lowest eigenvalue is uniformly bounded below by some $\lambda > 0$ depending only on M .

Moreover, using the definition of a $C^{k,\delta}$ -set and the fact that $R \lesssim r_0$, we deduce from (B.26) that

$$(B.32) \quad \sum_{r,s} |a_{r,s}|_{k-1,\delta,\tilde{\Omega} \cup T}^{(0)} \leq \Lambda$$

with Λ depending only on d, k, δ , and M . Similarly, for

$$(B.33) \quad b_r = - \sum_{s=1}^d \partial_s a_{sr} = \begin{cases} 0 & \text{if } r \leq d-1, \\ \Delta\Gamma & \text{if } r = d, \end{cases}$$

and $k \geq 2$, we have

$$(B.34) \quad \sum_r |b_r|_{k-2,\delta,\tilde{\Omega} \cup T}^{(1)} \leq \Lambda.$$

From Lemmas B.3 and B.4 we conclude that

$$(B.35) \quad \|\tilde{u}\|_{k,\delta,\tilde{\Omega} \cup T}^{(0)} \lesssim \|\tilde{u}\|_{0,\tilde{\Omega} \cup T}^{(0)} + \begin{cases} \|\tilde{f}\|_{0,\tilde{\Omega} \cup T}^{(2)} & \text{if } k = 1, \\ \|\tilde{f}\|_{k-2,\delta,\tilde{\Omega} \cup T}^{(2)} & \text{if } k \geq 2. \end{cases}$$

According to (B.26), the right side of (B.35) can be further bounded by a constant (depending only on d , k , δ , and M) times

$$\sup_{B_{2R}(a) \cap \Omega} |u| + \begin{cases} R^2 \sup_{B_{2R}(a) \cap \Omega} |f| & \text{if } k = 1, \\ \sum_{j=0}^{k-2} R^{j+2} \sup_{B_{2R}(a) \cap \Omega} |\partial^j f| \\ \quad + R^{k+\delta} \sup_{x,y \in B_{2R}(a) \cap \Omega} \frac{|\partial^{k-2} f(x) - \partial^{k-2} f(y)|}{|x-y|^\delta} & \text{if } k \geq 2. \end{cases}$$

We claim that the left side of (B.35) is bounded from below by a constant (depending only on d , k , δ , and M) times

$$\sum_{j=0}^k R^j \sup_{B_R(a) \cap \Omega} |\partial^j u| + R^{k+\delta} \sup_{x,y \in B_R(a) \cap \Omega} \frac{|\partial^k u(x) - \partial^k u(y)|}{|x-y|^\delta}.$$

The proof of the latter fact is similar to that of (B.26). Namely, for $x \in B_R(a) \cap \Omega$, one has

$$\text{dist}(\Phi(x), \partial\tilde{\Omega} \setminus T) \geq \frac{1}{2} \text{dist}(x, \partial(B_{2R}(a) \cap \Omega) \setminus (B_{2R}(a) \cap \partial\Omega)) \geq \frac{1}{2} R.$$

Moreover, factors of derivatives of Γ , which appear when computing derivatives of u in terms of derivatives of \tilde{u} , are handled as in the proof of (B.26). This completes the proof of the theorem in case $R_0 \leq r_1/4$ with r_1 defined in (B.17).

The case of larger R_0 is readily reduced to the previous case by covering the ball $B_R(a)$ with finitely many smaller balls of size $r_1/4$. As long as R_0/r_0 is bounded, this only modifies the constants in the bounds. \square

Appendix C Bounds on the Kernel of Functions of the Dirichlet Laplacian

In this appendix we will use the bounds in Appendix B, specifically Corollary B.2, to obtain estimates on derivatives of the integral kernel of various functions of the Dirichlet Laplacian Δ_Ω for $\Omega \subset \mathbb{R}^d$. We work in arbitrary dimension $d \geq 1$.

C.1 Simple Bounds

We recall [4, eq. (1.9.1)] that for any $x, y \in \Omega$, one has

$$(C.1) \quad 0 \leq e^{t\Delta_\Omega}(x, y) \leq e^{t\Delta_{\mathbb{R}^d}}(x, y) = (4\pi t)^{-d/2} e^{-(x-y)^2/(4t)}.$$

Therefore, by Bernstein's theorem we infer that for any completely monotone function f on $[0, \infty)$, we have

$$(C.2) \quad 0 \leq f(-\Delta_\Omega)(x, y) \leq f(-\Delta_{\mathbb{R}^d})(x, y) = \int_{\mathbb{R}^d} f(k^2) e^{ik \cdot (x-y)} \frac{dk}{(2\pi)^d}.$$

This bound is used in the main text multiple times, for instance with f given by $f(t) = t^{-1} e^{-t/K^2}$ and $f(t) = (t + K^2)^{-3}$.

To motivate the following, we shall first derive a more general but slightly worse bound on the diagonal $x = y$, assuming only that f is nonincreasing. Assuming that Ω is bounded (or more generally that the spectrum of $-\Delta_\Omega$ is discrete), we shall denote the eigenvalues of $-\Delta_\Omega$ (in increasing order and repeated according to their multiplicities) by e_n , and the corresponding eigenfunctions by φ_n . According to (C.1) we have for any $K > 0$

$$\sum_{e_n \leq K^2} |\varphi_n(x)|^2 \leq e^{tK^2} e^{t\Delta_\Omega}(x, x) \leq e^{tK^2} (4\pi t)^{-d/2}.$$

Optimizing in t yields

$$\sum_{e_n \leq K^2} |\varphi_n(x)|^2 \leq \left(\frac{e}{2\pi d}\right)^{d/2} K^d = \left(\frac{2e}{d}\right)^{d/2} \Gamma(1 + d/2) \int_{\{|k| \leq K\}} \frac{dk}{(2\pi)^d}.$$

Any nonincreasing function f with $\lim_{t \rightarrow \infty} f(t) = 0$ can be written as a superposition of characteristic functions as $f(t) = -\int_0^\infty \chi_{\{t \leq s\}} f'(s) ds$, and hence

$$(C.3) \quad \begin{aligned} \sum_n f(e_n) |\varphi_n(x)|^2 &= f(-\Delta_\Omega)(x, x) \\ &\leq \left(\frac{2e}{d}\right)^{d/2} \Gamma(1 + d/2) \int_{\mathbb{R}^d} f(k^2) \frac{dk}{(2\pi)^d} \end{aligned}$$

for nonincreasing functions.

C.2 Bounds on the Diagonal

We now use the same method to derive bounds on $\sum_n f(e_n) |\partial^\beta \varphi_n(x)|^2$. To do so we shall use Corollary B.2 to prove the following.

LEMMA C.1. *Assume that $\Omega \subset \mathbb{R}^d$ is a bounded, open $C^{k, \delta}$ set for some $k \geq 1$ and $0 < \delta < 1$, and let $R_0 > 0$. For any bounded function $g : \mathbb{R}_+ \rightarrow \mathbb{R}$ of*

compact support, any $\beta \in \mathbb{N}_0^d$ with $|\beta| \leq k$, and any $R \in (0, R_0)$,

$$(C.4) \quad \begin{aligned} & R^{2|\beta|} \sum_n g(e_n)^2 |\partial^\beta \varphi_n(x)|^2 \\ & \lesssim \sum_{j=0}^{|\beta|} \sup_{x' \in B_{R^2}(x) \cap \Omega} \sum_n g(e_n)^2 (R^2 e_n)^{2j} |\varphi_n(x')|^2 \end{aligned}$$

for all $x \in \Omega$.

PROOF. We proceed by induction in $|\beta|$. For $|\beta| = 0$, (C.4) obviously holds. Assume now $|\beta| \geq 1$. Pick a $\psi \in L^2(\Omega)$, and let $u = g(-\Delta_\Omega)\psi$. From Corollary B.2, we obtain for any $x \in \Omega$

$$(C.5) \quad R^{|\beta|} |\partial^\beta u(x)| \lesssim \sup_{B_R(x) \cap \Omega} |u(x')| + \sum_{\alpha: |\alpha| < |\beta|} R^{|\alpha|+2} \sup_{B_R(x) \cap \Omega} |\partial^\alpha \Delta_\Omega u(x')|.$$

Now

$$(C.6) \quad \begin{aligned} |u(x')| &= |g(-\Delta_\Omega)\psi(x')| = \left| \sum_n g(e_n) \langle \varphi_n | \psi \rangle \varphi_n(x') \right| \\ &\leq \left(\sum_n g(e_n)^2 |\varphi_n(x')|^2 \right)^{1/2} \|\psi\|_2 \end{aligned}$$

and similarly

$$(C.7) \quad |\partial^\alpha \Delta_\Omega u(x')| \leq \left(\sum_n g(e_n)^2 e_n^{2|\alpha|} |\partial^\alpha \varphi_n(x')|^2 \right)^{1/2} \|\psi\|_2.$$

By combining (C.5)–(C.7) and using the induction hypotheses for α with $|\alpha| < |\beta|$, we obtain the bound

$$\begin{aligned} & R^{2|\beta|} |\partial^\beta g(-\Delta_\Omega)\psi(x)|^2 \\ & \lesssim \|\psi\|_2^2 \sum_{j=0}^{|\beta|} \sup_{B_{2R}(x) \cap \Omega} \sum_n g(e_n)^2 (R^2 e_n)^{2j} |\varphi_n(x')|^2 \end{aligned}$$

valid for all $\psi \in L^2(\Omega)$. Since

$$(C.8) \quad \sup_\psi \|\psi\|_2^{-2} |\partial^\beta g(-\Delta_\Omega)\psi(x)|^2 = \sum_n g(e_n)^2 |\partial^\beta \varphi_n(x)|^2$$

the result follows. \square

We apply (C.4) with g the characteristic function of $\{e \leq K^2\}$ for some $K > 0$, $R = K^{-1}$, and $R_0 = e_1^{-1/2}$. This yields

$$(C.9) \quad \sum_{e_n \leq K^2} |\partial^\beta \varphi_n(x)|^2 \lesssim K^{2|\beta|} \sup_{B_{K^{-1}}(x) \cap \Omega} \sum_{e_n \leq K^2} |\varphi_n(x')|^2 \lesssim K^{2|\beta|+d}$$

where we have used (C.3) in the last step. More generally, we obtain for any nonincreasing function f with $\lim_{t \rightarrow \infty} t^{d/2+|\beta|} f(t) = 0$ that

$$\begin{aligned}
\sum_n f(e_n) |\partial^\beta \varphi_n(x)|^2 &= - \int_0^\infty \sum_{e_n \leq E} |\partial^\beta \varphi_n(x)|^2 f'(E) dE \\
&\lesssim - \int_0^\infty E^{d/2+|\beta|} f'(E) dE \\
&= \text{const} \int_0^\infty E^{d/2+|\beta|-1} f(E) dE \\
&= \text{const} \int_{\mathbb{R}^d} k^{2|\beta|} f(k^2) \frac{dk}{(2\pi)^d}.
\end{aligned}
\tag{C.10}$$

The validity of (C.9) is shown in [16, theorem 17.5.3] if Ω has C^∞ boundary. Following the proof there (which is based on regularity theory in L^2 -based Sobolev spaces), one sees that a certain finite number of derivatives is actually sufficient, but the result is not as precise as ours, which only requires $C^{|\beta|, \delta}$ regularity of the boundary.

C.3 Off-Diagonal Bounds

In this section we shall derive a bound on the derivatives of the kernel of certain functions of the Dirichlet Laplacian, valid even away from the diagonal. These bounds are much less general than the ones in the previous two subsections, however. For simplicity we only consider the particular class of functions needed in the main text, but the method obviously extends to other functions as well.

For $\Lambda > 0$ and $\ell > 0$, let

$$z_\ell(t) = t^{-\ell} (1 - e^{-t/\Lambda^2})^2.$$

LEMMA C.2. *Assume that $\Omega \subset \mathbb{R}^d$ is a bounded, open $C^{k, \delta}$ set for some $k \geq 1$ and $0 < \delta < 1$. For any $\beta \in \mathbb{N}_0^d$ with $|\beta| \leq k$ and $|\beta| < 2 + d/2$, and any $\ell \in (|\beta|, 2 + d/2)$ and $\Lambda > 0$, we have*

$$\begin{aligned}
|\partial_x^\beta z_\ell(-\Delta_\Omega)(x, y)| &\lesssim \\
\begin{cases} |x - y|^{2\ell - d - |\beta|} & \text{for } \ell < d/2, \\ \ln(1 + (\Lambda|x - y|)^{-1}) |x - y|^{-|\beta|} & \text{for } \ell = d/2, \\ \Lambda^{d-2\ell} |x - y|^{-|\beta|} & \text{for } \ell > d/2, \end{cases}
\end{aligned}
\tag{C.11}$$

for $\Lambda|x - y| \leq 1$, and

$$|\partial_x^\beta z_\ell(-\Delta_\Omega)(x, y)| \lesssim \Lambda^{-4} |x - y|^{2\ell - 4 - d - |\beta|}$$

for $\Lambda|x - y| \geq 1$.

PROOF. We use again Corollary B.2 above. A simple induction argument as in the proof of Lemma C.1 shows that

$$(C.13) \quad R^{|\beta|} |\partial_x^\beta z_\ell(-\Delta_\Omega)(x, y)| \lesssim \sum_{i=0}^{|\beta|} R^{2i} \sup_{x' \in B_R(x)} |z_{\ell-i}(-\Delta_\Omega)(x', y)|$$

for any $R > 0$ (smaller than some arbitrary, fixed value). To estimate the right side of (C.13), we write for $j > 0$

$$\begin{aligned} z_j(t) &= t^{-j} (1 - e^{-t/\Lambda^2})^2 \\ &= \frac{1}{\Gamma(j)} \int_0^\infty e^{-\lambda t} (\lambda^{j-1} - 2[\lambda - \Lambda^{-2}]_+^{j-1} + [\lambda - 2\Lambda^{-2}]_+^{j-1}) d\lambda \end{aligned}$$

where the term $[\lambda - \Lambda^{-2}]_+^{j-1}$ is understood as being 0 for $\lambda < \Lambda^{-2}$ even when $j < 1$, and likewise for $[\lambda - 2\Lambda^{-2}]_+^{j-1}$. In particular, from (C.1), we thus have

$$(C.14) \quad |z_j(-\Delta_\Omega)(x, y)| \leq \Lambda^{d-2j} f_j(\Lambda|x-y|)$$

with

$$(C.15) \quad f_j(t) = \frac{1}{\Gamma(j)(4\pi)^{d/2}} \int_0^\infty e^{-t^2/(4\lambda)} |\lambda^{j-1} - 2[\lambda - 1]_+^{j-1} + [\lambda - 2]_+^{j-1}| \lambda^{-d/2} d\lambda.$$

We note that

$$(C.16) \quad |\lambda^{j-1} - 2[\lambda - 1]_+^{j-1} + [\lambda - 2]_+^{j-1}| \lesssim \lambda^{j-3}$$

for $\lambda \geq 3$. Using this, one readily checks that as long as $0 < j < 2 + d/2$,

$$(C.17) \quad \begin{aligned} f_j(t) &\lesssim t^{2j-4-d} \quad \text{for } t \geq 1, \\ f_j(t) &\lesssim \begin{cases} 1 & \text{for } j > d/2, \\ \ln(2/t) & \text{for } j = d/2, \\ t^{2j-d} & \text{for } j < d/2, \end{cases} \quad \text{for } t \leq 1. \end{aligned}$$

We plug these bounds into (C.14) and choose $R = |x - y|/2$ in (C.13). (Note that $R \leq R_0$, as required for (C.13), where $R_0 = \text{diameter of } \Omega$.) For all $x' \in B_R(x)$, we then have $|x' - y| \geq |x - y|/2$, and hence (C.13), (C.14), and (C.17) imply the desired bounds (C.11) and (C.12) for this choice of R . \square

Recall the definition $u_{jk}(x) = \sup_{y \in \mathbb{R}^3} |p_j p_k |p|^{-4} w_{x+y}(y)|$ with

$$(C.18) \quad w_x(y) = z_{1/2}(-\Delta_\Omega)(x, y).$$

Applying the bounds (C.11) and (C.12), with $\ell = 5/2$, $d = 3$, and $|\beta| = 2$, we readily obtain

$$(C.19) \quad u_{jk}(x) \lesssim \min\{\Lambda^{-2}|x|^{-2}, \Lambda^{-4}|x|^{-4}\}.$$

The function $\min\{|x|^{-2}, |x|^{-4}\}$ is in $L^{6/5}(\mathbb{R}^3)$ and hence has finite Coulomb norm. By the Hardy-Littlewood-Sobolev inequality and scaling, it thus follows immediately that $\|u_{jk}\|_C \lesssim \Lambda^{-5/2}$, as claimed in (5.22).

Acknowledgments. Partial support through National Science Foundation Grant DMS-1363432 (R.L.F.) and the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 694227; R.S.), is acknowledged.

Bibliography

- [1] Alexandrov, A. S.; Devreese, J. T. *Advances in polaron physics*. Springer, Berlin-Heidelberg, 2010.
- [2] Allcock, G. R. On the polaron rest energy and effective mass. *Adv. Phys.* **5** (1956), no. 20, 412–451.
- [3] Allcock, G. R. Strong-coupling theory of the polaron. *Polarons and Excitons*, 45–70. Plenum Press, 1963.
- [4] Davies, E. B. *Heat kernels and spectral theory*. Cambridge Tracts in Mathematics, 92. Cambridge University Press, Cambridge, 1990.
- [5] Donsker, M. D.; Varadhan, S. R. S. Asymptotics for the polaron. *Comm. Pure Appl. Math.* **36** (1983), no. 4, 505–528. doi:10.1002/cpa.3160360408
- [6] Feliciangeli, D.; Seiringer, R. Uniqueness and non-degeneracy of minimizers of the Pekar functional on a ball. *SIAM J. Math. Anal.* **52** (2020), 605–622. doi:10.1137/19M126284X
- [7] Frank, R. L.; Lieb, E. H.; Seiringer, R. Symmetry of bipolaron bound states for small Coulomb repulsion. *Comm. Math. Phys.* **319** (2013), no. 2, 557–573. doi:10.1007/s00220-012-1604-y
- [8] Frank, R. L.; Lieb, E. H.; Seiringer, R.; Thomas, L. E. Ground state properties of multi-polaron systems. *XVIIth International Congress on Mathematical Physics*, 477–485. World Scientific, Hackensack, N.J., 2014.
- [9] Frank, R. L.; Schlein, B. Dynamics of a strongly coupled polaron. *Lett. Math. Phys.* **104** (2014), no. 8, 911–929. doi:10.1007/s11005-014-0700-7
- [10] Fröhlich, H. Theory of electrical breakdown in ionic crystals. *Proc. R. Soc. Lond. A* **160** (1937), no. 901, 230–241. doi:10.1098/rspa.1937.0106
- [11] Gerlach, B.; Löwen, H. Analytical properties of polaron systems or: Do polaronic phase transitions exist or not? *Rev. Mod. Phys.* **63** (1991), no. 1, 63–90. doi:10.1103/RevModPhys.63.63
- [12] Gilbarg, D.; Trudinger, N. S. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer, Berlin, 2001.
- [13] Griesemer, M.; Wunsch, A. Self-adjointness and domain of the Fröhlich Hamiltonian. *J. Math. Phys.* **57** (2016), no. 2, 021902, 15 pp. doi:10.1063/1.4941561
- [14] Gross, E. P. Particle-like solutions in field theory. *Ann. Physics* **19** (1962), 219–233. doi:10.1016/0003-4916(62)90216-6
- [15] Gross, E. P. Strong coupling polaron theory and translational invariance. *Ann. Physics* **99** (1976), no. 1, 1–29. doi:10.1016/0003-4916(76)90082-8
- [16] Hörmander, L. *The analysis of linear partial differential operators. III. Pseudo-differential operators*. Classics in Mathematics. Springer, Berlin, 2007. doi:10.1007/978-3-540-49938-1
- [17] Lenzmann, E. Uniqueness of ground states for pseudorelativistic Hartree equations. *Anal. PDE* **2** (2009), no. 1, 1–27. doi:10.2140/apde.2009.2.1
- [18] Lieb, E. H. Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation. *Studies in Appl. Math.* **57** (1976/77), no. 2, 93–105. doi:10.1002/sapm197757293
- [19] Lieb, E. H.; Loss, M. *Analysis*. Second edition. Graduate Studies in Mathematics, 14. American Mathematical Society, Providence, R.I., 2001. doi:10.1090/gsm/014

- [20] Lieb, E. H.; Thomas, L. E. Exact ground state energy of the strong-coupling polaron. *Comm. Math. Phys.* **183** (1997), no. 3, 511–519. doi:10.1007/s002200050040 Erratum-ibid. [21].
- [21] Lieb, E. H.; Thomas, L. E. Erratum: “Exact ground state energy of the strong-coupling polaron”. *Comm. Math. Phys.* **188** (1997), no. 2, 499–500. doi:10.1007/s002200050175
- [22] Lieb, E. H.; Yamazaki, K. Ground-state energy and effective mass of the polaron. *Phys. Rev.* **111** (1958), no. 3, 728–733. doi:10.1103/PhysRev.111.728
- [23] Møller, J. S. The polaron revisited. *Rev. Math. Phys.* **18** (2006), no. 5, 485–517. doi:10.1142/S0129055X0600267X
- [24] Mukherjee, C.; Varadhan, S. R. S. Identification of the Polaron measure in strong coupling and the Pekar variational formula. Preprint, 2018. arXiv:1812.06927 [math.PR]
- [25] Mukherjee, C.; Varadhan, S. R. S. Strong coupling limit of the Polaron measure and the Pekar process. Preprint, 2018. arXiv:1806.06865 [math.PR]
- [26] Nelson, E. Interaction of nonrelativistic particles with a quantized scalar field. *J. Math. Phys.* **5** (1964), 1190–1197. doi:10.1063/1.1704225
- [27] Pekar, S. I. *Untersuchung über die Elektronentheorie der Kristalle*. Akademie, Berlin, 1954.
- [28] Reed, M.; Simon, B. *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press, New York–London, 1978.
- [29] Spohn, H. Effective mass of the polaron: a functional integral approach. *Ann. Physics* **175** (1987), no. 2, 278–318. doi:10.1016/0003-4916(87)90211-9
- [30] Tjablikow, S. W. Adiabatische Form der Störungstheorie im Problem der Wechselwirkung eines Teilchens mit einem gequantelten Feld. *Abhandlungen aus der Sowjetischen Physik, Folge IV*, 54–68. Verlag Kultur und Fortschritt, Berlin, 1954.
- [31] Tod, P.; Moroz, I. M. An analytical approach to the Schrödinger-Newton equations. *Nonlinearity* **12** (1999), no. 2, 201–216. doi:10.1088/0951-7715/12/2/002
- [32] Wei, J.; Winter, M. Strongly interacting bumps for the Schrödinger-Newton equations. *J. Math. Phys.* **50** (2009), no. 1, 012905, 22 pp. doi:10.1063/1.3060169

RUPERT L. FRANK
 Mathematisches Institut
 Ludwig-Maximilians Universität
 München
 Theresienstr. 39
 80333 München
 GERMANY
 and
 Department of Mathematics
 California Institute of Technology
 Pasadena, CA 91125
 USA
 E-mail: r.frank@lmu.de,
 rlf Frank@caltech.edu

ROBERT SEIRINGER
 IST Austria (Institute of Science
 and Technology Austria)
 Am Campus 1
 3400 Klosterneuburg
 AUSTRIA
 E-mail: robert.seiringer@
 ist.ac.at

Received February 2019.