

Association of IceCube neutrinos with radio sources observed at Owens Valley and Metsähovi Radio Observatories

T. Hovatta^{1,2}, E. Lindfors¹, S. Kiehlmann^{3,4}, W. Max-Moerbeck⁵, M. Hodges⁶, I. Lioudakis¹, A. Lähteemäki^{2,7}, T. J. Pearson⁶, A. C. S. Readhead⁶, R. A. Reeves⁸, S. Suutarinen², J. Tammi², and M. Tornikoski²

¹ Finnish Centre for Astronomy with ESO, FINCA, University of Turku, Finland
e-mail: talvikki.hovatta@utu.fi, elilin@utu.fi

² Aalto University Metsähovi Radio Observatory, Metsähovintie 114, FI-02540 Kylmälä, Finland

³ Institute of Astrophysics, Foundation for Research and Technology-Hellas, GR-71110 Heraklion, Greece

⁴ Department of Physics, Univ. of Crete, GR-70013 Heraklion, Greece

⁵ Departamento de Astronomía, Universidad de Chile, Camino El Observatorio 1515, Las Condes, Santiago, Chile

⁶ Owens Valley Radio Observatory, California Institute of Technology, Pasadena, CA 91125, USA

⁷ Aalto University Department of Electronics and Nanoengineering, PO Box 15500, 00076 Aalto, Finland

⁸ CePIA, Astronomy Department, Universidad de Concepción, Casilla 160-C, Concepción, Chile

Received 21 September 2020; accepted

ABSTRACT

Context. Identifying the most likely sources for high-energy neutrino emission has been one of the main topics in high-energy astrophysics ever since the first observation of high-energy neutrinos by the IceCube Neutrino Observatory. Active galactic nuclei with relativistic jets, blazars, have been considered to be one of the main candidates due to their ability to accelerate particles to high energies.

Aims. We study the connection between radio emission and IceCube neutrino events using data from the Owens Valley Radio Observatory and Metsähovi Radio Observatory blazar monitoring programs.

Methods. We identify sources in our radio monitoring sample, which are positionally consistent with IceCube high-energy neutrino events. We estimate their mean flux density and variability amplitudes around the neutrino arrival time, and compare these with values from random samples to establish the significance of our results.

Results. We find radio source associations within our samples with 16 high-energy neutrino events detected by IceCube. Nearly half of the associated sources are not detected in the γ -ray energies, but their radio variability properties and Doppler boosting factors are similar to the γ -ray detected objects in our sample so that they could still be potential neutrino emitters. We find that the number of strongly flaring objects in our samples is unlikely to occur due to a random coincidence (at 2σ level), and in the case of OVRO samples, the sample of associated sources is on average at an active state compared to random samples.

Conclusions. Based on our results we conclude that although it is clear that not all neutrino events are associated with strong radio flaring blazars, when we see large amplitude radio flares in a blazar at the same time as a neutrino event, it is unlikely to happen by random coincidence.

Key words. neutrinos, BL Lacertae objects: general, quasars: general, galaxies: jets, radio continuum: galaxies

1. Introduction

Blazars, active galactic nuclei (AGNs) with the relativistic jet pointing close to our line of sight, are among the main candidate sources of astrophysical high energy neutrinos. The first γ -ray detection of blazar 3C 279 (Hartman et al. 1992) triggered the first estimates of neutrino fluxes from blazars assuming the γ -ray emission was of hadronic origin (Mannheim et al. 1992), and many followed (e.g. Mücke & Protheroe 2001).

The first observation of high energy neutrinos by the IceCube Neutrino Observatory (Aartsen et al. 2013; IceCube Collaboration 2013) immediately triggered searches for counterpart sources (e.g. Padovani & Resconi 2014), but no significant association with γ -ray blazars was found (Brown et al. 2015; Aartsen et al. 2017a). In September 2017, an event designated IceCube-170922A was observed. The best-fit reconstructed direction was at 0.1° from the sky position of the BL Lac object TXS 0506+056, which was flaring in the γ -ray and VHE

γ -ray bands (Tanaka et al. 2017; Mirzoyan 2017). The positional and temporal coincidences of the events were high, which suggests that the neutrino originated from this flaring blazar (IceCube Collaboration et al. 2018b). Several papers after the event showed that sufficient neutrino flux to explain the detection of the event could be produced within the source, even if the leptonic processes dominated the electromagnetic emission even in γ -rays with only sub-dominant contribution from cascade emission in VHE γ -rays (see e.g. Ansoldi et al. 2018; Petropoulou et al. 2020).

While the association of neutrino events and γ -ray emission has been getting plenty of attention, the connection has also proven to be very complicated. The regions that would host sufficiently dense external photon fields to produce high neutrino fluxes (such as observed from TXS 0506+056 in 2014–2015 (IceCube Collaboration et al. 2018a)) would also imply strong internal absorption of γ -rays (see e.g. Böttcher 2019; Reimer et al. 2019), and therefore no temporal coincidence would be ex-

pected. Indeed, no γ -ray flare was detected from TXS 0506+056 during the 2014-2015 neutrino burst (IceCube Collaboration et al. 2018a). Based on a sample of three sources, Kun et al. (2020) also suggest that the γ -ray emission may be suppressed, as they see a local decrease in the γ -ray flux during the time of the neutrino arrival.

It is also possible that not all blazar classes are equally "good" neutrino candidates (e.g. Neronov & Semikoz 2002). Blazars are classified as flat spectrum radio quasars (FSRQs) and BL Lacertae objects (BL Lacs) based on optical spectra. They can also be classified based on their synchrotron peak frequency. The low synchrotron peak sources (LSPs) have $\nu_S < 10^{14}$ Hz, the intermediate synchrotron peak sources (ISPs) 10^{14} Hz $< \nu_S < 10^{15}$ Hz and high synchrotron peak sources (HSPs) $\nu_S > 10^{15}$ Hz (Abdo et al. 2010). In FSRQs, the synchrotron peak is typically in the infrared regime (i.e. they are LSPs) while for BL Lacs it can be anywhere between infrared and hard X-rays (LBLs, IBLs and HBLs). Padovani et al. (2016) found the most significant connection with HSP sources. The most recent revision (with more data) suggested a connection with ISP and HSP sources (Giommi et al. 2020). On the other hand, Huber (2019) found the most significant connection (though only at 1.9σ level) with low and intermediate synchrotron peaked BL Lac objects (LBL/IBLs) and the connection to high synchrotron peaked sources had a much lower significance (only 0.5σ).

In this paper we investigate the connection between radio bright AGNs and astrophysical high energy neutrinos. In AGN jets, the radio emission is synchrotron self-absorbed close to the central engine and therefore the emission we observe originates several parsecs downstream the jet (Readhead et al. 1978; Blandford & Königl 1979). However, radio emission is a good proxy for the general jet activity, for example, radio flux densities are often seen to increase already before the flaring at higher energy bands begins (e.g. Lähteenmäki & Valtaoja 2003; León-Tavares et al. 2011). This could be, for example, due to a long-term increase in the total jet power. The increasing jet power would trigger current-driven kink instabilities and lead to strong magnetic dissipation and efficient particle acceleration (Nalewajko 2017), which is crucial also in neutrino production. Furthermore, even though the fourth catalogue of AGNs (4LAC) detected by *Fermi*-LAT γ -ray satellite includes 3207 objects (The Fermi-LAT collaboration 2019), not all blazars are γ -ray detected. For example, only 444 of the 1157 sources (38%) in the Candidate Gamma-Ray Blazar Survey (CGRaBS) sample (Healey et al. 2008) currently monitored at Owens Valley Radio Observatory, are included in the 4LAC catalog.

There have been some reports on the connection between radio-selected blazars inside IceCube neutrino positional error circles. Krauß et al. (2014) studied the six radio-brightest blazars within the error regions of PeV IceCube neutrino events. They found that the six sources alone were sufficient to explain the PeV neutrino flux. Kadler et al. (2016) discovered PKS B1424-418 flaring in the radio and γ -ray bands in temporal and positional coincidence with the IceCube neutrino IC35, but this neutrino event had a large positional error circle of 15.6° . The first systematic search of radio sources was performed only recently, by Plavin et al. (2020a), who investigated the association of neutrinos with radio bright AGNs using the VLBI based radio fundamental catalogue (RFC)¹ and RATAN-600 (Korolkov & Pariiskii 1979) data. They found that AGNs positionally associated with high-energy IceCube events typically have stronger parsec-scale cores compared to the rest of the sample. Moreover,

they found an average increase of radio emission at frequencies above 10 GHz around neutrino arrival times for several AGNs on the basis of RATAN-600 monitoring. In Plavin et al. (2020b) they further analyzed the IceCube point source likelihood map, including also lower energy events, against the RFC catalogue and found a 3σ significance for the correlation between them.

In this paper we investigate the connection between radio emission and neutrinos using the high cadence long-term monitoring data of the Owens Valley Radio Observatory (Richards et al. 2011) and Metsähovi Radio Observatory (Teräsraanta et al. 2004). Our paper is organized as follows. In Sect. 2, we describe the selection of the IceCube neutrino events and our radio samples. Our analysis methods and results are given in Sect. 3 and we discuss them in the context of other recent studies in Sect. 4. Our conclusions are drawn in Sect. 5.

2. Data

2.1. Neutrino events

High energy neutrinos can interact with ice creating high energy leptons via charged current interactions. For association with astrophysical sources the track-like events created by high energy muons are particularly important, as they have the smallest error circle around the arrival direction. The angular resolution can be as good as 0.4° (Aartsen et al. 2017a). The positional errors are different in right ascension (RA) and declination (DEC). There is also some level of uncertainty in the detector systematic uncertainties and therefore in the reconstruction errors, which is why we use the published upper limit of 1.0° (IceCube Collaboration 2013) in our analysis. However, following Plavin et al. (2020a), we estimate the optimal systematic uncertainty using our data (see Sect. 3).

While most of the events are associated with the atmospheric background, some high-energy events have a good chance of being of astrophysical origin. In general, events with energy > 200 TeV have likelihood $> 50\%$ of being of astrophysical origin (IceCube Collaboration et al. 2019). There are also indications that the observed neutrino spectrum hardens above 200 TeV (IceCube Collaboration et al. 2017), while below 200 TeV there could also be a contribution from the Galactic component (Palladino & Vissani 2016).

The IceCube Collaboration has published lists of high energy neutrino candidates (Aartsen et al. 2014; The IceCube Collaboration et al. 2015; Aartsen et al. 2016; IceCube Collaboration et al. 2017, 2018b) and real time alerts², see also https://icecube.wisc.edu/science/data/TXS0506_alerts. The high-energy neutrino candidates come from several different analysis chains: the muon tracks (MUONT), high energy starting events (HESEA), extremely high energy alerts (EHEA) and the GOLD and BRONZE events that replaced the HESEA and EHEA chains in mid 2019. These lists were used as a starting point in Plavin et al. (2020a) and the above described considerations led them to perform a cut in positional accuracy and energy. The cuts were performed at the 90% containment area on the celestial sphere $\Omega_{90} < 10 \text{ deg}^2$ and $E > 200$ TeV, and only track-like events were considered. For EHEA neutrino alerts, energies were not published, but by definition these events have extremely high energy (the pipeline is optimized for events with $E \geq 500$ TeV) (Aartsen et al. 2017c) so we include all of them in our event selection. We follow almost the same criteria as Plavin et al. (2020a) and therefore the list of neutrinos used in our work is almost identical to theirs.

¹ http://astrogeo.org/vlbi/solutions/rfc_2019c/

² <https://gcn.gsfc.nasa.gov/amon.html>

We have only two additional neutrino events in our list. One is the event on May 15, 2012 (MJD 56062) exactly at $E=200\text{TeV}$, which was excluded from Plavin et al. (2020a) because they used a non-inclusive cutoff $E > 200\text{ TeV}$, while we use $E \geq 200\text{ TeV}$. We also added another neutrino event to our sample from January 15, 2020 (MJD 58863). This event was a BRONZE alert and had an extremely large deposited charge in the detector, and was consequently reconstructed with a likely neutrino energy in excess of 6 PeV^3 . We did not consider neutrino events after May 2020 since for these events analyzing the radio activity around the neutrino event with a reasonable time window would be impossible. We also exclude one event at $\text{DEC} < -86^\circ$ because sources at this low declination are not included in OVRO or Metsähovi monitoring. In total our sample includes 57 neutrino events, which are listed in Table 1.

2.2. OVRO data and samples

The OVRO blazar monitoring program (Richards et al. 2011) has been continuously on-going since 2008. It uses the 40 m Telescope at Owens Valley Radio Observatory (37 d 14' 02" N, 118 d 16' 56" W). The OVRO 40 m uses two symmetric off-axis beams and a cryogenic receiver with a 15.0 GHz center frequency and 3 GHz bandwidth. The source is alternated between the two beams in an ON-ON fashion to remove atmospheric and ground contamination. The receiver was replaced in May 2014 with a dual-beam correlation receiver instead of the original Dicke switched receiver, providing better performance and higher time stability. Calibration is achieved using a temperature-stable diode noise source to remove receiver gain drifts and the flux density scale is determined from observations of 3C 286 assuming the Baars et al. (1977) value of 3.44 Jy at 15.0 GHz. The systematic uncertainty of about 5% in the flux density scale is not included in the error bars. Complete details of the reduction and calibration procedure are given in Richards et al. (2011).

Originally the monitoring included only sources from the CGRaBS sample, which is a catalog of AGN with radio and X-ray properties similar to EGRET γ -ray blazars (Healey et al. 2008). CGRaBS sample objects are derived from a parent population of flat-spectrum radio sources ($\alpha > -0.5$) with 4.8 GHz flux density $> 65\text{ mJy}$. In the OVRO monitoring 1158 sources at $\text{Dec} > -20^\circ$ were included. One faint source CGRaBS J1310+3233 was later removed from the sample because it is exactly 13 arcmin distant from another bright CGRaBS target CGRaBS J1310+3220, which corresponds to the separation of the two beams of the receiver, resulting in confusion in its fluxes. The final CGRaBS sample used in this paper consists of 1157 objects (hereafter CGRaBS sample).

In addition to the CGRaBS sample, the OVRO monitoring sample has subsequently been increased by adding γ -ray detected sources from the 1st *Fermi*-LAT AGN catalog (1LAC, Abdo et al. 2010) and 2LAC (Ackermann et al. 2011). Additionally, the source list includes various other AGN and blazars "of interest" so that the total number of monitored AGN is currently 1795 (hereafter all-AGN sample).

We note that the full OVRO sample is not statistically well defined, but based on a source-count distribution, Lioudakis et al. (2017) determined that it is complete down to a flux density limit of 350 mJy when using the maximum likelihood mean flux density at 15 GHz from Richards et al. (2014). The sample in

Richards et al. (2014) includes only objects in the CGRaBS catalog and the 1LAC catalog at $\text{DEC} > -20^\circ$, and excludes objects near the Galactic plane with Galactic latitude $|b| < 10^\circ$. There are 589 objects with maximum likelihood mean flux density greater than 350 mJy forming the flux density limited sample (hereafter OVRO-350mJy sample). However, we note that because the source-count distribution is not derived from a full sky survey, there are still individual sources brighter than this limit missing from our sample.

Plavin et al. (2020a) used the RFC catalog down to its completeness limit of 150 mJy of unresolved flux density at 8 GHz in their analysis (completeness also determined from a source-count distribution). For direct comparison with their results, we have cross-matched our full sample with the RFC and use the resulting 1156 objects in our sample (hereafter RFC-150mJy sample). Because the first IceCube neutrino event meeting our selection criteria is from 2009, in our analysis we use data from Jan 1, 2008 until Jul 1, 2020.

2.3. Metsähovi data and sample

A sample of over 1000 AGN is being monitored at Aalto University Metsähovi Radio Observatory in Finland (60 d 13' 04" N, 24 d 23' 35" E). The monitored sample is heterogeneous. The current high priority list contains over 400 sources where the aim is to observe them regularly, and weekly observations are being done for about a hundred sources, for some of which the data sets run for over 40 years. At the flux density limit of 2 Jy at $\text{DEC} > 0^\circ$ the sample can be considered complete (Valtaoja et al. 1992).

The Metsähovi 14-m telescope has a radome and Cassegrain optics. The measurements were made with a 1 GHz-band dual beam receiver centered at 36.8 GHz. The HEMT (high electron mobility transistor) front end operates at ambient temperature. The observations are Dicke switched ON-ON observations, alternating the source and the sky in each feed horn. A typical integration time to obtain one flux density data point is between 1200 and 1600 s. The detection limit of the telescope at 37 GHz is on the order of 0.2 Jy under optimal conditions, but it is highly weather-dependent. Data points with a signal-to-noise ratio < 4 are handled as non-detections.

The flux density scale is set by observations of the HII region DR21. Sources NGC 7027, 3C 274 and 3C 84 are used as secondary calibrators. A detailed description of the data reduction and analysis is given in Teräsraanta et al. (1998). The error estimate in the flux density includes the contribution from the measurement rms and the uncertainty of the absolute calibration.

We selected sources that have at least 6 years of data between Jan 1, 2008 and Feb 29, 2020. Additionally, we require that there are at least 10 detections (signal-to-noise ratio ≥ 4) during that time period. We also applied a cut on the flux density to exclude sources that are too close to the detection threshold, and only included sources that had a maximum flux $> 0.7\text{ Jy}$. Applying these criteria to the Metsähovi database resulted in 183 sources. The source with the lowest declination is at $\text{DEC} = -13.08^\circ$, which means that there are two neutrino events that are excluded from the comparison with the Metsähovi sample, when the uncertainties and the maximum systematic uncertainty of 1° in the neutrino event positions is accounted for.

3. Analyses and results

We have looked at three different metrics when studying the association of IceCube neutrinos with radio data.

³ https://gcn.gsfc.nasa.gov/notices_amon_g_b/133644_43767651.amon

Table 1. IceCube neutrino events at $E \geq 200$ TeV used in our analyses

Date	MJD	Type	RA (deg.)	RA Error (deg.)	DEC (deg.)	DEC Error (deg.)	Energy (TeV)	Reference
2009-08-13	55056	MUONT	29.51	+0.40	-0.38	1.23	+0.18 -0.22	480 1
2009-11-06	55141	MUONT	298.21	+0.53	-0.57	11.74	+0.32 -0.38	250 1
2010-06-23	55370	MUONT	141.25	+0.46	-0.45	47.80	+0.56 -0.48	260 1
2010-09-25	55464	MUONT	266.29	+0.58	-0.62	13.40	+0.52 -0.45	460 1
2010-10-09	55478	EHEA	331.09	+0.56	-0.72	11.10	+0.48 -0.58	- 1,2
2010-10-28	55497	EHEA	88.68	+0.54	-0.55	0.46	+0.33 -0.27	- 1,2
2010-11-13	55513	MUONT	285.95	+1.29	-1.50	3.15	+0.70 -0.63	520 1
2011-01-28	55589	EHEA	307.53	+0.82	-0.81	1.19	+0.35 -0.32	- 1,2
2011-03-04	55624	EHEA	116.37	+0.73	-0.73	-10.72	+0.57 -0.65	- 2
2011-05-21	55702	MUONT	235.13	+2.70	-1.76	20.30	+0.44 -0.62	300 1
2011-06-10	55722	MUONT	272.22	+1.23	-1.19	35.55	+0.28 -0.29	210 1
2011-07-14	55756	HESEA	67.86	+0.51	-0.72	40.32	+0.73 -0.25	253 2,3
2011-09-30	55834	EHEA	266.48	+2.09	-1.55	-4.41	+0.59 -0.86	- 2
2012-03-01	55987	EHEA	238.01	+0.60	-0.59	18.60	+0.46 -0.39	- 2
2012-05-15	56062	MUONT	198.74	+1.44	-1.09	31.96	+0.81 -0.58	200 1
2012-05-23	56070	EHEA	171.03	+0.81	-0.90	26.36	+0.49 -0.30	- 2
2012-08-07	56146	MUONT	330.10	+0.65	-0.82	1.57	+0.46 -0.42	260 1
2012-09-22	56192	EHEA	70.75	+1.56	-1.63	19.79	+1.37 -0.68	- 2
2012-10-11	56211	EHEA	205.22	+0.59	-0.65	-2.39	+0.51 -0.57	- 1,2
2012-10-26	56226	MUONT	169.61	+1.16	-1.11	28.04	+0.67 -0.66	750 1
2013-06-27	56470	HESEA	93.43	+0.80	-0.85	14.02	+0.72 -0.75	200 2
2013-08-17	56521	MUONT	224.89	+0.87	-1.19	-4.44	+1.21 -0.94	400 1
2013-09-07	56542	EHEA	129.81	+0.48	-0.28	-10.36	+0.36 -0.31	- 2
2013-10-14	56579	MUONT	32.94	+0.63	-0.62	10.20	+0.34 -0.49	390 1
2013-10-23	56588	EHEA	301.82	+1.10	-0.93	11.49	+1.19 -1.09	- 2
2013-12-04	56630	EHEA	289.16	+1.08	-0.94	-14.25	+0.91 -0.81	- 2
2014-01-08	56665	EHEA	344.53	+0.67	-0.48	1.57	+0.35 -0.32	- 2
2014-01-09	56666	EHEA	292.85	+0.87	-0.94	33.06	+0.50 -0.46	- 1,2
2014-02-03	56691	EHEA	349.54	+2.21	-1.97	-13.71	+1.23 -1.38	- 2
2014-06-09	56817	MUONT	106.26	+2.27	-1.90	1.29	+0.83 -0.74	340 1
2014-06-11	56819	EHEA	110.30	+0.66	-0.45	11.57	+0.14 -0.24	- 1,2
2014-09-23	56923	EHEA	169.72	+0.91	-0.86	-1.34	+0.73 -0.66	- 2
2015-01-27	57049	MUONT	100.48	+0.95	-1.87	4.56	+0.68 -0.50	210 1
2015-05-15	57157	MUONT	91.60	+0.16	-0.74	12.18	+0.37 -0.35	240 1
2015-07-14	57217	MUONT	325.50	+1.77	-1.46	26.10	+1.68 -1.85	300 4
2015-08-12	57246	EHEA	328.19	+1.01	-1.03	6.21	+0.44 -0.49	- 2,5
2015-08-31	57265	EHEA	54.85	+0.94	-0.98	33.96	+1.07 -1.19	- 2
2015-09-04	57269	MUONT	134.00	+0.39	-0.58	28.00	+0.47 -0.47	220 5
2015-09-23	57288	EHEA	103.27	+0.70	-1.36	3.88	+0.59 -0.71	- 2
2015-09-26	57291	EHEA	194.50	+0.76	-1.21	-4.34	+0.70 -0.95	- 2
2015-11-14	57340	MUONT	76.30	+0.75	-0.74	12.60	+0.61 -0.58	740 5
2015-11-22	57348	EHEA	262.18	+0.90	-1.21	-2.38	+0.73 -0.43	- 2
2016-01-28	57415	EHEA	263.40	+1.35	-1.18	-14.79	+0.99 -1.02	- 2
2016-03-31	57478	MUONT	15.60	+0.45	-0.58	15.60	+0.53 -0.60	380 4
2016-05-10	57518	EHEA	352.34	+1.63	-1.31	2.09	+0.99 -0.85	- 2
2016-07-31	57600	EHEA	214.50	+0.75	-0.75	-0.33	+0.75 -0.75	- 2,5
2016-08-06	57606	EHEA	122.81	+0.50	-0.50	-0.81	+0.50 -0.50	- 2
2016-12-10	57732	EHEA	46.58	+1.10	-1.00	14.98	+0.45 -0.40	- 2
2017-03-21	57833	EHEA	98.30	+1.20	-1.20	-15.02	+1.20 -1.20	- 2
2017-09-22	58018	EHEA	77.43	+0.95	-0.65	5.72	+0.50 -0.30	290 6
2017-11-06	58063	EHEA	340.00	+0.70	-0.50	7.40	+0.35 -0.25	- 7
2018-09-08	58369	EHEA	144.58	+1.55	-1.45	-2.13	+0.90 -1.20	- 7
2018-10-23	58414	EHEA	270.18	+2.00	-1.70	-8.57	+1.25 -1.30	- 7
2019-03-31	58573	HESEA	337.68	+0.23	-0.34	-20.70	+0.30 -0.48	1987 7
2019-05-03	58606	EHEA	120.28	+0.57	-0.77	6.35	+0.76 -0.70	- 7
2019-07-30	58694	GOLD	225.79	+1.28	-1.43	10.47	+1.14 -0.89	299 7
2020-01-15	58863	BRONZE	67.46	+0.34	-0.34	-14.63	+0.34 -0.34	6000 7

References. (1) Aartsen et al. (2016); (2) IceCube Collaboration (2018); (3) The IceCube Collaboration et al. (2015); (4) IceCube Collaboration et al. (2019); (5) IceCube Collaboration et al. (2017); (6) IceCube Collaboration et al. (2018b); (7) <https://gcn.gsfc.nasa.gov/amon.html>

1. Mean flux density of the associated sources.
2. Mean activity index of the associated sources.
3. Number of flaring sources in the sample.

In Sect. 3.1, we describe how we match the neutrino positions with our radio samples. Following Plavin et al. (2020a), we estimate the (unknown) systematic uncertainties in the IceCube event positions using our data. We also describe how we generate random Monte Carlo samples to obtain the chance coincidence probability in our analysis. In Sect. 3.2 we discuss the analysis of the mean flux density, while the mean activity index and the number of flaring sources in the sample are discussed in Sect. 3.3.

3.1. Position matching and systematic uncertainty estimation

As described in Plavin et al. (2020a), in order to match the neutrino positions with our samples, we must first translate the Ω_{90} error regions into two-dimensional coverage regions. This is done by multiplying the individual statistical errors in RA and DEC with the ratio of 90% quantiles of two- and one-dimensional Gaussian distributions: $\frac{\sqrt{-\log(1-0.9)}}{\text{erf}^{-1}(0.9)} \approx 1.3$ (Plavin et al. 2020a). This way we obtain regions around the neutrino events that are bounded by four quarters of ellipses.

Following Plavin et al. (2020a), we estimate the unknown additional systematic uncertainty in the neutrino positions using our data. We leave the systematic uncertainty $\Delta\psi$ free in our analysis, and scan over a range from 0.1° to 1.0° with a step of $\Delta\psi = 0.1^\circ$. We then add $\Delta\psi$ to the statistical uncertainties in RA and DEC in all directions.

For each $\Delta\psi$ we first find all the sources in our samples that fall within the elliptical region around the neutrino event (see Fig. 1 for an example). We then calculate the test statistic of interest (mean flux density or activity index) for each source. We repeat this step for random Monte Carlo samples. Similar to Plavin et al. (2020a), we generate random comparison samples by shifting the IceCube neutrino positions in RA while keeping the DEC constant. This way we reproduce the effect that the sensitivity of the instrument depends only on zenith angle (Aartsen et al. 2017b). We then match these random neutrino samples with our observed samples, and calculate the same test statistic (mean flux density and activity index) for each associated random source.

With this procedure we can estimate the distribution of the test statistic under the null hypothesis of no association between the neutrino events and the radio sources. We can then estimate, under the null hypothesis, the chance probability of obtaining a value of the test statistic equal or larger than the one obtained from the data as,

$$p = \frac{M + 1}{N + 1}, \quad (1)$$

where M is the number of random samples with a larger test statistic than in the real data, and N is the number of random samples (Davison & Hinkley 2013), which in our case is 10 000. This is given as the pre-trial p-value when reporting our results. We then select as the optimal systematic uncertainty the value that gives the smallest pre-trial p-value.

In order to account for the factor resulting from these multiple trials, we again follow Plavin et al. (2020a) and insert the calculation of the post-trial p-value into our Monte Carlo simulations. To do this, we treat each random sample as the real observation, and calculate how many times we obtain as small




Fig. 1. Example of the elliptical uncertainty region around the neutrino event on Aug 12, 2015 (MJD 57246) and the positions of the three associations found for it in the OVRO sample. The solid black line shows the statistical error ellipse of the neutrino position (shown with a black cross), where the quarters are asymmetric in all directions. The blue dashed line shows the additional systematic uncertainty $\Delta\psi = 0.6^\circ$ as found optimal for the mean flux density analysis in the CGRaBS sample.

pre-trial p-value by chance as is obtained for the real data. This will be M in Eq. 1, which we now use to obtain the post-trial p-value.

3.2. Mean flux density

The mean flux densities of the associated sources in the different samples are given in Table 2 for the OVRO samples and Table 3 for the Metsähovi sample. The optimal systematic uncertainty $\Delta\psi$ varies between 0.3° and 0.6° for the different samples due to the random nature of the process. These are, however, similar to values found by Plavin et al. (2020a). The optimal value and the number of associations within this limit are given in Table 4. When comparing the mean flux density values with random samples, we average the values of the individual sources within the sample of associated sources to obtain a single parameter for the sample. These are also given in Table 4, along with the resulting pre-trial and post-trial p-values. The OVRO and Metsähovi light curves of these sources are shown in Appendix A.

3.3. Time-Dependent analysis

3.3.1. Mean activity index

Because both OVRO and Metsähovi light curves are well sampled, we can estimate whether any of the associated blazars show flaring behavior during the arrival time of the neutrino. Following Plavin et al. (2020a), we use the activity index (A.I.), defined as the mean flux density within a time window around the neu-

Table 2. Mean flux density (Sect. 3.2) of OVRO AGN within the error regions of IceCube events

IceCube Event		AGN								
MJD	Category	J2000 Name	Other Name	RA	DEC	all	CGR	350	150	< S >
(1)	(2)	(3)	(4)	(deg.)	(deg.)	(7)	(8)	(9)	(10)	(Jy)
				(5)	(6)					(11)
55478.38058	EHEA	J2203+1007	B2201+098	330.8790	10.1283	X	X	...	X	0.13
		J2200+1030	TXS 2157+102	330.0330	10.5021	X	0.22
55834.44514	EHEA	J1743-0350	PKS 1741-038	265.9950	-3.8344	X	X	X	X	4.24
56062.96000	MUONT	J1310+3220	B1308+326	197.6190	32.3453	X	X	X	X	2.08
56070.57428	EHEA	J1125+2610	PKS 1123+264	171.4740	26.1722	X	X	X	X	0.88
56146.21000	MUONT	J2200+0234	GB6 B2158+0219	330.2060	2.5750	...	X	0.10
56579.91000	MUONT	J0211+1051	GB6 B0208+1037	32.8049	10.8594	X	X	X	X	0.90
56630.47006	EHEA	J1916-1519	B1914-154	289.2190	-15.3167	X	X	...	X	0.21
57246.75909	EHEA	J2151+0552	PKS 2149+056	327.9080	5.8703	X	X	X	X	0.40
		J2148+0657	PKS 2145+067	327.0230	6.9606	...	X	...	X	3.99
		J2151+0709	PKS 2149+069	327.8810	7.1572	...	X	...	X	0.84
57265.21782	EHEA	J0341+3352	TXS 0337+337	55.2916	33.8725	X	X	0.10
		J0336+3218 ^a	B0333+321	54.1254	32.3082	X	X	1.86
57291.90119	EHEA	J1256-0547	3C279	194.0470	-5.7892	X	X	X	X	18.02
57340.90000	MUONT	J0502+1338	PKS 0459+135	75.6384	13.6361	...	X	...	X	0.44
57415.18346	EHEA	J1724-1443	TXS 1721-146	261.1960	-14.7331	...	X	...	X	0.41
		J1733-1304	PKS 1730-130	263.2610	-13.0803	...	X	...	X	4.49
57833.31413	EHEA	J0631-1410	TXS 0629-141	97.8343	-14.1753	X	X	X	X	0.59
		J0630-1323	TXS 0628-133	97.7246	-13.3928	X	X	X	X	0.43
58018.87118	EHEA	J0509+0541	TXS 0506+056	77.3582	5.6931	X	X	X	X	0.69
58063.65254	EHEA	J2238+0724	GB6 B2235+0708	339.5430	7.4039	X	X	...	X	0.16
58694.74353	GOLD	J1504+1029	PKS 1502+106	226.1040	10.4942	X	X	X	X	2.07

Notes. Cols (5) and (6) give the coordinates of the associated AGN. Cols 7-10 indicate the sample where the match is found. All for all-AGN, CGR for CGRaBS, 350 for OVRO-350mJy, and 150 for RFC-150mJy samples. The optimal systematic uncertainty was 0.6° for the CGRaBS and RFC samples, and 0.3° for all-AGN and OVRO-350mJy samples. Col (11) gives the mean flux density of the source.

^(a) Although this source is brighter than 350 mJy, it is not included in the OVRO-350mJy sample, because it is not included in the sample of Richards et al. (2014), which we use to derive our flux-density limited sample.

Table 3. Mean flux density (Sect. 3.2) and activity index (Sect. 3.3) of Metsähovi AGN within the error regions of IceCube events

IceCube Event		AGN					
MJD	Category	J2000 Name	Other Name	RA	DEC	< S >	A.I.
(1)	(2)	(3)	(4)	(deg.)	(deg.)	(Jy)	(8)
				(5)	(6)	(7)	
55834.44514	EHEA	J1743-0350	PKS 1741-038	265.9953	-3.8346	3.61	1.36
56062.96000	MUONT	J1310+3220	B1308+326	197.6194	32.3455	1.76	1.28
		J1310+3233 ^a	B1308+328	197.7475	32.5596	0.49	0.81
56579.91000	MUONT	J0211+1051	GB6 B0208+1037	32.8049	10.8597	1.15	2.04
57265.21782	EHEA	J0336+3218	B0333+321	54.1255	32.3081	1.58	0.55
57291.90119	EHEA	J1256-0547	3C279	194.0465	-5.7893	19.07	0.89
58018.87118	EHEA	J0509+0541	TXS 0506+056	77.3579	5.6931	1.17	0.42
58694.74353	GOLD	J1504+1029	PKS1502+106	226.1041	10.4942	2.05	1.69

Notes. Cols (5) and (6) give the coordinates of the associated AGN. Col (7) gives the mean flux density and col (8) the activity index of the source. Sources marked with bold-face font in Col. 8 are those showing strong flaring activity (Sect. 3.3.2)

^(a) This source is the only one of the Metsähovi associations not included in the OVRO samples. It is included in the 4LAC catalog and its optical class is FSRQ and SED class LSP.

trino event divided by the mean flux density outside this time window. Plavin et al. (2020a) set the time window as a free parameter similar to the systematic uncertainty. They then selected the time window maximising the mean activity index of their real sample as the optimal time window, which they found to be 0.9 yrs for the RATAN-600 sample.

We explored this approach and we found that it does not work for either the OVRO or the Metsähovi data. The reason

is that with the well-sampled light curves from OVRO and Metsähovi, leaving the length of the time window free causes a few individual blazars with large flares around a neutrino event to dominate the optimal time window size. For example, in the OVRO data for TXS 0506+056 (J0509+0541) the optimal time window (ΔT) maximising the activity index is 5.6 yrs, simply because at this limit, the window around the neutrino event (calculated as $\pm\Delta T/2$) includes the peak of the large flare following

Table 4. Chance coincidence for the mean flux density analysis

Sample	N_{Source}	$\Delta\psi$ (deg)	N_{Assoc}	$\langle S \rangle$ (Jy)	pre-trial p	post-trial p
(1)	(2)	(3)	(4)	(5)	(6)	(7)
OVRO all-AGN	1795	0.3	15	2.19	0.012	0.025
CGRaBS	1157	0.6	20	2.06	0.015	0.037
OVRO-350mJy	589	0.3	10	3.03	0.061	0.137
RFC-150mJy	1156	0.6	20	2.15	0.035	0.079
Metsähovi	183	0.3	8	3.86	0.089	0.183

Notes. Col (1) indicates the sample studied and col (2) gives the number of sources in that sample. Col (3) gives the optimal systematic uncertainty for the sample, and col (4) the number of associations found using this systematic uncertainty. The pre-trial and post-trial p-values are given in cols (6) and (7), respectively. Post-trial values significant at the 2σ level are shown in bold-face font in Col (7).

the neutrino arrival (see Fig. A.1). We also found that the pre-trial p-value is very insensitive to the length of the time window with values between 0.3–0.4% for all window sizes between 0.4 and 3 yrs for the CGRaBS sample.

Because the optimal window size ΔT for each source clearly depends on how the neutrino event is positioned with respect to large flares, we think that ΔT is more related to the typical flaring time scales of blazars in radio wavelengths than directly to the possible link between neutrinos and radio flares. Therefore, we set the time window ΔT to the median structure function time scale from Hovatta et al. (2007), which can be taken as the typical flaring time scale of blazars. This value is 2.3 yrs at 15 GHz and 1.4 yrs at 37 GHz. In our analysis, we calculate the activity index using a time range of $\pm\Delta T/2$ around the neutrino event.

For the activity index analysis we find the optimal systematic uncertainty $\Delta\psi$ to be 0.4° for the OVRO samples and $\Delta\psi = 0.3^\circ$ for the Metsähovi sample. The activity index for the associated sources in our samples are tabulated in Tables 5 and 3 for the OVRO and Metsähovi samples, respectively.

3.3.2. Number of flaring sources

For our third analysis, we need an objective way to estimate the number of flaring sources. In general, any source with activity index greater than one could be considered to be in a flaring state. However, especially in fainter objects, it is possible to obtain values greater than one simply due to noise in the data. To estimate a threshold for flaring sources that accounts for the typical uncertainties in the OVRO and Metsähovi data, we simulate data of one window length with the same time sampling as the real data whose fluxes are varied around a fixed constant value by the typical uncertainty in our data. This way we get samples with no intrinsic variability, whose only variation is produced by the observing process. We then calculate the resulting A.I. as the mean during this time window over a fixed flux density limit (0.15 Jy for OVRO and 1 Jy for Metsähovi to match a typical fainter target in each sample). We repeat this 100 000 times to obtain a false detection rate.

The limit for obtaining a higher activity index due to noise in 1% of cases is $\text{A.I.} > 1.1$ in the OVRO data and $\text{A.I.} > 1.16$ in the Metsähovi data. A false detection rate of 0.01% is obtained for $\text{A.I.} > 1.29$ for the OVRO data and $\text{A.I.} > 1.71$ for the Metsähovi data. We use these limits to estimate the number of flaring sources in the samples. We then use these same limits for the random samples to obtain the probability of chance coincidence. The results from the comparison to random samples are summarized in Table 6 for the mean activity index and flaring source analysis.

4. Discussion

We have looked at the connection between our radio observations of blazars and IceCube neutrino events by first associating the high-energy neutrino events listed in Table 1 with our monitoring samples at OVRO and Metsähovi. In the following, we first discuss the number of associations in our samples, and the lack of potential counterparts in our samples based on other recent studies. We then discuss the implications of the small number of γ -ray detected sources among our associations. Finally, we discuss our results on the mean flux density and activity index analysis, and their implications for future searches on connection between neutrino events and blazars.

4.1. Number of associated sources

As can be seen from Tables 2, 3, and 5, the number of neutrino events with associated blazars in our samples is in general small, from seven to 16 events (12 – 28%), depending on the sample studied. Although we can expect that only $\sim 50\%$ of the 57 high-energy neutrinos in Table 1 are of astrophysical origin (IceCube Collaboration et al. 2019), the number of associations is still much smaller than that.

One potential explanation is the incompleteness of our monitoring samples. At OVRO, the monitoring began with the CGRaBS sample of 1157 objects, for which we find 14 associated events when using $\Delta\psi = 0.4^\circ$ as found optimal for the activity index analysis (Table 5), and this increases to 16 events when $\Delta\psi = 0.6^\circ$ from the mean flux density analysis is used (Table 2). The number of associated sources is slightly higher than the number of associated events (16 and 20 for the activity index and mean flux density analysis, respectively) because there can be multiple sources within the error ellipses. Interestingly, including all AGN monitored at OVRO does not change the associations drastically, although the number of objects in the sample increases to 1795. There are no additional neutrino events associated with OVRO sources and the increase in number of associated sources is simply due to the increase of sources within the same error regions.

This is interesting considering that most of the additional sources were included in the OVRO monitoring after they were detected by the Fermi-LAT. This may imply that γ -ray activity does not necessarily play a major role in the neutrino association, and indeed recent papers on the association between γ -ray and neutrino emission have been inconclusive (see also Sect. 4.2). Krauß et al. (2018) studied the γ -ray fluxes of all sources coincident with IceCube neutrino events at $E > 100$ TeV, and found no direct correlation between the γ -ray flux and expected neutrino flux. Franckowiak et al. (2020) also studied γ -ray detected

Table 5. Activity index (Sect. 3.3) of OVRO AGN within the error regions of IceCube events

IceCube Event		AGN								
MJD	Category	J2000 Name	Other Name	RA	DEC	all	CGR	350	150	A.I.
(1)	(2)	(3)	(4)	(deg.)	(deg.)	(7)	(8)	(9)	(10)	(11)
				(5)	(6)					
55478.38058	EHEA	J2200+1030 ^a	TXS 2157+102	330.0330	10.5021	X	X	...
		J2203+1007	B2201+098	330.8790	10.1283	X	X	...	X	0.98
55834.44514	EHEA	J1743-0350	PKS 1741-038	265.9950	-3.8344	X	X	X	X	1.20
56062.96000	MUONT	J1310+3220	B1308+326	197.6190	32.3453	X	X	X	X	1.46
56070.57428	EHEA	J1125+2610	PKS 1123+264	171.4740	26.1722	X	X	X	X	1.02
56579.91000	MUONT	J0211+1051	GB6 B0208+1037	32.8049	10.8594	X	X	X	X	1.85
56630.47006	EHEA	J1916-1519	B1914-154	289.2190	-15.3167	X	X	...	X	0.84
57246.75909	EHEA	J2151+0709	PKS 2149+069	327.8810	7.1572	X	X	X	X	0.84
		J2151+0552	PKS 2149+056	327.9080	5.8703	X	X	X	X	1.03
57265.21782	EHEA	J0341+3352	TXS 0337+337	55.2916	33.8725	X	X	1.29
		J0336+3218	B0333+321	54.1254	32.3082	X	X	0.70
57291.90119	EHEA	J1256-0547	3C279	194.0470	-5.7892	X	X	X	X	1.17
57340.90000	MUONT	J0502+1338	PKS 0459+135	75.6384	13.6361	X	X	X	X	1.63
57833.31413	EHEA	J0631-1410	TXS 0629-141	97.8343	-14.1753	X	X	X	X	0.95
		J0630-1323	TXS 0628-133	97.7246	-13.3928	X	X	X	X	1.49
58018.87118	EHEA	J0509+0541	TXS 0506+056	77.3582	5.6931	X	X	X	X	0.99
58063.65254	EHEA	J2238+0724	GB6 B2235+0708	339.5430	7.4039	X	X	...	X	0.96
58694.74353	GOLD	J1504+1029	PKS 1502+106	226.1040	10.4942	X	X	X	X	1.81

Notes. Cols (5) and (6) give the coordinates of the associated AGN. Cols 7-10 indicate the sample where the match is found. All for all-AGN, CGR for CGRaBS, 350 for OVRO-350mJy, and 150 for RFC-150mJy samples. The optimal systematic uncertainty $\Delta\psi$ was 0.4° for all the samples. Sources, for which the activity index is marked with bold-face font in Col. 11 are those showing strong flaring activity (Sect. 3.3.2).

^(a) This source was added to OVRO monitoring only in December 2011 after its inclusion in the 2LAC catalog, which is why there is no activity index available for it around the neutrino event in October 2010.

Table 6. Chance coincidence for the activity index analysis

Sample	N _{Source}	$\Delta\psi$ (deg)	N _{Assoc}	<A.I.>	p pre	p post	N _{flaring} (1%)	p pre	p post	N _{flaring} (0.01%)	p pre	p post
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
OVRO all-AGN	1795	0.4	18	1.19	0.002	0.007	8	0.354	0.514	6	0.026	0.049
CGRaBS	1157	0.4	16	1.22	0.003	0.010	8	0.087	0.166	6	0.007	0.017
OVRO-350mJy	589	0.4	12	1.29	0.006	0.020	7	0.016	0.033	5	0.003	0.006
RFC-150mJy	1156	0.4	17	1.18	0.008	0.028	7	0.174	0.310	5	0.021	0.045
Metsähovi	183	0.3	8	1.13	0.232	0.420	4	0.004	0.007	1	0.076	0.083

Notes. Col (1) indicates the sample studied and col (2) gives the number of sources in that sample. Col (3) gives the optimal systematic uncertainty $\Delta\psi$, which for each sample was found to be the same for all the A.I. analyses. Col (4) gives the number of associations found using this systematic uncertainty. The pre-trial and post-trial p-values are given in cols (6) and (7), respectively. Col (8) gives the number of flaring sources in the sample when 1% false-detection rate is used. The threshold is A.I. > 1.1 for the OVRO samples and A.I. > 1.16 for the Metsähovi sample. Col (11) is the same for 0.01% false-detection rate, which is A.I. > 1.29 for the OVRO samples and A.I. > 1.71 for the Metsähovi sample. Post-trial values significant at the 2σ level are shown in bold-face font in Cols (7) and (13).

sources associated with IceCube neutrino events at $E > 100$ TeV, and found that the results are consistent both with no correlation between the γ -ray and neutrino fluxes, as well as a linear correlation between the two. They also included the OVRO light curves of nine associated sources, and found no direct link between the radio flux density and neutrino arrival times, although this was not quantified.

Table 7 lists the optical classification, redshift, synchrotron peak frequency and luminosity, variability Doppler factor, and radio modulation index of the associated sources in the OVRO sample. The large majority of our 22 associations are FSRQs (15) with only five BL Lacs and two blazars of unknown type (BCU). This reflects the composition of our original CGRaBS sample, where 70% of the sources are FSRQs and only 11% are BL Lacs (Richards et al. 2014), and there is no contradic-

tion with results of Huber (2019) who found the strongest connection between LBLs and IBLs. Interestingly, one of our targets J2148+0657 has been identified in Lioudakis & Petropoulou (2020) as a probable neutrino candidate based on a proton-synchrotron model. Although two other sources are within the error ellipse of IceCube (Fig. 1), confirming J2148+0657 as a neutrino emitter would have interesting implications for our understanding of jet energetics.

As mentioned earlier, Giommi et al. (2020) found in their analysis an excess of ISP and HSP sources in the neutrino associations. We note that although the inclusion of the LAT-detected sources has increased the number of HSP sources in our monitoring sample, most of the sources are still LSP objects (Richards et al. 2014). Although Giommi et al. (2020) also included LSP and ISP objects in their study, they still only considered the γ -ray

Table 7. Properties of the associated OVRO sources

J2000 Name	4FGL Name	Opt. class	z	$\log(\nu_s)$ (Hz)	$\log(\nu F\nu_s)$ ($\text{erg cm}^{-2}\text{s}^{-1}$)	δ_{var}	mod. index
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J0211+1051	4FGL J0211.2+1051	BLL	0.2	14.2	-10.3	$8.4^{+6.0}_{-2.1}$	0.255
J0336+3218	4FGL J0336.4+3224	FSRQ	1.259	13.2	-11.0	$5.5^{+8.8}_{-0.8}$	0.275
J0341+3352		FSRQ	0.725	12.9	-12.6	$5.1^{+3.3}_{-4.1}$	0.227
J0502+1338	4FGL J0502.5+1340	BLL	0.35	13.0	-11.7	$15.8^{+5.4}_{-3.0}$	0.301
J0509+0541	4FGL J0509.4+0542	BLL	0.3365	14.6	-10.7	$14.7^{+3.0}_{-5.1}$	0.167
J0630-1323		FSRQ	1.021	13.6	-12.1	$32.3^{+21.2}_{-30.1}$	0.125
J0631-1410		FSRQ	1.017	12.6	-12.2	$0.6^{+2.7}_{-0.4}$	0.100
J1125+2610		FSRQ	2.341	12.3	-12.1	$56.8^{+27.0}_{-20.3}$	0.140
J1256-0547	4FGL J1256.1-0547	FSRQ	0.5361	12.7	-10.5	$11.6^{+1.8}_{-1.1}$	0.203
J1310+3220	4FGL J1310.5+3221	FSRQ	0.997	12.7	-11.2	$26.4^{+13.4}_{-16.6}$	0.244
J1504+1029	4FGL J1504.4+1029	FSRQ	1.839	12.7	-11.7	$17.7^{+6.4}_{-6.1}$	0.230
J1724-1443		BLL ^a	0.899	13.2	-12.0	$7.9^{+11.5}_{-4.0}$	0.172
J1733-1304	4FGL J1733.0-1305	FSRQ	0.902	12.6	-10.9	$21.5^{+20.3}_{-11.9}$	0.059
J1743-0350	4FGL J1744.2-0353	FSRQ	1.057	12.7	-11.2	$6.4^{+15.3}_{-2.5}$	0.056
J1916-1519	4FGL J1916.7-1516	BCU	...	12.8	-12.0	...	0.185
J2148+0657	4FGL J2148.6+0652	FSRQ	0.99	12.5	-10.7	$1.2^{+0.9}_{-0.4}$	0.201
J2151+0709		FSRQ	1.364	12.8	-12.1	$13.1^{+14.4}_{-4.7}$	0.115
J2151+0552		FSRQ	0.74	12.4	-12.2	$18.0^{+4.3}_{-4.8}$	0.030
J2200+1030	4FGL J2200.3+1029	BLL	0.172	13.0	-12.5	$3.4^{+6.3}_{-2.8}$	0.156
J2200+0234		FSRQ	1.323	13.1	-12.5	$31.1^{+4.1}_{-5.8}$	0.246
J2203+1007		BCU ^b	0.172	12.6	-12.5	$7.6^{+14.6}_{-7.3}$	0.156
J2238+0724		FSRQ	1.011	12.9	-12.0	$16.8^{+7.3}_{-9.7}$	0.347

Notes. Optical classifications in Col (3) are taken from the 4LAC catalog (The Fermi-LAT collaboration 2019) for the sources included in there (ones with name given in Col (2)), and from the 5BZCat catalog (Massaro et al. 2015) for the non-LAT sources unless otherwise stated. BCU stands for blazars of unknown optical class. Col (4) gives the redshift of the source. Cols (5) and (6) give the synchrotron peak frequency and peak luminosity, respectively, taken from 4LAC catalog or determined by ourselves for the non-LAT sources. Variability Doppler factors, given in Col (7) are taken from Liodakis et al. (2018) or determined by ourselves using the same methods. Radio modulation index determined from OVRO data are given in Col (8), and are taken from Liodakis et al. (2017) or determined by ourselves.

^(a) Classification taken from D'Abrusco et al. (2014) ^(b) Classification taken from Sowards-Emmerd et al. (2005)

detected sources when finding objects near the neutrino events, while our samples also include objects not detected in the γ -ray energies. In fact, ten out of the 22 associations listed in Table 7 have not been detected by Fermi-LAT (see also Sect. 4.2).

We took the SED parameters of our γ -ray detected associations from 4LAC (The Fermi-LAT collaboration 2019) and modeled the non γ -ray detected ones ourselves using the Space Science Data Center SED builder⁴. These are also given in Table 7. Indeed a large majority of the sources are LSPs, and there are only two ISPs (TXS 0506+056 and J0211+1051) and no HSPs. When comparing our neutrino sample with the 3HSP sample (Chang et al. 2019), which was used as the parent sample in Giommi et al. (2020), we find that there are eight possible neutrino associations with HSP sources that are missing from our radio monitoring sample.

Plavin et al. (2020a) found that radio sources selected based on their compact 8 GHz flux density and associated with neutrino events, have on average higher flux densities than randomly selected objects, suggesting a connection between compact radio emission and neutrino events. As discussed earlier, their sample is based on the RFC catalog, and they use a flux density limit of 150 mJy. Our sample has a large overlap with theirs, and we have only three additional associated sources, out of which two are on neutrino events for which they did not find

an association. One of these is J1310+3220, which is associated with a neutrino event on May 15, 2012 at energy $E = 200$ TeV, and as discussed earlier, was excluded from Plavin et al. (2020a) because of their energy limit. The other two sources (J2200+0234 and J0341+3352) have mean flux density of less than 150 mJy at 15 GHz, and are not included in the RFC sample at this limit. J0341+3352 is found in the same uncertainty region as 0333+321, which is included in Plavin et al. (2020a), while J2200+0234 is associated with a neutrino event on Aug 7, 2012 for which Plavin et al. (2020a) did not find any counterparts.

Although we have included the cross-match of the RFC sample and the OVRO monitoring sample in our analysis, our current monitoring sample does not include all potential counterparts from RFC, and we lack several of the associations listed in Plavin et al. (2020a). Cross-matching the full RFC catalog (above compact flux density of 150 mJy at 8 GHz and $\text{DEC} > -20^\circ$) of 2355 objects with our neutrino list using $\Delta\psi = 0.4^\circ$ would give us eight additional neutrino events bringing the number of associated events up to 24, and the total number of associated objects to 31.

One important point to note is that the sample of Plavin et al. (2020a), which is restricted to flux density greater than 150 mJy is missing most of the HSP sources that Giommi et al. (2020) include in their analysis, which are typically fainter in the radio band. In fact, only 11 out of 84 HSPs in Richards et al. (2014) have intrinsic mean flux density above 150 mJy at 15 GHz. This

⁴ <https://tools.ssd.csi.it/>

shows that selecting the potential candidates from samples generated at single energy ranges may not reveal the complete picture.

Because Plavin et al. (2020a) found a connection especially with the highest flux density objects, in our analysis we have also used higher flux density cutoffs in the radio band, which reduces the number of associations further. In the 350 mJy flux density limited OVRO sample of 589 objects (limit to which the full OVRO sample is complete) we find only 12 that are positionally coincident with the neutrino arrival directions. The higher flux density limit of the Metsähovi sample (only sources with maximum flux density > 700 mJy are included) is also clearly seen in the analysis, with only seven neutrino events with eight counterparts found with the optimal systematic uncertainty $\Delta\psi = 0.3^\circ$ (Table 3).

4.2. Connection with γ -ray emission

One significant feature on the list of associated OVRO sources is the large number of non- γ -ray emitting sources, 10 out of 22. In general, some significant differences in properties of non- γ -ray emitting and γ -ray emitting blazars have been found and these could be of relevance also to the expected neutrino fluxes, therefore we discuss these properties in the following.

Savolainen et al. (2010) found that γ -ray detected blazars had in general higher Doppler factors than non γ -ray detected ones. Later studies (Richards et al. 2014; Lister et al. 2015) confirmed that the non γ -ray blazars had smaller 15 GHz modulation indices (defined as the standard deviation of the flux density variations over the mean flux density) and lower apparent jet speeds, both suggesting lower Doppler boosting. Therefore we investigated modulation indices and variability Doppler factors of the γ -ray and non γ -ray detected sources in our sample. For some of the sources these were available in Liodakis et al. (2017, 2018) and for the missing ones these were calculated using the same methods as in those studies.

The modulation indices and variability Doppler factors for the sources are given in Table 7. We performed a Kolmogorov-Smirnov test and found that there are no significant differences between the Doppler factors (p-value 0.63) or 15 GHz modulation indices (p-value 0.48) of the non γ -ray and γ -ray sources in our sample. For the non γ -ray sources in our sample the mean modulation index is 0.179 and mean variability Doppler factor 18.9, while for the γ -ray sources they are 0.194 and 12.0. Comparing these to average modulation indices for γ -ray blazars $0.175^{+0.012}_{-0.011}$, and for non γ -ray blazars $0.099^{+0.004}_{-0.003}$ (Richards et al. 2014) and to median Doppler factors for the γ -ray detected ($\delta_{\text{var}} = 14$) and non-detected ($\delta_{\text{var}} = 8$) from Liodakis et al. (2018) demonstrates that the non γ -ray detected sources in our sample are actually as variable and as Doppler boosted as typical γ -ray blazars and from that point of view they are not less likely neutrino sources than an average γ -ray blazar.

As discussed in e.g., Richards et al. (2014) and Lister et al. (2015) the non γ -ray blazars also have lower synchrotron peak frequencies than the γ -ray detected blazars. Since our sample mainly consists of LSP sources, comparing our γ -ray and non- γ -ray detected sources is not of interest. We note however that our non γ -ray detected sources all have low ν_S values. The range is $\log(\nu_S) = 12.3 - 13.6$, with only 3 sources with $\log(\nu_S) \geq 13$, while e.g., Keenan et al. (2020) found the average for LSP and FSRQ sources to be ~ 13.4 . This of course largely explains why they are non γ -ray detected, with a typical $\sim 10^7$ between the peaks (Fossati et al. 1998), the second peak in the SED is not in the *Fermi*-LAT energy range. The nature of these sources is little

studied as most of the "blazar sequence" studies are concentrated on γ -ray detected blazars (e.g. Ghisellini et al. 2017). We note however that we also estimated the νF_ν of these sources from the SED fits (see Table 7). The non γ -ray sources in our sample are fainter than the γ -ray detected ones, but still have $L_{\text{peak}} = 10^{44.9} - 10^{46.8}$ erg/s, so these sources are not low luminosity sources in general.

The actual conditions and predictions of the neutrino rates from the sources discussed in our paper would require detailed modelling and this is beyond the scope of the current paper. We simply note that there is no obvious reason to exclude any of the sources as a possible neutrino emitter. This is also in line with the discussion in Plavin et al. (2020b). They did not find reasons to disfavor the non γ -ray AGNs as neutrino sources in terms of target photon fields or the accelerated protons.

4.3. Mean flux density and variability of the radio emission

As discussed earlier, although we have used several subsets of the OVRO monitoring sample to look at the connection between radio emission and IceCube neutrino events, the number of associated sources is very similar in the different samples. Therefore, it is not surprising that the results for these different samples are in general consistent, as can be seen from Tables 4 and 6.

Unlike Plavin et al. (2020a), we do not see a highly significant difference between the mean flux density of the associated sources compared to the other sources in our samples (Table 4). Only in case of the OVRO all-AGN sample and CGRaBS sample, our results are indicative at the 2σ level of possibly higher mean flux densities in the sources positionally coincident with the neutrino events. Especially if we limit the analysis to the brightest blazars (RFC-150mJy, OVRO-350mJy and Metsähovi) samples, there is no significant difference between the mean flux densities of the associated objects and other objects in the samples.

However, when we look at the radio variability of these sources, our results are suggestive of a connection between the neutrino events and radio flares (Table 6). Regardless of the OVRO sample used, when we average the activity index of the associated sources, we find that the post-trial chance probability for obtaining these mean values is less than 3%. For the CGRaBS sample, which we consider statistically well defined, this is 1%. This is slightly smaller than 5%, which Plavin et al. (2020a) found for the sources in the RATAN-600 monitoring program, although it is still not significant at the 3σ level.

For the Metsähovi sample, we do not find a significant effect when we look at the average of the activity indices, but the chance probability to find four flaring blazars ($A.I. > 1.16$) coincident with neutrino arrival times is only 0.7% (post-trial). This value is obtained when using the 1% false-detection limit for defining the threshold for flaring blazars. If we use the 0.01% limit of $A.I. > 1.71$, there is only one source in our sample above that limit, also making the chance probability much higher (8.3% post-trial).

In the case of the OVRO samples, the case with flaring blazars is the opposite, i.e. our results are mostly not significant when the false-detection limit of 1% ($A.I. > 1.10$ for the OVRO data) is used, while using a limit of 0.01% ($A.I. > 1.29$) the post-trial p-values are less than 5% in all samples. The most significant results are found for the brightest objects (OVRO-350mJy sample) with a post-trial p-value of 0.6%, when there are five sources indicating strong flaring. These sources are not necessarily the brightest sources in the sample, and therefore they are

not the same objects that Plavin et al. (2020a) indicated as the most likely neutrino emitting sources based on their mean flux density.

We note that the number of strongly flaring blazars in both OVRO and Metsähovi samples is very small and the significance of the analysis is somewhat sensitive to the false-detection limits used because inclusion or exclusion of individual sources changes the results. Moreover, the number of strongly flaring blazars is very small compared to the number of IceCube neutrino events even if 50% of them are not astrophysical. Therefore it is clear that not all neutrino events happen when a cospatial blazar shows strong radio activity. However, when we see strong radio activity in a blazar and a neutrino event at the same time, it is unlikely to be a chance coincidence. This is consistent with the association of PKS B1424-418 with the high-energy neutrino IC35 by Kadler et al. (2016) who found the source to be flaring in both γ -ray and radio bands at the time of the neutrino arrival. Based on a sample of four possible neutrino associations and their OVRO light curves (including J0506+056 and J1504+1029 from our sample), Weber (2020) also note that the neutrino arrival times seem to coincide with flaring in the radio band.

Finally, although the activity index is a simple way to estimate the flaring state of a blazar, it is not ideal when using light curves that are several years long and well-sampled. For example, in the source J1743-0350 the coincident neutrino arrives during the rising part of a large outburst (Fig. A.2) but because the light curve includes other flares of similar strength, the overall mean flux density of the light curve is high, resulting in a smaller activity index. Another similar case is J1125+2610 (Fig. A.1). Even in the case of TXS 0506+056 (J0509+0541, Fig. A.1) the activity index does not indicate flaring because the radio outburst continues after the neutrino arrival.

The activity index and mean flux density in the radio bands are also not appropriate metrics for associating the HSP sources because in addition to being faint in the radio frequencies, the HSP sources are also less variable (Richards et al. 2014). Other ways to combine the radio and γ -ray catalogs should be investigated, in addition to more sophisticated methods for comparing radio flares to neutrino arrival times, but these are beyond the scope of this paper.

4.4. Comparison between OVRO and Metsähovi associations

All but one of the Metsähovi associations (1308+328, which was eliminated from the OVRO sample, see Sect. 2.2) are also included in the OVRO associations. We can thus compare the light curves and activity index values in the two different frequencies. In general, the light curves at 37 GHz show much faster variations than at 15 GHz (e.g., Hovatta et al. 2007), which is also seen when comparing the light curves of the associated sources. As discussed in the previous subsection, there are four associations in the Metsähovi sample that show strong radio flares at the time of the neutrino arrival. One of these is J1743-0350, which did not meet the formal criteria of strong flares in the OVRO data as discussed above. In the Metsähovi light curve (Fig. A.4), it can be seen that the neutrino arrival time coincides with an extremely fast increase of the radio flux density at 37 GHz. At 15 GHz the substructures of the flares are not as clearly visible, but also there the neutrino arrival time coincides with a fast increase of flux density. The other three flaring sources in the Metsähovi sample, are also identified as strongly flaring sources in the OVRO light curves, and overall the activity index values between the two data sets are similar.

We also note that while there is a large fraction of non- γ -ray detected sources in the OVRO associations (see Sect. 4.2), all the Metsähovi associations are γ -ray detected. This is mainly due to the smaller number of objects monitored regularly at Metsähovi, and the higher flux density limit we have used in our analysis. As can be seen in Table 2, all the non γ -ray detected sources have $\bar{S}_{15\text{GHz}} < 0.9\text{ Jy}$.

5. Conclusions

We have studied the possible connection between high-energy neutrinos and blazars by comparing our radio monitoring catalogs from OVRO and Metsähovi with positions of IceCube neutrino events. The number of associations we find is in general small, only with 12 – 28% of the 57 neutrino events, depending on the radio sample used. The small number of associations can be largely explained by incomplete parent samples of radio sources used in this study and is not in contradiction with the recent studies of Plavin et al. (2020a); Plavin et al. (2020b) associating the observed astrophysical neutrino flux to radio bright AGN.

Our associated sources are largely LSP FSRQs and a large fraction of the OVRO associations are not γ -ray detected (see Sect. 4.2). This reflects the composition of the OVRO monitoring sample, which largely consists of FSRQs and non γ -ray detected sources so that our study is not suitable for identifying the most likely class of sources associated with neutrinos.

We find that the luminosity and radio variability characteristics, including variability Doppler boosting factors, of the non γ -ray detected associations in our sample are indistinguishable from the γ -ray detected associations. Because these sources are mostly luminous FSRQs, they are also potential neutrino emitters, which are often ignored in studies concentrating only on γ -ray detected sources.

When studying the radio emission and variability of the associated sources, we find no difference in the mean flux density of the associated sources compared to random control populations. However, when radio variability amplitudes are studied, we find that on average, the associated sources in the OVRO sample are in a more active state at the time of the neutrino arrival. This is especially clear when largest flares are studied (both in OVRO and Metsähovi samples) and we conclude that when we see strong radio activity in a blazar and a neutrino event at the same time, it is unlikely to be a random coincidence.

This is in line with the results of recent studies on connection of blazar γ -ray flares and production of detectable neutrino fluxes. It is evident that the fluence of even the brightest keV-GeV flares is not sufficient to result in a likely detection of neutrinos and the only way to increase this likelihood is to increase the duration of the flare (Oikonomou et al. 2019; Kreter et al. 2020). Strong radio flares have typical duration of 1-2 years (see Section 3.3.) and are often accompanied with long-duration flaring at higher energies. Therefore they are natural candidates to be associated with detectable neutrino emission and we will study this connection in more detail in future works.

References

- Aartsen, M. G., Abbasi, R., Abdou, Y., et al. 2013, Phys. Rev. Lett., 111, 021103
- Aartsen, M. G., Abraham, K., Ackermann, M., et al. 2017a, ApJ, 835, 45
- Aartsen, M. G., Abraham, K., Ackermann, M., et al. 2016, ApJ, 833, 3
- Aartsen, M. G., Abraham, K., Ackermann, M., et al. 2017b, ApJ, 835, 151
- Aartsen, M. G., Ackermann, M., Adams, J., et al. 2017c, Astroparticle Physics, 92, 30

Acknowledgements

We thank A. Plavin, Y. Kovalev and M. Petropoulou for useful discussions. The OVRO 40 m program was supported by NASA grants NNG06GG1G, NNX08AW31G, NNX11A043G, and NNX13AQ89G from 2006 to 2016 and NSF grants AST-0808050, and AST-1109911 from 2008 to 2014. T. H. was supported by the Academy of Finland projects 317383, 320085, and 322535. E. L. was supported by the Academy of Finland projects 317636 and 320045. S.K. acknowledges support from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme under grant agreement No. 771282. W.M. acknowledges support from ANID projects Basal AFB-170002 and PAI79160080. Handling of catalogs in the paper was partially done using TOPCAT (Taylor 2005).

- Aartsen, M. G., Ackermann, M., Adams, J., et al. 2014, *Phys. Rev. Lett.*, 113, 101101
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2010, *ApJ*, 715, 429
- Ackermann, M., Ajello, M., Allafort, A., et al. 2011, *ApJ*, 743, 171
- Ansoldi, S., Antonelli, L. A., Arcaro, C., et al. 2018, *ApJ*, 863, L10
- Baars, J. W. M., Genzel, R., Pauliny-Toth, I. I. K., & Witzel, A. 1977, *A&A*, 61, 99
- Blandford, R. D. & Königl, A. 1979, *ApJ*, 232, 34
- Böttcher, M. 2019, *Galaxies*, 7, 20
- Brown, A. M., Adams, J., & Chadwick, P. M. 2015, *MNRAS*, 451, 323
- Chang, Y. L., Arsioli, B., Giommi, P., Padovani, P., & Brandt, C. H. 2019, *A&A*, 632, A77
- D'Abusco, R., Massaro, F., Paggi, A., et al. 2014, *The Astrophysical Journal Supplement Series*, 215, 14
- Davison, A. C. & Hinkley, D. V. 2013, *Bootstrap Methods and their Application*, Cambridge Series in Statistical and Probabilistic Mathematics (Cambridge University Press)
- Fossati, G., Maraschi, L., Celotti, A., Comastri, A., & Ghisellini, G. 1998, *MNRAS*, 299, 433
- Franckowiak, A., Garrappa, S., Paliya, V., et al. 2020, *ApJ*, 893, 162
- Ghisellini, G., Righi, C., Costamante, L., & Tavecchio, F. 2017, *MNRAS*, 469, 255
- Giommi, P., Glauch, T., Padovani, P., et al. 2020, *MNRAS*, 497, 865
- Hartman, R. C., Bertsch, D. L., Fichtel, C. E., et al. 1992, *ApJ*, 385, L1
- Healey, S. E., Romani, R. W., Cotter, G., et al. 2008, *ApJS*, 175, 97
- Hovatta, T., Tornikoski, M., Lainela, M., et al. 2007, *A&A*, 469, 899
- Huber, M. 2019, in *International Cosmic Ray Conference*, Vol. 36, 36th International Cosmic Ray Conference (ICRC2019), 916
- IceCube Collaboration. 2013, *Science*, 342, 1242856
- IceCube Collaboration. 2018, DOI:10.21234/B4KS6S
- IceCube Collaboration, Aartsen, M. G., Ackermann, M., et al. 2018a, *Science*, 361, eaat1378
- IceCube Collaboration, Aartsen, M. G., Ackermann, M., et al. 2019, arXiv e-prints, arXiv:1907.11699
- IceCube Collaboration, Aartsen, M. G., Ackermann, M., et al. 2018b, *Science*, 361, 147
- IceCube Collaboration, Aartsen, M. G., Ackermann, M., et al. 2017, arXiv e-prints, arXiv:1710.01191
- Kadler, M., Krauß, F., Mannheim, K., et al. 2016, *Nature Physics*, 12, 807
- Keenan, M., Meyer, E. T., Georganopoulos, M., Reddy, K., & French, O. J. 2020, arXiv e-prints, arXiv:2007.12661
- Korolkov, D. V. & Pariiskii, I. N. 1979, *S&T*, 57, 324
- Krauß, F., Deoskar, K., Baxter, C., et al. 2018, *A&A*, 620, A174
- Krauß, F., Kadler, M., Mannheim, K., et al. 2014, *A&A*, 566, L7
- Kreter, M., Kadler, M., Krauß, F., et al. 2020, arXiv e-prints, arXiv:2009.00125
- Kun, E., Bartos, I., Tjus, J. B., et al. 2020 [arXiv:2009.09792]
- Lähteenmäki, A. & Valtaoja, E. 2003, *ApJ*, 590, 95
- León-Tavares, J., Valtaoja, E., Tornikoski, M., Lähteenmäki, A., & Nieppola, E. 2011, *A&A*, 532, A146
- Liodakis, I., Hovatta, T., Huppenkothen, D., et al. 2018, *ApJ*, 866, 137
- Liodakis, I., Pavlidou, V., Hovatta, T., et al. 2017, *MNRAS*, 467, 4565
- Liodakis, I. & Petropoulou, M. 2020, *ApJ*, 893, L20
- Lister, M. L., Aller, M. F., Aller, H. D., et al. 2015, *ApJ*, 810, L9
- Mannheim, K., Stanev, T., & Biermann, P. L. 1992, *A&A*, 260, L1
- Massaro, E., Maselli, A., Leto, C., et al. 2015, *Ap&SS*, 357, 75
- Mirzoyan, R. 2017, *The Astronomer's Telegram*, 10817, 1
- Mücke, A. & Protheroe, R. J. 2001, in *International Cosmic Ray Conference*, Vol. 3, International Cosmic Ray Conference, 1153
- Nalewajko, K. 2017, *Galaxies*, 5, 64
- Neronov, A. Y. & Semikoz, D. V. 2002, *Phys. Rev. D*, 66, 123003
- Oikonomou, F., Murase, K., Padovani, P., Resconi, E., & Mészáros, P. 2019, *MNRAS*, 489, 4347
- Padovani, P. & Resconi, E. 2014, *MNRAS*, 443, 474
- Padovani, P., Resconi, E., Giommi, P., Arsioli, B., & Chang, Y. L. 2016, *MNRAS*, 457, 3582
- Palladino, A. & Vissani, F. 2016, *ApJ*, 826, 185
- Petropoulou, M., Murase, K., Santander, M., et al. 2020, *ApJ*, 891, 115
- Plavin, A., Kovalev, Y. Y., Kovalev, Y. A., & Troitsky, S. 2020a, *ApJ*, 894, 101
- Plavin, A. V., Kovalev, Y. Y., Kovalev, Y. A., & Troitsky, S. V. 2020b, Submitted to *ApJ*[arXiv:2009.08914]
- Readhead, A. C. S., Cohen, M. H., Pearson, T. J., & Wilkinson, P. N. 1978, *Nature*, 276, 768
- Reimer, A., Böttcher, M., & Buson, S. 2019, *ApJ*, 881, 46
- Richards, J. L., Hovatta, T., Max-Moerbeck, W., et al. 2014, *MNRAS*, 438, 3058
- Richards, J. L., Max-Moerbeck, W., Pavlidou, V., et al. 2011, *ApJS*, 194, 29
- Savolainen, T., Homan, D. C., Hovatta, T., et al. 2010, *A&A*, 512, A24
- Sowards-Emmerd, D., Romani, R. W., Michelson, P. F., Healey, S. E., & Nolan, P. L. 2005, *The Astrophysical Journal*, 626, 95
- Tanaka, Y. T., Buson, S., & Kocevski, D. 2017, *The Astronomer's Telegram*, 10791, 1
- Taylor, M. B. 2005, in *Astronomical Society of the Pacific Conference Series*, Vol. 347, *Astronomical Data Analysis Software and Systems XIV*, ed. P. Shopbell, M. Britton, & R. Ebert, 29
- Teräsranta, H., Achren, J., Hanski, M., et al. 2004, *A&A*, 427, 769
- Teräsranta, H., Tornikoski, M., Mujunen, A., et al. 1998, *A&AS*, 132, 305
- The Fermi-LAT collaboration. 2019, arXiv e-prints, arXiv:1905.10771
- The IceCube Collaboration, Aartsen, M. G., Abraham, K., et al. 2015, arXiv e-prints, arXiv:1510.05223
- Valtaoja, E., Lähteenmäki, A., & Teräsranta, H. 1992, *A&AS*, 95, 73
- Weber, P. 2020, Bachelor Thesis, Julius-Maximilians-Universität Würzburg, Germany

**Appendix A: OVRO 15 GHz and Metsähovi 37 GHz
light curves of the associated sources**




Fig. A.1. OVRO light curves of the associated sources. The neutrino arrival time is shown with a blue solid line, while the blue dashed lines indicate the window of 2.3 yrs around the neutrino event.




Fig. A.2. OVRO light curves of the associated sources. The neutrino arrival time is shown with a blue solid line, while the blue dashed lines indicate the window of 2.3 yrs around the neutrino event.




Fig. A.3. OVRO light curves of the associated sources. The neutrino arrival time is shown with a blue solid line, while the blue dashed lines indicate the window of 2.3 yrs around the neutrino event.




Fig. A.4. Metsähovi light curves of the associated sources. The neutrino arrival time is shown with a blue solid line, while the blue dashed lines indicate the window of 1.4 yrs around the neutrino event.