A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: Catalysis-Enabled Concise and Stereoselective Total Synthesis of the Tricyclic Prostaglandin D2 Metabolite Methyl Ester

Authors: Mingji Dai, Hunter S. Sims, Pedro de Andrade Horn, Ryota Isshiki, Melissa Lim, Yan Xu, and Robert H. Grubbs

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.202115633

Link to VoR: <https://doi.org/10.1002/anie.202115633>

COMMUNICATION

Catalysis-Enabled Concise and Stereoselective Total Synthesis of the Tricyclic Prostaglandin D₂ Metabolite Methyl Ester

Hunter S. Sims,^[a] Pedro de Andrade Horn,^[a] Ryota Isshiki,^{[a][c]} Melissa Lim,^[a] Yan Xu,^[b] Robert H. Grubbs,^[b] and Mingji Dai^{*[a]}

[a] H. S. Sims, P. de Andrade Horn, R. Isshiki, M. Lim, and Prof. Dr. M. Dai
Department of Chemistry and Center for Cancer Research, Purdue University, West Lafayette, Indiana 47907, USA
E-mail: mj Dai@purdue.edu

[b] Dr. Y. Xu, Prof. Dr. R. H. Grubbs
The Arnold and Mabel Beckman Laboratory of Chemical Synthesis, Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, USA

[c] R. Isshiki
Department of Applied Chemistry, Waseda University, Shinjuku, Tokyo 162-0041, Japan

Supporting information for this article is given via a link at the end of the document

Abstract: A concise and stereoselective total synthesis of the clinically relevant tricyclic prostaglandin D₂ metabolite (tricyclic-PGDM) methyl ester in racemic form was accomplished in 8 steps from a readily available known cyclopentene-diol derivative. The synthesis features a nickel-catalyzed Ueno-Stork-type dicarbofunctionalization to generate two consecutive stereocenters, a palladium-catalyzed carbonylative spirocyclization to build the core oxaspirolactone, and a Z-selective cross metathesis to introduce the (Z)-3-butenate side chain, a group challenging to introduce through traditional Wittig protocols and troublesome for the two previous total syntheses. A general Z-selective cross metathesis protocol to construct (Z)-β,γ-unsaturated esters was also developed that has broad functional group tolerance and high stereoselectivity. Additionally, our synthesis already accumulated 75 mg of valuable material for an ¹⁸O tricyclic-PGDM based assay used in clinical settings for inflammation.

Prostaglandins are signaling molecules that play a pivotal role in numerous biological pathways. A range of chemical and physical stimuli can activate their biosynthesis, which progresses through an enzymatic cascade involving phospholipase-mediated release of arachidonic acid (AA) from the membrane. Upon release, AA is transformed by cyclooxygenase (COX) enzymes to prostaglandin H₂ (PGH₂), a common intermediate that gives rise to the four principal prostaglandins by tissue specific isomerases.^[1] Prostaglandins serve a crucial regulatory function in the acute inflammatory response. However, upregulation of these pathways spontaneously or extending beyond an initial inflammation event can be detrimental and has been linked to multiple diseases.^[2] In addition, other unique non-inflammatory roles of prostaglandins are continuing to emerge further demonstrating that these scaffolds hold a privileged position as biological mediators.^[3] Thus, it is no surprise that the development of sensitive detection methods for prostaglandins has been an active research area. By virtue of their short half-lives, most current methods rely on the detection of metabolites that can be traced to the prostaglandin of interest.^[4] Notably, following its recognition as the major urinary metabolite of prostaglandin D₂ (PGD₂, **1**), tricyclic-PGDM (**2**) was used as a valuable indicator for PGD₂ overproduction.^[5] Using ¹⁸O-labelling, Roberts and colleagues at Vanderbilt University

developed an assay for tricyclic-PGDM quantification which has been a valuable tool in clinical settings (Figure 1).^[6] However, limited material is a constraint that has restricted this assay from becoming more prevalent since it relies on the fully synthetic tricyclic-PGDM methyl ester (**3**).

Figure 1. Clinical use of tricyclic-PGDM methyl ester and its synthesis.

The prostaglandin molecules and their metabolites have garnered a tremendous amount of interest and efforts from the synthetic community.^[7] The complexity of tricyclic-PGDM methyl ester (**3**) arises from the four contiguous stereocenters on the central cyclopentane ring, the labile oxaspirolactone, and the (Z)-3-butenate side chain. In 1988, Taber and coworkers surmounted these challenges achieving the first synthesis of (±)-**3** in 17 steps from the Corey lactone (**4**, prepared in 9 steps), with

COMMUNICATION

their final step giving 5 mg of **3** for the clinical assay (Figure 1).^[8] In their synthesis, the (Z)-3-butenolate side chain proved to be troublesome to install on a hemi-acetal intermediate, with β -phosphoranylidene esters being noncompatible with the Wittig reaction. They overcame this issue by using an orthoester phosphorane.^[9] To complete the synthesis, an acid-catalyzed cyclization of a hydroxy ketoacid was used to install the oxaspirolactone. More recently, to help replenish the assay Sulikowski and coworkers developed the second synthesis of ($\pm$)-**3** from the known α -iodoenone **5** (prepared in 6 steps) in 13 steps, with their final step yielding 1 mg of **3**.^[10] Their route featured a Lindlar reduction of a β -alkynyl ester to install the (Z)-3-butenolate side chain, albeit in low yield, and an acid-promoted spiroketalization to afford the oxaspirolactone. Herein, we report an efficient synthesis of ($\pm$)-**3** (75 mg) with three transition metal (Ni, Pd, and Ru)-catalyzed transformations to build strategic C-C bonds and key ring systems.

A. Pd-catalyzed carbonylative spirocyclization of hydroxycyclopropanols

B. Retrosynthetic analysis

Figure 2. Carbonylative spirocyclization and retrosynthetic analysis.

The clinical importance of tricyclic-PGDM methyl ester (**3**) and its intriguing structure, particularly the oxaspirolactone moiety, attracted our attention. We previously developed a palladium-catalyzed carbonylative spirocyclization to synthesize oxaspirolactones from readily available hydroxycyclopropanols (**8**→**9**, Figure 2A), and have used this method to efficiently synthesize a multitude of complex natural products.^[11] To this end, we were inspired to establish an efficient carbonylative lactonization route to **3** and provide sufficient amount of material to support the clinical assays. Retrosynthetically (Figure 2B), we reasoned that the (Z)-3-butenolate side chain would need to be incorporated late in the synthesis to ensure its integrity. While the Wittig protocol with an orthoester phosphorane employed by Taber et al. gave 50% yield, the Z:E selectivity was only moderate (4:1) and additional orthoester deprotection and esterification steps were required.^[8] Compound **12** is a known intermediate in

the Sulikowski synthesis, however, six more steps (5% overall yield) including a Lindlar reduction were required to incorporate the (Z)-3-butenolate side chain.^[10] Apparently, a direct Z-selective cross metathesis between **12** and methyl 3-butenolate **13** would be ideal to install the side chain, but with foreseeable challenges. While significant advances have been made in Z-selective cross metathesis,^[12] β,γ -unsaturated esters including **13** still remain as challenging substrates. Therefore, it is not surprising that no such methods for (Z)- β,γ -unsaturated esters have been disclosed thus far. Olefins with a proximal chelating group, such as esters and amides, are particularly challenging substrates. β,γ -Unsaturated esters are among the worst substrates because the ester can chelate on the metal center after carbenoid formation (cf. **A**) forming a stable five-membered metallocycle. Furthermore, the cross-metathesis product tends to undergo double bond isomerization to the E-product or migration to the α,β -unsaturated ester. However, the recent Z-selective cross metathesis results between allylic-substituted olefins^[13] or acrylamides^[14] and common terminal olefins are encouraging. Despite the challenges, a direct Z-selective cross metathesis between **12** and **13** to complete the synthesis of **3** in just one step is attractive and worth developing, which may result in a generalized method for (Z)- β,γ -unsaturated ester preparation. The cross-metathesis precursor **12** could be prepared via olefination of the hemi-acetal derived from hydrolysis of acetal **14**. The latter could be prepared from hydroxycyclopropanol **15** using the palladium-catalyzed carbonylative spirocyclization to generate its oxaspirolactone moiety. Hydroxycyclopropanol **15** could be generated by the Kulinkovich reaction of lactone **16**, which we anticipated to arise from a Ueno-Stork^[15] type tandem radical cyclization of iodo-acetal **17**. The radical cyclization precursor **17** could be assembled from the readily available known mono-protected diol **6**.^[16]

Our synthesis commenced with the multi-decagram preparation of known compound **6** in six steps in racemic form from cyclopentadiene (Scheme 1). Enantioselective synthesis of **6** with an enzymatic hydrolysis^[17] or a Noyori reduction was reported as well.^[18] Subsequent conversion to iodo-acetal **18**, silyl ether deprotection, and attachment of the acrylate side chain then gave **17**, setting the stage for the key Ueno-Stork type cyclization. Unfortunately, tricyclic lactone **16** could not be isolated. Complex mixtures were obtained under various tin-based Ueno-Stork cyclization conditions, and the acrylate tether was cleaved when palladium, nickel and other transition metals were used to promote a dicarbofunctionalization due to its allylic nature.^[19] We presumed the terminal 6-endo-trig cyclization was unfeasible for this substrate, and therefore proceeded to an intermolecular variant utilizing intermediate **18** and methyl acrylate. Gratifyingly, with modifications to the original Stork conditions,^[19a] product **20** was obtained in 61% yield and excellent stereoselectivity at the newly formed stereogenic centers on up to 10 g scales. While efficient and scalable, the organotin toxicity and complications with *t*-BuOH during the workup led us to explore more environmentally benign conditions. Accordingly, we developed a novel nickel-catalyzed protocol to obtain dicarbofunctionalization product **20** in 59% yield.^[20] While the yield slightly decreased (46%) on larger scale, the procedure was operationally convenient. Next, the Kulinkovich reaction^[21] and TBS-ether deprotection proceeded smoothly on multi-gram scale to give cyclopropanol **15**, which was unambiguously confirmed by x-ray analysis (CCDC 2089935).^[22]

The unsuccessful one-carbon homologation of **23** prompted us to rethink our route, and we surmised the olefination would likely be more feasible without the sensitive oxaspirolactone present. Accordingly, cyclopropanol **15** was chosen as an appropriate candidate, since it would not require us to extensively modify our route. We suspected that the cyclopropanol would

COMMUNICATION

likely survive mild acetal hydrolysis conditions, however, we anticipated the methylenation to be a formidable challenge. To this end, hydrolysis of **15** proceeded smoothly in dilute HCl (0.05 M) to give hemi-acetal **25**, but as expected the olefination was not as straightforward. Finally, with careful monitoring of reaction temperature and time, the methylenation product **26** was obtained in acceptable yield on fairly large scales. Moving forward, the carbonylative spirocyclization was once again reliable. Oxaspirolactone **12** was delivered as a single stereoisomer due to the anomeric effect and in good yield despite the potential competition of a 5-*exo-trig* oxypalladation and other unproductive pathways.

Our next task was to find a suitable Z-selective cross metathesis condition to complete the total synthesis in just one step.^[25] We reasoned a cross metathesis using the commercially available **Ru-Z-Mes** as catalyst and methyl 3-butenolate (**13**) as the partner would be a suitable starting point. After extensive investigations, we found that about 10–15% (90% Z) of the cross-metathesis product could be obtained with the commercially available **Ru-Z-Mes**, but the results were not consistent. We further learned that re-purification of the **Ru-Z-Mes** was beneficial and increased the yield to 35%. To further enhance the yield and Z selectivity, the **Ru-Z-DIPP** catalyst was evaluated and found to be superior to the **Ru-Z-Mes** catalyst. Final product **3** was produced in better yield (52%) and higher selectivity (>95% Z). Interestingly, while both **Ru-Z-Mes** and **Ru-Z-DIPP** usually perform best in THF, only negligible yields (<10%) were obtained for this case. The reaction could also be conducted at 100-mg scale to give 38% (>95% Z) yield of **3** at a lower catalyst loading (5 mol%) and less amount of **13** (5 equiv.).

Table 1. Substrate scope for the Z-selective cross metathesis with **13**.

With the total synthesis of **3** secured, we next sought to generalize the Z-selective cross metathesis with the challenging β,γ -unsaturated ester substrate **13** (Table 1). Various terminal olefins with a wide range of functional groups were investigated. Under the aforementioned conditions (15 mol% **Ru-Z-DIPP**, 8 equiv of **13**, 40 °C), although cross metathesis products were obtained in good yield, the Z/E selectivity dropped slightly (90%–92% Z). Gratifyingly, lower the catalyst loadings (5 mol% **Ru-Z-DIPP**) and temperature (30 °C) gave satisfactory yields and excellent Z selectivity (>95%) for a range of olefin substrates. Notably, free alcohol, TBS ether, alkyl bromide, ester, and boronate are all tolerated and cross metathesis product **28g** is a

known precursor for synthesis of the beetle pheromone Ferrulactone II.^[26]

In summary, a concise synthesis of tricyclic-PGDM methyl ester (**3**) was achieved in 8 steps from readily available known compound **6**. The synthesis features a Z-selective cross metathesis to form the challenging (Z)- β,γ -unsaturated ester, a palladium-catalyzed carbonylative spirocyclization to build the oxaspirolactone moiety, and a nickel-catalyzed Ueno-Stork-type dicarbofunctionalization to form two key C–C bonds and stereocenters. A general cross-metathesis protocol for (Z)- β,γ -unsaturated esters was established. Through this route, we have accumulated over 75 mg of pure **3**. We anticipate this material will be useful for the tricyclic-PGDM assay and further its availability as a clinical tool.

Acknowledgements

This work was supported by NSF 2102022. The NIH CA023168 is acknowledged for supporting shared NMR resources to Purdue Center for Cancer Research. H.S. gratefully acknowledges support from the Purdue Drug Discovery Training Program (NIH T32 GM125620). The XRD data was collected on a new single crystal X-ray diffractometer supported by the NSF through the Major Research Instrumentation Program under Grant No. CHE 1625543.

Conflicts of interest

The authors declare no conflicts of interest.

Keywords: prostaglandins • carbonylation • olefin metathesis • nickel catalysis • total synthesis

- [1] G. A. FitzGerald, *Nat. Rev. Drug Discov.*, **2003**, 2, 879–890.
- [2] E. Ricciotti, G. A. FitzGerald, *Arterioscler. Thromb. Vasc. Biol.* **2011**, 31, 986–1000.
- [3] For selected examples: a) Y. Urade, O. Hayaishi, *Sleep Med. Rev.*, **2011**, 15, 411–418; b) M. L. Lee, H. Matsunaga, Y. Sugiura, T. Hayasaka, I. Yamamoto, T. Ishimoto, D. Imoto, M. Suematsu, N. Iijima, K. Kimura, S. Diano, C. Toda, *Nat. Commun.*, **2021**, 12, 2330; c) A. R. Palla, M. Ravichandran, Y. X. Wang, L. Alexandrova, A. V. Yang, P. Kraft, C. A. Holbrook, C. M. Schürch, A. T. V. Ho, H. M. Blau, *Science*, **2021**, 371, eabc8059.
- [4] D. Tsikas, *J. Chromatogr. B Biomed. Sci. Appl.*, **1998**, 717, 201–245.
- [5] T. E. Liston, L. J. Roberts, *J. Biol. Chem.*, **1985**, 260, 13172–13180.
- [6] J. D. Morrow, C. Prakash, J. A. Awad, T. A. Duckworth, W. E. Zackert, I. A. Blair, J. A. Oates, L. J. Roberts, *Anal. Biochem.*, **1991**, 193, 142–148.
- [7] a) S. Das, S. Chandrasekhar, J. S. Yadav, R. Grée, *Chem. Rev.* **2007**, 107, 3286–3337; b) H. H. Peng, F. E. Chen, *Org. Biomol. Chem.* **2017**, 15, 6281–6301.
- [8] C. Prakash, S. Saleh, L. J. Roberts, I. A. Blair, D. F. Taber, *J. Chem. Soc. Perkin Trans. 1*, **1988**, 2821–2826.
- [9] E. J. Corey, K. Shimoi, *J. Am. Chem. Soc.*, **1983**, 105, 1662–1664.
- [10] J. R. Kimbrough, Z. Austin, G. L. Milne, G. A. Sulikowski, *Org. Lett.*, **2019**, 21, 10048–10051.
- [11] a) D. C. Davis, K. L. Walker, C. Hu, R. N. Zare, R. M. Waymouth, M. Dai, *J. Am. Chem. Soc.*, **2016**, 138, 10693–10699; b) K. Ma, X. Yin, M. Dai, *Angew. Chem. Int. Ed.*, **2018**, 57, 15209–15212; *Angew. Chem.*, **2018**, 130, 15429–15432; c) X. Yin, K. Ma, Y. Dong, M. Dai, *Org. Lett.*, **2020**, 22, 5001–5004.

COMMUNICATION

- [12] a) K. Endo, R. H. Grubbs, *J. Am. Chem. Soc.*, **2011**, *133*, 8525–8527; b) B. K. Keitz, K. Endo, P. R. Patel, M. B. Herbert, R. H. Grubbs, *J. Am. Chem. Soc.*, **2012**, *134*, 693–699; c) L. E. Rosebrugh, M. B. Herbert, V. M. Marx, B. K. Keitz, R. H. Grubbs, *J. Am. Chem. Soc.*, **2013**, *135*, 1276–1279; d) S. J. Meek, R. V. O'Brian, J. Llaveria, R. R. Schrock, A. H. Hoveyda, *Nature*, **2011**, *471*, 461–466; e) T. J. Mann, A. W. H. Speed, R. R. Schrock, A. H. Hoveyda, *Angew. Chem. Int. Ed.*, **2013**, *52*, 8395–8400; *Angew. Chem.*, **2013**, *125*, 8553–8558; f) M. J. Koh, R. K. M. Khan, S. Torker, M. Yu, M. S. Mikus, A. H. Hoveyda, *Nature*, **2015**, *517*, 181–186; g) G. Occhipinti, F. R. Hansen, K. W. Törnroos, V. R. Jensen, *J. Am. Chem. Soc.*, **2013**, *135*, 3331–3334.
- [13] B. L. Quigley, R. H. Grubbs, *Chem. Sci.*, **2014**, *5*, 501–506.
- [14] Y. Xu, J. J. Wong, A. E. Samkian, J. H. Ko, S. Chen, K. N. Houk, R. H. Grubbs, *J. Am. Chem. Soc.*, **2020**, *142*, 20987–20993.
- [15] For a comprehensive review on the Ueno-Stork reaction see: X. J. Salom-Roig, F. Dénès, P. Renaud, *Synthesis*, **2004**, *12*, 1903–1928.
- [16] A. K. Ghosh, C. X. Xu, H. L. Osswald, *Tetrahedron Lett.*, **2015**, *56*, 3314–3317.
- [17] D. R. Deardorff, C. Q. Windham, C. L. Craney, *Org. Synth.*, **1996**, *73*, 25.
- [18] G. Singh, A. Meyer, J. Aubé, *J. Org. Chem.*, **2014**, *79*, 452–458.
- [19] a) G. Stork, P. M. Sher, H. L. Chen, *J. Am. Chem. Soc.*, **1986**, *108*, 6384–6385; b) L. Firmansjah, G. C. Fu, *J. Am. Chem. Soc.*, **2007**, *129*, 11340–11341; c) G. Z. Wang, R. Shang, W. M. Cheng, Y. Fu, *J. Am. Chem. Soc.*, **2017**, *139*, 18307–18312; d) J. D. Nguyen, E. M. D'Amato, J. M. R. Narayanam, C. R. J. Stephenson, *Nat. Chem.*, **2012**, *4*, 854–859; e) H. Kim, C. Lee, *Org. Lett.*, **2011**, *13*, 2050–2053. For a review: X. Qi, T. Diao, *ACS Catal.*, **2020**, *10*, 8542–8556.
- [20] For a review: X. Qi, T. Diao, *ACS Catal.*, **2020**, *10*, 8542–8556.
- [21] The reaction was unsuccessful under Kulinkovich's originally reported catalytic conditions: a) O. G. Kulinkovich, S. V. Sviridov, D. A. Vasilevski, *Synthesis*, **1991**, 234; when using stoichiometric amounts of $\text{CITi}(\text{O}i\text{Pr})_3$, the reaction proceeded smoothly: b) J. S. Kingsbury, E. J. Corey, *J. Am. Chem. Soc.*, **2005**, *127*, 13813–13815; also see reference 11b.
- [22] CCDC 2089935 contains the supplementary crystallographic data for compound **15**.
- [23] X. Cai, W. Liang, M. Liu, X. Li, M. Dai, *J. Am. Chem. Soc.*, **2020**, *142*, 13677–13682.
- [24] a) H. Lebel, V. Paquet, C. Proulx, *Angew. Chem. Int. Ed.*, **2001**, *40*, 2887–2890; *Angew. Chem.*, **2001**, *113*, 2971–2974; b) H. Lebel, V. Paquet, *J. Am. Chem. Soc.*, **2004**, *126*, 320–328.
- [25] a) M. B. Herbert, V. M. Marx, R. L. Pederson, R. H. Grubbs, *Angew. Chem. Int. Ed.*, **2013**, *52*, 310–314; *Angew. Chem.*, **2013**, *125*, 328–332; b) J. S. Cannon, R. H. Grubbs, *Angew. Chem. Int. Ed.*, **2013**, *52*, 9001–9004; *Angew. Chem.*, **2013**, *125*, 9171–9174; c) S. M. Bronner, M. B. Herbert, P. R. Patel, V. M. Marx, R. H. Grubbs, *Chem. Sci.*, **2014**, *5*, 4091–4098; d) S. L. Mangold, D. J. O'Leary, R. H. Grubbs, *J. Am. Chem. Soc.*, **2014**, *136*, 12469–12478; For reviews see: e) M. B. Herbert, R. H. Grubbs, *Angew. Chem. Int. Ed.*, **2015**, *54*, 5018–5024; *Angew. Chem.*, **2015**, *127*, 5104–5110; f) T. P. Montgomery, A. M. Johns, R. H. Grubbs, *Catalysts*, **2017**, *7*, 87. For examples in total synthesis see: g) W. J. Chung, J. S. Carlson, D. K. Bedke, C. D. Vanderwal, *Angew. Chem. Int. Ed.*, **2013**, *52*, 10052–10055; *Angew. Chem.*, **2013**, *125*, 10236–10239; h) J. Li, T. S. Ahmed, C. Xu, B. M. Stoltz, R. H. Grubbs, *J. Am. Chem. Soc.*, **2019**, *141*, 154–158; i) J. Li, B. M. Stoltz, R. H. Grubbs, *Org. Lett.*, **2019**, *21*, 10139–10142; j) M. De Léséleuc, É. Godin, S. Parisien-Collette, A. Lévesque, S. K. Collins, *J. Org. Chem.*, **2016**, *81*, 6750–6756.
- [26] a) K. Mori, H. Tomioka, *Liebigs Ann. Chem.*, **1992**, 1011–1017; b) A. S. Pawar, S. Sankaranarayanan, S. Chattopadhyay, *Tetrahedron Asymmetry*, **1995**, *6*, 2219–2226.

COMMUNICATION

Entry for the Table of Contents

The clinically important tricyclic prostaglandin D₂ metabolite (PGDM) methyl ester was synthesized in eight steps from a readily available known compound. The synthesis features Z-selective cross-metathesis to form the challenging Z β,γ -unsaturated ester, palladium-catalyzed carbonylative spirocyclization to build the oxaspirolactone moiety, and nickel-catalyzed Ueno–Stork-type dicarbofunctionalization to form two key C–C bonds and stereocenters.

Institute and/or researcher Twitter usernames: Institute Twitter: @PurdueChemistry; Researcher Twitter: @MingjiDai