

The California-Kepler Survey. X. The Radius Gap as a Function of Stellar Mass, Metallicity, and Age.

ERIK A. PETIGURA,¹ JAMES G. ROGERS,² HOWARD ISAACSON,^{3,4} JAMES E. OWEN,² ADAM L. KRAUS,⁵
JOSHUA N. WINN,⁶ MASON G. MACDOUGALL,¹ ANDREW W. HOWARD,⁷ BENJAMIN FULTON,⁸ MOLLY R. KOSIAREK,⁹
LAUREN M. WEISS,¹⁰ AIDA BEHMARD,¹¹ AND SARAH BLUNT¹²

¹*Department of Physics & Astronomy, University of California Los Angeles, Los Angeles, CA 90095, USA*

²*Astrophysics Group, Imperial College London, Blackett Laboratory, Prince Consort Road, London SW7 2AZ, UK*

³*Department of Astronomy, University of California Berkeley, Berkeley CA 94720, USA*

⁴*Centre for Astrophysics, University of Southern Queensland, Toowoomba, QLD, Australia*

⁵*Department of Astronomy, The University of Texas at Austin, Austin, TX 78712, USA*

⁶*Princeton University, Princeton, NJ 08540, USA*

⁷*Department of Astronomy, California Institute of Technology, Pasadena, CA 91125, USA*

⁸*NASA Exoplanet Science Institute/Caltech-IPAC, MC 314-6, 1200 E. California Blvd., Pasadena, CA 91125, USA*

⁹*Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064, USA*

¹⁰*Department of Physics, University of Notre Dame, Notre Dame, IN 46556, USA*

¹¹*Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125, USA*

¹²*Department of Astronomy, California Institute of Technology, Pasadena, CA, USA*

ABSTRACT

In 2017, the California-*Kepler* Survey (CKS) published its first data release (DR1) of high-resolution optical spectra of 1305 planet hosts. Refined CKS planet radii revealed that small planets are bifurcated into two distinct populations: super-Earths (smaller than $1.5 R_{\oplus}$) and sub-Neptunes (between 2.0 and $4.0 R_{\oplus}$), with few planets in between (the “Radius Gap.”) Several theoretical models of the Radius Gap predict variation with stellar mass, but testing these predictions are challenging with CKS DR1 due to its limited M_{\star} range of 0.8–1.4 M_{\odot} . Here, we present CKS DR2 with 411 additional spectra and derived properties focusing on stars of 0.5–0.8 M_{\odot} . We found the Radius Gap follows $R_p \propto P^m$ with $m = -0.10 \pm 0.03$, consistent with predictions of XUV- and core-powered mass-loss mechanisms. We found no evidence that m varies with M_{\star} . We observed a correlation between the average sub-Neptune size and M_{\star} . Over 0.5 to 1.4 M_{\odot} , the average sub-Neptune grows from 2.1 to 2.6 R_{\oplus} , following $R_p \propto M_{\star}^{\alpha}$ with $\alpha = 0.25 \pm 0.03$. In contrast, there is no detectable change for super-Earths. These M_{\star} - R_p trends suggests that protoplanetary disks can efficiently produce cores up to a threshold mass of M_c , which grows linearly with stellar mass according to $M_c \approx 10 M_{\oplus} (M_{\star}/M_{\odot})$. There is no significant correlation between sub-Neptune size and stellar metallicity (over -0.5 to $+0.5$ dex) suggesting a weak relationship between planet envelope opacity and stellar metallicity. Finally, there is no significant variation in sub-Neptune size with stellar age (over 1 to 10 Gyr), which suggests that the majority of envelope contraction concludes after ~ 1 Gyr.

Keywords: catalogs — stars: abundances — stars: fundamental parameters — stars: spectroscopic

1. INTRODUCTION

The key legacy of NASA’s *Kepler* mission is its sample of over 4000 extrasolar planets. These planets were discovered from precise, nearly continuous photometry obtained over four years of roughly 200,000 stars. Importantly, the *Kepler* planet population covers a wide range of sizes and orbital periods; sizes range from super-Jupiter to sub-Mercury and orbital periods extend from less than a day to more than a year (Thompson et al. 2018). The *Kepler* exoplanet census has, and will con-

tinue to, shed light on the diverse outcomes of planet formation.

The distribution of *Kepler* planets encodes key aspects of planet formation physics including the growth of solid cores, the accretion (and loss) of gaseous envelopes, and the prevalence of orbital migration. Gaining insights into these processes requires detailed knowledge of host stars. Until 2017, the properties of the vast majority of *Kepler* planet host stars were based on broadband photometry, which limited the accuracy of star and planet

properties. Importantly, photometric stellar radii R_* were uncertain at the $\sim 40\%$ level (Brown et al. 2011).

To address these limitations, our group conducted the California-*Kepler* Survey (CKS; Petigura et al. 2017), a spectroscopic survey of 1305 planet hosting stars observed by *Kepler* (technically “*Kepler* Objects of Interest” or “KOIs”). CKS consisted of several overlapping samples, the largest of which was magnitude-limited and included 960 planet hosts brighter than *Kepler*-band magnitude $m_{\text{Kep}} = 14.2$. Using these spectra, we measured T_{eff} , $\log g$, and $[\text{Fe}/\text{H}]$ and refined stellar M_* , R_* , and age along with planet size R_p and incident stellar flux S_{inc} .

CKS stellar properties enabled several insights into the occurrence and properties of planets as a function of stellar metallicity (Petigura et al. 2018a), stellar age (Berger et al. 2018a), planet multiplicity (Weiss et al. 2018), stellar obliquity (Winn et al. 2017), and other properties. One such insight flowed directly from the improvement in R_* errors from 40% to 10% — the radius distribution of small planets is bimodal (Fulton et al. 2017). This result has been confirmed in samples of planets with even smaller radius uncertainties that have leveraged asteroseismic or parallax constraints (Van Eylen et al. 2018; Fulton & Petigura 2018; Berger et al. 2018b; Petigura 2020).

This Radius Gap was predicted by several groups who considered the effect of photoevaporation on planetary envelopes by X-ray and extreme ultraviolet (XUV) radiation (Lopez & Fortney 2013; Owen & Wu 2013; Jin et al. 2014; Chen & Rogers 2016). Other processes that inject energy into planet envelopes may also produce the gap such as the luminosity from cooling cores (Ginzburg et al. 2018; Gupta & Schlichting 2019) or the heat from accreting planetesimals (Chatterjee & Chen 2018; Zeng et al. 2019; Wyatt et al. 2020).

Projecting the planet population along the axis of stellar mass should help illuminate which processes sculpt the distribution of small planets. For example, photoevaporation models predict that small planets will lose their envelopes out to lower bolometric flux levels around low-mass stars due to increased $F_{\text{xuv}}/F_{\text{bol}}$.

Such studies are challenging with CKS DR1 because it spans a limited range of $M_* \approx 0.8\text{--}1.4 M_\odot$. There are several reasons for this: (1) the *Kepler* selection function favored FGK stars (Batalha et al. 2010); (2) most of the *Kepler* stars below $0.8 M_\odot$ are fainter than $m_{\text{Kep}} = 14.2$; (3) the spectroscopic techniques used in Petigura et al. (2017) could not return reliable parameters for stars cooler than $T_{\text{eff}} = 4700$ K. Despite the limited M_* range, analyses of this dataset indicated an increase in the average size of sub-Neptunes with increasing stel-

lar mass (Fulton & Petigura 2018; Petigura 2020). In addition, Wu (2019) and Berger et al. (2020b) found a positive $M_*\text{--}R_p$ correlation for sub-Neptunes.

The $T_{\text{eff}} > 4700$ K boundary in CKS poses an additional challenge for interpreting trends with M_* . This boundary corresponds to $M_* > 0.65 M_\odot$ and $M_* > 0.80 M_\odot$ at $[\text{Fe}/\text{H}] = -0.4$ dex and $+0.4$ dex, respectively, the upper and lower ends of the CKS metallicity distribution. Owen & Murray-Clay (2018) discussed how this correlation acts as a confounding factor for interpreting both mass and metallicity trends in CKS DR1.

Here, we present a follow-on survey designed to expand the stellar mass range of the CKS sample. We constructed a sample of planet hosts spanning $0.5\text{--}1.4 M_\odot$ (§2) and gathered Keck/HIRES spectra of those absent from CKS DR1. New spectra of 411 hosts are presented here as CKS DR2 (§3). We analyzed the combined DR1/DR2 samples to derive temperatures, metallicities, and limits on stellar companions, which we combined with astrometric and photometric measurements to derive stellar masses, radii, and ages (§4). With updated stellar properties, we refined the precision and purity of the planet catalog (§5). We analyzed both the distribution of detected planets (§6) and the underlying occurrence rate (§7). We offer some comparisons to planet formation theory (§8) and provide a brief summary and conclusion (§9).

2. SAMPLE SELECTION

Our main goal was to measure planet occurrence as a function of stellar mass. This requires a sample of planets drawn from a well-defined sample of stars spanning a broad range in stellar mass. Before describing our selection function in detail, we first orient the reader with an overview of our survey design and practical considerations. When we designed our survey in 2018, our goal was to increase the number of low-mass stars with CKS-quality parameters given a fixed telescope allocation of 15 Keck/HIRES nights. Simply extending the magnitude-limited survey beyond $m_{\text{Kep}} = 14.2$ would have included many stars with $T_{\text{eff}} \approx 5000\text{--}6500$ K that were already well-sampled by DR1 (see Figure 1). Observing all targets down to $m_{\text{Kep}} = 16.0$ in the manner described in §3 would have required ~ 75 nights and was impractical. Instead, we adopted a selection function and observing strategy that favored cool stars but mitigated magnitude-dependent effects. The two mitigation strategies are described in detail in §2–5 but briefly involved selecting single isolated stars and collecting spectra of homogeneous quality.




Figure 1. Panel (a): Gray points show effective temperature and apparent magnitude m_{Kepler} of stars observed by *Kepler*; red points are stars that pass the selection criteria described in §2. Panel (b): same as (a), but for planet hosts. Panel (c): same as (a) but showing *Gaia* absolute *G*-band magnitude M_G and $B_p - R_p$ color. The boundaries of our main-sequence filter are shown as red-dashed lines. Panel (d): same as (c) but for planet hosts.

We began with 184933 stars in the union of the following three catalogs: the *Kepler* project Data Release 25 (DR25) stellar properties catalog (Mathur et al. 2017, M17 hereafter), the *Gaia* DR2 catalog (Gaia Collaboration et al. 2018), and the Berger et al. (2020b) catalog of stellar properties (B20 hereafter). We then applied the following filters:

1. *Stellar brightness and effective temperature.* Our sample consists of two magnitude-limited components. If $T_{\text{eff}} = 5000\text{--}6500$ K (as measured by M17), we required $m_{\text{Kepler}} < 14.2$; if $T_{\text{eff}} = 3000\text{--}5000$ K, we required $m_{\text{Kepler}} < 16.0$. A total of 77243 stars remained.
2. *Main-sequence stars.* We selected main-sequence stars using their position in the *Gaia* color-magnitude diagram. Stars were included if they reside within the boundaries shown in Figure 1. A total of 47502 stars remained.
3. *Single stars: Gaia source catalog.* If a transit is observed within a *Kepler* aperture containing more

than one star, it is often not possible to identify the star hosting the transiting object. Whichever star is the host, the transit depth is diluted by the neighboring star(s) and the inferred planet radius will be biased to smaller values. To ensure high-precision R_p , we worked to exclude targets that are diluted by $>10\%$, which would result in a R_p error of $>5\%$. We excluded stars where the cumulative *G*-band flux contribution from all *Gaia*-identified neighboring stars within 4 arcsec exceeded 10%. Companions outside of this limit are resolved as separate sources in the KIC, which had an average FWHM of 2.5 arcsec (Brown et al. 2011). The *Kepler* project removed the diluting flux from such sources during the photometric extraction (see the discussion of the “crowding metric” in Stumpe et al. 2012). A total of 46468 stars remained.

4. *Single stars: Gaia astrometric noise.* Companions within 1 arcsec are often not resolved as separate sources in *Gaia* DR2, but can be identified us-

ing the residuals to the astrometric fits. *Gaia* Renormalized Unit Weight Error (RUWE) is conceptually similar to the square root of the reduced χ^2 and is close to unity when the astrometric data are consistent with parallax and proper motion only, with no additional variation due to any companions (Lindgren 2018). Analyses of stars with high-resolution imaging have demonstrated that requiring $\text{RUWE} < 1.2$ removes companions with $\Delta m_G < 3$ mag and separations of $\rho = 0.1\text{--}1$ arcsec (Bryson et al. 2020, Wood et al. 2021, Kraus et al. in preparation). A total of 39423 stars remained.

We cross-referenced these stars against the Thompson et al. (2018) catalog of KOIs, T18 hereafter, which we accessed from the NASA Exoplanet Archive (NEA; Akeson et al. 2013). This catalog was derived from the final planet search conducted by the *Kepler* project (DR25). Among our filtered stellar sample, T18 lists 39423 stars hosting 2373 KOIs. We subjected the KOIs to the following additional filters:

5. *Candidate reliability.* T18 dispositioned KOIs using a fully automated procedure called the “Robovetter” that performed a series of tests on a suite of data quality metrics designed to identify false positives due to data anomalies and eclipsing binaries. When setting the Robovetter thresholds, T18 emulated the human classification of Threshold Crossing Event Review Team (TCERT). Robovetter dispositioned all KOIs as either “False Positive” or “Planet Candidate.” Some KOIs fall near the boundary between the two dispositions. T18 quantified the proximity of each KOI to this boundary through Monte Carlo simulations. T18 reported a “disposition score” which is the fraction of Monte Carlo simulations classified as planet candidates. We required “Planet Candidate” status¹ with a disposition score² exceeding 75%. Finally, we removed a handful of known false positives that cleared the Robovetter that are listed in the DR25 supplemental table.³ A total of 1425 KOIs orbiting 1003 host stars remained.
6. *Candidate signal-to-noise ratio (S/N).* Low S/N candidates are more likely to be false positives (see, e.g., Bryson et al. 2020) and yield lower overall precision on R_p/R_\star . The *Kepler* project re-

ported S/N as a “Multiple Event Statistic” (MES; Jenkins 2002).⁴ We required $\text{MES} > 10$. A total of 1250 KOIs orbiting 891 host stars remained.

7. *Transit Modeling.* To compute planet properties, we used posterior samples of transit parameters derived from Markov Chain Monte Carlo (MCMC). A small fraction of the T18 KOIs were not modeled using MCMC.⁵ Most of these are dispositioned as false positives. We removed a handful of candidates that passed the previous cuts. A total of 1246 KOIs orbiting 888 host stars remained.

Figure 1 shows the application of these filters to the field star and planet samples. In this paper, we refer to these 888 stars as the “CKS Extended Mass” (CXM) sample.

3. SPECTROSCOPIC OBSERVATIONS

We worked to compile a homogeneous spectroscopic catalog of all CXM stars from new and archival observations. All spectra were obtained with Keck/HIRES (Vogt et al. 1994) to ensure common systematics in our analysis. Our goal was to gather spectra with $R \geq 60,000$ and $\text{S/N} \geq 20 \text{ pix}^{-1}$ on blaze at 5500 \AA .

The inventory of DR1/DR2 spectra and the CXM host star sample are shown in Figure 2. By design, there was substantial overlap between the 1305-star DR1 sample and the CXM sample (628/888 spectra). The DR1 spectra have $R \geq 60,000$ and $\text{S/N} \geq 45 \text{ pix}^{-1}$ and thus met our spectral quality goals. The 411 spectra in DR2 are a mixture of 81 archival HIRES spectra taken before 2018 and 330 new spectra taken by our team since then. For the archival spectra, all have $R \geq 60,000$ and all but 4 had $\text{S/N} \geq 20 \text{ pix}^{-1}$. An additional 260/888 CXM stars have spectra in DR2.

Below, we describe the observations of the post-2018 DR2 spectra, which were executed as part of a NASA Key Strategic Mission Support program (PI: Petigura). Our strategy was identical to that of CKS DR1 except we exposed to a lower S/N. We used the standard setup of the California Planet Search (CPS; Howard et al. 2010). We observed stars through the “C2” slit with sky-projected dimensions of $0.86 \text{ arcsec} \times 14.0 \text{ arcsec}$, which achieves a spectral resolution of $R \approx 60,000$. We used the exposure meter to attain a designed S/N which is measured at the peak of the blaze function at $\lambda = 5500 \text{ \AA}$.

Given the magnitude-limited nature of DR1, most DR2 stars had $m_{\text{Kep}} = 14.2\text{--}16.0$ mag. It would have

¹ NEA data column: *koi_pdisposition*

² NEA data column: *koi_score*

³ NEA data column: *koi_pdisposition*

⁴ NEA data column: *koi_max_mult_ev*

⁵ NEA data column: *koi_fittype*

been prohibitively expensive to observe these stars to the same S/N level as DR1 (45 pix^{-1}), so we used the exposure meter to achieve $S/N = 20 \text{ pix}^{-1}$ for stars in this magnitude range. In §4, we demonstrate that the additional photon-limited uncertainties are smaller than other sources of uncertainty in the extracted parameters.

For faint targets, the sky background is occasionally comparable to the stellar spectrum depending on moon separation, moon phase, and cloud cover. In our spectral reduction, we removed the sky background (see Batalha et al. 2011 for details), but it still contributes to exposure meter counts. Thus, at fixed m_V and exposure meter setting, additional sky background would cause exposures to terminate early. While at the telescope, we used a custom script to compute the sky contribution in realtime and adjusted the exposure meter setting such that we would reach the desired S/N *after* sky subtraction.

We reduced our spectra according to the standard practices of CPS. We removed the blaze function by dividing our spectra by a composite spectrum of spectrally flat standards compiled by Clubb et al. (2018). One component of our spectroscopic analysis involved registering each spectrum to a common wavelength scale. We used the SpecMatch-Empirical code (Yee et al. 2017) that includes a ladder of reference spectra that have been registered against the National Solar Observatory solar atlas (Kurucz et al. 1984). Figure 3 shows the Mg I b region for representative spectra over a range of T_{eff} . The deblazed spectra are available via the Keck Observatory Archive⁶ the CFOP website,⁷ and a website maintained by our team.⁸ We have also made available the standard observatory-frame wavelength solution applicable to every spectrum, which is accurate to within one reduced pixel or 1.2 km/s. The registered spectra are also available at the above repositories.

4. HOST STAR CHARACTERIZATION

4.1. Spectroscopy

We derived stellar parameters from our spectra using two related and complementary codes: SpecMatch-Synthetic and SpecMatch-Empirical. Both codes were designed to yield high quality parameters even at moderate or low S/N. A detailed description of SpecMatch-Synthetic is available in Petigura (2015) and the code’s application to CKS DR1 is given in Petigura et al. (2017). In brief, the code generates a synthetic spec-




Figure 2. Venn diagram illustrating the number of host stars belonging to various samples described in this paper. CXM—888 stars in extended stellar mass sample described in §2 that forms the basis of the exoplanet demographic work described in §6 and §7. DR1—1305 stars from (Petigura et al. 2017), DR2—411 stars with new spectra presented in this work.

trum by interpolating within a set of library spectra computed over a grid of T_{eff} , $\log g$, and $[\text{Fe}/\text{H}]$. It then applies line broadening kernels that account for the instrumental profile, stellar rotation, and macroturbulence. The library spectra were computed by Coelho et al. (2005) under the assumption of local thermodynamic equilibrium (LTE) using the Castelli & Kurucz (2004) model atmospheres. The code derives T_{eff} , $\log g$, $[\text{Fe}/\text{H}]$, and $v \sin i$ using non-linear optimization of an l_2 metric, which is evaluated over five spectral segments with a combined spectral bandwidth of 380 Å between 5200 Å and 6260 Å.

The uncertainties of SpecMatch-Synthetic parameters have been extensively vetted for stars having $T_{\text{eff}} = 4700\text{--}6500$ K. Errors are dominated by systematic uncertainties which are $\sigma(T_{\text{eff}}) = 100$ K, $\sigma(\log g) = 0.10$ dex, and $\sigma([\text{Fe}/\text{H}]) = 0.06$ dex. Poisson uncertainties are smaller than systematic uncertainties even at low S/N because the code fits a wide spectral bandpass and combines information from many lines. At our lowest S/N (20 pix^{-1}), photon-limited errors are $\sigma(T_{\text{eff}}) = 34$ K, $\sigma(\log g) = 0.05$ dex, and $\sigma([\text{Fe}/\text{H}]) = 0.02$ dex, respectively, below the systematic floor. Above 6500 K, the quality of the derived parameters degrades due to the reduced number of lines and substantial rotational broadening. As T_{eff} drops below 4700 K, the number of lines in SpecMatch-Synthetic wavelength regions increases dramatically due to molecular features (of which MgH is the most prominent). The synthetic spectra do not accurately reproduce these complex spectra and the derived parameters suffer.

⁶ <http://www2.keck.hawaii.edu/koa/public/koa.php>

⁷ <http://cfop.ipac.caltech.edu>

⁸ <http://astro.caltech.edu/~howard/cks/>




Figure 3. The black lines show a segment of HIRES spectra containing the Mg I b triplet for seven representative spectra observed by our team to illustrate the spectral resolution $R \geq 60,000$ and $S/N \geq 20 \text{ pix}^{-1}$ (see §3). The red lines are SpecMatch-Empirical fits (see §4.1).

By design, the CXM sample included many stars cooler than 4700 K so we needed another method that is better suited to cool stars. In preparation for this project, Yee et al. (2017) developed SpecMatch-Empirical, which sidesteps the difficulties of cool star spectral synthesis. In brief, SpecMatch-Empirical compares a target spectrum to a spectral library of 404 stars with precise and accurate T_{eff} , R_{\star} , and $[\text{Fe}/\text{H}]$ measured through a combination of the following techniques: interferometry, asteroseismology, line-by-line LTE spectral synthesis, and spectrophotometry. The code identifies the top five matches based on an l_2 metric. The target spectrum is then fit using a linear combination of these five spectra over a broad 1000 Å spectral region spanning 5000 Å to 6000 Å. The extracted parameters are a weighted average of the input parameters. For the representative spectra in Figure 3 we have over-plotted the best-fit spectra.

Yee et al. (2017) validated the precision and accuracy of SpecMatch-Empirical across the HR diagram and found that quality of the derived parameters varies in this space due to the relative density of library spectra. For stars having $T_{\text{eff}} = 3500\text{--}5000$ K and $[\text{Fe}/\text{H}]$ between -0.5 and $+0.5$ dex, the code achieves uncertainties of $\sigma(T_{\text{eff}}) = 60$ K and $\sigma([\text{Fe}/\text{H}]) = 0.12$ dex. Because this code analyzes 1000 Å of spectrum, it is robust to photon-limited uncertainties at low S/N. For our lowest S/N spectra at 20 pix^{-1} , Poisson uncertainties amount to $\sigma(T_{\text{eff}}) = 6$ K and $\sigma([\text{Fe}/\text{H}]) = 0.004$ dex and thus contribute negligibly to the overall error budget.

Table 1 lists our adopted stellar parameters for the 1716 star union of DR1 and DR2. Given the relative strengths and weakness of the two codes, we split our sample into two groups at $T_{\text{eff}} = 4800$ K, as measured by SpecMatch-Empirical. We report SpecMatch-Synthetic parameters for the 1350 stars above this threshold and SpecMatch-Empirical for the 366 below.

In the following section, we combine our spectroscopic T_{eff} and $[\text{Fe}/\text{H}]$ measurements with astrometric and photometric constraints to characterize the stellar hosts. Before proceeding, we note that the input spectra have a bimodal distribution of S/N and consider whether this could bias the DR1 and DR2 stellar parameters relative to one another. As explained above, the photon-limited T_{eff} and $[\text{Fe}/\text{H}]$ errors at $S/N = 20 \text{ pix}^{-1}$ are less than half the systematic errors. Any S/N dependent offsets are washed out by the larger systematic uncertainties.

4.2. Stellar Properties

We derived stellar masses, radii, and ages using our spectroscopic measurements of T_{eff} and $[\text{Fe}/\text{H}]$, along with astrometric and photometric constraints. Our methodology closely follows that of Fulton & Petigura (2018). We give a brief summary below, noting differences where relevant.

We measured R_{\star} using the isclassify package in its “direct” mode. This code evaluates R_{\star} from the Stefan–Boltzmann law,

$$R_{\star} = \left(\frac{L_{\text{bol}}}{4\pi\sigma_{\text{sb}}T_{\text{eff}}^4} \right)^{1/2}, \quad (1)$$

where L_{bol} is the bolometric stellar luminosity and σ_{sb} is the Stefan-Boltzmann constant. L_{bol} is directly related to bolometric magnitude M_{bol} , which may be expressed as

$$M_{\text{bol}} = m - \mu - A + BC, \quad (2)$$

where m is the apparent magnitude, μ is the distance modulus, A is the line-of-sight extinction, and BC is the bolometric correction. We summarize each input below along with their typical uncertainties. Given the discontinuity in our sample selection at $m_{\text{Kep}} = 14.2$ described in §2, we report median uncertainties above and below this value.

1. *Apparent magnitude.* We used 2MASS K -band photometric measurements because dust extinction is less severe in the infrared. We used 2MASS m_K which have a median precision of 0.02 mag for $m_{\text{Kep}} < 14.2$ and 0.03 mag for $m_{\text{Kep}} > 14.2$.
2. *Extinction.* We accounted for K -band extinction A using the 3D dust map of Green et al. (2019), an update to the Green et al. (2018) map used in Fulton & Petigura (2018). Median A is 0.01 mag for both bright and faint subsets.
3. *Distance modulus.* We used *Gaia* DR2 parallaxes and applied a correction of +0.053 mas to account for a known systematic offset in the *Kepler* field (Zinn et al. 2018). The parallaxes have a median precision of 1.2% for $m_{\text{Kep}} < 14.2$ and 1.8% for $m_{\text{Kep}} > 14.2$. *Isoclassify* handles the conversion between parallax and distance modulus using a Bayesian framework.
4. *Bolometric correction.* *Isoclassify* interpolates over a grid of bolometric corrections computed by the MESA Isochrones and Stellar Tracks project (MIST v1.2, Choi et al. 2016; Dotter 2016; Paxton et al. 2011, 2013, 2015). Uncertainties in T_{eff} dominate the uncertainty of the K -band bolometric correction and a 60 K uncertainty translates to a ≈ 0.03 mag error on BC .

Table 1 lists our measured R_{\star} , and Figure 4 shows the distribution of fractional R_{\star} uncertainties. The median fractional uncertainty is 4.0% for $m_{\text{Kep}} < 14.2$ and 4.4% for $m_{\text{Kep}} > 14.2$.

We also ran *isoclassify* in its “grid” mode with the same constraints. In this mode, the code queries the MIST isochrones which provide various stellar properties over a grid of M_{\star} , $[\text{Fe}/\text{H}]$, and age. The grid extends to 20 Gyr to minimize grid edge-effects, which can bias the parameter posteriors. At each grid point, *isoclassify*

computes the likelihood that point is consistent with the input constraints.

Marginalized values are determined by direct integration. We list $M_{\star, \text{iso}}$, $R_{\star, \text{iso}}$, $\rho_{\star, \text{iso}}$, and $\text{age}_{\star, \text{iso}}$ in Table 1. The “iso” subscript indicates that these parameters are forced to be consistent with MIST isochrones. The median fractional uncertainty for $M_{\star, \text{iso}}$ is 3.8% for $m_{\text{Kep}} < 14.2$ and 3.7% for $m_{\text{Kep}} > 14.2$; for $\rho_{\star, \text{iso}}$ it is 8.7% for $m_{\text{Kep}} < 14.2$ and 9.7% for $m_{\text{Kep}} > 14.2$. A handful of stars have $\text{age}_{\star, \text{iso}} > 14$ Gyr because the grid extends to 20 Gyr for reasons explained above. Age uncertainties vary widely across the HR diagram and are discussed later in §6.

5. PLANET CATALOG

In the previous section, we computed stellar properties in the combined DR1/DR2 dataset. For the remainder of the paper, we focus on the 1246 planets associated with the 888 star CXM subsample (see Figure 2). In this section, we first compute planet properties given our updated stellar properties and then apply an additional set of quality checks to produce a high-reliability sample of planets for demographic work in later sections.

5.1. Planet properties

We computed R_p from our R_{\star} (listed in Table 1) and T18’s modeling of *Kepler* light curves. We multiplied our posterior samples on R_{\star} by the MCMC-derived posterior samples on R_p/R_{\star} (available at the NEA). Median fractional error on R_p/R_{\star} is 3.9% for $m_{\text{Kep}} < 14.2$ and 4.8% for $m_{\text{Kep}} > 14.2$. Uncertainties in R_{\star} and R_p/R_{\star} are independent and propagated into R_p , and we quote the 16th, 50th, and 84th posterior quantiles in Table 2. The median fractional precision in R_p is 5.6% for $m_{\text{Kep}} < 14.2$ and 6.3% for $m_{\text{Kep}} > 14.2$. Figure 4 shows the fractional precision of R_p , R_{\star} , and R_p/R_{\star} . For most of the sample, R_{\star} errors are comparable to or larger than R_p/R_{\star} errors. However, for a small subset, R_p/R_{\star} errors dominate.

A note about the provenance of R_p/R_{\star} : There are a number of published catalogs of R_p/R_{\star} from *Kepler* project, of which T18 is the most recent. The NEA table corresponding to the T18 catalog lists the “best-fit” R_p/R_{\star} , not the posterior medians. Petigura (2020) found that these best-fit values contribute noise to R_p/R_{\star} from the non-linear optimization and recommended using posterior medians which are a more robust estimator.

We compared our planet radii to T18, by computing the ratio of the radii $r = R_{p, \text{us}}/R_{p, \text{T18}}$. There was a negligible systematic offset $\text{mean}(r) = 1.00$ but a substantial dispersion $\text{RMS}(r) = 20\%$. The dispersion is dominated by uncertainties in the T18 R_p .

Table 1. Stellar Properties

KOI	m_K	π	T_{eff}	[Fe/H]	$v \sin i$	prov	R_*	$M_{*,\text{iso}}$	$R_{*,\text{iso}}$	$\rho_{*,\text{iso}}$	$\text{age}_{*,\text{iso}}$	π_{spec}	SB2	CXM
	mag	mas	K	dex	km s^{-1}		R_{\odot}	M_{\odot}	R_{\odot}	g/cc	Gyr	mas		
1	9.85	4.67	5857	0.02	1.3	syn	1.04	1.02	1.03	0.91	3.7	4.76	1	1
2	9.33	2.96	6403	0.22	5.1	syn	2.00	1.50	1.99	0.19	1.8	3.72	1	1
3	7.01	26.50	4619	0.16	...	emp	0.78	0.76	0.75	1.79	11.2	23.80	1	1
4	10.20	1.29	5948	-0.27	38.0	syn	3.26	1.45	3.22	0.04	2.0	0.93	3	0
6	10.99	2.13	6356	0.05	11.8	syn	1.29	1.23	1.28	0.57	1.5	2.24	1	0

NOTE—Properties of 1716 planet hosting stars from the union of the DR1 and DR2 datasets. m_K is 2MASS K -band apparent magnitude and π is *Gaia* DR2 parallax. T_{eff} , [Fe/H], and $v \sin i$ were derived by one of two methods listed in the ‘prov’ column: ‘syn’—SpecMatch-Synthetic and ‘emp’—SpecMatch-Empirical. R_* is our adopted stellar radius from the Stefan-Boltzmann law. Stellar properties with the ‘iso’ subscript incorporate constraints from the MIST isochrones. ‘SB2’ encodes the limits on spectroscopic binaries (SB2s) using the *ReaMatch* code (Kolbl et al. 2015): 1—No detection of SB2 with $\Delta m_V \lesssim 5$ mag and $\Delta RV \gtrsim 12$ km s^{-1} ; 2— $T_{\text{eff}} < 3500$ K, star unfit for ReaMatch; 3— $v \sin i > 10$ km s^{-1} , star unfit for ReaMatch; 4—Ambiguous detection; 5—Obvious detection. Note: *ReaMatch* computes T_{eff} and $v \sin i$ independently from SpecMatch-Synthetic and SpecMatch-Empirical. The ‘CXM’ flag is 1 if the star is in the 888 star CXM subset. Median uncertainties are as follows: m_K —0.02 mag; π —1.4%; T_{eff} —100 K, [Fe/H]—0.06 dex; $v \sin i$ —1 km s^{-1} ; R_* —3.9%; $M_{*,\text{iso}}$ —3.7%; $R_{*,\text{iso}}$ —2.6%; $\rho_{*,\text{iso}}$ —8.9%; $\text{age}_{*,\text{iso}}$ —0.24 dex; π_{spec} —16%. Table 1 is published in its entirety with uncertainties in machine-readable format in the online journal.

Table 2. Planet Properties

Planet	P	R_p/R_*	T	R_p	$T_{\text{max,circ}}$	a	S_{inc}	samp
	d	%	hr	R_{\oplus}	hr	au	S_{\oplus}	
K00001.01	2.5	12.39	1.29	14.04	2.83	0.037	814.85	0
K00002.01	2.2	7.52	3.55	16.42	4.58	0.039	3855.68	1
K00003.01	4.9	5.80	2.23	4.92	2.84	0.053	80.94	1
K00007.01	3.2	2.45	3.84	4.07	4.29	0.046	1082.71	1
K00010.01	3.5	9.21	2.75	15.51	4.47	0.050	1262.15	1

NOTE—Properties of 1246 planets orbiting the 888 star CXM sample. Orbital period P , planet-to-star radius ratio R_p/R_* , and transit duration T were measured from *Kepler* photometry by Thompson et al. (2018). R_p follows from R_p/R_* and R_* . The expected duration of a centrally transiting object on a circular orbit $T_{\text{max,circ}}$, semi-major axis a , and incident bolometric flux S_{inc} are determined from Kepler’s Third Law and the Stefan-Boltzmann law. The ‘samp’ flag is 1 if planet is in curated sample of 970 planets described at the end of §5. Median uncertainties are as follows: P —3 ppm; R_p/R_* —4.3%; T —2.3%, R_p —5.7%; $T_{\text{max,circ}}$ —2.9%; a —1.2%; S_{inc} —9.2%. Table 2 is published in its entirety with uncertainties in machine-readable format in the online journal.

Two additional quantities were relevant to our goal of precise planet radii: the observed transit duration T and the expected duration of centrally transiting planet on a circular orbit $T_{\text{max,circ}}$. Here, T refers to time between the midpoints of first/second contact and third/fourth contact. In the T18 modeling, transit shape was set by the following free parameters: period P , transit epoch T_0 , impact parameter b , mean stellar density assuming a circular orbit $\rho_{*,\text{circ}}$, and R_p/R_* . We derived T and

its uncertainties according to

$$T = 2.036 \text{ hr} (1 - b^2)^{1/2} \left(\frac{P}{1 \text{ d}} \right)^{1/3} \left(\frac{\rho_{*,\text{circ}}}{1 \text{ g cm}^{-3}} \right)^{-1/3} \quad (3)$$

using the T18 MCMC chains. We computed $T_{\text{max,circ}}$ according to

$$T_{\text{max,circ}} = 2.036 \text{ hr} \left(\frac{P}{1 \text{ d}} \right)^{1/3} \left(\frac{\rho_{*,\text{iso}}}{1 \text{ g cm}^{-3}} \right)^{-1/3} \quad (4)$$

Both T and $T_{\text{max,circ}}$ are listed in Table 2 along with semi-major axis a and incident stellar flux S_{inc} .




Figure 4. Fractional precision of R_p , R_* , and R_p/R_* for the planets in the CXM sample.

5.2. Refining the planet sample

While the initial CXM sample was optimized for high-reliability planets (§2), we incorporated additional diagnostics from our spectra to further increase sample purity. We first filtered on the following stellar properties:

1. *Rotation rate.* Our spectroscopic codes are unreliable at $v \sin i > 20 \text{ km s}^{-1}$ (Petigura et al. 2017). We excluded 17 stars above this threshold which all had $T_{\text{eff}} > 6500 \text{ K}$ as measured by B20. A total of 1227 KOIs orbiting 871 hosts remained.
2. *Spectroscopic parallax.* Following Fulton & Petigura (2018), we used isoclassify to compute a “spectroscopic parallax” constrained by our T_{eff} , $[\text{Fe}/\text{H}]$, m_K , and the MIST models. If spectroscopic and trigonometric parallaxes differ significantly, we conclude that the input parameters used to compute R_* are inconsistent due to unresolved binaries or other effects. We removed 9 stars where these parallaxes differed by 4σ or more. We chose a 4σ over a 3σ threshold because the probability of a 3σ event is 3×10^{-3} and thus we expect ≈ 3 such events in a sample of $\approx 10^3$ stars. The probability of a 4σ event 6×10^{-5} so a single such event is unlikely. A total of 1215 KOIs orbiting 862 hosts remained.
3. *Secondary spectra.* We removed 1 star where we identified a secondary set of spectral lines using the methodology of Kolbl et al. (2015), which is sensitive to binaries having $\Delta m_V \lesssim 5 \text{ mag}$ and $\Delta v \gtrsim 12 \text{ km s}^{-1}$. A total of 1214 KOIs orbiting 861 hosts remained.

Next, we filtered based on the following planet properties:

4. *Impact parameter.* We removed grazing and high-impact parameter planets by excluding planets where the median of the b posterior exceeded 0.8. Petigura (2020) found, however, that for most *Kepler* light curves, this cut is ineffective at excluding high- b transits because b is nearly unconstrained. For such transits, R_p/R_* is biased by 10–20% due to the strong b - R_p/R_* covariance. However, $T/T_{\text{max,circ}}$ is an effective proxy for b . In addition to our explicit filter on b , we also required $T/T_{\text{max,circ}} > 0.6$ in order to exclude transits with $b > 0.8$. While planets with eccentric orbits that transit near periastron also produce low $T/T_{\text{max,circ}}$, Petigura (2020) showed that fewer than 20% of *Kepler* planets with $T/T_{\text{max,circ}} < 0.6$ have $b < 0.8$. A total of 973 KOIs orbiting 704 hosts remained.
5. *Radius precision.* We required fractional errors on R_p of 20% or less. A total of 970 KOIs orbiting 703 hosts remained.

The properties of the stars that passed these cuts are shown in Figure 5. They span a mass range of $M_* \approx 0.5$ – $1.4 M_\odot$.

6. DISTRIBUTION OF DETECTED PLANETS

From here on, we describe and interpret the features in our curated sample of 970 planets. In this section, we treat the population of detected planets before turning to the occurrence distribution in §7. We explore how the Radius Gap correlates with P , S_{inc} , M_* and age in §6.1. We then inspect how the super-Earth and sub-Neptune populations vary with stellar mass in §6.2.

6.1. Dependence of the Radius Gap on period, flux, stellar mass, and stellar metallicity.

Figure 6 shows several views of our planet sample. In the four panels, the y-axes show updated R_p , and the x-axes show P , S_{inc} , M_* , and $[\text{Fe}/\text{H}]$. The contours show the relative density of detected planets in these domains using a Gaussian kernel density estimator (KDE), and the KDE bandwidths are specified in the caption. Figure 7 is analogous to Figure 6 but with R_p restricted to 1–4 R_\oplus for a detailed view of the Radius Gap.

The P - R_p distribution and closely related S_{inc} - R_p distribution of *Kepler* planets have been extensively studied in previous works. We begin by remarking on how several known features that appear in our extended stellar mass sample. Most *Kepler* planets fall into one of two size categories: “super-Earths” and “sub-Neptunes.” For definitiveness, we adopt the following size limits for the two planet classes: $R_p = 1.0$ – $1.7 R_\oplus$ and $R_p = 1.7$ – $4.0 R_\oplus$. The Radius Gap separating the populations




Figure 5. Properties of host stars in the filtered CXM sample described in §5.2. Panel (a): Stellar mass and metallicity. Panel (b): Stellar radius and effective temperature. The color scale conveys the isochrone age uncertainty in dex (half the difference between the 84th and 16th percentiles of the age posterior). Panel (c): Stellar metallicity and age. Panel (d): Stellar mass and age where the color scale is equivalent to panel (b). Several stars have ages exceeding 14 Gyr for reasons we explain in §4.2. We discuss correlations between mass, metallicity, and age in §6.2.

was observed by [Fulton et al. \(2017\)](#) in CKS DR1 ($M_{\star} \approx 0.8\text{--}1.3 M_{\odot}$), [Van Eylen et al. \(2018\)](#) in a sample of stars with asteroseismic detections ($M_{\star} \approx 1.0\text{--}1.4 M_{\odot}$), and in other subsequent works. We again resolve the Radius Gap, despite the broader M_{\star} range of the CXM sample ($M_{\star} \approx 0.5\text{--}1.4 M_{\odot}$).

The distribution of super-Earths is centered at shorter periods and higher incident fluxes than the distribution of sub-Neptunes. Compared to super-Earths, there are few sub-Neptunes with $P < 3$ days or $S_{\text{inc}} > 300 S_{\oplus}$. This paucity of sub-Neptunes is not a selection effect because larger planets are easier to detect. This zone of low sub-Neptune occurrence is sometimes referred to as the Sub-Neptune Desert (see, e.g., [Szabó & Kiss 2011](#); [Beaugé & Nesvorný 2013](#); [Lundkvist et al. 2016](#)). In

§7, we quantify the sharpness of this boundary and explore its dependence on stellar mass. The relative lack of super-Earths at $P > 30$ days and $S_{\text{inc}} < 10 S_{\oplus}$ is closely tied survey completeness, also addressed in §7.

We wish to characterize the slope of the Radius Gap in P - R_p and S_{inc} - R_p space because it is a metric by which to test theoretical predictions. However, fitting a parameterized description of the Radius Gap is challenging since it involves characterizing the *absence* of planets. [Van Eylen et al. \(2018\)](#) used a support vector classification (SVC) scheme to find the line which maximized the distance between the super-Earth and sub-Neptune populations. The SVC scheme struggles for our sample because the gap is not devoid of planets. (Recently, [David et al. 2021](#) investigated how regular-




Figure 6. Panel (a): sizes and orbital periods of detected planets after applying the filters described in §5.2. The contours show a Gaussian kernel density estimate (KDE) of the number density of planets in this space. The kernel bandwidth is 0.029 dex in R_p and 0.18 dex in P as shown by the symbol labelled ‘bw’. The ‘uncert’ symbol shows the median R_p uncertainty (P uncertainties are smaller than the line width). Panels (b–d): same as (a), except that x-axis is S_{inc} , M_* , and $[\text{Fe}/\text{H}]$ and the kernel bandwidths along the horizontal axes are 0.30, 0.061, and 0.1.

ization can assist the SVC identification of the Radius Gap in the CKS sample.)

We adopted a different approach starting with relative number density of detected planets in the P - R_p plane $d^2N/d\log R_p d\log P$ as measured by our KDE. This quantity is shown as contours in Figure 7a. We then found the minimum density along 100 vertical lines spanning $\log R_p = 0.15$ – 0.35 spaced uniformly in $\log P$ over 0.5 to 1.5. We then fit the train of 100 minima with following power-law

$$R_p(P) = R_{p,0} \left(\frac{P}{10 \text{ days}} \right)^m, \quad (5)$$

where the slope m and intercept $R_{p,0}$ are free parameters. To be clear, $R_{p,0}$ should be interpreted as the midpoint of the Radius Gap at 10-day orbital periods. We chose 10 days as a convenient reference point near the middle of the super-Earth and sub-Neptune popula-

tions shown in Figure 7a. To determine uncertainties we generated 1000 bootstrap resamples with replacement of the planet population shown in Figure 7a (Press 2002). For each bootstrapped population, we recomputed the planet number density with our KDE, the train of minima, and the best-fitting power-law. We found $m = d\log R_p/d\log P = -0.11^{+0.02}_{-0.02}$ and $R_{p,0} = 1.84^{+0.03}_{-0.03} R_{\oplus}$, which reflect the 16th, 50th, and 84th percentiles. Figure 7 shows the credible gap models. We explored the sensitivity of our method to different period ranges by perturbing the upper and lower boundaries by ± 0.25 dex. The derived parameters were consistent to 1σ . We also explored the sensitivity of our method to our adopted KDE period bandwidth. A very wide bandwidth will flatten the Radius Gap (i.e. bias $|m|$ toward smaller values). We repeated our analysis with a much smaller bandwidth of 0.04 dex and found that m and $R_{p,0}$ changed by less than 1σ .




Figure 7. A zoomed-in version of Figure 6 highlighting the Radius Gap, which is visible in all projections. In the M_* projection, the typical sub-Neptune becomes larger with M_* , while the super-Earths remain the same size. The bands are power-law fits to Radius Gap (see §6.1).

We performed similar analyses to fit the Radius Gap as a function of S_{inc} , M_* , and $[\text{Fe}/\text{H}]$. Table 3 lists the parametric models used in the fits and the credible range of parameters. Figure 7 shows the range of credible gap models. We observed a positive slope with incident flux ($d \log R_p / d \log S_{\text{inc}} = 0.06^{+0.01}_{-0.01}$), a positive slope with stellar mass ($d \log R_p / d \log M_* = 0.18^{+0.08}_{-0.07}$), and no significant correlation with metallicity ($d \log R_p / d [\text{Fe}/\text{H}] = 0.01^{+0.05}_{-0.06}$).

We explored how the P - R_p distribution of planets changes with M_* . The left panels of Figure 8 shows this distribution for three different bins of stellar mass with the following boundaries: 0.5, 0.7, 1.0, and 1.4 M_\odot . The super-Earth and sub-Neptune populations are distinct in each bin, but their locations vary with M_* . As M_* increases, the population of super-Earths and sub-Neptunes separates; the super-Earth population is confined to $R_p < 1.7 R_\oplus$ while the sub-Neptunes grow in size. We measured the slope and intercept of the Ra-

dius Gap in these three bins. They are given in Table 3 and are consistent to 2σ .

We also explored how the S_{inc} - R_p distribution changes with M_* . Figure 9 is analogous to Figure 8 except it shows the S_{inc} - R_p distribution for three bins of stellar mass. The credible models describing the Radius Gap in this space are consistent to 2σ .

6.2. Dependence of planet size on stellar mass

In the previous section, we measured the slope of the Radius Gap in the R_p - M_* plane and noted that it increases with M_* . Inspecting Figure 7c, we see qualitatively that the Radius Gap trend driven by the fact that the typical sub-Neptune grows with stellar mass while the typical super-Earth does not. Here, we quantify these trends and explore possible confounding factors.

We first measured the sub-Neptune R_p - M_* correlation by selecting all sub-Neptunes from the population shown in Figure 7 based on their radii ($R_p = 1.7$ – $4.0 R_\oplus$). We




Figure 8. Left panels are analogous to Figure 7a, except we have split the planet sample into three bins of stellar mass. Right panels show the occurrence rate density of planets $df/d \log P/d \log R_p$. The maximum occurrence rate density increases with decreasing M_* (note different color scales). Regions of low planet detectability, defined as $n_{\text{trial}} < 50$, are gray. The typical sub-Neptune size grows with M_* , while super-Earths have a nearly constant size. The bands show fits to the Radius Gap.

then fit a power-law

$$R_p = R_{p,0} \left(\frac{M_*}{M_\odot} \right)^\alpha. \quad (6)$$

We used the Levenberg-Marquardt algorithm as implemented in the *lmfit* Python package to find the best-fit parameters and measured uncertainties via 1000 bootstrap resamples. We detected a positive correlation between M_* and sub-Neptune size at 8σ significance $\alpha = 0.25 \pm 0.03$. An identical analysis for the super-

Earths ($R_p = 1.0 - 1.7 R_\oplus$) revealed no correlation $\alpha = 0.02 \pm 0.03$.

Before interpreting these trends with M_* , we must consider confounding effects of other stellar properties that are correlated with M_* . (We consider effects from M_* -dependent survey completeness in §7.2). Stellar metallicity and age are both concerns because they are astrophysically correlated with mass. This M_* -[Fe/H]-age correlation is tied to the chemical enrichment of the galaxy and stellar lifespans. A star's main-sequence lifetime is a strong function of its mass;




Figure 9. Same as Figure 8 except the x-axis is incident bolometric flux.

$t_{\text{ms}} \sim 10 \text{ Gyr} (M_{\star}/M_{\odot})^{-2.5}$. Massive stars are younger on average and formed when the galaxy was more enriched in metals, as shown in Figures 5a and 5c. Figure 5b highlights the variable uncertainties of our age measurements which range from ≈ 0.1 dex for stars that have evolved off the main sequence, to ≈ 0.5 dex for stars with $T_{\text{eff}} \lesssim 5500$ K, i.e. indistinguishable from the zero-age main-sequence.

We quantified possible metallicity and age effects by including them in our regression:

$$R_p = R_{p,0} \left(\frac{M_{\star}}{M_{\odot}} \right)^{\alpha} \left(\frac{N_{\text{Fe}}}{N_{\text{Fe},\odot}} \right)^{\beta} \left(\frac{\text{age}}{5 \text{ Gyr}} \right)^{\gamma}. \quad (7)$$

When we allowed α and β to vary and fixed γ to 0, we found $\beta = -0.01 \pm 0.02$ and -0.00 ± 0.02 for sub-Neptunes and super-Earths, i.e. no significant R_p -[Fe/H] correlation. Finally, we considered the effects of age by allowing γ to vary as well. We restricted our analysis to stars with $T_{\text{eff}} > 5500$ K where $\text{age}_{\star,\text{iso}}$ is not prior-dominated. We found $\gamma = -0.01 \pm 0.03$ and 0.02 ± 0.02 for the sub-Neptunes and super-Earths, i.e. no R_p -age correlation.

Based on these fits, we concluded that stellar mass rather than metallicity or age is the property most closely linked to sub-Neptune size variation. We acknowledge two additional caveats to the above analysis

Table 3. Power-law fits to the Radius Gap

Fit	M_\star	m	$R_{p,0}$	Dist
$R_p = R_{p,0} \left(\frac{P}{10 \text{ days}} \right)^m$	0.5–1.4	$-0.11^{+0.02}_{-0.02}$	$1.84^{+0.03}_{-0.03}$	D
	0.5–0.7	$-0.12^{+0.06}_{-0.04}$	$1.78^{+0.05}_{-0.06}$	D
	0.7–1.0	$-0.13^{+0.07}_{-0.06}$	$1.74^{+0.09}_{-0.08}$	D
	1.0–1.4	$-0.06^{+0.02}_{-0.02}$	$1.93^{+0.04}_{-0.04}$	D
	0.5–0.7	$-0.06^{+0.04}_{-0.05}$	$1.71^{+0.05}_{-0.04}$	O
	0.7–1.0	$-0.10^{+0.07}_{-0.05}$	$1.70^{+0.09}_{-0.10}$	O
	1.0–1.4	$-0.05^{+0.03}_{-0.03}$	$1.92^{+0.06}_{-0.05}$	O
$R_p = R_{p,0} \left(\frac{S_{\text{inc}}}{100 S_\oplus} \right)^m$	0.5–1.4	$0.06^{+0.01}_{-0.01}$	$1.86^{+0.04}_{-0.03}$	D
	0.5–0.7	$0.07^{+0.02}_{-0.04}$	$2.00^{+0.11}_{-0.19}$	D
	0.7–1.0	$0.10^{+0.04}_{-0.04}$	$1.86^{+0.07}_{-0.07}$	D
	1.0–1.4	$0.05^{+0.01}_{-0.01}$	$1.85^{+0.04}_{-0.04}$	D
	0.5–0.7	$0.06^{+0.02}_{-0.03}$	$1.93^{+0.11}_{-0.13}$	O
	0.7–1.0	$0.08^{+0.04}_{-0.04}$	$1.81^{+0.08}_{-0.06}$	O
	1.0–1.4	$0.05^{+0.01}_{-0.01}$	$1.84^{+0.04}_{-0.04}$	O
$R_p = R_{p,0} \left(\frac{M_\star}{1 M_\odot} \right)^m$	0.5–1.4	$0.18^{+0.08}_{-0.07}$	$1.86^{+0.03}_{-0.03}$	D
$R_p = m[\text{Fe}/\text{H}] + R_{p,0}$	0.5–1.4	$0.01^{+0.05}_{-0.06}$	$1.82^{+0.05}_{-0.03}$	D

NOTE—We fit the Radius Gap in the *detected* planet population in §6.1 as function of orbital period, incident stellar flux, stellar mass, and stellar metallicity (see Figure 7). We repeated the orbital period and incident flux fits for narrower ranges of stellar mass (see Figures 8 and 9). We fit the Radius Gap in the de-biased planet *occurrence* distribution in §7.1 (see Figures 8 and 9). We elaborate on the following columns: ‘Fit’—the functional form of the fit; ‘ M_\star ’—range of stellar masses; ‘Dist’—indicates whether the fit was of the distribution of planet detections ‘D’ or occurrence ‘O’.

which we address in later sections. First, we fit the population of detected planets which includes survey completeness effects that are addressed in §7.2. Second, orbital period is another possible confounding variable if the *shape* of the period distribution varies with stellar mass. Fortunately, as we show in §7.3 the period distribution is nearly independent of stellar mass up to a normalization constant.

7. DE-BIASED DISTRIBUTION OF PLANETS

We have characterized the location and slope of the Radius Gap and its dependence on stellar mass. However, one must exercise caution when connecting these trends to planet formation models because the distributions of planets have been filtered through *Kepler*’s selection function. In this section, we remove selection effects to characterize variation in the planet population with stellar mass.

In §7.1, we reexamine the slope of the Radius Gap as a function of period and incident flux (first treated in §6.2) and find consistent results after accounting for selection effects. In §7.2, we measure the sizes of super-Earths and

sub-Neptunes (as in §6.2) and observe the same qualitative trends after removing selection effects. In §7.3, we characterize how the period and incident flux distributions of super-Earths and sub-Neptunes vary with stellar mass.

7.1. Dependence of the Radius Gap on stellar mass

We return to the P - R_p distribution of planets shown in Figure 8. We wish to remove *Kepler*’s observational biases to measure the planet occurrence rate density (ORD), $d^2f/d \log P d \log R_p$. One may integrate the ORD over a specified domain in the P - R_p plane to derive the number of planets per star residing in that domain.

For every detected planet in a sample of n_\star stars, a large number are missed due to non-transiting geometries or insufficient photometric S/N. We account for both effects here using the inverse detection efficiency method (IDEM, see [Fulton & Petigura 2018](#)). Each detected planet i with (P, R_p) represents $w_i = 1/\langle p_{\text{tr}} p_{\text{det}} \rangle$ total planets where $\langle p_{\text{tr}} p_{\text{det}} \rangle$ is the product of transit and detection probability averaged over the parent stellar population. We computed $\langle p_{\text{tr}} p_{\text{det}} \rangle$ over a grid of (P, R_p) . The probability that a randomly inclined planet with semi-major axis a will transit with $b < 0.8$ is $p_{\text{tr}} = 0.8R_\star/a$, assuming circular orbits. We computed p_{tr} for each star using the B20 R_\star and deriving a from P and the B20 M_\star using Kepler’s Third Law.

We characterized the recovery rate p_{det} following the procedure described in [Christiansen et al. \(2020\)](#). These authors injected a suite of 146,295 synthetic transits into the raw *Kepler* pixel-level data and searched for transits using the same pipeline used to produce the T18 catalog. The output was a list of injections that were either successfully recovered or not. This list may be used to determine the average recovery rate for any *Kepler* sample of interest. We selected the subset of injections with $P = 1 - 300$ days into our parent stellar population described in §2 and measured the recovery rate as a function of expected MES, a close relative of transit S/N. We modeled the completeness curve as a Γ cumulative distribution function

$$p_{\text{det}}(\text{MES}) = \frac{c}{b^a \Gamma(a)} \int_0^{\text{MES}} t^{a-1} e^{-t/b} dt. \quad (8)$$

The best-fit model had the following coefficients $(a, b, c) = (26.1, 0.320, 0.941)$ and is shown in Figure 10. We set the recovery rate to zero when $\text{MES} < 10$ to incorporate our filter described in §2. We evaluated MES for each star based on the putative planet’s transit depth, transit duration, and the star’s tabulated photometric noise (Combined Differential Photometric Precision; [Jenkins et al. 2010](#)).

With our calculated weights w , we may now compute the total number of planets within a given region of the P - R_p plane by summing the weights of the planets within that region, $\sum_i w_i$, or compute the occurrence rate (number of planets per star) via $(1/n_\star) \sum_i w_i$.

The IDEM, while mathematically straightforward, has some drawbacks particularly when p_{det} is small. In brief, Hsu et al. (2018) showed that the IDEM is a biased estimator at low p_{det} because the weights w_i are computed assuming no uncertainties in R_p . Hsu et al. (2018) proposed a less biased estimator using Approximate Bayesian Computation (ABC). We avoided these complications by restricting our analysis to regions where $p_{\text{det}} > 25\%$, where differences between the IDEM and ABC amount to $< 1\sigma$ (Fulton & Petigura 2018).

We measured the ORD in the three bins of stellar mass introduced in §6.2 via

$$\frac{d^2 f}{d \log P d \log R_p} = \frac{1}{n_\star} \sum_i w_i k(P - P_i, R_p - R_{p,i}). \quad (9)$$

where k is a 2D gaussian with a bandwidth of $\log(2) = 0.3$ dex in P and $\log(1.05) = 0.02$ dex in R_p . The results are shown in the right-hand panels of Figure 8. The value at each (P, R_p) is the number of planets per star in a 1 dex \times 1 dex interval centered at (P, R_p) . It is convenient to scale this value to smaller intervals because the ORD varies significantly over 1 dex in P and R_p . For example, to read off the number of planets per star within a box having the dimensions of the kernel FWHM, multiply the ORD by $2.355 \times 0.3 \times 2.355 \times 0.02 = 0.033$.

We do not plot occurrence in regions of low-planet detectability, defined to be where $n_\star \langle p_{\text{tr}} p_{\text{det}} \rangle < 50$. The ORD ranges from ≈ 0 –4 which corresponds to ≈ 0 –0.1 planets per star within the kernel FWHM. For (P, R_p) values where $n_\star \langle p_{\text{tr}} p_{\text{det}} \rangle = 50$, there are effectively 50 stars where such a planet could be detected, and we would expect 0–5 detected planets within the kernel FWHM. At this point Poisson fluctuations become unmanageable.

We fit a power-law to the Radius Gap for each mass bin. Our methodology, power-law parameterization, fitting domain, and uncertainty analysis are identical to that presented in §6, except that we fit ORD rather than density of detected planets. The credible range of values is given in Table 3. The measured slope and intercepts are consistent at 1σ to those measured using planet detections only.

7.2. Dependence of planet size on stellar mass

In §6.2, we noted a positive R_p - M_\star correlation for sub-Neptunes that was not mirrored by super-Earths. However, we must consider the possibility that such cor-

relations (or lack thereof) could stem from the *Kepler* selection function. To do so, we first divided the CXM planet hosts and parent stellar population into our three M_\star bins from §6.2. In order to treat hosts and non-hosts equally, we used the B20 M_\star which are available for all stars.

Returning to our binned planet population, we computed the occurrence-weighted mean planet size for sub-Neptunes having $P < 100$ days and display them in Figure 11. Fitting the power-law given by Equation 6 to the three binned mean size measurements shown in Figure 11 yields an index $\alpha = 0.23 \pm 0.04$ and an intercept of $R_{p,0} = 2.47 \pm 0.03 R_\oplus$ (uncertainties from bootstrap resampling). These values are consistent with our fits to the population of detected planets (§6.2). For the sub-Neptunes, the weights w are dominated by geometrical corrections which are independent of R_p . Planets with larger P have larger w and are more strongly weighted than in §6.2. However, given the lack of a significant tilt in the P - R_p distribution, the average value is similar.

Corrections stemming from low detectability are larger for the super-Earths. While there is no obvious tilt of the super-Earth population in Figure 7c, completeness effects could conspire to produce a flat dependence when there is actually a positive slope in the underlying population. We computed the occurrence-weighted mean planet size for super-Earths with $P < 30$ days and found $\alpha = -0.01 \pm 0.03$ and $R_{p,0} = 1.34 \pm 0.01$, i.e. no significant correlation. We interpret the different R_p - M_\star correlations super-Earths and sub-Neptunes and compare our results to previous literature measurements in §8.

7.3. Orbital period and incident flux distributions

One key feature of the exoplanet population is the paucity of sub-Neptunes with $P \lesssim 10$ days, i.e. the “Sub-Neptune Desert.” The edge of this desert is significant in both the core and XUV-powered models. It represents a boundary, inside of which core-dominated planets (i.e. $M_c \gg M_{\text{env}}$) cannot retain H/He envelopes and are completely stripped.

Here, we characterize the edge of the Sub-Neptune Desert in terms of period and flux and note how it changes with stellar mass. One can see qualitatively from Figures 8 and 9 that the edge of the Sub-Neptune Desert occurs at a similar P for all mass bins but varies in S_{inc} by $\sim 10\times$. We first considered the edge in period space in our three bins of stellar mass introduced in §6.2. Quantitatively measuring this boundary in P by visually inspecting Figures 8 is suboptimal for a number of reasons. First, the contours do not convey uncertainties. Second, the Gaussian KDE used to generate the 2D dis-




Figure 10. Points show the recovery rate of the *Kepler* planet detection pipeline of simulated planets injected by Christiansen et al. (2020) as a function of multiple event statistic (MES). The curve shows the parametric model we used to account for pipeline incompleteness in our occurrence calculations described in §7




Figure 11. Mean size of super-Earths with $P < 30$ days and sub-Neptunes with $P < 100$ days. Over $M_* = 0.5\text{--}1.4 M_\odot$, the average sub-Neptune grows from $2.1 R_\oplus$ to $2.6 R_\oplus$ with a slope of $\alpha = d \log R_p / d \log M_* = 0.23 \pm 0.04$. The mean super-Earth has a nearly constant size of $1.35 R_\oplus$ with $\alpha = -0.01 \pm 0.03$. Wu (2019) reported $\alpha = 0.25$ for both populations and is shown by the dashed lines. Our sub-Neptune results are consistent with this index, but our super-Earth results are inconsistent at 8σ .

tributions smears out sharp features in the P distributions. Third, the Sub-Neptune Desert is the paucity of short-period Sub-Neptunes *relative* to long-period Sub-Neptunes, but the contours show *absolute* occurrence which also varies with stellar mass. These shortcomings motivated a different method, which we describe below.

Within each M_* bin, modeled the ORD as

$$\frac{df}{dPdR_p} = f_* \lambda(P, R_p; \boldsymbol{\theta}) \quad (10)$$

over a rectangular domain of P and R_p . Here, f_* is the mean number of planets per star within the P - R_p domain. In principle, λ may be any function that integrates to unity over the same domain. We consider parametric functions described by a vector of parameters $\boldsymbol{\theta}$ that we will constrain through Bayesian inference. In short, the free parameters f_* and $\boldsymbol{\theta}$ control the normalization and shape of the ORD, respectively.

Following previous works, we modeled the detected population of planets as a realization of an inhomogeneous Poisson point process (see, e.g., Rogers & Owen 2021, and references therein). The log-likelihood of observing the full sample is given by

$$\ln \mathcal{L} = -\Lambda + \sum_i \ln [f_* \lambda(P_i, R_{p,i})] \quad (11)$$

where

$$\Lambda = n_* f_* \iint \eta(P, R_p) \lambda(P, R_p) dP dR_p. \quad (12)$$

is the expected number of detected planets given f_* and $\boldsymbol{\theta}$. Here, $\eta = \langle p_{\text{tr}} p_{\text{det}} \rangle$.

For simplicity, we assumed that the period and radius distributions are independent, i.e. $\lambda(P, R_p) \propto \lambda(P) \lambda(R_p)$. We also approximated the R_p distribution log-uniform within the R_p boundaries, i.e. $\lambda(R_p) \propto 1/R_p$. We precomputed

$$\eta(P) = \frac{1}{\Delta \ln R_p} \int \eta(P, R_p) d \ln R_p \quad (13)$$

to reduce Λ to a 1D integral

$$\Lambda = n_* f_* \int \eta(P) \lambda(P) dP. \quad (14)$$

Following Rogers & Owen (2021) we described the period distribution with a smooth broken power-law.

$$\lambda(P) = \frac{C}{(P/P_0)^{-k_1} + (P/P_0)^{-k_2}}, \quad (15)$$

where C is the constant of proportionality that enforces normalization.⁹ In this parameterized description, P_0 is the knee of the distribution; for $P \ll P_0$ the distribution approaches a power-law with an index of k_1 ; for $P \gg P_0$ the index is k_2 .

We first fit the sub-Neptune distribution for our three bins of stellar mass over $P = 1\text{--}300$ days. Our goal was to characterize the knee and slope of the occurrence period distribution. For consistency between the subsamples, we fixed $k_2 = -1$, i.e. log-uniform occurrence for $P \gg P_0$. We maximized the likelihood (Equation 11) using the L-BFGS-B algorithm and explored the credible range of parameters using the MCMC sampler of [Goodman & Weare \(2010\)](#). We imposed log-uniform priors on f_\star and P_0 and a linear prior on k_1 . We sampled the posterior with 8 walkers for 5,000 steps each and discarded the first 1,000 as burn-in. The chains for all parameters were at least 50 times longer than the integrated autocorrelation time, indicating convergence. Table 4 lists the 16, 50, 84 percentiles of our posterior samples. Figure 12 shows the posterior probability density of f_\star and P_0 for different M_\star bins.

Figure 13 shows the best-fit and credible range of models for our three M_\star bins. They are similar except for an overall shift in the normalization. For the low-, medium-, and high-mass bins, the breakpoint P_0 is at $7.0^{+1.4}_{-1.1}$, $9.7^{+2.0}_{-1.5}$, $8.6^{+1.6}_{-1.2}$ days, respectively, consistent to 2σ ; the power-law indices k_1 agree to 1σ and are $1.7^{+0.5}_{-0.4}$, $1.5^{+0.4}_{-0.3}$, $2.2^{+0.6}_{-0.5}$; we observed a steady decrease in the absolute number of planets per star from 0.5 to $1.4 M_\odot$ of $1.37^{+0.19}_{-0.18}$, to $0.79^{+0.08}_{-0.08}$, to $0.41^{+0.04}_{-0.04}$.

The similarity of P_0 for the different mass bins is noteworthy because it indicates a key boundary in the planet formation process that is stationary over a wide range of stellar mass. Naturally, stationary P_0 implies a variable $S_{\text{inc},0}$ given the relationship between M_\star and bolometric luminosity. To quantify this boundary in flux space, we again characterized the distribution as a smooth broken power-law,

$$\lambda(S_{\text{inc}}) = \frac{C}{(S_{\text{inc}}/S_{\text{inc},0})^{-k_1} + (S_{\text{inc}}/S_{\text{inc},0})^{-k_2}}, \quad (16)$$

but fixed k_1 to -1 . Over the three M_\star bins, we observed factor of 9 increase in the breakpoint $S_{\text{inc},0}$ from 27^{+6}_{-6} , to 70^{+16}_{-14} , to $244^{+64}_{-55} S_\oplus$.

⁹ The normalization constant C is

$$C = \left[\frac{P_0 u^{k_1+1}}{k_1+1} {}_2F_1 \left(1, \frac{k_1+1}{k_1-k_2}; \frac{k_1+1}{k_1-k_2} + 1; -u^{k_1-k_2} \right) \right]_{u=\frac{P_1}{P_0}}^{u=\frac{P_2}{P_0}}$$

where ${}_2F_1$ is the hypergeometric function.

We also characterized the period distribution of super-Earths. Our modeling was identical to that of the sub-Neptunes except we only included planets out to 30 days due to low completeness at longer orbital periods. Here, the breakpoint P_0 increases with M_\star from $2.6^{+0.8}_{-0.4}$, to $3.3^{+0.8}_{-0.5}$, to $5.2^{+1.3}_{-0.9}$ days. The breakpoints in flux $S_{\text{inc},0}$ occur at 89^{+35}_{-31} , 265^{+84}_{-76} , and $615^{+159}_{-152} S_\oplus$. Figure 14 contrasts the super-Earth and sub-Neptune breakpoints. The super-Earth breakpoint is shifted to smaller orbital periods and higher incident fluxes for all stellar mass bins. We interpret these shifts in §8.5.

8. DISCUSSION

8.1. Positive M_\star - R_p correlation suggests massive stars make massive planet cores

One of the striking features of Figure 7c is the divergence of the super-Earth and sub-Neptune populations with stellar mass: the sub-Neptunes grow larger while the super-Earths do not. When we modeled the size dependence of each population as a power-law $R_p \propto M_\star^\alpha$, we found $\alpha = 0.25 \pm 0.03$ and $\alpha = 0.02 \pm 0.03$, respectively. The slope of the Radius Gap in M_\star - R_p space (displayed in Figure 7) lies between those two values $m = 0.18^{+0.08}_{-0.07}$.

A positive sub-Neptune R_p - M_\star correlation has been noted in several previous works. [Fulton & Petigura \(2018\)](#) observed such a trend over a narrower stellar mass range of $\approx 0.8\text{--}1.3 M_\odot$. Later, [Wu \(2019\)](#) stitched together the [Fulton & Petigura \(2018\)](#) sample with a compilation of ~ 100 *Kepler* and *K2* planets with hosts of $M_\star \lesssim 0.7 M_\odot$ from [Newton et al. \(2015\)](#) and [Dressing et al. \(2017\)](#) and reported $\alpha = 0.23\text{--}0.35$, consistent with what we found.

In contrast to the sub-Neptunes, we observe a flat super-Earth R_p - M_\star dependence, $\alpha = -0.01 \pm 0.03$. This result differs from that of [Wu \(2019\)](#) who reported $\alpha = 0.23\text{--}0.35$. As a point of comparison, Figure 11 shows our occurrence-weighted mean super-Earth size with an $\alpha = 0.25$ dependence overplotted; such a strong relationship is ruled out at 8σ significance. The differences between the two works may stem from the fact that [Wu \(2019\)](#) used an add-mixture of planet samples with heterogeneously derived properties.

[Berger et al. \(2020a\)](#) offers a second point of comparison. They derived properties of 2956 KOIs spanning $M_\star \approx 0.6\text{--}1.4 M_\odot$ that host 3898 confirmed/candidate planets. They measured the Radius Gap slope in M_\star - R_p space and found $m = 0.26^{+0.21}_{-0.16}$, which is consistent with our measured value, albeit with larger uncertainties.

What causes the super-Earth size to be independent of stellar mass while the sub-Neptune size grows with stellar mass? In our discussion below, we assume plan-

Table 4. Fits to the planet period and flux distributions

Size	M_\star	Period			Flux		
		f_\star	P_0	k_1	f_\star	$S_{\text{inc},0}$	k_2
SE	0.5–0.7	$0.42^{+0.07}_{-0.06}$	$2.6^{+0.8}_{-0.4}$	$2.4^{+1.1}_{-1.0}$	$0.33^{+0.06}_{-0.05}$	89^{+35}_{-31}	$-3.1^{+0.4}_{-0.5}$
	0.7–1.0	$0.22^{+0.03}_{-0.02}$	$3.3^{+0.8}_{-0.5}$	$1.5^{+0.6}_{-0.5}$	$0.26^{+0.03}_{-0.03}$	265^{+84}_{-76}	$-2.6^{+0.3}_{-0.3}$
	1.0–1.4	$0.18^{+0.02}_{-0.02}$	$5.2^{+1.3}_{-0.9}$	$1.9^{+0.7}_{-0.5}$	$0.28^{+0.04}_{-0.03}$	615^{+159}_{-152}	$-3.0^{+0.4}_{-0.5}$
SN	0.5–0.7	$1.37^{+0.19}_{-0.18}$	$7.0^{+1.4}_{-1.1}$	$1.7^{+0.5}_{-0.4}$	$1.13^{+0.16}_{-0.14}$	27^{+6}_{-6}	$-3.2^{+0.3}_{-0.3}$
	0.7–1.0	$0.79^{+0.08}_{-0.08}$	$9.7^{+2.0}_{-1.5}$	$1.5^{+0.4}_{-0.3}$	$0.76^{+0.08}_{-0.08}$	70^{+16}_{-14}	$-3.1^{+0.3}_{-0.3}$
	1.0–1.4	$0.41^{+0.04}_{-0.04}$	$8.6^{+1.6}_{-1.2}$	$2.2^{+0.6}_{-0.5}$	$0.52^{+0.06}_{-0.05}$	244^{+64}_{-55}	$-3.0^{+0.3}_{-0.4}$

NOTE—In §7.3, we modeled both the period and flux distribution of super-Earths (SE) and sub-Neptunes (SN) for different bins of stellar mass. We used a smooth broken power-law to model the period distribution $df/dP \propto ((P/P_0)^{-k_1} + (P/P_0)^{-k_2})^{-1}$. Here, f_\star is the mean number of planets per star over the following period ranges: $P = 1\text{--}30$ days (SE); $P = 1\text{--}300$ days (SN). The power-law indices below and above the breakpoint P_0 are k_1 and k_2 , respectively. In our modeling, k_2 is fixed to -1 , i.e. $df/d\log P \approx \text{constant}$ for $P \gg P_0$. Similarly, we also used a smooth broken power-law to model the flux distribution. Here, k_1 was fixed to -1 .

ets between $1\text{--}4 R_\oplus$ are Earth-composition bodies that may also include a H/He envelope. This assumption is supported by the growing body of mass measurements of planets in this size range (see, e.g., [Weiss & Marcy 2014](#); [Sinukoff 2018](#)). Most planets smaller than $1.5 R_\oplus$ have densities consistent with Earth composition bodies; most planets larger than $2.0 R_\oplus$ are too low-density to be rock/iron bodies and are consistent with having H/He envelopes of $f_{\text{env}} \gtrsim 1\%$ by mass.

With this two-component model, both the XUV- and core-powered models produce the Radius Gap. In the 1D, energy-limited, XUV-powered models of [Owen & Wu \(2017\)](#) the mass loss-timescale $\tau_{\text{xuv}} = M/\dot{M}$ is a strong function of f_{env} and the local XUV flux. This timescale is maximized when the size of the core and thickness of the envelope are approximately equal, i.e. $R_p \approx 2 R_c$. This occurs at $f_{\text{env},2} \approx 3\%$. Planets with less envelope experience accelerating mass-loss; those with more experience decelerating mass-loss. Planets are herded toward $f_{\text{env}} = f_{\text{env},2}$ or $f_{\text{env}} = 0$ (bare cores).

In the XUV-model, the fate of a planet’s envelope depends the time-integrated XUV exposure $\mathcal{F}_{\text{xuv}} = \int F_{\text{xuv}} dt$. This quantity is strongly correlated with orbital period and weakly correlated with stellar mass ([Owen & Wu 2017](#); [Lopez & Rice 2018](#)). At fixed orbital period, planets around low-mass stars receive less bolometric flux F_{bol} , but $F_{\text{xuv}}/F_{\text{bol}}$ is higher and stars stay XUV-active longer. [McDonald et al. \(2019\)](#) found $\mathcal{F}_{\text{xuv}} \propto F_{\text{bol}} M_\star^{-3}$. Applying following relationships, $P^2 \propto a^3/M_\star$ (Kepler’s Third Law), $F_{\text{bol}} \propto L_{\star,\text{bol}}/a^2$, $L_{\star,\text{bol}} \propto M_\star^4$ ([Kippenhahn & Weigert 1990](#), for FGK and early M stars), we find $\mathcal{F}_{\text{xuv}} \propto M_\star^{0.33} P^{-1.33}$.

If the XUV model is correct and the initial planet population is independent of M_\star , we expect only minor sub-Neptune size variability with M_\star . We can get a rough estimate the expected $R_p\text{--}M_\star$ dependence at fixed P by considering the amount of energy needed to unbind a $f_{\text{env}} \approx 3\%$ envelope

$$E_{\text{unb}} \sim GM_c M_{\text{env}}/R_c \propto M_c^{1.75}. \quad (17)$$

We have approximated the rocky mass-radius relationship as $R_c \propto M_c^{0.25}$. The XUV energy received by the planet is approximately

$$E_{\text{rec}} \sim F_{\text{xuv}} \pi R_p^2 \propto F_{\text{xuv}} M_c^{0.5} \propto M_\star^{0.33} M_c^{0.5}. \quad (18)$$

Equating the two energies yields $M_c \propto M_\star^{0.26}$ or $R_c \propto R_p \propto M_\star^{0.06}$, i.e. $\alpha \approx 0.06$.

In the core-powered model of [Gupta & Schlichting \(2020\)](#), planets with $f_{\text{env}} < 5\%$ have enough thermal energy in their cores to unbind their envelopes and become bare rocky planets. Thus, this process also works to clear the $P\text{--}R_p$ plane of planets with f_{env} of a few percent or less. The fate of a planet’s envelope depends on the mass-flux at the Bondi radius R_B , and planets with smaller R_B lose mass faster. Since $R_B \propto M_c/T_{\text{eq}}$, planets more readily lose their envelopes when T_{eq} (or equivalently S_{inc}) is large, or when M_c is small. If the core-powered model is correct and the initial planet population is independent of M_\star , we expect significant changes with M_\star because S_{inc} is a strong function of M_\star . [Gupta & Schlichting \(2020\)](#) predicted $\alpha \approx 0.33$.

It is tempting to attribute the larger sub-Neptunes around massive stars as a direct result of higher mass cores. One may, for example, adopt a monotonically increasing mass-radius relationship (e.g. [Weiss & Marcy](#)




Figure 12. Panel (a): posterior probability density of the mean number of sub-Neptunes per star f_* with $P = 1\text{--}300$ days for three stellar mass bins. Dashed lines show the 25%, 50%, and 75% quartiles. Panel (b) shows the period breakpoint P_0 . Panel (c) shows the flux breakpoint $S_{\text{inc},0}$. Panels (d–f) same as (a–c) but for super-Earths. The mean occurrence in panel (d) corresponds to $P = 1\text{--}30$ days. For both classes of planets, there is a decline in total occurrence with increasing M_* . For the sub-Neptunes, P_0 ranges from $\approx 7.0\text{--}9.6$ days, consistent to 40%; for the super-Earths P_0 is $\approx 2.7\text{--}5.1$ days, consistent to a factor of two. For the sub-Neptunes, the breakpoint $S_{\text{inc},0}$ increases by a factor of 9 from $27_{-6}^{+6} S_{\oplus}$ to $244_{-55}^{+64} S_{\oplus}$ between the low- and high-mass bin. For the super-Earths, the breakpoint also increases with stellar mass from $89_{-31}^{+35} S_{\oplus}$ to $615_{-152}^{+159} S_{\oplus}$, a factor of 7 increase.

2014), invert it, and infer higher mass sub-Neptunes around more massive stars. However, there is significant astrophysical scatter about the mean mass-radius relationship (Wolfgang et al. 2016). Planets of $2\text{--}3 R_{\oplus}$ have a nearly uniform distribution in mass from $5\text{--}20 M_{\oplus}$. In short: radius is not a good predictor of mass. It could also be true that faster or more efficient gas accretion occurs in the protoplanetary disks of massive stars because of differences in gas surface density, ionization structure, or other properties.

With these caveats in mind, we *can* qualitatively explain the increase in sub-Neptune sizes and constant super-Earth sizes in both XUV- and core-powered models as a consequence of more massive cores around more

massive stars. We first consider photoevaporation in a toy model shown in Figure 15. Here, the cores have a log-uniform mass distribution up to a maximum cut-off mass of $M_{c,\text{max}}$. The maximum core mass is proportional to stellar mass, $M_{c,\text{max}} \propto M_*$. For simplicity, we show two populations of host stars: (1) where $M_{*,1} \approx 0.6 M_{\odot}$ and $M_{c,\text{max},1} \approx 10 M_{\oplus}$, and (2) where $M_{*,2} \approx 1.2 M_{\odot}$ where $M_{c,\text{max},2} \approx 20 M_{\oplus}$. At the time of disk dispersal ($t \sim 10$ Myr), the cores have accreted H/He envelopes with a broad range of $f_{\text{env}} \approx 0.3\text{--}30\%$ and have a broad size distribution $R_p \approx 2\text{--}10 R_{\oplus}$.

After the period of high XUV activity ($t \gtrsim 100$ Myr), cores up to a critical mass $M_{c,\text{crit}}$ are stripped bare and the largest super-Earth is given by $R_{\text{se,max}}/R_{\oplus} \approx$




Figure 13. Panel (a): period distribution of sub-Neptunes for three bins of stellar mass. The points show the number of planets per star in period bins that are 0.25 dex wide. We modeled the occurrence rate density df/dP as a smooth broken power-law which we fit to the unbinned planet population over $P = 1\text{--}300$ days (see §7.3). The bands show the credible range of $df/d \log P \times 0.25$ dex. The height of each curve at a particular P is the number of planets per star within a 0.25 dex period interval centered at P . Panel (b): same as (a) except for super-Earths where the fits extend to 30 days. Panel (c–d): same as (a–b) except showing incident flux distribution. The vertical axis has the following interpretation: number of planets per star within a 0.5 dex flux interval.

$(M_{c,\text{crit}}/M_{\oplus})^{0.25}$. Cores above $M_{c,\text{crit}}$ retain their envelopes and mass-loss concentrates envelope fractions to $f_{\text{env}} \approx 3\%$. The smallest sub-Neptunes have cores that barely retained their envelopes and obey $R_{\text{sn},\text{min}}/R_{\oplus} \approx 2(M_{c,\text{crit}}/M_{\oplus})^{0.25}$; the largest sub-Neptunes have an $f_{\text{env}} \approx 3\%$ envelope atop a $M_{c,\text{max}}$ core and obey $R_{\text{sn},\text{max}}/R_{\oplus} \approx 2(M_{c,\text{max}}/M_{\oplus})^{0.25}$.

This toy model reproduces some, but not all, of the observed trends. $M_{c,\text{max}} \propto M_{\star}$ so $R_{\text{sn},\text{max}} \propto M_{\star}^{0.25}$, which is consistent with the observed trend. Moreover, approximating $\mathcal{F}_{\text{xuv}} \propto M^{0.33}$ as constant over our mass range predicts constant $R_{\text{se},\text{max}}$ for different M_{\star} .

The fact that $R_{\text{se},\text{max}} \approx 1.7 R_{\oplus}$ implies $M_{c,\text{crit}} \approx 8 M_{\oplus}$ and $R_{\text{sn},\text{min}} \approx 3.4 R_{\oplus}$. Given our assumed $M_{c,\text{max}}$,

we find $R_{\text{sn},\text{max}} \approx 3.6 R_{\oplus}$ and $4.2 R_{\oplus}$. While these have the appropriate dependence on M_{\star} , they exceed the upper envelope of the observed sub-Neptunes. Note, however, that these sizes are at $t \sim 100$ Myr and observed planets are $t \approx 1\text{--}10$ Gyr and will contract as they radiate away their heat of formation. Gupta & Schlichting (2020) found that the radius of the radiative-convective boundary (RCB) shrinks as $d \log R_p / d \log t \approx -0.1$. So over $\log t = 8$ to 9.7 we expect planets to shrink by 0.17 dex or 33%. The largest sub-Neptunes would shrink to $2.2 R_{\oplus}$ and $2.8 R_{\oplus}$ for the $M_{\star,1} \approx 0.6 M_{\odot}$ and $M_{\star,2} \approx 1.2 M_{\odot}$ populations, respectively, in rough agreement with observations. However, we note that we observed no age dependence in the mean sub-Neptune




Figure 14. Panels (a–c): period distribution of super-Earths (green) and sub-Neptunes (blue) for different bins of stellar mass. Points and curves are the same as shown in Figure 13 but grouped according to stellar mass as opposed to planet size. The vertical bands show the range of credible period breakpoints P_0 . For all stellar mass bins, P_0 is smaller for super-Earths than for sub-Neptunes. Panels (d–f): same as (a–c), but x-axis is incident stellar flux. The flux breakpoint $S_{\text{inc},0}$ increases with stellar mass, but the relative offset between the two populations is nearly constant.

size over $\log t = 9\text{--}10$, and comment on this discrepancy later.

Given the similarities between the core- and XUV-powered models, this toy model may be easily adapted to the core-powered model. The key difference is that the time of rapid mass loss is $t \lesssim 1$ Gyr as opposed to $t \lesssim 100$ Myr.

To summarize, the $R_p\text{--}M_*$ correlation observed for super-Earths and sub-Neptunes is broadly consistent with maximum core-mass that increases with M_* . The $R_p \propto M_*^{0.25}$ trend suggests a $M_c \propto M_*$ relationship. A linear relationship between core mass and host star mass has also been suggested by Wu (2019) and Berger et al. (2020a).

Why should core mass increase with host star mass? One explanation could be that the core formation process is dust-limited and that the available dust mass M_{dust} grows with stellar mass. Pascucci et al. (2016) derived M_{dust} in protoplanetary disks from ALMA observations at ~ 1 mm and found a super-linear relationship with stellar mass $M_{\text{dust}} \propto (M_*)^{1.3\text{--}1.9}$, with a large

0.8 dex dispersion about this relationship. It is not clear, however, if core-formation is dust-limited due to the difficulties inferring the total inventories of solids from mm emission. Wu (2019) offered a different interpretation, positing that planet cores form at a “thermal mass” M_{th} , the mass at which a planet’s Hill sphere equals the disk scale height. $M_{\text{th}} \propto M_*^{11/8}$, which is close to the inferred $M_c \propto M_*$ dependence.

8.2. Absence of size-metallicity trend suggests weak connection between envelope opacity and stellar metallicity

Models of sub-Neptunes depend on the envelope opacity κ , which sets the cooling rate, location of the RCB, and thus the planet’s size. Higher opacity envelopes cool and contract more slowly.

In their core-powered models, Gupta & Schlichting (2020) assumed $\kappa \propto Z$, where Z is the bulk stellar metallicity. To isolate the effects of metallicity, they constructed a population of planets with M_* fixed at $1.0 M_\odot$, $[\text{Fe}/\text{H}]$ ranging from -0.5 to $+0.5$, and evolved them for 3 Gyr. The final population is shown in their




Figure 15. A toy model to explore how a correlation between host star mass M_\star and planet core mass M_c impacts super-Earth and sub-Neptune sizes when photoevaporation drives mass-loss. Panels (a–c) show the distributions of M_c , envelope fraction f_{env} , and planet size R_p at the time of disk dispersal ($t \sim 10$ Myr). Here, low-mass stars have a lower maximum core mass than high-mass stars: for $M_\star \approx 0.6 M_\odot$, $M_{c,\text{max},1} \approx 10 M_\oplus$; for $M_\star \approx 1.2 M_\odot$, $M_{c,\text{max},2} \approx 20 M_\oplus$. Planets have a broad distribution of envelope fractions $f_{\text{env}} \approx 0.3\text{--}30\%$ and a broad distribution of sizes. Panels (d–f) show the same properties after the period of rapid photoevaporation ($t \sim 100$ Myr). Cores up to M_{crit} have been stripped bare. We have approximated the time-integrated XUV flux to be independent of M_\star , so M_{crit} is also independent of M_\star (see §8.1). Photoevaporation bifurcates the f_{env} distribution into two modes with $f_{\text{env}} \approx 0\%$ and $f_{\text{env}} \approx 3\%$. The largest super-Earths have $R_{\text{crit}}/R_\oplus \approx (M_{\text{crit}}/M_\oplus)^{0.25}$ so R_{crit} is independent of M_\star . The smallest sub-Neptunes barely retained their envelopes and have $R_p \approx 2 R_{\text{crit}}$. The largest sub-Neptune size is set by a $f_{\text{env}} \approx 3\%$ envelope atop a $M_{c,\text{max}}$ core or $R_{\text{max}} = 2(M_{c,\text{max}}/M_\oplus)^{0.25}$. Since $M_{c,\text{max}} \propto M_\star$, $R_{\text{max}} \propto M_\star^{0.25}$. The model is equally applicable to the core-powered framework except mass is lost over ~ 1 Gyr timescales.

Figure 9 is reproduced here in Figure 16. In the models, sub-Neptune size increases with stellar metallicity as $R_p \propto Z^\beta$, with $\beta = 0.1$. In contrast, in §6.2 we measured $\beta = -0.01 \pm 0.02$, which rules out the predicted $\beta = 0.1$ at 4σ significance.

For visual comparison, we reproduced Figure 7d in Figure 16 with a $\beta = 0.1$ relationship over-plotted. The CXM distribution is peaked near $[\text{Fe}/\text{H}] = 0.0$, but no tilt is observable. To visually enhance the tails of the metallicity distribution, we normalized the density by the integrated density along vertical columns. Both super-Earths and sub-Neptunes show no significant size-metallicity dependence.

Our results suggests that the dominant opacity at the RCB does not track stellar metallicity. The tension between the observations and the Gupta & Schlichting

(2020) model does not reflect an insurmountable obstacle for core-powered theory, but does suggest a modification to the opacity treatment. We also note that the XUV-powered models of Owen & Wu (2017) also assumed a $\kappa \propto Z$ relationship.

8.3. Age effects

After the period of rapid mass-loss, we expect the sizes of the sub-Neptunes to decrease due to the Kelvin-Helmholtz mechanism. In §6.2, we found $R_p \propto (\text{age}/5 \text{ Gyr})^\gamma$ with $\gamma = -0.01 \pm 0.03$, consistent with no R_p -age collocation. This flat age dependence is interesting given that several studies have noted planets younger than 100 Myr seem to have inflated envelopes. In a recent compilation of nine planets with cluster-based ages under 100 Myr by Bouma et al. (2020), all were between 4 and $10 R_\oplus$. Rizzuto et al. (2018) and David




Figure 16. Panel (a): predicted planet sizes as a function of stellar metallicity in the core-powered models of Gupta & Schlichting (2020). Here, M_* and age were fixed at $1.0 M_\odot$ and 3 Gyr. Panel (c): the relative number density of *detected* planets (not occurrence). Panel (e): same as (c), except we have normalized the number density by its integral along columns of constant $[\text{Fe}/\text{H}]$ to accentuate features at low/high $[\text{Fe}/\text{H}]$. In the Gupta & Schlichting (2020) models, the typical sub-Neptune is expected to grow with metallicity according to $R_p \propto Z^\beta$, with $\beta = 0.1$. We observed $\beta = -0.01 \pm 0.02$ and show the predicted $\beta = 0.1$ dependence for comparison in panels (c) and (e). The lack of detectable variation in sub-Neptune size with metallicity when such a trend is predicted in the core-powered models points to missing or incomplete physics, perhaps involving the treatment of envelope opacity (see §8.2). Panel (b): same as panel (a) except that M_* and $[\text{Fe}/\text{H}]$ were fixed to $1.0 M_\odot$ and 0.0 dex and age was allowed to vary. Panel (d) and (f): same as panel (c) and (e) except for age. The sub-Neptunes are expected to shrink with time according to $R_p \propto t^\gamma$, with $\gamma = -0.1$. While we observed $\gamma = -0.01 \pm 0.03$, we could not conclusively rule out a $\gamma = -0.1$ dependence given the age uncertainties (see §8.3).

et al. (2019) made a similar observation among earlier compilations.

Planets in this size range are intrinsically rare among field-age stars. As a point of reference, for $P < 100$ days, Petigura et al. (2018b) measured an occurrence rate of 0.04 planets per star with $R_p = 4.0\text{--}11.3 R_\oplus$ vs. 0.64 planets per star with $R_p = 1\text{--}4 R_\oplus$ among stars with $M_* \approx 0.8\text{--}1.3 M_\odot$. Yet V1298Tau, a 20–30 Myr, solar-mass star hosts *four* planets with sizes ranging from 5 to

$10 R_\oplus$ (David et al. 2019). Planets in this size range are even rarer among low-mass stars. Dressing & Charbonneau (2013) measured an occurrence rate of 0.007 planets per star with $R_p = 4.0\text{--}11.3 R_\oplus$ and $P < 50$ days in a sample of stars with $M_* \approx 0.4\text{--}0.6 M_\odot$. Yet K2-33 is a 5–10 Myr, $0.4 M_\odot$ star that hosts a $6 R_\oplus$ planet (David et al. 2016).

Given the evidence for inflated planets younger than 100 Myr in the literature combined with the flat R_p -age

relationship observed here for ages spanning 1 to 10 Gyr, we conclude that large-scale evolution in the size of sub-Neptune envelopes concludes by ~ 1 Gyr.

The envelopes may continue to contract after ~ 1 Gyr, but the R_p -age trends may be too subtle to detect given our sample size and age uncertainties. For example, [Gupta & Schlichting \(2020\)](#) isolated the age effect in the core-powered context by constructing a planet population with M_\star and $[\text{Fe}/\text{H}]$ held fixed at $1.0 M_\odot$ and 0.0 dex and evolved them from 1 to 10 Gyr (their Figure 10, reproduced here in Figure 16). In this model, the planets obeyed a $\gamma = -0.1$ relationship. However, errors in the measured ages will conspire to flatten such a trend. We explored this effect by injecting a $\gamma = -0.1$ dependence into the observed sub-Neptune population, perturbing their ages by their uncertainties, and refitting our power-law model. We found $\gamma = -0.06$, which is consistent with our measured value of $\gamma = -0.01 \pm 0.03$ at the 2σ level. Thus, we cannot rule out the subtle $\gamma = -0.1$ dependence predicted by [Gupta & Schlichting \(2020\)](#) given our current sample size and age uncertainties.

8.4. Slope of the Radius Gap in period-radius space

While a number of theoretical models can produce the Radius Gap, they do not produce identical slopes in the P - R_p plane. Thus the slope of the Radius Gap has emerged as a potential discriminant between different models. In the [Owen & Wu \(2017\)](#) photoevaporation model, the maximum size of a super-Earth decreases with orbital period. The upper envelope is set by the size of a rocky core that is barely stripped by the integrated XUV exposure. Slightly more massive cores retain their envelopes and remain sub-Neptunes.

At larger orbital periods, \mathcal{F}_{XUV} decreases and the cores that can be completely stripped have lower masses and smaller sizes. Assuming a fixed energy-limited efficiency factor $\eta = 10\%$, [Owen & Wu \(2017\)](#) predicted that the bottom of the Radius Gap will follow $R_p \propto P^m$ with $m = -0.25$. Realistically, η should decline as mass increases since escaping gas has more time to cool as it climbs out of a deeper potential well. Adopting $\eta = 10\% (v_{\text{esc}}/15 \text{ km s}^{-1})^{-2}$ yields a shallower slope $m = -0.16$.

In the core-powered framework of [Gupta & Schlichting \(2019\)](#), the fate of an envelope around a planet with $P \gtrsim 8$ days depends on the cooling timescale $t_{\text{cool}} = E/L$ relative to the envelope mass-loss timescale $t_{\text{loss}} = M_{\text{env}}/\dot{M}$. Envelopes that escape faster than they cool $t_{\text{loss}} < t_{\text{cool}}$ are lost completely. The mass-loss timescale is depends on \dot{M} at the Bondi radius, and $t_{\text{loss}} \propto M_{\text{env}} R_s^2 c_s \rho_{\text{rcb}} \exp(GM_p/c_s^2 R_{\text{rcb}})$ where R_s

is the sonic point and c_s is the sound speed. The cooling timescale may be computed by dividing the combined thermal and gravitational energy of the envelope and core by the luminosity evaluated at the RCB. The exponential behavior of t_{loss} results in the dividing line $t_{\text{loss}} = t_{\text{cool}}$ being set by $GM_p/c_s^2 R_{\text{rcb}} = \text{constant}$, which implies $R_p \propto P^{-0.11}$ or $m = -0.11$.

We emphasize that the values of m predicted by the core- and XUV-powered models discussed above have all incorporated assumptions about the properties of planet cores such as the distribution of masses, orbital periods, and bulk compositions. [Rogers et al. \(2021\)](#) showed how changing these input parameters can alter the predicted m . *However, both the core- and XUV-models categorically predict a negative slope.*

Models that delay the accretion of gas until the nebula is nearly or completely dissipated predict an increasing slope with orbital period ([Lee et al. 2014](#); [Lee & Chiang 2016](#); [Lopez & Rice 2018](#)). Here, the core mass is determined by the solid surface density profile within a feeding zone that is proportional to the Hill radius $R_H \propto a(M_p/M_\star)^{1/3}$. A [Hayashi \(1981\)](#) profile $\Sigma(a) \propto a^{-3/2}$ yields $M_p \propto a^{0.6} M_\star^{-0.5}$ or $R_p \propto P^{0.11} M_\star^{-0.09}$ or $m = 0.11$.

For the full CXM sample, we measured $m = -0.11^{+0.02}_{-0.02}$. [Van Eylen et al. \(2018\)](#) fit the gap in a sample of 117 planets orbiting 75 stars with asteroseismic detections and found $m = -0.09^{+0.02}_{-0.04}$. Their population of stellar hosts most closely resembles a combination of our middle and high mass bins, for which we measured $m = -0.13^{+0.07}_{-0.06}$ and $-0.06^{+0.02}_{-0.02}$, which straddle the [Van Eylen et al. \(2018\)](#) value. [Martinez et al. \(2019\)](#) performed an independent analysis of CKS DR1 spectra including parallax constraints and found $m = -0.11 \pm 0.02$. Despite differences in planets used, planet/star parameter provenance, and slope measurement technique, our measurements of slope are consistent with the [Van Eylen et al. \(2018\)](#) and [Martinez et al. \(2019\)](#) measurements and with the core- and XUV-powered model predictions.

Shifting our attention to low-mass stars, we resolved the Radius Gap among the 185 planets in our low stellar mass bin $M_\star = 0.5\text{--}0.7 M_\odot$. We measured $m = -0.12^{+0.06}_{-0.04}$, consistent with that of the full sample. As a point of comparison, [Cloutier & Menou \(2020\)](#) measured the occurrence of planets in a sample of stars having $T_{\text{eff}} < 4700$ K drawn from *Kepler* and *K2* (275 and 53 planets). The host stars were $M_\star = 0.08\text{--}0.93 M_\odot$, but most were between 0.5 and $0.8 M_\odot$. They resolved the Radius Gap and measured a positive slope in P - R_p space of $m = 0.058 \pm 0.022$. They interpreted this as a signature of gas-poor formation channels around lower

mass stars. However, the Radius Gap is only visible in their full 328 planet sample, but not when they restrict their analysis to the 126 planets around $M_\star = 0.08\text{--}0.65 M_\odot$ hosts (their Figure 12). Our measured slope is $\approx 4\sigma$ lower than Cloutier & Menou (2020). We found no evidence of a change of slope and thus do not favor an alternate formation pathway for planets around $M_\star = 0.5\text{--}0.7 M_\odot$ stars. Van Eylen et al. (2021) arrived at a similar conclusion after analyzing a sample of 27 planets orbiting stars with $T_{\text{eff}} < 4000$ K.

Before concluding this section, we wish to remark on some of the challenges associated with using slope of the Radius Gap as a signpost of formation. Note that our uncertainties on m are *larger* than Martinez et al. (2019) even though we resolved the valley at higher contrast (compare our Figure 7 to their Figure 12). One may understand this through the following limiting case: Consider super-Earth/sup-Neptune populations separated by a gap completely devoid of planets. In such a scenario, selecting the minima used to fit the Radius Gap in §6.1 is ill-posed because there is no longer a single minimum R_p . Thus, a broad range of $R_{p,0}$ and m are allowed. *Thus, the statistic for assessing model predictions has the undesirable quality of becoming more uncertain as observational uncertainties improve.*

Moreover, there is no standard approach to fitting the absence of planets. For example, Van Eylen et al. (2018) used a support vector machine scheme, Berger et al. (2020a) used the ‘gapfit’ code that subtracts off a trial $R_p \propto P^m$ relationship and evaluates a 1D KDE of the residuals, and Martinez et al. (2019) and this work fit a train of minima computed along 1D projections through the P - R_p plane. All approaches rely on an *ad hoc* smoothing parameter which determines the relative influence of the few planets near the edge of the gap relative to the many planets far from the gap on the fit. These differences in fitting method are less troublesome in theoretical studies which can model limitless numbers of planets and do not contend with complications like false positive contamination or mischaracterized measurement uncertainties.

Recently, Rogers & Owen (2021) introduced a different framework for evaluating the agreement between formation theory and the observed planet population that does not rely on measuring the slope of the gap. Here, a model planet population is subjected to a specified set of physical processes, e.g. XUV- or core-powered mass-loss. The synthetic population is transformed into an occurrence rate density distribution. This is then converted into a number rate density distribution by accounting for properties of the parent stellar population, losses due to non-transiting planets, and losses due to

pipeline incompleteness. In this framework, one evaluates the likelihood that the observed planets are a realization of an inhomogeneous Poisson point process with the specified number rate density. One may then optimize the model parameters and characterize their uncertainties with using standard techniques. Such a treatment of the CXM sample would be interesting, but is beyond the scope of this paper.

8.5. Period-flux distribution of planets

In §7.3, we characterized the period distribution of super-Earths and sub-Neptunes dN/dP in three bins of stellar mass. We parameterized these distributions as smooth broken power-laws with variable breakpoints P_0 . From $M_\star = 0.5\text{--}1.4 M_\odot$, the sub-Neptune falloff occurs at a nearly constant $P_0 \approx 10$ days. Similarly, when we modeled the flux distribution, dN/dS_{inc} , we found that $S_{\text{inc},0}$ increases by nearly an order of magnitude from ≈ 20 to $200 S_\oplus$.

Under the assumptions that planet cores are (1) uniformly distributed in log-period and (2) have masses that are uncorrelated with M_\star , the above observation would be strong evidence for XUV-powered and against core-powered mass-loss. As we explain in §8.1, $\mathcal{F}_{\text{XUV}} \propto M_\star^{0.33}$, and thus only varies by 40% over our full mass stellar range. Thus the critical \mathcal{F}_{XUV} that strips the typical sub-Neptune maps to P_0 that is constant with M_\star . Fixed P_0 corresponds to $S_{\text{inc},0}$ that grows with M_\star . Under the same assumptions, $S_{\text{inc},0}$ should be constant with M_\star in the core-powered model.

Neither of the above assumptions, however, appear to be correct. The steep decline in super-Earths for $P \lesssim P_0 \approx 5$ days suggests rocky cores struggle to form inside this boundary. Several mechanisms may be responsible, including magnetospheric truncation of the inner disk (e.g., Lee & Chiang 2017) or a pressure trap due a silicate sublimation front (e.g., Flock et al. 2019). Moreover, the M_\star - R_p correlation discussed in §8.1 suggests that massive stars make massive cores.

Therefore, we cannot rule out the core-powered model based on the dependence of P_0 and $S_{\text{inc},0}$ on M_\star . In this framework, stars that are $1.0\text{--}1.4 M_\odot$ efficiently produce sub-Neptunes when $S_{\text{inc}} \lesssim S_{\text{inc},0} \approx 200 S_\oplus$. Stars that are $0.5\text{--}0.7 M_\odot$ only efficiently produce sub-Neptunes when $S_{\text{inc}} \lesssim S_{\text{inc},0} \approx 20 S_\oplus$. The lack of sub-Neptunes in the $S_{\text{inc}} \approx 20\text{--}200 S_\oplus$ range is not due to envelope stripping but instead due to the absence of suitable cores. Inspecting Figure 13, we see that the occurrence rate density of super-Earths declines by a factor of five over $S_{\text{inc}} = 20\text{--}200 S_\oplus$.

There is still considerable uncertainty in the distribution of core masses and further uncertainty in how that

distribution varies with P and M_\star . Given these unknowns, both core- and XUV-powered frameworks appear sufficiently flexible to match the observations. Additional mass measurements of sub-Neptunes as a function of P - S_{inc} - M_\star should help to constrain the models since R_p and M_p together constrain both core mass and envelope fraction.

9. SUMMARY AND CONCLUSION

Here, we provide a brief summary of our results and point toward future improvements to this work:

- We augmented the 1305-star CKS DR1 spectral library with DR2 containing 411 new spectra.
- We performed a homogeneous analysis of the combined DR1 and DR2 samples and derived T_{eff} , $[\text{Fe}/\text{H}]$, $v \sin i$, M_\star , R_\star , and age.
- We constructed a curated sample of 970 planets orbiting 703 stars with updated star/planet properties.
- We resolved the Radius Gap and projected the planet population as a function of P , S_{inc} , M_\star , $[\text{Fe}/\text{H}]$, and age.
- The Radius Gap in the P - R_p plane follows $R_p \propto P^m$ with $m = -0.10 \pm 0.03$, consistent with both core-powered and XUV-powered models but inconsistent with gas-poor formation models.
- We observed no significant change in m over 0.5–1.4 M_\odot .
- Sub-Neptunes tend to be larger around higher mass stars and follow $R_p \propto (M_\star/M_\odot)^\alpha$ with $\alpha = 0.25 \pm 0.03$. The super-Earths exhibit no measurable M_\star dependence $\alpha = 0.02 \pm 0.03$. Taken together, these trends are consistent with a core-mass distribution that scales linearly with stellar mass $M_c \propto M_\star$.
- The average sub-Neptune is not measurably larger around higher metallicity stars, disfavoring a simple linear relationship between envelope opacity and stellar metallicity $\kappa \propto Z$.
- The average sub-Neptune does not measurably shrink over 1–10 Gyr. Given the large radii observed among stars younger than 100 Myr, we conclude that the majority of planet radius contraction concludes by ~ 1 Gyr.
- The period distribution of sub-Neptunes has a breakpoint P_0 in M_\star . This is consistent with the

predictions of photoevaporation models. However, core-powered models may still be viable given uncertainties in the underlying distribution of planet cores.

The bifurcation of small planets in to two distinct populations was one of the most intriguing results from the *Kepler* mission. This feature appears most consistent with models where close-in planets acquire and lose H/He envelopes. XUV radiation from young stars or heat from cooling cores are leading candidates for the energy source that powers this mass loss. While we worked to identify the dominant mass-loss process, we found both mechanisms are flexible enough to match our observations.

We look forward to additional observational and theoretical work that could help discriminate between these two theories. With *Gaia*, it will soon be possible to extend the analysis presented here to a larger sample of planets (2–3 \times). Future *Gaia* data releases will include low- and high-resolution spectra for nearly all *Kepler* stars. These data will increase the upper M_\star limit since rapid rotation above 6500 K does not wash out broadband SED information. *Gaia*-based refinements in T_{eff} and $[\text{Fe}/\text{H}]$ will be especially useful in determining masses and ages for additional early-type stars. However, increases in sample size from *Gaia* may not settle the core-powered vs. XUV-powered question. An analysis of synthetic planet populations by Rogers et al. (2021) suggested that $\gtrsim 5,000$ planets are needed to do so.

We found that heuristic descriptions of the Radius Gap such as its slope in the P - R_p plane were neither straightforward to compute nor straightforward to compare to model predictions, which are based on a previous occurrence analysis. The disconnect between the demographic analysis and the population modeling leads to concerns of self-inconsistency. The modeling approach of Rogers & Owen (2021) offers a firmer statistical basis for comparing models and observations. This approach can also naturally accommodate realistic non-independent distributions of host star properties like mass, metallicity, and age.

Aside from improved host star properties, additional planet observables should shed light on the mass-loss process. Mass measurements of transiting planets spanning P , S_{inc} , $[\text{Fe}/\text{H}]$, and age will help constrain the core-mass distribution which contributes to the remaining flexibility of both models since mass and radius together constrain core mass and envelope fraction. Finally, direct measurements of mass-loss as a function of the same properties would be especially valuable. Outflows from a handful of planets have been recently de-

tected using the He-10830 Å line, and Gupta & Schlichting (2021) have highlighted a number of planets that may actively be experiencing mass loss in the core-powered framework.

Software: The software used in this analysis is available at <https://github.com/petigura/CKS-Cool>. We used the following additional packages: *Astropy* (Astropy Collaboration et al. 2013), *emcee* (Foreman-Mackey et al. 2013), *isoclassify* (Huber et al. 2017; Berger et al. 2020b), *Lmfit* (Newville et al. 2014), *Matplotlib* (Hunter 2007), *Numpy* (van der Walt et al. 2011), *Scipy* (Jones et al. 2001), *H5py* (<https://doi.org/10.5281/zenodo.5585380>), *Pandas* (McKinney 2010), *Seaborn* (Waskom 2021), and *Xarray* (Hoyer & Hamman 2017)

Facilities: Keck:I (HIRES), Kepler

ACKNOWLEDGMENTS

We thank Natalie Batalha, Konstantin Batygin, Akash Gupta, Daniel Huber, Eric Lopez, Hilke Schlichting, Joshua Schlieder, Samuel Yee, and Jon Zink for valuable conversations that improved this manuscript.

Data presented herein were obtained at the W. M. Keck Observatory from telescope time allocated to the National Aeronautics and Space Administration through the agency’s scientific partnership with the California

Institute of Technology and the University of California. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation

Kepler was competitively selected as the tenth NASA Discovery mission. Funding for this mission is provided by the NASA Science Mission Directorate. We thank the Kepler Science Office, the Science Operations Center, Threshold Crossing Event Review Team (TCERT), and the Follow-up Observations Program (FOP) working group for their work on all steps in the planet discovery process ranging from selecting target stars to curating planet catalogs.

E.A.P. acknowledges support from the following sources: a Hubble Fellowship grant HST-HF2-51365.001-A awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc. for NASA under contract NAS 5-26555; a NASA Keck PI Data Award (80NSSC20K0308), administered by the NASA Exoplanet Science Institute; a NASA Astrophysics Data Analysis Program (ADAP) grant (80NSSC20K0457); and the Alfred P. Sloan Foundation.

The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has long had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

REFERENCES

- Akeson, R. L., Chen, X., Ciardi, D., et al. 2013, *PASP*, 125, 989, doi: [10.1086/672273](https://doi.org/10.1086/672273)
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Batalha, N. M., Borucki, W. J., Koch, D. G., et al. 2010, *ApJL*, 713, L109, doi: [10.1088/2041-8205/713/2/L109](https://doi.org/10.1088/2041-8205/713/2/L109)
- Batalha, N. M., Borucki, W. J., Bryson, S. T., et al. 2011, *ApJ*, 729, 27, doi: [10.1088/0004-637X/729/1/27](https://doi.org/10.1088/0004-637X/729/1/27)
- Beaugé, C., & Nesvorný, D. 2013, *ApJ*, 763, 12, doi: [10.1088/0004-637X/763/1/12](https://doi.org/10.1088/0004-637X/763/1/12)
- Berger, T. A., Howard, A. W., & Boesgaard, A. M. 2018a, *ApJ*, 855, 115, doi: [10.3847/1538-4357/aab154](https://doi.org/10.3847/1538-4357/aab154)
- Berger, T. A., Huber, D., Gaidos, E., & van Saders, J. L. 2018b, *ApJ*, 866, 99, doi: [10.3847/1538-4357/aada83](https://doi.org/10.3847/1538-4357/aada83)
- Berger, T. A., Huber, D., Gaidos, E., van Saders, J. L., & Weiss, L. M. 2020a, *AJ*, 160, 108, doi: [10.3847/1538-3881/aba18a](https://doi.org/10.3847/1538-3881/aba18a)
- Berger, T. A., Huber, D., van Saders, J. L., et al. 2020b, *AJ*, 159, 280, doi: [10.3847/1538-3881/159/6/280](https://doi.org/10.3847/1538-3881/159/6/280)
- Bouma, L. G., Hartman, J. D., Brahm, R., et al. 2020, *AJ*, 160, 239, doi: [10.3847/1538-3881/abb9ab](https://doi.org/10.3847/1538-3881/abb9ab)
- Brown, T. M., Latham, D. W., Everett, M. E., & Esquerdo, G. A. 2011, *AJ*, 142, 112, doi: [10.1088/0004-6256/142/4/112](https://doi.org/10.1088/0004-6256/142/4/112)
- Bryson, S., Coughlin, J., Batalha, N. M., et al. 2020, *AJ*, 159, 279, doi: [10.3847/1538-3881/ab8a30](https://doi.org/10.3847/1538-3881/ab8a30)
- Castelli, F., & Kurucz, R. L. 2004, *ArXiv Astrophysics e-prints*
- Chatterjee, S., & Chen, H. 2018, *ApJ*, 852, 58, doi: [10.3847/1538-4357/aa9e05](https://doi.org/10.3847/1538-4357/aa9e05)
- Chen, H., & Rogers, L. A. 2016, *ApJ*, 831, 180, doi: [10.3847/0004-637X/831/2/180](https://doi.org/10.3847/0004-637X/831/2/180)
- Choi, J., Dotter, A., Conroy, C., et al. 2016, *ApJ*, 823, 102, doi: [10.3847/0004-637X/823/2/102](https://doi.org/10.3847/0004-637X/823/2/102)
- Christiansen, J. L., Clarke, B. D., Burke, C. J., et al. 2020, *AJ*, 160, 159, doi: [10.3847/1538-3881/abab0b](https://doi.org/10.3847/1538-3881/abab0b)
- Cloutier, R., & Menou, K. 2020, *AJ*, 159, 211, doi: [10.3847/1538-3881/ab8237](https://doi.org/10.3847/1538-3881/ab8237)

- Clubb, K., Howard, A. W., Isaacson, H., Marcy, G. W., & Wright, J. T. 2018, *Research Notes of the American Astronomical Society*, 2, 44, doi: [10.3847/2515-5172/aab7e9](https://doi.org/10.3847/2515-5172/aab7e9)
- Coelho, P., Barbuy, B., Meléndez, J., Schiavon, R. P., & Castilho, B. V. 2005, *A&A*, 443, 735, doi: [10.1051/0004-6361:20053511](https://doi.org/10.1051/0004-6361:20053511)
- David, T. J., Petigura, E. A., Luger, R., et al. 2019, *ApJL*, 885, L12, doi: [10.3847/2041-8213/ab4c99](https://doi.org/10.3847/2041-8213/ab4c99)
- David, T. J., Hillenbrand, L. A., Petigura, E. A., et al. 2016, *Nature*, 534, 658, doi: [10.1038/nature18293](https://doi.org/10.1038/nature18293)
- David, T. J., Contardo, G., Sandoval, A., et al. 2021, *AJ*, 161, 265, doi: [10.3847/1538-3881/abf439](https://doi.org/10.3847/1538-3881/abf439)
- Dotter, A. 2016, *ApJS*, 222, 8, doi: [10.3847/0067-0049/222/1/8](https://doi.org/10.3847/0067-0049/222/1/8)
- Dressing, C. D., & Charbonneau, D. 2013, *ApJ*, 767, 95, doi: [10.1088/0004-637X/767/1/95](https://doi.org/10.1088/0004-637X/767/1/95)
- Dressing, C. D., Newton, E. R., Schlieder, J. E., et al. 2017, *ApJ*, 836, 167, doi: [10.3847/1538-4357/836/2/167](https://doi.org/10.3847/1538-4357/836/2/167)
- Flock, M., Turner, N. J., Mulders, G. D., et al. 2019, *A&A*, 630, A147, doi: [10.1051/0004-6361/201935806](https://doi.org/10.1051/0004-6361/201935806)
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306, doi: [10.1086/670067](https://doi.org/10.1086/670067)
- Fulton, B. J., & Petigura, E. A. 2018, *ArXiv e-prints*. <https://arxiv.org/abs/1805.01453>
- Fulton, B. J., Petigura, E. A., Howard, A. W., et al. 2017, *AJ*, 154, 109, doi: [10.3847/1538-3881/aa80eb](https://doi.org/10.3847/1538-3881/aa80eb)
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1, doi: [10.1051/0004-6361/201833051](https://doi.org/10.1051/0004-6361/201833051)
- Ginzburg, S., Schlichting, H. E., & Sari, R. 2018, *MNRAS*, 476, 759, doi: [10.1093/mnras/sty290](https://doi.org/10.1093/mnras/sty290)
- Goodman, J., & Weare, J. 2010, *Communications in Applied Mathematics and Computational Science*, 5, 65, doi: [10.2140/camcos.2010.5.65](https://doi.org/10.2140/camcos.2010.5.65)
- Green, G. M., Schlafly, E., Zucker, C., Speagle, J. S., & Finkbeiner, D. 2019, *ApJ*, 887, 93, doi: [10.3847/1538-4357/ab5362](https://doi.org/10.3847/1538-4357/ab5362)
- Green, G. M., Schlafly, E. F., Finkbeiner, D., et al. 2018, *ArXiv e-prints*. <https://arxiv.org/abs/1801.03555>
- Gupta, A., & Schlichting, H. E. 2019, *MNRAS*, 487, 24, doi: [10.1093/mnras/stz1230](https://doi.org/10.1093/mnras/stz1230)
- . 2020, *MNRAS*, 493, 792, doi: [10.1093/mnras/staa315](https://doi.org/10.1093/mnras/staa315)
- . 2021, *MNRAS*, 504, 4634, doi: [10.1093/mnras/stab1128](https://doi.org/10.1093/mnras/stab1128)
- Hayashi, C. 1981, *Progress of Theoretical Physics Supplement*, 70, 35, doi: [10.1143/PTPS.70.35](https://doi.org/10.1143/PTPS.70.35)
- Howard, A. W., Johnson, J. A., Marcy, G. W., et al. 2010, *ApJ*, 721, 1467, doi: [10.1088/0004-637X/721/2/1467](https://doi.org/10.1088/0004-637X/721/2/1467)
- Hoyer, S., & Hamman, J. 2017, *Journal of Open Research Software*, 5, doi: [10.5334/jors.148](https://doi.org/10.5334/jors.148)
- Hsu, D. C., Ford, E. B., Ragozzine, D., & Morehead, R. C. 2018, *AJ*, 155, 205, doi: [10.3847/1538-3881/aab9a8](https://doi.org/10.3847/1538-3881/aab9a8)
- Huber, D., Zinn, J., Bojsen-Hansen, M., et al. 2017, *ApJ*, 844, 102, doi: [10.3847/1538-4357/aa75ca](https://doi.org/10.3847/1538-4357/aa75ca)
- Hunter, J. D. 2007, *Computing In Science & Engineering*, 9, 90, doi: [10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)
- Jenkins, J. M. 2002, *ApJ*, 575, 493, doi: [10.1086/341136](https://doi.org/10.1086/341136)
- Jenkins, J. M., Caldwell, D. A., Chandrasekaran, H., et al. 2010, *ApJL*, 713, L87, doi: [10.1088/2041-8205/713/2/L87](https://doi.org/10.1088/2041-8205/713/2/L87)
- Jin, S., Mordasini, C., Parmentier, V., et al. 2014, *ApJ*, 795, 65, doi: [10.1088/0004-637X/795/1/65](https://doi.org/10.1088/0004-637X/795/1/65)
- Johnson, J. A., Petigura, E. A., Fulton, B. J., et al. 2017, *AJ*, 154, 108, doi: [10.3847/1538-3881/aa80e7](https://doi.org/10.3847/1538-3881/aa80e7)
- Jones, E., Oliphant, T., Peterson, P., et al. 2001, *SciPy: Open source scientific tools for Python*. <http://www.scipy.org/>
- Kippenhahn, R., & Weigert, A. 1990, *Stellar Structure and Evolution*
- Kolbl, R., Marcy, G. W., Isaacson, H., & Howard, A. W. 2015, *AJ*, 149, 18, doi: [10.1088/0004-6256/149/1/18](https://doi.org/10.1088/0004-6256/149/1/18)
- Kurucz, R. L., Furenlid, I., Brault, J., & Testerman, L. 1984, *Solar flux atlas from 296 to 1300 nm*
- Lee, E. J., & Chiang, E. 2016, *ApJ*, 817, 90, doi: [10.3847/0004-637X/817/2/90](https://doi.org/10.3847/0004-637X/817/2/90)
- . 2017, *ApJ*, 842, 40, doi: [10.3847/1538-4357/aa6fb3](https://doi.org/10.3847/1538-4357/aa6fb3)
- Lee, E. J., Chiang, E., & Ormel, C. W. 2014, *ApJ*, 797, 95, doi: [10.1088/0004-637X/797/2/95](https://doi.org/10.1088/0004-637X/797/2/95)
- Lindegren, L. 2018, *Re-normalising the astrometric chi-square in Gaia DR2, GAIA-C3-TN-LU-LL-124-01*, Tech. rep., Lund Observatory
- Lopez, E. D., & Fortney, J. J. 2013, *ApJ*, 776, 2, doi: [10.1088/0004-637X/776/1/2](https://doi.org/10.1088/0004-637X/776/1/2)
- Lopez, E. D., & Rice, K. 2018, *MNRAS*, 479, 5303, doi: [10.1093/mnras/sty1707](https://doi.org/10.1093/mnras/sty1707)
- Lundkvist, M. S., Kjeldsen, H., Albrecht, S., et al. 2016, *Nature Communications*, 7, 11201, doi: [10.1038/ncomms11201](https://doi.org/10.1038/ncomms11201)
- Martinez, C. F., Cunha, K., Ghezzi, L., & Smith, V. V. 2019, *ApJ*, 875, 29, doi: [10.3847/1538-4357/ab0d93](https://doi.org/10.3847/1538-4357/ab0d93)
- Mathur, S., Huber, D., Batalha, N. M., et al. 2017, *ApJS*, 229, 30, doi: [10.3847/1538-4365/229/2/30](https://doi.org/10.3847/1538-4365/229/2/30)
- McDonald, G. D., Kreidberg, L., & Lopez, E. 2019, *ApJ*, 876, 22, doi: [10.3847/1538-4357/ab1095](https://doi.org/10.3847/1538-4357/ab1095)
- McKinney, W. 2010, in *Proceedings of the 9th Python in Science Conference*, ed. S. van der Walt & J. Millman, 51 – 56
- Newton, E. R., Charbonneau, D., Irwin, J., & Mann, A. W. 2015, *ApJ*, 800, 85, doi: [10.1088/0004-637X/800/2/85](https://doi.org/10.1088/0004-637X/800/2/85)

- Newville, M., Stensitzki, T., Allen, D. B., & Ingargiola, A. 2014, LMFIT: Non-Linear Least-Square Minimization and Curve-Fitting for Python, doi: [10.5281/zenodo.11813](https://doi.org/10.5281/zenodo.11813).
<https://doi.org/10.5281/zenodo.11813>
- Owen, J. E., & Murray-Clay, R. 2018, MNRAS, 480, 2206, doi: [10.1093/mnras/sty1943](https://doi.org/10.1093/mnras/sty1943)
- Owen, J. E., & Wu, Y. 2013, ApJ, 775, 105, doi: [10.1088/0004-637X/775/2/105](https://doi.org/10.1088/0004-637X/775/2/105)
- . 2017, ApJ, 847, 29, doi: [10.3847/1538-4357/aa890a](https://doi.org/10.3847/1538-4357/aa890a)
- Pascucci, I., Testi, L., Herczeg, G. J., et al. 2016, ApJ, 831, 125, doi: [10.3847/0004-637X/831/2/125](https://doi.org/10.3847/0004-637X/831/2/125)
- Paxton, B., Bildsten, L., Dotter, A., et al. 2011, ApJS, 192, 3, doi: [10.1088/0067-0049/192/1/3](https://doi.org/10.1088/0067-0049/192/1/3)
- Paxton, B., Cantiello, M., Arras, P., et al. 2013, ApJS, 208, 4, doi: [10.1088/0067-0049/208/1/4](https://doi.org/10.1088/0067-0049/208/1/4)
- Paxton, B., Marchant, P., Schwab, J., et al. 2015, ApJS, 220, 15, doi: [10.1088/0067-0049/220/1/15](https://doi.org/10.1088/0067-0049/220/1/15)
- Petigura, E. A. 2015, PhD thesis, University of California, Berkeley
- . 2020, AJ, 160, 89, doi: [10.3847/1538-3881/ab9fff](https://doi.org/10.3847/1538-3881/ab9fff)
- Petigura, E. A., Howard, A. W., Marcy, G. W., et al. 2017, AJ, 154, 107, doi: [10.3847/1538-3881/aa80de](https://doi.org/10.3847/1538-3881/aa80de)
- Petigura, E. A., Marcy, G. W., Winn, J. N., et al. 2018a, AJ, 155, 89, doi: [10.3847/1538-3881/aaa54c](https://doi.org/10.3847/1538-3881/aaa54c)
- Petigura, E. A., Crossfield, I. J. M., Isaacson, H., et al. 2018b, AJ, 155, 21, doi: [10.3847/1538-3881/aa9b83](https://doi.org/10.3847/1538-3881/aa9b83)
- Press, W. H. 2002, Numerical recipes in C++ : the art of scientific computing, ed. Press, W. H.
- Rizzuto, A. C., Vanderburg, A., Mann, A. W., et al. 2018, Zodiacal Exoplanets in Time (ZEIT). VIII. A Two-planet System in Praesepse from K2 Campaign 16, doi: [10.3847/1538-3881/aadf37](https://doi.org/10.3847/1538-3881/aadf37)
- Rogers, J. G., Gupta, A., Owen, J. E., & Schlichting, H. E. 2021, arXiv e-prints, arXiv:2105.03443.
<https://arxiv.org/abs/2105.03443>
- Rogers, J. G., & Owen, J. E. 2021, MNRAS, 503, 1526, doi: [10.1093/mnras/stab529](https://doi.org/10.1093/mnras/stab529)
- Sinukoff, E. 2018, PhD thesis, University of Hawaii, Manoa
- Stumpe, M. C., Smith, J. C., Van Cleve, J. E., et al. 2012, PASP, 124, 985, doi: [10.1086/667698](https://doi.org/10.1086/667698)
- Szabó, G. M., & Kiss, L. L. 2011, ApJL, 727, L44, doi: [10.1088/2041-8205/727/2/L44](https://doi.org/10.1088/2041-8205/727/2/L44)
- Thompson, S. E., Coughlin, J. L., Hoffman, K., et al. 2018, ApJS, 235, 38, doi: [10.3847/1538-4365/aab4f9](https://doi.org/10.3847/1538-4365/aab4f9)
- van der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, Computing in Science Engineering, 13, 22, doi: [10.1109/MCSE.2011.37](https://doi.org/10.1109/MCSE.2011.37)
- Van Eylen, V., Agentoft, C., Lundkvist, M. S., et al. 2018, MNRAS, 479, 4786, doi: [10.1093/mnras/sty1783](https://doi.org/10.1093/mnras/sty1783)
- Van Eylen, V., Astudillo-Defru, N., Bonfils, X., et al. 2021, MNRAS, 507, 2154, doi: [10.1093/mnras/stab2143](https://doi.org/10.1093/mnras/stab2143)
- Vogt, S. S., Allen, S. L., Bigelow, B. C., et al. 1994, 2198, 362
- Waskom, M. L. 2021, Journal of Open Source Software, 6, 3021, doi: [10.21105/joss.03021](https://doi.org/10.21105/joss.03021)
- Weiss, L. M., & Marcy, G. W. 2014, ApJL, 783, L6, doi: [10.1088/2041-8205/783/1/L6](https://doi.org/10.1088/2041-8205/783/1/L6)
- Weiss, L. M., Marcy, G. W., Petigura, E. A., et al. 2018, AJ, 155, 48, doi: [10.3847/1538-3881/aa9ff6](https://doi.org/10.3847/1538-3881/aa9ff6)
- Winn, J. N., Petigura, E. A., Morton, T. D., et al. 2017, AJ, 154, 270, doi: [10.3847/1538-3881/aa93e3](https://doi.org/10.3847/1538-3881/aa93e3)
- Wolfgang, A., Rogers, L. A., & Ford, E. B. 2016, ApJ, 825, 19, doi: [10.3847/0004-637X/825/1/19](https://doi.org/10.3847/0004-637X/825/1/19)
- Wood, M. L., Mann, A. W., & Kraus, A. L. 2021, arXiv e-prints, arXiv:2106.09040.
<https://arxiv.org/abs/2106.09040>
- Wu, Y. 2019, ApJ, 874, 91, doi: [10.3847/1538-4357/ab06f8](https://doi.org/10.3847/1538-4357/ab06f8)
- Wyatt, M. C., Kral, Q., & Sinclair, C. A. 2020, MNRAS, 491, 782, doi: [10.1093/mnras/stz3052](https://doi.org/10.1093/mnras/stz3052)
- Yee, S. W., Petigura, E. A., & von Braun, K. 2017, ApJ, 836, 77, doi: [10.3847/1538-4357/836/1/77](https://doi.org/10.3847/1538-4357/836/1/77)
- Zeng, L., Jacobsen, S. B., Sasselov, D. D., et al. 2019, Proceedings of the National Academy of Science, 116, 9723, doi: [10.1073/pnas.1812905116](https://doi.org/10.1073/pnas.1812905116)
- Zinn, J. C., Pinsonneault, M. H., Huber, D., & Stello, D. 2018, ArXiv e-prints. <https://arxiv.org/abs/1805.02650>