

MID-INFRARED SPECTRAL DIAGNOSIS OF SUBMILLIMETER GALAXIES

ALEXANDRA POPE,^{1,2,3} RANGA-RAM CHARY,⁴ DAVID M. ALEXANDER,⁵ LEE ARMUS,⁴ MARK DICKINSON,⁶
DAVID ELBAZ,⁷ DAVID FRAYER,⁴ DOUGLAS SCOTT,¹ AND HARRY TEPLITZ⁴

Received 2007 August 22; accepted 2007 November 19

ABSTRACT

We present deep mid-IR spectroscopy with *Spitzer* of 13 SMGs in the GOODS-N field. We find strong PAH emission in all of our targets, which allows us to measure mid-IR spectroscopic redshifts and place constraints on the contribution from star formation and AGN activity to the mid-IR emission. In the high-S/N composite spectrum, we find that the hot dust continuum from an AGN contributes at most 30% of the mid-IR luminosity. Individually, only 2/13 SMGs have continuum emission dominating the mid-IR luminosity; one of these SMGs, C1, remains undetected in the deep X-ray images but shows a steeply rising continuum in the mid-IR indicative of a Compton-thick AGN. We find that the mid-IR properties of SMGs are distinct from those of 24 μm -selected ULIRGs at $z \sim 2$; the former are predominantly dominated by star formation, while the latter are a more heterogeneous sample with many showing significant AGN activity. We fit the IRS spectrum and the mid-IR to radio photometry of SMGs with template SEDs to determine the best estimate of the total IR luminosity from star formation. While many SMGs contain an AGN as evinced by their X-ray properties, our multiwavelength analysis shows that the total IR luminosity, L_{IR} , in SMGs is dominated by star formation. We find that high-redshift SMGs lie on the relation between L_{IR} and $L_{\text{PAH},6.2}$ (or $L_{\text{PAH},7.7}$ or $L_{\text{PAH},11.3}$) that has been established for local starburst galaxies. This suggests that PAH luminosity can be used as a proxy for the SFR in SMGs. SMGs are consistent with being a short-lived cool phase in a massive merger where the AGN does not appear to have become strong enough to heat the dust and dominate the mid- or far-IR emission.

Subject headings: galaxies: active — galaxies: evolution — galaxies: starburst — infrared: galaxies — submillimeter — techniques: spectroscopic

Online material: color figures

1. INTRODUCTION

The *Cosmic Background Explorer (COBE)* revealed that the extragalactic background light at infrared (IR) wavelengths was comparable to that at optical wavelengths, indicating that half of the star formation and active galactic nucleus (AGN) emission, averaged over cosmic time, is obscured by dust (e.g., Puget et al. 1996; Hauser & Dwek 2001). In the local universe, the volume density of dusty, IR-luminous galaxies is insufficient to account for this background. Therefore, the density of such galaxies must evolve strongly with redshift (for a review see, e.g., Lagache et al. 2005). In fact, imaging observations with the *Infrared Space Observatory (ISO)* and the *Spitzer Space Telescope* (hereafter *Spitzer*) have shown that about 70% of the comoving star formation rate density (SFRD) at $0.5 < z < 3$ is obscured by dust (Chary & Elbaz 2001, hereafter CE01; Le Floc'h et al. 2005; Pérez-González et al. 2005).

In the local universe, we refer to these dusty galaxies as luminous infrared galaxies [LIRGs; $11 \leq \log L_{\text{IR}}(L_{\odot}) < 12$] and ultraluminous infrared galaxies [ULIRGs; $\log L_{\text{IR}}(L_{\odot}) \geq 12$], where L_{IR} is the total IR luminosity from 8 to 1000 μm (for a review see Sanders & Mirabel 1996). It is now becoming clear

that IR-luminous galaxies are a heterogeneous population with selection functions at different mid- and far-IR wavelengths revealing different subsets of this population (Daddi et al. 2005; Lutz et al. 2005b; Yan et al. 2005; Pope et al. 2006).

Perhaps the most enigmatic members of the high-redshift ULIRG population are the submillimeter galaxies (SMGs). These galaxies were initially discovered in the late 1990s with the Submillimeter Common-User Bolometer Array (SCUBA; Holland et al. 1999) on the James Clerk Maxwell Telescope (JCMT) and then later with the Max-Planck Millimeter Bolometer (MAMBO) camera on the Institut de Radio Astronomie Millimétrique (IRAM) 30 m telescope (for a review see Blain et al. 2002). They have proven to be difficult to study, due to the large beam size of submillimeter observations and the difficulty in finding counterparts at other wavelengths (e.g., Ivison et al. 2002; Pope et al. 2005). Nevertheless, it has been found that they are very massive systems (Swinbank et al. 2004; Borys et al. 2005; Greve et al. 2005; Tacconi et al. 2006) at $z \sim 2$ (Chapman et al. 2005; Pope et al. 2006; Aretxaga et al. 2007), which are thought to be connected to the most massive present-day galaxies via an evolutionary sequence (Lilly et al. 1999). A crucial aspect that needs to be explored in SMGs is what powers their extreme luminosities: AGN activity or star formation?

The mid-IR spectral regime is a particularly good probe of what is producing the total IR luminosity since it has contributions from both star formation and AGNs each with a unique signature at these wavelengths. The 5–15 μm mid-IR spectral energy distribution (SED) can be decomposed into three main components (e.g., Sajina et al. 2007): (1) emission features from what are thought to be polycyclic aromatic hydrocarbon (PAH) molecules (Puget & Leger 1989; Allamandola et al. 1999), (2) power-law (or warm blackbody) emission, and (3) extinction characterized by prominent silicate absorption features (e.g., Draine 2003). The

¹ Department of Physics and Astronomy, University of British Columbia, Vancouver, BC V6T 1Z1, Canada.

² Current address: National Optical Astronomy Observatory, Tucson, AZ 85719; pope@noao.edu.

³ *Spitzer* Fellow.

⁴ *Spitzer* Science Center, MS 220-6, California Institute of Technology, Pasadena, CA 91125.

⁵ Department of Physics, Durham University, Durham DH1 3LE, UK.

⁶ National Optical Astronomy Observatory, Tucson, AZ 85719.

⁷ Laboratoire AIM, CEA/DSM-CNRS-Université Paris Diderot, DAPNIA/Service d'Astrophysique, CEA Saclay, 91191 Gif-sur-Yvette Cedex, France.

first is understood to be powered entirely by star formation, while the second is predominantly emission from the AGN (Genzel et al. 1998). The second component can also contain contributions from hot dust (very small grains) present in the most energetic H II regions (e.g., Tran et al. 2001). The third component can be found in both starbursts and AGNs since it just requires a dust screen around hot dust emission.

Measuring the contribution from ULIRGs to the global star formation rate (SFR) requires dissecting their IR emission into contributions from starbursts and AGNs. Mid-IR spectroscopy with *ISO* made it possible to decompose the various contributors in local ULIRGs with the result that they are predominantly starburst powered, although they also show signs of AGN activity (e.g., Genzel et al. 1998; Lutz et al. 1999; Rigopoulou et al. 1999; Sturm et al. 2000; Tran et al. 2001). In particular, *ISO* studies of local ULIRGs showed that at low luminosities the ULIRGs are powered mainly by a starburst, but at the very highest luminosities ($L_{\text{IR}} \gtrsim 3 \times 10^{12} L_{\odot}$) the dominant emission source is the AGN (Lutz et al. 1998; Tran et al. 2001). From this one might expect that SMGs, with their extreme IR luminosities ($L_{\text{IR}} \sim 6 \times 10^{12} L_{\odot}$; Pope et al. 2006), are primarily AGN dominated.

The *Spitzer* Infrared Spectrograph (IRS; Houck et al. 2004), with its sensitivity and spectral resolution, can extend this effort outside the local universe, allowing much larger samples to be observed to much fainter flux limits. The IRS can detect emission features and continuum in ULIRGs out to $z \sim 4$ (Valiante et al. 2007). The three mid-IR spectral components listed above have been fitted to the *Spitzer* IRS spectra of ULIRGs at various redshifts to determine the AGN contribution to the mid-IR luminosity (e.g., Sajina et al. 2007). Although X-ray imaging is very effective at identifying the presence of an AGN in a galaxy, it is subject to obscuration in extreme, Compton-thick sources. The X-ray luminosity provides a direct measurement of the power of the AGN at X-ray energies, but it is necessary to make assumptions about the geometry of the obscuring dust to determine the bolometric luminosity of the AGN (e.g., Alexander et al. 2005). Furthermore, AGN indicators at UV and optical wavelengths can be very obscured in dusty SMGs (Swinbank et al. 2004; Chapman et al. 2005). Mid-IR spectroscopy, by directly detecting the thermally reradiated emission from the obscuring dust, provides one of the most effective ways to identify AGNs, including Compton-thick sources, and measure their contribution to the IR luminosity (e.g., Lutz et al. 2004).

Two previous studies have presented IRS spectroscopy of SMGs. Menéndez-Delmestre et al. (2007) presented IRS spectra of five radio-detected, spectroscopically confirmed SMGs. They found that four SMGs with lower redshifts ($z < 1.5$) showed PAH emission and that a composite spectrum of these four sources fits well to a scaled M82 spectrum plus a power-law component. They conclude that these systems host both star formation and AGN activity. Their fifth source at $z = 2.4$ was found to have a more AGN-type spectrum. Valiante et al. (2007) discussed IRS spectra of 13 bright SMGs, 9 of which are detected with the IRS (the first two spectra from this sample were presented in Lutz et al. 2005a). This sample was more representative of the redshift distribution of SMGs. They also found that SMGs are likely to be mainly starburst powered. Sajina et al. (2007) used IRS spectra to study a sample of *Spitzer* 24 μm -selected galaxies at $z \sim 2$. This sample was selected to have $S_{24 \mu\text{m}} > 900 \mu\text{Jy}$ and very red 24 μm -to-8 μm and 24 μm -to- R colors (Yan et al. 2007). A total of 75% of this sample consists of continuum-dominated sources, with some weak PAH emission indicative of star formation activity. They conclude that these sources display mid-IR spectra similar to those of the most luminous local ULIRGs. Follow-up studies

of these galaxies have shown that they are not generally (sub)-millimeter-detected sources ($S_{1.2 \text{ mm}} \lesssim 1.5 \text{ mJy}$ for 33/40 sources; Lutz et al. 2005b).

In this paper we present *Spitzer* IRS spectra of a sample of SMGs in GOODS-N that have extensive multiwavelength information from X-ray to radio wavelengths. This allows us to produce full SEDs and to calculate the bolometric luminosity. Our *Spitzer* IRS observations go substantially deeper ($S_{24 \mu\text{m}} > 200 \mu\text{Jy}$) than previous work. This enables a study of the typical blank-field submillimeter galaxy population that has $S_{850 \mu\text{m}} > 2 \text{ mJy}$. The main goal of our program is to decompose the mid-IR spectra into star formation and AGN components, in order to infer how much of the total IR luminosity is coming from each. From the mid-IR spectroscopy we are also able to obtain spectroscopic redshifts and measure the luminosities and equivalent widths of individual PAH lines to investigate various line diagnostics. We compare our measurements with line strengths in other star-forming galaxies to put constraints on the evolution of SMGs and the relationship between star formation and black hole growth.

This paper is formatted as follows. Our *Spitzer* observations are described in § 2, and the data analysis is explained in § 3. Section 4 presents the main results of the IRS spectroscopy including redshifts, spectral decomposition, and AGN classification. In § 5 we fit the full SEDs of these SMGs to determine their total IR luminosities. We explore correlations between the PAH and IR luminosities in § 6. We discuss our results in the context of the role that SMGs play in galaxy evolution in § 7. Finally, our conclusions are listed in § 8. We include two appendices that discuss the noise properties of our *Spitzer* IRS spectra and include notes on individual sources.

All magnitudes in this paper use the AB system. We use a standard cosmology with $H_0 = 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$, and $\Omega_{\Lambda} = 0.7$.

2. OBSERVATIONS

The *Spitzer* IRS observations presented here were taken as part of *Spitzer* program GO-20456 (PI: R. Chary). The full sample consists of high-redshift SMGs, AGNs, and optically faint 24 μm sources in the Great Observatories Origins Deep Survey—North (GOODS-N; Dickinson et al. 2003) field. The GOODS-N field is one of the most extensively studied regions of the sky, with deep data existing across all wave bands, including *Chandra* 2 Ms X-ray observations (Alexander et al. 2003); deep *Hubble Space Telescope* (*HST*) optical imaging using the Advanced Camera for Surveys (ACS) in four bands, B_{435} , V_{606} , i_{775} , and z_{850} (Giavalisco et al. 2004); and deep *Spitzer* imaging at five IR wavelengths (M. Dickinson et al. 2008, in preparation). More recently, GOODS-N has been surveyed with IRS peak-up observations at 16 μm (Teplitz et al. 2005) and MIPS imaging at 70 μm (Frayer et al. 2006). In addition to the extensive space-based imaging campaign, many ground-based programs have also targeted this field for imaging and spectroscopy, making it one of the best fields for investigating the role of AGNs and star formation in high-redshift galaxies. Here we present the IRS spectroscopy for the SMGs from this project. Results from the other sources will be presented in other papers.

Our sample of SMGs was primarily selected from sources detected in the GOODS-N SCUBA “supermap” with secure multi-wavelength counterparts (Borys et al. 2003; Pope et al. 2006; Pope 2007). We imposed a 24 μm flux cut of $S_{24} > 200 \mu\text{Jy}$ in order to ensure that we achieve adequate signal-to-noise ratio (S/N) on the continuum in the IRS spectra. Roughly half of the secure counterparts in Pope et al. (2006) meet this 24 μm flux cut, which yields a sample of 10 IRS targets (after removing sources that are

TABLE 1
COORDINATES AND FLUX DENSITIES OF IRS TARGETS

SMG Name	SMG ID	R.A. (J2000.0)	Decl. (J2000.0)	S_{16} (mJy)	S_{24} (mJy)	S_{70} (mJy)
SMM J123653.1+621120	GN31	12 36 53.22	62 11 16.7	0.301 ± 0.007	0.367 ± 0.006	<3.5
SMM J123635.5+621238	GN26	12 36 34.51	62 12 40.9	0.992 ± 0.012	0.446 ± 0.005	13.9 ± 1.8
SMM J123701.2+621147	GN17	12 37 01.59	62 11 46.2	0.209 ± 0.006	0.710 ± 0.008	<3.5
SMM J123622.6+621629	C2	12 36 22.66	62 16 29.5	0.098 ± 0.006	0.414 ± 0.007	<3.5
SMM J123555.1+620901	C3	12 35 55.13	62 09 01.6	...	0.374 ± 0.009	<3.5
SMM J123711.1+621325	GN39a	12 37 11.37	62 13 31.1	0.097 ± 0.005	0.537 ± 0.009	<3.5
SMM J123711.9+621331	GN39b	12 37 11.97	62 13 25.8	0.051 ± 0.005	0.225 ± 0.007	<3.5
SMM J123621.3+621711	GN07	12 36 21.27	62 17 08.1	0.048 ± 0.007	0.370 ± 0.011	<3.5
SMM J123618.7+621553	GN06	12 36 18.33	62 15 50.4	0.036 ± 0.006	0.330 ± 0.008	<3.5
SMM J123600.2+621047	C1	12 36 00.16	62 10 47.3	0.478 ± 0.011	1.220 ± 0.008	5.5 ± 1.2
SMM J123618.8+621008	GN05	12 36 19.13	62 10 04.3	0.044 ± 0.007	0.215 ± 0.006	<3.5
SMM J123707.7+621411	GN19	12 37 07.19	62 14 08.0	<0.02	0.280 ± 0.015	<3.5
SMM J123616.6+621520	GN04	12 36 16.11	62 15 13.5	0.060 ± 0.006	0.303 ± 0.007	<3.5

NOTE.—Units of right ascension are hours, minutes, and seconds, and units of declination are degrees, arcminutes, and arcseconds.

already observed with the IRS in other programs). To expand our sample slightly, we also included three SCUBA sources from Chapman et al. (2005). These three sources are not detected at $>3.5\sigma$ in the supermap, but they are detected in SCUBA photometry observations with a lower statistical significance.

Our 13 SMG IRS targets have 850 and 24 μm flux densities of 2–10 and 200–1200 μJy , respectively. This sample is expected to be representative of the submillimeter population discovered in existing blank-field 850 μm imaging. There is no evidence supporting the idea that SMGs that are fainter than 200 μJy at 24 μm are fundamentally different besides lying at slightly higher redshifts (e.g., Pope et al. 2006). We list the positions and mid-IR fluxes for these 13 SMGs in Table 1. SMG IDs starting with “GN” and “C” are from the SCUBA supermap (Pope et al. 2006; Pope 2007) and Chapman et al. (2005), respectively, and the 850 μm fluxes can be found in these papers. The majority of the SMGs are undetected in the deep *Spitzer* 70 μm images (see also Huynh et al. 2007). A total of 11/13 SMGs have optical spectroscopic redshifts, while the other 2 have photometric redshifts. The redshift distribution of our sample is consistent with that of all SMGs (Chapman et al. 2005; Pope et al. 2006). Figure 1 shows multi-wavelength images, which span roughly the width of the IRS slit, of our targets. In some cases there are several optical or IRAC sources in the aperture. For GN04, GN07, and GN19, the two IRAC sources are both expected to be associated with the submillimeter emission since both components are at the same redshift (see Pope et al. 2006). The IRS spectra of these three SMGs will have contributions from both components. For the remaining sources, the 8, 16, and 24 μm images show that only our primary targets are producing the mid-IR emission that we will detect with the IRS.

The *Spitzer* IRS observations are taken in spectral staring mode, which observed the target at two nod positions within the slit. We observed in low resolution ($R = \lambda/\Delta\lambda \sim 100$) using the Short-Low 1 (SL1; 7.4–14.5 μm), Long-Low 2 (LL2; 14.0–21.3 μm), and/or Long-Low 1 (LL1; 19.5–38.0 μm) orders, depending on the redshift of the source. We chose the wavelength coverage in order to include one or more of the main PAH features at 6.2, 7.7, and 11.3 μm . Integration times varied depending on the flux of the target at 24 μm . We used blue peak-up acquisition on nearby, isolated, bright Two Micron All Sky Survey (2MASS) stars. Our observations were taken in 2006 April and May.

Table 2 shows our list of targets and integration times.

3. DATA ANALYSIS

We started with the S14.0.0 *Spitzer* IRS pipeline data,⁸ which produce two-dimensional (2D) basic calibrated data (BCD) files. Note that our targets are not resolved with *Spitzer* at these wavelengths and therefore can be treated as point sources.

At mid-IR wavelengths, the dominant source of noise is the sky background, mostly from zodiacal light. Since we have long integrations of very faint targets, additional data reduction steps are required to alleviate the detector and sky noise. The first step is to identify and clean the rogue pixels. There are a number of hot pixels in the IRS arrays ($\sim 15\%$). Rogue pixel masks were created from the campaign masks and from examining each 2D spectral image individually. We used IRSCLEAN⁹ to replace the values of the hot pixels by extrapolating from the surrounding pixels. Note that IRSCLEAN fails when there is a group of several hot pixels together, so we made sure to exclude these from the spectral extraction. The fraction of pixels in the 2D files that remain unusable after IRSCLEAN is only $\sim 2\%$.

The next step is to remove the latent charge buildup on the array. In long IRS integrations, it has been found that latent charge builds up with time, despite the detectors being reset at the end of each integration (see Teplitz et al. 2007). This buildup is different depending on the wavelength (y position on the 2D spectral image).¹⁰ We found charge buildup on the arrays for integrations of more than 1 hr in all LL observations. For the SL1 observations, we only saw the latent charge buildup for a few targets, even though all of these observations were greater than 1 hr. The latent charge buildup was fitted by a polynomial and the temporal component subtracted off.

We explore several methods for removing the sky background, including (1) subtracting the average spectrum at each nod position, (2) subtracting individual observations from opposite nod positions and then averaging, (3) subtracting a “supersky” created from all files from that astronomical observation request (AOR), and (4) subtracting a supersky created from all our data taken in a given campaign. In all methods, we mask out all sources that fall within the IRS slit identified from the deep 24 μm

⁸ See <http://ssc.spitzer.caltech.edu/irs/dh/>.

⁹ See http://ssc.spitzer.caltech.edu/archanalyst/contributed/irsclean/IRSCLEAN_MASK.html.

¹⁰ See http://ssc.spitzer.caltech.edu/irs/documents/irs_ultra-deep-memo.pdf.




FIG. 1.—Images of SMG targets, in order of increasing redshift. Images are $10'' \times 10''$ (roughly the size of the aperture of the IRS observations) at B , i , and 3.6, 8.0, 16, and 24 μm from left to right. Data are not available in some of these wave bands for source C3 since it is outside the *HST* GOODS-N coverage. GN04, GN07, GN19, GN39a, and GN39b have two sources within the slit and that contribute flux in the IRS spectra (see text for details). For the other sources, the 8, 16, and 24 μm images show that only the primary targets are bright in the mid-IR.

imaging of the GOODS-N field. The supersky subtraction was carried out with and without normalizing to the median in each image. The third method of subtracting a normalized supersky created from all observations from that AOR gave the lowest residual sky noise (and thus highest S/N) in the final spectrum; therefore, this is what we used to remove the sky from all our observations.

After the sky was removed from the individual data files, we then co-added them using a clipped median. The co-adds were checked again for persisting hot pixels and cleaned once more using IRSCLEAN. These steps produced two 2D spectra for each target, one for each nod position.

We used the *Spitzer* IRS Custom Extraction (SPICE¹¹) to extract the one-dimensional (1D) spectra at each nod position. We used a narrow extraction window of 2 pixels ($\sim 10''$ for LL observations), which was constant as a function of wavelength. We found that this produced higher S/N spectra than using an extraction window of increasing width, starting with 2 pixels at the blue end. Using the optimal extraction option in SPICE, we found comparable results to the 2 pixel, fixed width, extraction. For each science spectrum, we also extracted a residual sky spectrum, offset from any sources, to provide a level of the error in our final

¹¹ See <http://ssc.spitzer.caltech.edu/postbcd/spice.html>.




FIG. 1—Continued

TABLE 2
Spitzer IRS OBSERVATIONS

SMG ID	INTEGRATION TIME			Total (hr)
	SL1 ($\times 240$ s)	LL2 ($\times 120$ s)	LL1 ($\times 120$ s)	
GN31.....	...	35×2	...	2.3
GN26.....	15×2	20×2	$[12 \times 2]$	3.3 [4.1]
GN17.....	20×2	12×2	...	3.5
C2.....	45×2	3.0
C3.....	...	31×2	40×2	4.7
GN39a.....	...	20×2	28×2	3.2
GN39b.....	...	$[20 \times 2]^a$	$[28 \times 2]$	[3.2]
GN07.....	...	35×2	45×2	5.3
GN06.....	...	35×2	55×2	6.0
C1.....	...	6×2	10×2	1.1
GN05.....	55×2	3.7
GN19.....	...	$[20 \times 2]$	$[28 \times 2]$	[3.2]
GN04.....	70×2	4.7

^a Sources with time in square brackets are secondary sources observed in the slits of the primary targets.

spectrum as a function of wavelength. The uncertainty is calculated as the standard deviation of 10 pixels surrounding each wavelength in the residual sky spectrum. For each source, the 1D spectra at each nod position were then averaged together to produce the final spectrum for each source. Further details about the final noise in our IRS spectra are given in Appendix A.

In order to calibrate the final IRS spectra, we reduced standard star calibrator observations from the same campaign as our observations, in exactly the same way that we reduced the science observations. We then extracted the spectrum for the calibrator using our 2 pixel width and the default SPICE width in order to determine the calibration factor as a function of wavelength. This calibration was verified against the broadband photometry at 8, 16, and $24 \mu\text{m}$, and we found very good agreement (see Fig. 2).

3.1. Spectral Line Measurements

In order to measure the individual PAH line luminosities, we used the IDEA (ISAP¹²) environment in the SMART¹³ software

¹² The ISO Spectral Analysis Package (ISAP) is a joint development by the LWS and SWS Instrument Teams and Data Centers. Contributing institutes are CESR, IAS, IPAC, MPE, RAL, and SRON.

¹³ SMART was developed by the IRS Team at Cornell University and is available through the *Spitzer* Science Center at Caltech.




FIG. 2.—Calibration of IRS spectra compared to the MIPS and peak-up photometry. MIPS photometry is from the GOODS 24 μm imaging (R. Chary et al. 2008, in preparation), and the 16 μm peak-up coverage is from Teplitz et al. (2005). Most of the 16 μm IRS fluxes are systematically higher because roughly half of the 16 μm filter falls outside of the LL2 IRS coverage and so the IRS flux is overestimated, since the spectrum actually drops off at lower wavelengths. These sources are indicated as the open circles. For sources with SL1 and LL2 observations, there should be no such discrepancy, which we confirm. The filled circle in the right panel that is significantly above the line is GN39b, whose IRS spectrum is known to contain a contribution ($\frac{1}{5}$ of the flux) from GN39a.

package (Higdon et al. 2004). For each of the four main PAH lines (6.2, 7.7, 8.6, and 11.3 μm), we isolated a region around each line and simultaneously fitted the line and continuum. Both the center and the width of the line were allowed to vary. All measured line widths were consistent within the errors with the line widths measured in IRS spectra of local star-forming galaxies (Smith et al. 2007). Note that the errors we calculated from the residual sky were propagated in the fitting of individual lines. All SMG line measurements used in this paper, including 3 σ upper limits when the line is undetected in the IRS spectrum, are listed in Table 5 below.

The biggest uncertainty in comparing PAH line luminosities from different samples comes from differences in how the spectral lines are measured, or more specifically, how the continuum is estimated for each line. This can lead to variations in the line flux and equivalent width of up to a factor of 4 (see Sajina et al. 2007; Smith et al. 2007). Our method of measuring the line strengths gives similar results to the cubic spline continuum fits performed in Brandl et al. (2006; see next paragraph). This method is known to underestimate the flux of the PAH lines (particularly the 7.7 μm line) compared to methods that fit all PAH lines simultaneously (e.g., Sajina et al. 2007; Smith et al. 2007) since the continuum we adopt is higher. Therefore, we took care to measure the PAH lines in any comparison samples in the exact same way as we did for the SMGs.

In this paper we compare the SMG PAH line luminosities to those of other galaxy populations. Brandl et al. (2006) presented IRS spectra of 22 local starburst galaxies with IR luminosities ranging from 5×10^9 to $5 \times 10^{11} L_{\odot}$. These galaxies are known to have little or no AGN contributing to their IR luminosity. Because the choice for estimating the continuum under the PAH lines makes a significant difference to the results, we have measured the lines for the starburst galaxies in exactly the same way as we did for the SMGs. Another interesting comparison sample

is the local ULIRG galaxies. Armus et al. (2007) presented IRS spectroscopy of 10 ULIRGs from the Bright Galaxy Sample (BGS; $S_{60} > 5.4$ mJy; Soifer et al. 1987), and the IRS spectra for a sample of over 100 ULIRGs were recently presented in Desai et al. (2007) and Farrah et al. (2007). The local ULIRGs show a broad range of mid-IR properties and a mix of AGN and starburst mid-IR emission. However, we did not have access to the IRS spectra for the full sample of local ULIRGs in order to measure the line luminosities using our method; therefore, we could not include them in this paper.

We measured the PAH lines in 10 of the Brandl et al. (2006) starburst galaxies to verify that our method is consistent with their measurements. On average the luminosities of the 6.2 and 11.3 μm lines we measure are consistent with the values listed in Brandl et al. (2006), but the 7.7 μm line luminosity is 1.5 times larger on average from our measurements. This is probably because we are using a straight line for the continuum as opposed to a spline. We have corrected the measurements of the 7.7 μm line luminosity for the Brandl et al. (2006) starbursts for this factor of 1.5 for all plots in this paper. For the Brandl et al. (2006) starbursts we also applied a correction to the line measurements to account for the fractional flux found within the IRS apertures (see Table 2 of Brandl et al. 2006) when comparing to the unresolved SMGs. The error on the line fluxes for the Brandl et al. (2006) starbursts is dominated by the calibration error, which is on the order of 10% (less than the size of the symbols in most of the plots in this paper).

4. RESULTS

In this section we use the mid-IR spectra to explore the nature of SMGs. We start by using the identification of PAH lines to estimate redshifts. We then proceed to decompose the mid-IR spectra into starburst and AGN components, starting first with a composite SMG spectrum since the S/N is higher and then




FIG. 3.—IRS spectra of SMGs. The solid black lines are the raw (unsmoothed) IRS spectra, while the shaded region is the associated 1σ noise from the sky background. The red dashed lines show the best-fit SED, which is made up of an extincted power-law component (*blue dashed line*; consistent with zero in most cases) and an extincted starburst component (*green dashed line*). Recall that the extinction curve is not monotonic in wavelength and contains various features, the most notable being the $9.7\ \mu\text{m}$ silicate feature.

proceeding to decompose each individual SMG spectrum. We then use the information from the spectral decomposition to classify the fractional contribution of AGNs to the mid-IR luminosity and compare this with other AGN indicators such as optical spectroscopy and X-ray emission. These results are then used to guide the fitting of the full IR SED to constrain the total IR luminosity in § 5.

Figure 3 shows the rest-frame (unsmoothed) IRS spectra for the 13 SMGs. PAH emission lines are present in all sources. Despite the relatively low S/N in the spectra, all SMGs seem to show very similar mid-IR spectral shapes with the exception of C1.

While this SMG clearly shows PAH emission, it is superimposed on a steeply rising continuum. We discuss this source further in Appendix B.

4.1. Redshifts

Given the presence of PAH features in all of our sources, we are able to extract redshifts from the IRS spectra. In order to calculate the redshift and its associated error from the IRS data, we follow the following steps. We assumed the following rest-frame wavelengths for the main PAH features: 6.22 , 7.71 , 8.61 , and $11.27\ \mu\text{m}$ (Draine & Li 2007). For each PAH line, we find the




FIG. 3—Continued

center of the Gaussian fit to the line from SMART and calculate the redshift. The redshift for each source is calculated by averaging the redshift estimates from each PAH line. The total error in the redshift is the quadratic sum of the deviation due to the spread in the redshift from the different PAH lines and the centroid uncertainty. For sources whose spectra only contain one PAH line, the error is just the centroid uncertainty of that line. The deviation in the redshift from each line is calculated as $1\sigma = [\sum_i (z_i - \langle z \rangle)^2 / (n - 1)]^{1/2}$. We verified the centroid uncertainties by performing Monte Carlo simulations. We take a Gaussian of the same height and width as each PAH feature and run 1000 trials where we add noise that has the same standard deviation as the noise in the real data. For each trial, we fit the data to get the line center and keep track of the difference between the input line

center and the measured line center. The centroid error is estimated from the standard deviation of this difference for all 1000 trials. We repeat this for each source. With the exception of GN26, whose spectrum has a very high S/N, the centroid uncertainties for individual PAH features range from 0.08 to 0.25 μm , which corresponds to 0.5–1.5 pixels. This is roughly an order of magnitude higher than what we would expect from the instrument based on the pixel size in the case of no noise. The total error in the IRS redshift ranges from 0.01 to 0.05 for these SMGs.

Table 3 lists the new IRS redshifts and the previous optical, spectroscopic or photometric, redshifts. The new IRS mid-IR redshifts are compared with the optical redshifts in Figure 4. Two of our 13 sources seem to have very different redshifts as determined from IRS and optical spectroscopy (C2 and GN06, which we




FIG. 3—Continued

TABLE 3
MEASURED IRS REDSHIFTS COMPARED TO OPTICAL SPECTROSCOPIC AND PHOTOMETRIC REDSHIFTS

SMG ID	i mag (AB) ^a	z_{optical}^b	$z_{\text{IRS}} \pm 1 \sigma$	Comment
GN31.....	21.8	0.935 ^c	0.93 \pm 0.03	
GN26.....	22.7	1.219 ^d	1.23 \pm 0.01	
GN17.....	27.7	1.72 ^e	1.73 \pm 0.01	New IRS redshift
C2.....	25.5	2.466 ^{d,f}	1.79 \pm 0.04	Inconsistent
C3.....	23.5 ^g	1.875 ^d	1.88 \pm 0.02	
GN39a.....	25.9	1.996 ^f	1.98 \pm 0.01	
GN39b.....	26.4	1.992 ^d	1.99 \pm 0.04	
GN07.....	27.8/23.9 ^h	1.988, ^d 1.992 ^f	1.99 \pm 0.02	
GN06.....	27.8	1.865 ^d	2.00 \pm 0.03	Inconsistent
C1.....	24.8	1.994, ^d 2.002 ^f	2.01 \pm 0.05	
GN05.....	24.9	2.60 ^e	2.21 \pm 0.03	New IRS redshift
GN19.....	25.4/>28 ^h	2.484, ^d 2.490 ^f	2.48 \pm 0.03	
GN04.....	26.2/>28 ^h	2.578 ^d	2.55 \pm 0.01	

^a The i magnitudes are from the GOODS ACS observations unless otherwise noted.

^b Optical redshifts are spectroscopic unless otherwise noted.

^c Cowie et al. (2004).

^d Chapman et al. (2005).

^e Pope et al. (2006), photometric redshift.

^f Swinbank et al. (2004).

^g Capak et al. (2004).

^h These SMGs have two counterparts within the IRS aperture.

discuss in Appendix B). Since the mid-IR and submillimeter wavelengths both trace dust emission, we believe that the IRS redshifts presented here are correct for the galaxies that are causing the submillimeter emission.

4.2. Spectral Decomposition of SMG Composite Spectrum

We can study the ensemble properties of the SMG sample by averaging together the individual mid-IR spectra. When plotted




FIG. 4.—Comparison of IRS spectroscopic redshifts and previous optical redshifts. All optical redshifts are spectroscopic except for the two sources indicated by open symbols. There appear to be two optical spectroscopic redshifts that disagree with our new IRS redshifts. In these cases we believe that the IRS redshift is correct for the SMG, since the mid-IR emission is more likely to trace the submillimeter emission than the optical/UV emission.

in rest-frame luminosity units, the spectra for all SMGs in this sample show very little dynamic range. Therefore, we are able to average the spectra without having to normalize each at a specific wavelength, which can introduce additional biases in the composite spectrum. We exclude C1 from the composite spectrum, since it shows a very different spectral shape from all the other SMGs (see Appendix B). The error on the composite spectrum is calculated from combining the errors on the individual flux measurements. We restrict the final SMG composite spectrum to areas where there are more than three data files present at that wavelength, which results in a wavelength coverage of $\sim 5\text{--}12 \mu\text{m}$ in the rest frame. Figure 5 shows our composite spectrum of 12 SMGs calculated from averaging the individual spectra (excluding C1).

We detect the 6.2, 7.7, and 11.3 μm emission lines, with a hint of the 8.6 μm emission line in the composite spectrum. The composite spectrum also shows a very flat underlying continuum. In order to quantify the fractional contribution of star formation and AGN activity to the emission at these wavelengths, we fit our composite spectrum with a model containing the three components discussed in § 1. The PAH emission is fitted using two different templates: (1) the mid-IR spectrum of M82 (Förster Schreiber et al. 2003) and (2) the starburst composite template from Brandl et al. (2006). M82 is a prototypical star-forming galaxy, and its mid-IR spectrum is expected to be dominated by emission from star-forming regions. The starburst template from Brandl et al. (2006) is a composite of 13 local starburst galaxies with bright flux densities and without a strong AGN component. As we show below, these templates are composed almost entirely of the sum of the individual PAH features. The AGN emission is characterized by a power law with both the normalization and slope as free parameters. For the extinction, we obtain τ_ν from the Draine (2003) extinction curves and applied it to both the PAH and continuum components separately. The extinction curve is not just monotonic in wavelength and contains silicate absorption features, the most notable being at 9.7 μm . Our model, F_ν , can be expressed as

$$F_\nu = c_1 \nu^{-c_2} e^{-c_3 \tau_\nu} + c_4 f_\nu e^{-c_5 \tau_\nu}, \quad (1)$$




FIG. 5.—Composite IRS spectrum of 12 SMGs (excluding C1). This SMG composite spectrum (*black line*) has been smoothed by a Gaussian with an FWHM of $0.12 \mu\text{m}$ (rest frame), which corresponds to the average resolution of the IRS. The shaded region shows the 1σ uncertainty. *Left*: The red line is the best-fit model (eq. [1]), which is composed of a PAH template (*green line*) and a power-law continuum component (*blue line*), each with extinction applied to them. *Right*: The composite is fitted using individual PAH line profiles and a continuum, each with extinction. In either case, we find that the PAH emission dominates the mid-IR luminosity.

where f_ν is the PAH template. We performed a χ^2 minimization fit for each of the two PAH templates and solved simultaneously for c_1 , c_2 , c_3 , c_4 , and c_5 .

The best-fit model, F_ν , is shown in the left panel of Figure 5 as the red dashed line. The blue dot-dashed and green dotted lines show the individual contributions from the continuum and PAH components, respectively. The best-fit PAH template was the starburst composite from Brandl et al. (2006) although M82 gave very similar results. With this PAH template, the extinction needed is $\tau_{9.7} \sim 1$. The slope of the power law is very shallow ($c_2 \sim 0.2$). The continuum component accounts for 30% of the luminosity at these wavelengths ($5\text{--}11.5 \mu\text{m}$). This implies that the mid-IR luminosity of SMGs is dominated by star formation with a maximum mid-IR AGN contribution of 30%.

To be clear about the origin of the emission in the PAH template, we also fit the SMG composite spectrum to individual PAH line profiles and a continuum with extinction. We include the PAH lines at 5.70 , 6.22 , 7.42 , 7.60 , 7.85 , 8.33 , 8.61 , 11.23 , and $11.33 \mu\text{m}$, each modeled as a Drude profile (e.g., Draine & Li 2007) where the height of each line is treated as a free parameter. The resulting best-fit model shown in the right panel of Figure 5 is very similar to the best-fit model that includes a PAH template and continuum (*left panel*).

In Figure 6 we show the SMG IRS composite spectrum compared to the IRS spectra of $z \sim 2$ *Spitzer* $24 \mu\text{m}$ -selected ULIRGs from Sajina et al. (2007) and the local ULIRG Arp 220. While Arp 220 is often considered the typical local analog to high-redshift SMGs, it appears to have more silicate absorption and less 6.2 and $11.3 \mu\text{m}$ PAH emission than the average SMG. Recall that the sample in Sajina et al. (2007) consists of red $24 \mu\text{m}$ -selected galaxies with $S_{24} > 0.9$ mJy, which is much brighter than most of the SMG population. The majority of the ULIRGs in the *Spitzer*-selected sample have weak PAH emission and are AGN dominated; the green and red lines in Figure 6 are composite spectra of the weak-PAH (i.e., AGN dominated) *Spitzer* $24 \mu\text{m}$ -selected ULIRGs with $\tau_{9.7} > 1$ and $\tau_{9.7} < 1$, respectively. The




FIG. 6.—SMG composite spectrum compared to those other ULIRGs. The SMG composite spectrum is the black line, and the shaded region shows the 1σ error. The colored histograms are high-redshift *Spitzer* $24 \mu\text{m}$ -selected ULIRGs observed with the IRS from Sajina et al. (2007; S07); the green and red lines are composite spectra of the weak-PAH sources for $\tau_{9.7} > 1$ and $\tau_{9.7} < 1$, respectively, and the blue line is the composite spectrum for the strong-PAH sources. The light blue line is the local ULIRG Arp 220 (Charmandaris et al. 1999; Fischer et al. 1999). All lines have been normalized at $\sim 7 \mu\text{m}$. The numbers in the legend in parentheses indicate the number of sources in each composite spectrum.

TABLE 4

CLASSIFICATION OF MID-IR SPECTRA COMPARED WITH CLASSIFICATIONS BASED ON X-RAY IMAGING AND UV/OPTICAL SPECTROSCOPY

SMG ID	CLASSIFICATION ^a			MID-IR CONTINUUM ^d
	X-Ray ^b	UV/Op ^c	Mid-IR	
GN31.....	U	...	SB	<45
GN26.....	SB	SB	SB	<15
GN17.....	SB	...	SB	<29
C2.....	AGN	SB	SB	<34
C3.....	AGN	SB	SB+AGN	48
GN39a.....	oAGN	SB	SB	<35
GN39b.....	oAGN	SB	SB	<10
GN07.....	SB	SB	SB+weak AGN	18
GN06.....	SB	SB	SB+AGN	47
C1.....	SB	SB	AGN+weak SB	82
GN05.....	U	...	SB	<7
GN19.....	oAGN	SB	SB	<14
GN04.....	AGN	SB	AGN+SB	61

^a oAGN=obscured AGN, SB=starburst, U=undetected.

^b Classified as AGN if detected in the hard X-ray band.

^c Classification from Chapman et al. (2005) and/or Swinbank et al. (2004).

^d Percentage of mid-IR flux that comes from continuum emission.

blue line is the composite spectrum for the strong-PAH *Spitzer* 24 μm -selected ULIRGs and is very similar to the SMG composite. In particular, the relative line strengths are consistent. The strong-PAH *Spitzer*-selected sources represent only a small fraction of the Sajina et al. (2007) sample, and these sources tend to have a higher detection rate in the (sub)millimeter than the rest of their sample (Sajina et al. 2008). From Yan et al. (2007) and Sajina et al. (2007), the surface density of the $z \sim 2$ *Spitzer* 24 μm -selected ULIRGs is 10.4 deg^{-2} , and the strong-PAH subsample of these sources has a surface density of 2.6 deg^{-2} . The surface density of SMGs with $S_{850} > 4 \text{ mJy}$ (i.e., $L_{\text{IR}} \gtrsim 5 \times 10^{12} L_{\odot}$; Pope et al. 2006) is 844 deg^{-2} (Coppin et al. 2006). While most submillimeter-selected galaxies are much fainter at 24 μm , the strong-PAH *Spitzer* 24 μm -selected ULIRGs appear to be a very small subsample of the SMG population.

4.3. Spectral Decomposition of Individual Galaxies

Given the good agreement between our SMG composite spectrum and the spectrum of starburst galaxies, we proceed to fit the individual SMG mid-IR spectra to the model given in equation (1). For each of the 13 SMGs we solve for the five parameters in equation (1) and plot the best-fit model in Figure 3 as the red dashed line. The separate starburst and AGN components including extinction are shown as the green and blue lines. For most SMGs, the continuum component is zero and not plotted in Figure 3. We note that the spectrum of GN26 is even lower than the model at long wavelengths, which confirms that the AGN component must be negligible in this source.

From our model fits, we derived the fraction of the mid-IR luminosity that comes from the continuum component. The continuum component provides an upper limit to the AGN contribution since hot dust present in the most energetic H II regions can also contribute to the continuum. When the best-fit model does not include a continuum component, we calculate an upper limit by conservatively assuming that all of the emission around 6 μm is coming from a continuum component. We used this fraction to classify the mid-IR spectrum, and the results are listed in Table 4. Eight of the 13 SMGs are clearly starburst-dominated sources (continuum component $\lesssim 50\%$ of the mid-IR luminosity),

and only 2/13 SMGs are AGN dominated (continuum component $\gtrsim 50\%$ of the mid-IR luminosity) in the mid-IR. The remaining sources have a combination of PAH and continuum emission in the mid-IR. The results for the individual galaxies are in agreement with the composite spectrum, which showed a 30% AGN contribution on average.

4.4. AGN Classification

Alexander et al. (2005) showed that deep X-ray observations are a powerful tool for identifying the presence of AGNs in SMGs. They conclude that the majority of SMGs host an AGN, but that the bolometric output from SMGs is dominated by star formation because of the low ratio of X-ray luminosity to IR luminosity. They also stress that some fraction of SMGs will be AGN dominated. Following the same method as Alexander et al. (2005), we list the X-ray classification in Table 4, where a hard X-ray detection tells us that the X-ray emission is dominated by AGN activity.

We find that the X-ray classification gives a higher fraction of AGN sources (6/13) than the mid-IR spectra (2/13) for SMGs with $S_{24} > 200 \mu\text{Jy}$. For the X-ray-classified AGN sources, the X-ray emission expected from star formation (e.g., Bauer et al. 2002) is less than that observed, indicating that there is an AGN present. Our IRS observations show that this AGN is not important to the mid-IR luminosity. On the other hand, the line widths and line ratios in the UV/optical spectra more often indicate that these SMGs are starburst sources (Swinbank et al. 2004; Chapman et al. 2005) and appear to be insensitive to the AGN activity in many sources.

C1 is classified as a starburst from the X-ray data but shows a very strong AGN in the mid-IR. This SMG is likely to host a very obscured AGN, the kind of system that can help explain the unresolved portion of the hard X-ray background (e.g., Brandt & Hasinger 2005; Daddi et al. 2007). See Appendix B for more discussion about C1. Combining the X-ray imaging with mid-IR spectroscopy provides the best census of AGNs in SMGs.

In Pope et al. (2006), we found that most SMGs showed evidence for a stellar bump in the four IRAC channels. We found only four sources that had a good fit to a simple power law from IRAC+24 μm photometry. Only one of these is in our IRS sample, GN04, since the others are too faint for spectroscopy at 24 μm . We find that the IRAC SED, the mid-IR spectrum, and X-ray classification all converge on the conclusion that there is an energetically important AGN in GN04. In contrast, the UV/optical spectra of this SMG did not reveal the presence of an AGN in it.

Another approach for separating sources that contain a mid-IR AGN component is through the use of *Spitzer* color-color diagrams. Ivison et al. (2004) proposed that the $S_8/S_{4.5}$ versus S_{24}/S_8 color plane could be used to identify AGN-dominated SMGs. The IRS spectra have allowed us to decompose the starburst and AGN components in SMGs; therefore, we are in a position to test the plot as a diagnostic. In Figure 7 we plot this *Spitzer* color-color diagram. The dashed and dotted lines show the positions of Mrk 231 (mid-IR spectrum from Rigopoulou et al. [1999] spliced with a fit to the *Infrared Astronomical Satellite* [IRAS] photometry) and M82 (Förster Schreiber et al. 2003), respectively, as a function of redshift (redshift is indicated by the numbers along the track). Based on these broadband colors, AGN-dominated sources are expected to lie along the Mrk 231 tracks, while starburst-dominated systems should have similar colors to M82. Sajina et al. (2005) showed that while AGN-dominated sources do lie on this AGN sequence, there are also just as many high-redshift PAH-dominated sources with high extinction in this region. The large filled circles are the SMGs from this study for which we




FIG. 7.—*Spitzer* color-color diagram as an AGN diagnostic (see also Ivison et al. 2004). The dashed and dotted lines show the positions of Mrk 231 (an AGN) and M82 (a starburst), respectively, as a function of redshift (redshift is indicated by the numbers along the track). The large filled circles are the SMGs from this study for which we have IRS spectra. The two SMGs (C1 and GN04) that have a $>50\%$ contribution from continuum (AGN) emission to the mid-IR luminosity based on the IRS spectra (see Table 4) both lie off the M82 starburst sequence. Based on our IRS results, SMGs that lie in the light shaded region are starburst (SB) dominated in the mid-IR. The crosses indicate sources that are classified as X-ray AGNs (Alexander et al. 2005). The smaller filled circles are the rest of the SMGs from Pope et al. (2006). Using our IRS results, the fraction of all SMGs that are starburst dominated in the mid-IR (i.e., are found in the shaded region) is $\geq 80\%$. The small open triangles are the *Spitzer* $24\ \mu\text{m}$ -selected ULIRGs from Sajina et al. (2007) with $z > 1$; most of these sources are outside the starburst region defined for SMGs. [See the electronic edition of the *Journal* for a color version of this figure.]

have IRS spectra. Interestingly, both SMGs (C1 and GN04) that have a $>50\%$ contribution from continuum (AGN) emission to the mid-IR luminosity based on the IRS spectra (see Table 4) lie off the starburst sequence. Based on our IRS results, SMGs that lie in the light shaded region of Figure 7 are starburst dominated in the mid-IR. SMGs outside the shaded region likely have a combination of starburst and AGN emission in the mid-IR or lie at the highest redshifts ($z \geq 4$).

This division between starburst and AGN-dominated sources based on IRS spectra agrees with the broadband colors. The crosses indicate sources that are classified as X-ray AGNs (Alexander et al. 2005). Many of these lie along the starburst sequence and have IRS spectra that are starburst dominated. While the X-ray emission may be telling us that an AGN is present in the system, it does not seem to be a very good diagnostic of whether the AGN contributes to the bolometric luminosity. Sources that have AGN-dominated mid-IR spectra, like C1, clearly show very extreme colors in this diagram; however, there are not enough of these sources to characterize their allowable positions in this color-color space. The smaller filled circles are the rest of the SMGs from the GOODS-N submillimeter supermap from Pope et al. (2006). Extrapolating our IRS results, the fraction of all SMGs that are starburst dominated in the mid-IR is $\sim 80\%$. The small open triangles are the $z > 1$ bright $24\ \mu\text{m}$ ULIRGs from Sajina et al. (2007); very few of these sources lie within the starburst region defined for SMGs, indicating that they have a wider range of mid-IR spectra.




FIG. 8.—IR SED of source C2. This figure demonstrates our template fitting to the full SED from mid-IR to radio wavelengths. The dotted line is the CE01 template that best matches the submillimeter flux, while the dot-dashed line is the CE01 template that best matches the 16 and $24\ \mu\text{m}$ fluxes. Both of these models are clearly poor fits to the other data points and have a higher total IR luminosity than the best fit to all the available data. The double-dot-dashed line is the best fit to all the data points $>7\ \mu\text{m}$ rest frame where we have allowed the CE01 templates to scale. While this provides a good fit to the mid-IR spectrum and the submillimeter flux, the radio flux and the total IR luminosity are underestimated. The dashed line is a scaled CE01 template with additional extinction from the Draine (2003) extinction curve that provides the best fit to our data. We require both a scaling of local templates (in this case CE01) and additional extinction to fit the SED of high-redshift SMGs. [See the electronic edition of the *Journal* for a color version of this figure.]

5. FULL SED FITS

In Pope et al. (2006), we fit the $24\ \mu\text{m}$, $850\ \mu\text{m}$, and radio photometry to a suite of models to determine the total IR luminosity and dust temperatures. We had used the CE01 templates and modified them by adding additional extinction from the Draine (2003) models. The CE01 templates were developed to be representative of local galaxies; therefore, they contain an inherent luminosity-temperature (L - T) relation. When fitting, we had allowed the models to scale so that we could solve for both the luminosity and the temperature independently (i.e., a model with high luminosity but cool average dust temperature is allowed). We had found that SMGs fit best to scaled-up versions of lower luminosity templates that had cooler average dust temperatures than local ULIRGs. In those fits the short-wavelength part of the spectrum was constrained only by the $24\ \mu\text{m}$ flux. From the present study it is apparent that the $24\ \mu\text{m}$ flux of SMGs will vary considerably with redshift, due to the presence of the PAH lines. Therefore, we have repeated the full SED fitting now that we have the mid-IR spectra.

Figure 8 demonstrates why we need to allow the CE01 templates to scale (i.e., luminosity and temperature are both variables) and the need for an additional extinction term. Here we plot the mid-IR to radio SED for one of our SMG sources, C2. First, we simply plot the CE01 template that best matches the submillimeter flux only as the dotted line and the CE01 template that best matches the 16 and $24\ \mu\text{m}$ fluxes as the dot-dashed line. For these two models we emphasize that the CE01 templates have not been allowed to scale; therefore, they follow the local luminosity-temperature relation. Because SMGs are so luminous at mid-IR and submillimeter wavelengths, the dotted and dot-dashed models have a warm average dust temperature ($\sim 50\ \text{K}$). The triple-dot-dashed line is the best fit to all the photometry




FIG. 9.—Mid-IR to radio SEDs of SMGs. The IRS spectrum (*thick line*) plus $70\ \mu\text{m}$, submillimeter, and radio photometry (*filled circles*) are used in the fit. The open squares indicate the photometry at 16 and $24\ \mu\text{m}$. The dashed line is the best-fit modified CE01 template, while the dot-dashed line is Mrk 231 normalized to the AGN fraction of the mid-IR given in Table 4. We used the latter to determine the contribution of the AGN component to L_{IR} . The horizontal dotted lines indicate the region within which L_{IR} is calculated. [See the electronic edition of the Journal for a color version of this figure.]

between rest-frame $7\text{--}300\ \mu\text{m}$ where we have allowed the CE01 templates to scale. While this provides a good fit to the mid-IR spectrum and the submillimeter flux, the radio flux and L_{IR} are underestimated compared to our best fit. We know that the CE01 templates need to be scaled up to match the photometry of SMGs; however, as we scale them up, we are introducing a greater dust mass that must then cause more extinction. The dashed line is a scaled CE01 template with additional extinction from the Draine (2003) extinction curve that provides the best fit to our data. We require both a scaling of local templates (in this case CE01) and additional extinction to fit the SED of high-redshift SMGs.

We thus fit the IRS spectrum plus $70\ \mu\text{m}$, submillimeter, and radio photometry to the CE01 templates plus Draine (2003) extinction for each of the 13 SMGs in this paper.¹⁴ We solve for the three free parameters (luminosity, average dust temperature, and extinction) and find the template with the lowest χ^2 . Figure 9 shows the full SED fits. In general, the resulting fits are very good. The thick line and filled circles are the points used in the fit,

¹⁴ For the GN39 double system, we split the total submillimeter flux according to the radio flux of each component (note that this is the same as splitting by $24\ \mu\text{m}$ flux).




FIG. 9—Continued

while the open symbols show the other photometry points (mostly 16 and 24 μm) that were not used in the fits but agree well with the best-fit SED. We required a scaling of 10–100 in luminosity to fit the local CE01 templates to the SMG photometry. Since high-redshift SMGs can be well fitted by modified CE01 templates that contain an inherent radio-IR correlation, this suggests that SMGs also follow a similar radio-IR correlation to local galaxies (see also Kovács et al. 2006).

We integrated the full SEDs in Figure 9 from 8 to 1000 μm to get the total IR luminosity. Values of L_{IR} are listed in Table 5 and are in good agreement with the values from Pope et al. (2006).

By simply scaling to the wavelength where the SED is a maximum as an estimate of the average dust temperature, we obtain a

median of 32 K for these 13 SMGs. This is consistent with previous estimates of dust temperatures in SMGs (Chapman et al. 2005; Kovács et al. 2006; Pope et al. 2006). We also found a similar trend between luminosity and temperature as found in Chapman et al. (2005), which is offset to cooler temperatures from the local correlation. Knowing the shape of the far-IR SED and the total luminosity, we can also estimate the dust mass: following the relation given in Dunne et al. (2000), the median dust mass for this sample is $2.4 \times 10^8 M_{\odot}$. This is roughly an order of magnitude higher than dust masses observed in galaxies locally although the local galaxies studied are generally much less luminous.

In order to quantify the effect of the extinction term on the models, we calculate L_{IR} without the extinction. The median amount




FIG. 9—Continued

TABLE 5
PAH STRENGTHS AND INFRARED LUMINOSITIES

SMG ID	PAH Luminosity ($10^9 L_\odot$)			PAH Equivalent Width ^a (μm)			L_{IR} ($10^{12} L_\odot$)		
	6.2 μm	7.7 μm	11.3 μm	6.2 μm	7.7 μm	11.3 μm	SB	AGN	Total
GN31.....	...	3.4 ± 1.2	0.32 ± 0.11	...	>0.17	<0.10	0.27
GN26.....	12.2 ± 1.2	42.9 ± 5.1	8.1 ± 1.4	0.38 ± 0.04	1.00 ± 0.13	1.18 ± 0.54	>3.5	<0.1	3.6
GN17.....	13.0 ± 4.1	0.22 ± 0.07	>1.3	<0.7	2.0
C2.....	...	31.1 ± 6.3	7.0 ± 1.8	...	1.08 ± 0.30	1.00 ± 0.29	>2.8	<0.5	3.3
C3.....	<19.2	9.2 ± 2.7	7.9 ± 2.7	<0.50	0.25 ± 0.08	0.45 ± 0.16	5.2	0.8	6.0
GN39a.....	13.9 ± 4.9	39.2 ± 12.9	<36.3	0.66 ± 0.26	0.86 ± 0.30	<0.86	>4.2	<0.8	5.0
GN39b.....	12.3 ± 5.6	23.1 ± 5.1	<29.4	1.09 ± 0.65	0.59 ± 0.14	<1.14	>2.7	<0.1	2.8
GN07.....	12.5 ± 2.6	17.6 ± 3.2	9.9 ± 1.8	1.10 ± 0.38	0.41 ± 0.08	1.58 ± 0.47	6.7	0.2	6.9
GN06.....	18.0 ± 1.5	26.3 ± 4.4	10.5 ± 2.2	0.83 ± 0.10	0.93 ± 0.16	2.95 ± 1.38	5.7	0.7	6.4
C1.....	6.3 ± 1.5	59.1 ± 12.6	<18.4	0.05 ± 0.01	0.54 ± 0.14	<0.18	5.4	4.2	9.6
GN05.....	...	12.9 ± 4.3	0.44 ± 0.15	...	>3.9	<0.1	4.0
GN19.....	12.5 ± 3.0	32.5 ± 9.0	...	0.59 ± 0.20	0.77 ± 0.22	...	>11.9	<0.4	12.3
GN04.....	<23.7	43.3 ± 9.9	...	<0.45	0.63 ± 0.15	...	4.5	2.1	6.6

^a The equivalent widths are all in rest frame.

that the extinction term changes L_{IR} by is 13% (0%–20%), and so models that do not include the extinction term will incorrectly determine L_{IR} by around 13%. We find an average value of $\tau_{9.7} \sim 1$ and do not see any correlations between the extinction and L_{IR} for the SMGs given the error bars.

For sources that we found to contain an AGN component in the mid-IR, we calculate the contribution to L_{IR} from the AGN. Using *ISO* spectra, Tran et al. (2001) found that the ratio of $L_{5-10 \mu\text{m}}/L_{\text{IR}}$ was about 2.5 times greater for AGNs than for starbursts; i.e., as you extrapolate from mid-IR to far-IR wavelengths, AGNs become less important bolometrically than starbursts. In order to quantify this for our sample, we scale the Mrk 231 (an AGN whose L_{IR} is known to be dominated by AGN emission and not star formation; Armus et al. 2007) template to the AGN fraction (listed in Table 4) of the mid-IR luminosity and then integrate the SED from 8 to 1000 μm to get the contribution to L_{IR} from the AGN component. For sources that have an upper limit to the AGN contribution, we plot the scaled Mrk 231 in Figure 9 at this limit. The values of $L_{\text{IR}}^{\text{AGN}}$ from integrating under the curves and $L_{\text{IR}}^{\text{SB}} = L_{\text{IR}} - L_{\text{IR}}^{\text{AGN}}$ are given in Table 5. As seen in Figure 9, only C1 has enough AGN contribution in the mid-IR to make a significant contribution to L_{IR} . Therefore, the intense IR luminosities seen in SMGs are powered almost entirely by star formation. In § 7 we discuss how this luminosity can be converted to give us the SFR.

6. PAH LUMINOSITIES

PAH line profiles have broad wings that contribute a significant fraction of the line flux. PAH line fluxes, therefore, need to be consistently measured in a uniform way before comparisons between local sources and high-redshift sources can be performed. We have measured line fluxes of all sources consistently, including the comparison sample of low-redshift starbursts (Brandl et al. 2006), using the technique described in § 3.1.

The PAH line luminosities and equivalent widths of the SMGs, including 3σ upper limits when the line is undetected, are listed in Table 5. The 6.2 μm line is the cleanest to measure and use for diagnostics since it is the least affected by silicate extinction and line blending. We can see that $L_{\text{PAH},7.7}$ for the SMGs varies by over an order of magnitude, while $L_{\text{PAH},6.2}$ and $L_{\text{PAH},11.3}$ have much smaller dynamic ranges of a factor of a few. While this might seem odd, we note that only a subset (40%–60%) of our

sample have 6.2 and 11.3 μm spectral coverage, whereas 92% have coverage of the 7.7 μm emission line. Furthermore, the redshift range for the sources with 6.2 and 11.3 μm spectral coverage is very small, and therefore we expect a small dynamic range in PAH luminosity. The sources that are dominated by an AGN in the mid-IR, as classified by our IRS spectra (C1 and GN04), both show suppressed 6.2 μm emission relative to their 7.7 μm emission.

Figure 10 shows the relative strengths of PAH lines compared to local starbursts (Brandl et al. 2006). The top panels show $L_{\text{PAH},7.7}$ as a function of $L_{\text{PAH},6.2}$ (top left panel) and $L_{\text{PAH},11.3}$ (top right panel). The smaller bottom panels show the residual scatter after removing the fit shown in the top panels; the two larger error bars quantify the 1σ scatter seen in the starbursts and SMGs. The outlying source in the left panel of Figure 10 is C1, the SMG with the largest AGN contribution.

While the SMGs do not show enough dynamic range in PAH luminosity to pick out any strong correlations within the population, they do fall on the relation between various PAH lines established for local starburst galaxies. The dashed lines in Figure 10 are the best fits to the local starbursts only (squares). We find essentially the same relation if we fit the local starbursts and the SMGs, indicating that the PAH luminosities in SMGs are simply scaled up (albeit by several orders of magnitude) from less luminous starburst galaxies. The best-fit relations from fitting both the SMGs and the local starburst galaxies are

$$\log L_{\text{PAH},7.7} = (1.0 \pm 0.1) + (0.94 \pm 0.01) \log L_{\text{PAH},6.2}, \quad (2)$$

$$\log L_{\text{PAH},7.7} = (0.5 \pm 0.1) + (1.00 \pm 0.01) \log L_{\text{PAH},11.3}. \quad (3)$$

The SMG C1 appears to lie off this relation with an excess of 7.7 μm PAH emission for a given $L_{\text{PAH},6.2}$ or $L_{\text{PAH},11.3}$. Rigopoulou et al. (1999) found suppressed 6.2 μm emission relative to 7.7 μm emission in *ISO* spectra of local ULIRGs compared to starbursts and conclude that this is primarily because of extinction. This suggests that, on average, the SMG C1 considered here may contain greater extinction or a more dominant AGN component than most SMGs and starburst galaxies since an AGN continuum will dilute the strength of the weaker 6.2 and 11.3 μm features relative to the strong 7.7 μm line.

Figure 11 shows the equivalent widths of the 6.2 and 7.7 μm PAH lines. The vertical dashed line at $\text{EW}_{\text{PAH},6.2} = 0.2$ is often




FIG. 10.—Plot comparing the PAH line luminosities of SMGs presented in this paper (*circles*) to that of local starbursts (*squares*; Brandl et al. 2006). All line measurements have been made in a consistent way (see § 3.1). The 1σ error bars are shown on all points (the 10% errors on the starbursts are similar in size to the symbols). The top panels show $L_{\text{PAH},7.7}$ as a function of $L_{\text{PAH},6.2}$ (*left*) and $L_{\text{PAH},11.3}$ (*right*), where the dashed lines show the best fit to the starburst galaxies only. The similar relationship seen in starbursts and SMGs between PAH line strengths indicates that SMGs are starburst dominated and that PAH line strengths do not vary significantly with redshift. The smaller bottom panels show the residual scatter after removing the fit shown in the top panels; the larger error bars quantify the 1σ scatter seen in the starbursts and SMGs, respectively. [See the electronic edition of the Journal for a color version of this figure.]

used to separate starburst-dominated and AGN-dominated sources (Armus et al. 2007; Sajina et al. 2007). With the exception of C1, all of our sources with measurable $6.2\ \mu\text{m}$ PAH emission have $\text{EW}_{\text{PAH},6.2} > 0.2$, which indicates that they are starburst dominated. We have again plotted the local starbursts for comparison. We found that the values in Brandl et al. (2006) for $\text{EW}_{\text{PAH},7.7}$ were on average 1.6 times lower than our measurements of the




FIG. 11.—Rest-frame equivalent widths of main PAH features. Again the SMGs are denoted by the circles and the local starbursts (Brandl et al. 2006) by the squares. All measurements have been made in a consistent way (see § 3.1). Measurements of both the 6.2 and $7.7\ \mu\text{m}$ PAH lines are not available for all SMGs. C1 stands out again as having a very small $6.2\ \mu\text{m}$ EW, below the values of $\text{EW}_{\text{PAH},6.2} = 0.2$ (*vertical dashed line*), which is often used to separate AGN-dominated and starburst-dominated systems. [See the electronic edition of the Journal for a color version of this figure.]

starburst $\text{EW}_{\text{PAH},7.7}$, due to differences in the way the lines were measured. Therefore, we have revised all the Brandl et al. (2006) points upward in this plot. The starbursts cover a small area of this plot, while the SMGs are spread over a larger region, although with much larger error bars. Our current data do not have sufficient S/N to determine if there is any genuine spread or correlation among the EWs; however, our error bars are sufficiently small to exclude equivalent widths of less than 0.2 in either PAH line (with the exception of C1). Our SMGs show similar equivalent widths to the PAH-dominated ULIRGs from the *Spitzer* $24\ \mu\text{m}$ -selected sample in Sajina et al. (2007), if we decrease their values of $\text{EW}_{\text{PAH},7.7}$ by a factor of 4 to account for the different methods for measuring the EWs. With $\text{EW}_{\text{PAH},6.2}$ values of ≥ 0.5 and $\tau_{9.7} \sim 1$, high-redshift SMGs lie in region 2C of the diagnostic plot in Spoon et al. (2007). Sources in this region are classified as PAH dominated, and, not surprisingly, M82 also falls in this region.

Utilizing *ISO* spectra of starbursts, ULIRGs, and AGNs, Genzel et al. (1998) used the ratio of the $7.7\ \mu\text{m}$ PAH line flux to the continuum flux, l/c , as a diagnostic for AGN contribution in the mid-IR emission. In general, galaxies with $l/c \geq 1$ are classified as starbursts and those with $l/c < 1$ are AGN dominated. As we stress in § 3.1, this type of analysis is strongly dependent on the specific choice made for the continuum fitting procedure. Nevertheless, we calculate $l/c \sim 2.3$ for the composite SMG and note that all SMGs in our sample satisfy the $l/c \geq 1$ criterion for starburst-dominated sources. This is true even for C1, which clearly shows a strongly rising continuum indicative of the presence of an AGN. Based on Figure 11, the $6.2\ \mu\text{m}$ PAH line may be more useful in weeding out AGN-dominated sources.

In Figure 12 we plot the PAH line luminosities as a function of L_{IR} for SMGs and local starbursts (*large panels*). For the SMGs, we plot the $L_{\text{IR}}^{\text{SB}}$ coming from star formation (i.e., after subtracting the component from the AGN). Errors in L_{IR} for the SMGs have been estimated from the error in the submillimeter flux only, since this dominates (assuming fixed CE01 templates). In all panels, the dashed line is the best fit to the local starburst galaxies; the smaller panels show the residual scatter around this




FIG. 12.—Correlations between L_{IR} and PAH luminosities. The dashed lines show the best-fit relations for the local starburst galaxies only; the smaller panels show the residual scatter around the best-fit relations. The SMGs clearly lie on the relations established by local starburst galaxies. [See the electronic edition of the *Journal* for a color version of this figure.]

best-fit relation. The best-fit relation for the starburst galaxies only is consistent with the fit to both the starburst galaxies and the SMGs. The best-fit relation for both the SMGs and starburst galaxies is given by

$$\log L_{\text{PAH},6.2} = (-2.7 \pm 0.1) + (1.01 \pm 0.01) \log L_{\text{IR}}, \quad (4)$$

$$\log L_{\text{PAH},7.7} = (-1.5 \pm 0.1) + (0.95 \pm 0.01) \log L_{\text{IR}}, \quad (5)$$

$$\log L_{\text{PAH},11.3} = (-1.6 \pm 0.2) + (0.91 \pm 0.01) \log L_{\text{IR}}. \quad (6)$$

The derived parameters are all within 1σ of those found by fitting only the starbursts. The slopes are close to 1, indicating a direct proportionality between L_{IR} and the PAH line luminosities.

The first thing to note is that the SMGs lie on the relation established for local starburst galaxies extrapolated to very high luminosities. This is what we expect if both are dominated by the same emission mechanism, namely, star formation. Brandl et al. (2006) showed that the PAH line luminosities are correlated with the IR luminosities within their starburst sample. Schweitzer et al. (2006) showed that a similar correlation exists in a sample of QSOs due to star formation in them. Peeters et al. (2004) showed a correlation between L_{FIR} and $L_{\text{PAH},6.2}$ for a range of scales of star formation in local massive star-forming regions to normal and starburst galaxies. The fact that the SMGs lie on this relation demonstrates the reliability of our full SED fitting to derive the total IR luminosity and confirms that SMGs do have incredibly high IR luminosities powered primarily by star formation. Note that C1 does not stick out as an outlier in these plots, since we have removed the AGN component of its L_{IR} .

The second thing that stands out in Figure 12 is that there is significant scatter in this relation for both the low-redshift starbursts and the high-redshift SMGs. In the smaller panels the two larger error bars quantify the 1σ scatter seen in the starbursts and SMGs. The scatter is similar for the SMGs and starbursts and is considerably higher than would be implied by the measurement uncertainty. Other properties of individual objects, such as geometry of the starburst region, spatial distribution of the dust, viewing angle relative to a merging event, clumpiness of the star formation, etc., are likely playing a role in this scatter.

We found no correlations between the IR luminosity and equivalent widths of the PAH lines in SMGs. This is consistent with what was found in Brandl et al. (2006) for the local starburst galaxies and shows that both the mid-IR continuum and the PAH strength scale together with increasing L_{IR} . We also find no correlation between X-ray luminosity and L_{IR} (or the PAH luminosities) for the SMGs, a correlation that would have been expected had the IR emission been dominated by an AGN.

Using the Kennicutt (1998) relation between L_{IR} and SFR, we plot the corresponding SFR on the top axis of Figure 12. The Kennicutt (1998) relation between L_{IR} and SFR assumes a Salpeter (1955) initial mass function and applies to starbursts with ages less than 100 Myr. Assuming that both the PAH and far-IR emission are tracing the star formation, we can use the Kennicutt (1998) relation and the above PAH line luminosity-to- L_{IR} correlations to determine the relation between SFR and the PAH line luminosity. We use the $6.2 \mu\text{m}$ PAH line since it is the least affected by the silicate absorption and line blending. We find that given the PAH luminosity, one can calculate the SFR using

$$\text{SFR} (M_{\odot} \text{ yr}^{-1}) \simeq 10^{-7} L_{\text{PAH},6.2} (L_{\odot}). \quad (7)$$

Similar (although perhaps slightly less accurate) relations can be found for $L_{\text{PAH},7.7}$ and $L_{\text{PAH},11.3}$ using equation (4).

7. DISCUSSION

We have shown from the mid-IR spectra of high-redshift SMGs that these systems are starburst dominated and contain only a small contribution from the AGN. Despite the fact that X-ray observations suggest that most SMGs harbor an AGN, only a small fraction of SMGs (2/13) contain an AGN that contributes significantly to the mid-IR luminosity. When extrapolated to the total IR luminosity, all SMGs are dominated (>50% of L_{IR}) by star formation and not AGN activity.

From Table 5 we see that the median $L_{\text{IR}}^{\text{SB}}$ for our sample is $4 \times 10^{12} L_{\odot}$, which corresponds to an SFR of around $700 M_{\odot} \text{ yr}^{-1}$. This is an extremely high rate of star formation, capable of creating a massive $10^{11} M_{\odot}$ galaxy in only 10^8 yr. Coupled with the low number density of SMGs compared to other massive galaxies at high redshift, this leads us to suspect that SMGs are a short-lived phase in the evolution of massive galaxies.

SMGs are only one of several populations of high-redshift galaxies that are ultraluminous in the IR. Many $z \sim 2$ galaxies selected in deep $24 \mu\text{m}$ surveys also qualify as ULIRGs (e.g., Daddi et al. 2005; Yan et al. 2005, 2007). These other ULIRGs are generally not detected at submillimeter wavelengths. What then are the fundamental differences between various populations of high-redshift ULIRGs?

One of the main factors that could cause a bifurcation in the population of ULIRGs at high redshift is the presence or absence of a bolometrically significant AGN. The evolutionary scenario proposed by Sanders et al. (1988) shows massive galaxies going through several main stages en route to becoming a massive elliptical. It starts with an IR-luminous phase most likely triggered by a massive merger. During this stage of intense star formation, the AGN is also growing. As the AGN becomes larger, it begins to feed back on the galaxy, eventually quenching the star formation completely by blowing off all the dust and gas. Thus begins the QSO phase, where the AGN is free to dominate the emission. Eventually the QSO consumes its gas and we end up with a quiescent massive elliptical galaxy. In this scenario, the submillimeter emission will be at a maximum during the initial IR-luminous phase (Granato et al. 2004; Springel et al. 2005). However, one uncertainty in this scenario is the timescale. In particular, how long is the IR-luminous phase and how long does it take the AGN to develop?

Since most ULIRGs at $z \sim 2$ show a mix of AGN and starburst activity (e.g., Houck et al. 2005; Sajina et al. 2007), they fit into this evolutionary scenario at the stage where the AGN has developed but before it has had a chance to expel all the gas and dust, since the IR luminosities are high. Since SMGs show less of an AGN contribution, it is likely that they represent a slightly earlier phase of this scenario, before the AGN has time to develop fully and is significant to the IR luminosity. This is also consistent with the finding that SMGs have cool dust temperatures (Pope et al. 2006), partly because the star formation is taking place on extended spatial scales and partly because the AGN is not yet energetically important for heating the dust. Other high-redshift ULIRGs, like those selected at $24 \mu\text{m}$ (e.g., Sajina et al. 2007), should have warmer dust temperatures (or have overestimated IR luminosities); otherwise, they would be detected in the submillimeter. Therefore, they likely represent the stage after the SMG phase in this evolutionary scenario.

For local ULIRGs, there is evidence that shows that the luminosity of CO emission decreases as the merger progresses (Rigopoulou et al. 1999), which is consistent with decreasing submillimeter emission in the later stages of the merger. This idea is also consistent with morphological studies that show that SMGs are often tidally disrupted systems (Chapman et al. 2003;

Conselice et al. 2003; Pope et al. 2005). Alternatively, SMGs and other high-redshift ULIRGs might appear different because of different merger progenitors.

Since the number density of SMGs is much less than that of all $24 \mu\text{m}$ -selected IR-luminous galaxies, then this phase in the evolutionary sequence must be shorter. We can make a rough estimate for the duration of the submillimeter-luminous phase. In the 2 Gyr between $1.5 < z < 3$ there are roughly 20 SMGs and 500 $24 \mu\text{m}$ -selected galaxies (with $L_{\text{IR}} > 10^{11} L_{\odot}$; see Chary et al. 2004) within a portion of the GOODS-N field. Assuming that all IR-luminous $24 \mu\text{m}$ -selected galaxies will undergo a submillimeter-luminous stage during this period of time, we calculate the duration of the submillimeter-luminous phase to be on the order of 10^8 yr.¹⁵ As mentioned above, this timescale is what is needed to create a galaxy with a stellar mass of $\sim 10^{11} M_{\odot}$ (Borys et al. 2005) at a rate of $\sim 1000 M_{\odot} \text{ yr}^{-1}$. This timescale is consistent with those derived from gas masses of SMGs (Greve et al. 2005).

8. CONCLUSIONS

Spitzer IRS spectroscopy has been obtained for a sample of 13 SMGs brighter than $200 \mu\text{Jy}$ at $24 \mu\text{m}$. This has effectively confirmed the identification of these SMG counterparts and in some cases has provided a new redshift estimate. The SMGs show strong PAH emission and, on average, only a small continuum contribution from hot dust.

We have explored several diagnostics from the mid-IR spectra of SMGs to determine the level of AGNs contributing to the luminosity at these wavelengths. All of them seem to converge on a picture in which SMGs are starburst-dominated systems with at most a 30% contribution from an AGN at mid-IR wavelengths. Their mid-IR spectra are similar to a scaled spectrum of local starburst galaxies like M82.

The classification of an SMG as AGN or starburst dominated from the mid-IR often disagrees with the classification from the X-rays. While X-ray observations are better suited for determining the presence of an AGN in an SMG, the mid-IR spectra can determine how important the AGN is to the total IR luminosity since it is directly detecting the hot dust emission and not subject to details about the geometry of the obscuring material.

Full IR SED fits to the mid-IR, far-IR, and radio data show that SMGs are best fitted by scaled up versions of local IR-luminous galaxy templates with additional extinction from the Draine (2003) extinction curves. These models have cool dust temperatures ($T = 32$ K) and high L_{IR} values that imply SFRs of $\sim 700 M_{\odot} \text{ yr}^{-1}$ assuming a Salpeter initial mass function.

SMGs lie on the local relation between L_{IR} and $L_{\text{PAH},6.2}$ (or $L_{\text{PAH},7.7}$ or $L_{\text{PAH},11.3}$), which means that the PAH line flux can be used to estimate the SFRs in these systems, albeit with large uncertainties. Equivalent widths, which have been the focus of some other studies, are much less useful in this regard.

SMGs appear to be an earlier phase in the evolution of massive galaxies than other high-redshift ULIRGs. The average high-redshift ULIRG shows warmer dust and less PAH emission. It is therefore likely that the AGN in SMGs has not yet grown strong enough to heat the dust substantially and destroy the PAH molecules.

We have been able to put constraints on the contribution from AGN activity to the rest-frame 5–12 μm mid-IR emission and have extrapolated this to far-IR wavelengths. Without more data

¹⁵ It is not clear what fraction of $24 \mu\text{m}$ -selected galaxies will undergo a submillimeter-luminous phase since SMGs are known to be massive and could represent only the most extreme galaxies. Nevertheless, this calculation provides a useful reality check on the expected duration of the SMG phase.

from 30–100 μm rest frame, it is difficult to directly probe the emission at far-IR wavelengths. We will be able to model this part of the spectrum in more detail once data from the *Herschel Space Observatory*¹⁶ and SCUBA-2 (Holland et al. 2006) become available. Details of the SMG phase will be understood only after the spatially resolved spectroscopy capabilities of ALMA¹⁷ become available.

We are grateful to the referee for helpful suggestions which improved this paper. We thank Anna Sajina for helpful discussions and for providing composite IRS spectra of their high-redshift *Spitzer* 24 μm -selected ULIRGs. We are very grateful to

Bernhard Brandl for providing the IRS spectra of local starburst galaxies and Benjamin Magnelli for providing information from the completeness simulations of the 70 μm images. Support for this work was provided by NASA through the *Spitzer Space Telescope* Fellowship Program, through a contract issued by the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA. This work was supported by the Natural Sciences and Engineering Research Council of Canada and the Canadian Space Agency. D. M. A. acknowledges the Royal Society for support. This work is based on observations made with the *Spitzer Space Telescope*, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. The IRS was a collaborative venture between Cornell University and Ball Aerospace Corporation funded by NASA through the Jet Propulsion Laboratory and Ames Research Center.

¹⁶ See <http://herschel.esac.esa.int/>.

¹⁷ See <http://www.eso.org/projects/alma/>.

APPENDIX A

SPITZER IRS NOISE

Figure 13 shows the final rms of our spectra as a function of the integration time and spectral order. Values from the *Spitzer* Science Center sensitivity calculator, SPEC-PET,¹⁸ are given for comparison as the dashed lines. We confirm that the sensitivity goes as $\sigma(\text{mJy}) = at(\text{s})^{-1/2}$ even for very long integrations. The scatter in the measured values is probably dominated by the accuracy of the sky subtraction in these crowded fields. Given the scatter, our observed sensitivities are consistent with those from SPEC-PET. The decrease in sensitivity in the SL1 order could be related to our choice of 240 s ramps in fields with backgrounds $>20 \text{ MJy sr}^{-1}$ at 24 μm , which results in incorrect droop correction (see the *Spitzer* Observer's Manual).

APPENDIX B

NOTES ON INDIVIDUAL SOURCES

Here we describe several of our sources in more detail. Sources that are not discussed here are relatively straightforward in terms of interpreting their mid-IR spectra.

¹⁸ See <http://ssc.spitzer.caltech.edu/tools/specpet/>.




FIG. 13.—Observed 1σ rms as a function of integration time for our IRS staring mode observations. All observations from our *Spitzer* GO2-20456 program are shown here. The dashed lines show the predicted values from SPEC-PET for each order. We confirm that the sensitivity goes as $\sigma \propto t^{-1/2}$ even for very long integrations. The scatter in the measured values is probably dominated by the accuracy of the sky subtraction in these crowded fields. Given the scatter, our observed sensitivities are consistent with those from SPEC-PET. [See the electronic edition of the *Journal* for a color version of this figure.]

B1. GN06

The optical and IRS redshifts disagree for this SMG. The IRS redshift is based on three PAH features, 6.2, 7.7, and 11.3 μm , and the optical spectroscopic redshift of 1.865 is inconsistent with the positions of these emission features, even within the errors. The optical spectroscopic redshift of GN06 comes from Chapman et al. (2005), where they note that the optical galaxy was offset from the radio counterpart by about an arcsecond. From the images of GN06 in Figure 1, we see that there are two optical knots, both of which are offset from the counterpart that is detected in the radio and mid-IR. For this source, we conclude that the optical and mid-IR spectra are picking up different galaxies and the submillimeter emission is coming from the mid-IR-detected galaxy.

B2. C2

C2 has the largest discrepancy between the IRS and UV/optical spectroscopic redshifts. The IRS redshift for this source is based on the 7.7 and 11.3 μm PAH features, and there is also a hint of the 8.6 μm line. The optical spectroscopic redshift would have placed the 7.7 μm PAH line right in the middle of this spectrum; therefore, it is inconsistent with the IRS spectrum. The optical spectroscopic redshift comes from detections of both $\text{H}\alpha$ and $\text{Ly}\alpha$ emission (Swinbank et al. 2004; Chapman et al. 2005). The images in Figure 1 show an extended optical galaxy. This is a curious source; however, since the mid-IR is more likely to trace the submillimeter emission, we conclude that the IRS redshift is correct for this SMG.

B3. C1

This SMG is not detected in the deep *Chandra* 2 Ms imaging of GOODS-N. Furthermore, the optical spectrum from Swinbank et al. (2004) clearly shows $\text{H}\alpha$, N II , and the S II doublet, and the low N II -to- $\text{H}\alpha$ ratio is used to classify this SMG as a starburst. However, the mid-IR spectrum of C1 clearly shows a steeply rising continuum indicative of an AGN emission. This is also apparent from the extreme *Spitzer* colors shown in Figure 7. The mid-IR spectrum also shows PAH emission, and the ratio of PAH luminosity to IR luminosity for C1 is significantly lower than those seen in local starbursts. We conclude that this SMG harbors a Compton-thick AGN that accounts for 44% of the total IR luminosity. This is the only SMG that contains a bolometrically significant AGN. Interestingly, C1 has not been detected in CO (Greve et al. 2005). From the X-ray flux limit, we can place a limit on the column density of $N_{\text{H}} \gg 10^{24} \text{ cm}^{-2}$. C1 is relatively bright at 24 μm (1.2 mJy), suggesting that these types of obscured AGNs at high redshift should be detectable in even shallow *Spitzer* surveys.

B4. GN39 DOUBLE SYSTEM

One further SMG deserves special discussion, and that is GN39. This SMG has two radio/MIPS counterparts separated by 8'', which we refer to as GN39a and GN39b (note that this submillimeter system was first presented in Chapman et al. 2001). Both are confirmed to lie at $z = 1.996$ (Swinbank et al. 2004; Chapman et al. 2005); therefore, it is likely that they are both contributing to the submillimeter emission. The separation corresponds to 70 kpc at this redshift.

Our IRS observations picked up both counterparts, and, with a narrow extraction window, we were able to extract a separate spectrum of each component. Some of the flux from GN39a falls into the spectral extraction window of GN39b; however, a comparison of the 24 μm flux from the IRS spectra to that from the MIPS 24 μm image confirms that only $\frac{1}{3}$ of the flux in the IRS spectrum of GN39b is from GN39a. The extracted spectra confirm that these two sources are indeed at the same redshift, and both spectra look very similar to each other and to the rest of the SMGs, supporting the idea that they are both submillimeter emitters. Furthermore, both components are X-ray detected and classified as obscured AGNs by Alexander et al. (2005), although the two IRS spectra show a completely starburst-dominated system at mid-IR wavelengths.

Since these two SMGs are at the same redshift, we know that they must be associated. Given the physical distance between them, larger than those seen in most local ULIRGs (Murphy et al. 1996), we could be seeing the early stage in the merger of two massive galaxies (Springel et al. 2005). Note that GN04, GN07, and GN19 also have double radio counterparts, which are thought to be at the same redshift. However, in these sources, the two components are separated by only 3'' and therefore do not resolve into two separate 24 μm sources. Thus, the IRS spectra for these sources contain contributions from both components.

REFERENCES

- Alexander, D. M., Bauer, F. E., Chapman, S. C., Smail, I., Blain, A. W., Brandt, W. N., & Ivison, R. J. 2005, *ApJ*, 632, 736
 Alexander, D. M., et al. 2003, *AJ*, 126, 539
 Allamandola, L. J., Hudgins, D. M., & Sandford, S. A. 1999, *ApJ*, 511, L115
 Aretxaga, I., et al. 2007, *MNRAS*, 379, 1571
 Armus, L., et al. 2007, *ApJ*, 656, 148
 Bauer, F. E., Alexander, D. M., Brandt, W. N., Hornschemeier, A. E., Vignali, C., Garmire, G. P., & Schneider, D. P. 2002, *AJ*, 124, 2351
 Blain, A. W., Smail, I., Ivison, R. J., Kneib, J.-P., & Frayer, D. T. 2002, *Phys. Rep.*, 369, 111
 Borys, C., Chapman, S., Halpern, M., & Scott, D. 2003, *MNRAS*, 344, 385
 Borys, C., Smail, I., Chapman, S. C., Blain, A. W., Alexander, D. M., & Ivison, R. J. 2005, *ApJ*, 635, 853
 Brandl, B. R., et al. 2006, *ApJ*, 653, 1129
 Brandt, W. N., & Hasinger, G. 2005, *ARA&A*, 43, 827
 Capak, P., et al. 2004, *AJ*, 127, 180
 Chapman, S. C., Blain, A. W., Smail, I., & Ivison, R. J. 2005, *ApJ*, 622, 772
 Chapman, S. C., Richards, E. A., Lewis, G. F., Wilson, G., & Barger, A. J. 2001, *ApJ*, 548, 147
 Chapman, S. C., Windhorst, R., Odewahn, S., Yan, H., & Conselice, C. 2003, *ApJ*, 599, 92
 Charmandaris, V., et al. 1999, *Ap&SS*, 266, 99
 Chary, R., & Elbaz, D. 2001, *ApJ*, 556, 562 (CE01)
 Chary, R., et al. 2004, *ApJS*, 154, 80
 Conselice, C. J., Chapman, S. C., & Windhorst, R. A. 2003, *ApJ*, 596, L5
 Coppin, K., et al. 2006, *MNRAS*, 372, 1621
 Cowie, L. L., Barger, A. J., Hu, E. M., Capak, P., & Songaila, A. 2004, *AJ*, 127, 3137
 Daddi, E., et al. 2005, *ApJ*, 631, L13
 ———. 2007, *ApJ*, 670, 173
 Desai, V., et al. 2007, *ApJ*, 669, 810
 Dickinson, M., et al. 2003, in *The Mass of Galaxies at Low and High Redshift*, ed. R. Bender & A. Renzini (Berlin: Springer), 324

- Draine, B. T. 2003, *ARA&A*, 41, 241
- Draine, B. T., & Li, A. 2007, *ApJ*, 657, 810
- Dunne, L., Eales, S., Edmunds, M., Ivison, R., Alexander, P., & Clements, D. L. 2000, *MNRAS*, 315, 115
- Farrah, D., et al. 2007, *ApJ*, 667, 149
- Fischer, J., et al. 1999, *Ap&SS*, 266, 91
- Förster Schreiber, N. M., et al. 2003, *A&A*, 399, 833
- Frayser, D. T., et al. 2006, *ApJ*, 647, L9
- Genzel, R., et al. 1998, *ApJ*, 498, 579
- Giavalisco, M., et al. 2004, *ApJ*, 600, L93
- Granato, G. L., De Zotti, G., Silva L., Bressan, A., & Danese, L. 2004, *ApJ*, 600, 580
- Greve, T. R., et al. 2005, *MNRAS*, 359, 1165
- Hauser, M. G., & Dwek, E. 2001, *ARA&A*, 39, 249
- Higdon, S. J. U., et al. 2004, *PASP*, 116, 975
- Holland, W. S., et al. 1999, *MNRAS*, 303, 659
- . 2006, *Proc. SPIE*, 6275, 45
- Houck, J. R., et al. 2004, *ApJS*, 154, 18
- . 2005, *ApJ*, 622, L105
- Huynh, M. T., Pope, A., Frayer, D. T., & Scott, D. 2007, *ApJ*, 659, 305
- Ivison, R. J., et al. 2002, *MNRAS*, 337, 1
- . 2004, *ApJS*, 154, 124
- Kennicutt, R. C. 1998, *ARA&A*, 36, 189
- Kovács, A., Chapman, S. C., Dowell, C. D., Blain, A. W., Ivison, R. J., Smail, I., & Phillips, T. G. 2006, *ApJ*, 650, 592
- Lagache, G., Puget, J.-L., & Dole, H. 2005, *ARA&A*, 43, 727
- Le Floc'h, E., et al. 2005, *ApJ*, 632, 169
- Lilly, S. J., Eales, S. A., Gear, W. K. P., Hammer, F., Le Fèvre, O., Crampton, D., Bond, J. R., & Dunne, L. 1999, *ApJ*, 518, 641
- Lutz, D., Genzel, R., Kunze, D., Rigopoulou, D., Spoon, H. W. W., Sturm, E., Tran, D., & Moorwood, A. F. M. 1999, *Ap&SS*, 266, 85
- Lutz, D., Maiolino, R., Spoon, H. W. W., & Moorwood, A. F. M. 2004, *A&A*, 418, 465
- Lutz, D., Spoon, H. W. W., Rigopoulou, D., Moorwood, A. F. M., & Genzel, R. 1998, *ApJ*, 505, L103
- Lutz, D., Valiante, E., Sturm, E., Genzel, R., Tacconi, L. J., Lehnert, M. D., Sternberg, A., & Baker, A. J. 2005a, *ApJ*, 625, L83
- Lutz, D., Yan, L., Armus, L., Helou, G., Tacconi, L. J., Genzel, R., & Baker, A. J. 2005b, *ApJ*, 632, L13
- Menéndez-Delmestre, K., et al. 2007, *ApJ*, 655, L65
- Murphy, T. W., Jr., Armus, L., Matthews, K., Soifer, B. T., Mazzarella, J. M., Shupe, D. L., Strauss, M. A., & Neugebauer, G. 1996, *AJ*, 111, 1025
- Peeters, E., Spoon, H. W. W., & Tielens, A. G. G. M. 2004, *ApJ*, 613, 986
- Pérez-González, P. G., et al. 2005, *ApJ*, 630, 82
- Pope, A. 2007, Ph.D. thesis, Univ. British Columbia
- Pope, A., Borys, C., Scott, D., Conelice, C., Dickinson, M., & Mobasher, B. 2005, *MNRAS*, 358, 149
- Pope, A., et al. 2006, *MNRAS*, 370, 1185
- Puget, J.-L., Abergel, A., Bernard, J.-P., Boulanger, F., Burton, W. B., Desert, F.-X., & Hartmann, D. 1996, *A&A*, 308, L5
- Puget, J. L., & Leger, A. 1989, *ARA&A*, 27, 161
- Rigopoulou, D., Spoon, H. W. W., Genzel, R., Lutz, D., Moorwood, A. F. M., & Tran, Q. D. 1999, *AJ*, 118, 2625
- Sajina, A., Lacy, M., & Scott, D. 2005, *ApJ*, 621, 256
- Sajina, A., Yan, L., Armus, L., Choi, P., Fadda, D., Helou, G., & Spoon, H. 2007, *ApJ*, 664, 713
- Sajina, A., et al. 2008, *ApJ*, submitted
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Sanders, D. B., & Mirabel, I. F. 1996, *ARA&A*, 34, 749
- Sanders, D. B., Soifer, B. T., Elias, J. H., Madore, B. F., Matthews, K., Neugebauer, G., & Scoville, N. Z. 1988, *ApJ*, 325, 74
- Schweitzer, M., et al. 2006, *ApJ*, 649, 79
- Smith, J. D. T., et al. 2007, *ApJ*, 656, 770
- Soifer, B. T., Sanders, D. B., Madore, B. F., Neugebauer, G., Danielson, G. E., Elias, J. H., Lonsdale, C. J., & Rice, W. L. 1987, *ApJ*, 320, 238
- Spoon, H. W. W., Marshall, J. A., Houck, J. R., Elitzur, M., Hao, L., Armus, L., Brandl, B. R., & Charmandaris, V. 2007, *ApJ*, 654, L49
- Springel, V., Di Matteo, T., & Hernquist, L. 2005, *MNRAS*, 361, 776
- Sturm, E., Lutz, D., Tran, D., Feuchtgruber, H., Genzel, R., Kunze, D., Moorwood, A. F. M., & Thornley, M. D. 2000, *A&A*, 358, 481
- Swinbank, A. M., Smail, I., Chapman, S. C., Blain, A. W., Ivison, R. J., & Keel, W. C. 2004, *ApJ*, 617, 64
- Tacconi, L. J., et al. 2006, *ApJ*, 640, 228
- Teplitz, H. I., Charmandaris, V., Chary, R., Colbert, J. W., Armus, L., & Weedman, D. 2005, *ApJ*, 634, 128
- Teplitz, H. I., et al. 2007, *ApJ*, 659, 941
- Tran, Q. D., et al. 2001, *ApJ*, 552, 527
- Valiante, E., Lutz, D., Sturm, E., Genzel, R., Tacconi, L. J., Lehnert, M. D., & Baker, A. J. 2007, *ApJ*, 660, 1060
- Yan, L., et al. 2005, *ApJ*, 628, 604
- . 2007, *ApJ*, 658, 778