

KECK DEIMOS SPECTROSCOPY OF A *GALEX* UV-SELECTED SAMPLE FROM THE MEDIUM IMAGING SURVEY¹

RYAN P. MALLERY,² R. MICHAEL RICH,² SAMIR SALIM,² TODD SMALL,³ STEPHANE CHARLOT,^{4,5} MARK SEIBERT,³
TED WYDER,³ TOM A. BARLOW,³ KARL FORSTER,³ PETER G. FRIEDMAN,³ D. CHRISTOPHER MARTIN,³
PATRICK MORRISSEY,² SUSAN G. NEFF,⁶ DAVID SCHIMINOVICH⁷, LUCIANA BIANCHI,⁸ JOSÉ DONAS,⁹
TIMOTHY M. HECKMAN,¹⁰ YOUNG-WOOK LEE,¹¹ BARRY F. MADORE,¹² BRUNO MILLIARD,⁹
ALEX S. SZALAY,¹⁰ BARRY Y. WELSH,¹³ AND SUKYOUNG YI¹¹

Received 2006 January 25; accepted 2006 July 25

ABSTRACT

We report results from a pilot program to obtain spectroscopy for objects detected in the *Galaxy Evolution Explorer* (*GALEX*) Medium Imaging Survey (MIS). Our study examines the properties of galaxies detected by *GALEX* fainter than the Sloan Digital Sky Survey (SDSS) spectroscopic survey. This is the first study to extend the techniques of Salim and coworkers to estimate stellar masses, star formation rates (SFRs), and the b (star formation history) parameter for star-forming galaxies out to $z \sim 0.7$. We obtain redshifts for 50 *GALEX* MIS sources reaching NUV = 23.9 (AB mag) having counterparts in the SDSS Data Release 4 (DR4). Of our sample, 43 are star-forming galaxies with $z < 0.7$, 3 have emission-line ratios indicative of active galactic nuclei with $z < 0.7$, and 4 objects with $z > 1$ are QSOs, 3 of which are not previously cataloged. We compare our sample to a much larger sample of $\sim 50,000$ matched *GALEX*/SDSS galaxies with SDSS spectroscopy; while our survey is shallow, the optical counterparts to our sources reach ~ 3 mag fainter in SDSS r than the SDSS spectroscopic sample. We use emission-line diagnostics for the galaxies to determine that the sample contains mostly star-forming galaxies. The galaxies in the sample populate the blue sequence in the NUV $- r$ versus M_r color-magnitude diagram. The derived stellar masses of the galaxies range from 10^8 to $10^{11} M_\odot$, and derived SFRs are between 10^{-1} and $10^2 M_\odot \text{ yr}^{-1}$. Our sample has SFRs, luminosities, and velocity dispersions that are similar to the samples of faint compact blue galaxies studied previously in the same redshift range by Koo and collaborators, Guzmán and collaborators, and Phillips and collaborators. However, our sample is ~ 2 mag fainter in surface brightness than the compact blue galaxies. We find that the star formation histories for a majority of the galaxies are consistent with a recent starburst within the last 100 Myr.

Subject headings: galaxies: active — galaxies: high-redshift — galaxies: starburst — ultraviolet: galaxies

1. INTRODUCTION

Following the successful launch and early operations of the *Galaxy Evolution Explorer* (*GALEX*) satellite in 2003, we decided to undertake an initial spectroscopic assay of sources detected by *GALEX* using the DEIMOS multiobject spectrograph (Faber et al. 2003) on the Keck II telescope. Although *GALEX* data have been matched to relatively deep optical surveys (Schiminovich et al. 2005), our study is the first to extend the modeling technique of Salim et al. (2005) from $z = 0.25$ to $z \sim 0.5$.

The power of the *GALEX* Medium Imaging Survey (MIS) to select interesting objects is noteworthy. We recall that the MIS observations are 1500 s in duration, imaging FUV and NUV simultaneously (Martin et al. 2005a; Morrissey et al. 2005). The *GALEX* satellite has investigated the ultraviolet universe to

$z \sim 1$ with the goal of determining, among other things, the star formation rate (SFR) of galaxies in the local universe, $z \lesssim 0.25$ (Salim et al. 2005, 2007; Martin et al. 2005b; Wyder et al. 2005; Treyer et al. 2005), and the evolution of the global star formation density out to $z \sim 1$ (Schiminovich et al. 2005). The ultimate goal of such work along with other surveys and observations of galaxies at similar and higher redshifts is to constrain the baryonic physics of galaxy formation and evolution. Questions still remain as to how the SFR depends on different factors such as environment, mass, and morphology (Martin et al. 2005a). Previous work deriving SFRs with a UV-selected sample was carried out by Sullivan et al. (2000), who derived SFRs both from UV luminosities at 2000 Å and H α luminosities from optical spectra for a sample of galaxies at redshifts of $0 < z < 0.4$ detected in the UV by the balloon-borne telescope FOCA (Milliard

¹ Some of the data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California, and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation.

² Department of Physics and Astronomy, University of California, Los Angeles, CA 90095-1562.

³ California Institute of Technology, MC 405-47, 1200 East California Boulevard, Pasadena, CA 91125.

⁴ Max-Planck Institut für Astrophysik, D-85748 Garching, Germany.

⁵ Institut d'Astrophysique de Paris, CNRS, 98 bis boulevard Arago, F-75014 Paris, France.

⁶ Laboratory for Astronomy and Solar Physics, NASA Goddard Space Flight Center, Greenbelt, MD 20771.

⁷ Department of Astronomy, Columbia University, New York, NY 10027.

⁸ Center for Astrophysical Sciences, The Johns Hopkins University, 3400 North Charles Street, Baltimore, MD 21218.

⁹ Laboratoire d'Astrophysique de Marseille, BP 8, Traverse du Siphon, 13376 Marseille Cedex 12, France.

¹⁰ Department of Physics and Astronomy, The Johns Hopkins University, Homewood Campus, Baltimore, MD 21218.

¹¹ Center for Space Astrophysics, Yonsei University, Seoul 120-749, Korea.

¹² Space Sciences Laboratory, University of California, Berkeley, 601 Campbell Hall, Berkeley, CA 94720; and Observatories of the Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA 91101.

¹³ Space Sciences Laboratory, University of California, Berkeley, 601 Campbell Hall, Berkeley, CA 94720.




FIG. 4.—Emission-line diagnostic diagram first used by Baldwin et al. (1981). The shaded two-dimensional histogram corresponds to the *GALEX* IR1.1/SDSS DR2 spectroscopic sample. The dashed curves taken from Kewley et al. (2001) and Kauffmann et al. (2003) show the distinction between sources with emission-line flux coming from AGN (above) and sources with emission line flux coming from H II regions (below). Out of the 14 objects from our DEIMOS sample with detections of H β , [O III], H α , and [N II], only three have emission-line ratios consistent with emission due to an AGN.

and errors for the emission lines were measured in IDL using the MPFIT function. The continuum of each spectrum was first fit by a polynomial, and then the emission lines were simultaneously fit with Gaussians. Table 1 lists the objects, their equivalent widths, and flux ratios of [O III] λ 5007/H β and [N II] λ 6584/H α for objects with a 3σ measurement of at least one of the above lines. We did not perform a reddening correction to the fluxes, since the standard procedure requires either detections of both H α and H β or a measurement of the radio continuum (Osterbrock 1989). The effect of this on the flux ratios of [O III] λ 5007/H β and [N II] λ 6584/H α is thought to be negligible due to the small wavelength separation of the emission lines. In the DEIMOS sample we only have detections of both H α and H β for 18 objects, all at $z < 0.3$, and chose for consistency not to perform corrections on any of the spectra. We note that the foreground Galactic extinction, $E(B - V)$, calculated from the dust maps of Schlegel et al. (1998) is ~ 0.05 mag for the field.

Fourteen objects have measured fluxes of H β , [O III] λ 5007, H α , and [N II] λ 6584. Figure 4 shows the emission-line diagnostic of Baldwin et al. (1981) used to distinguish star-forming galaxies and type II active galactic nuclei (AGNs). The solid curve is the boundary between star-forming galaxies and AGNs determined through modeling of starburst galaxy spectra by Kewley et al. (2001). A star-forming galaxy with only $\sim 20\%$ of the optical emission-line flux due to an AGN would lie above the star-forming/AGN boundary (Kewley et al. 2001). The dashed curve is the more stringent demarcation used by Kauffmann et al. (2003) to distinguish between AGNs and star-forming galaxies in SDSS. The shaded histogram shows the distribution of emission-line ratios for a sample of 51,000 SDSS/*GALEX* objects (see § 4). The curve of Kauffmann et al. (2003) distinguishes between the two different sources of emission-line flux in this large sample, with the AGN occupying the parameter space to the right of the distribution of star-forming galaxies. In the DEIMOS sample only three galaxies of the 14 have emission-line flux ratios above this curve, implying that some portion of the flux is due to an AGN, although it is still within the errors that the emission-line flux for two of these objects results entirely from star-forming regions inside the host galaxies. Objects can also be AGNs if $\log [O III]/H\beta > 1$ or




FIG. 5.—SDSS r -magnitude distribution of the DEIMOS spectroscopic sample plotted with the SDSS r -magnitude distribution of the matched *GALEX* IR1.1/SDSS DR2 spectroscopic sample (shaded histogram). The DEIMOS-matched spectroscopic sample mean r -magnitude (21) is 3 mag fainter than the mean of the SDSS spectroscopic sample.

$\log [N II]/H\alpha > 0.3$. Another 13 objects also have only detections of H β and [O III] λ 5007. Of the 13, one has a flux ratio of $\log [O III]/H\beta > 1$, indicative of an AGN. Sullivan et al. (2000) found a similar ratio of galaxies classified as star-forming/AGN from examination of emission lines in their sample of UV-selected galaxies.

4. MAGNITUDE AND COLOR DISTRIBUTION

The DEIMOS sample is ~ 3 mag deeper in r than the SDSS spectroscopic sample. Figure 5 shows a histogram of the SDSS r magnitudes for the sample plotted with the SDSS r magnitude for *GALEX* MIS sources in IR1.1 with matches in the SDSS DR2 spectroscopic sample. The IR1.1/DR2 sample consists of $\sim 51,000$ galaxies at redshifts $0.005 < z < 0.25$ with derived star formation histories (SFHs) by SED fitting (see § 5). The IR1.1/DR2 histogram is normalized to the size of the DEIMOS galaxy sample. The IR1.1/DR2 sample is mostly bounded in r by the SDSS spectroscopic survey limits, $r < 17.7$ (Strauss et al. 2002).

Using the redshifts determined for the DEIMOS sample we calculate the absolute r magnitude, M_r , of galaxies in the sample, k -corrected to the mean redshift of the IR1.1/DR2 sample, $z = 0.1$; k -corrections for all bandpasses were calculated using the publicly available code of Blanton et al. (2003), version 4.1.4. In Figure 6 we construct a color-magnitude diagram (CMD) of $NUV - r$ versus M_r . We overplot our sample onto the IR1.1/DR2 sample, plotted as a shaded contour plot. In the figure the diamonds correspond to DEIMOS objects with spectroscopically measured redshifts at $z < 0.25$, and the crosses correspond to objects with spectroscopically measured redshifts at $z > 0.25$. The contours enclose 40% and 80% of the objects in the IR1.1/DR2 sample.

The CMD of the IR1.1/DR2 sample clearly shows the bimodality of galaxies seen by *GALEX* in the nearby universe, $z < 0.25$, with distribution peaks at blue $NUV - r$ colors of ~ 3 and at red $NUV - r$ colors of ~ 6 (Wyder et al. 2005). We henceforth call these the blue and red sequences. The galaxies in the blue sequence are generally late-type in morphology and have spectra with emission line ratios indicative of star formation (Salim et al. 2005; Brinchmann et al. 2004), while the galaxies in the red sequence typically have absorption-line spectra and lower star formation rates than blue-sequence galaxies. The majority of the objects in the DEIMOS sample lie in the region of the diagram




FIG. 6.—CMD of M_r vs. $NUV - r$ plotted for the sample of *GALEX*/SDSS objects with DEIMOS spectroscopy plotted with the entire *GALEX* (IR1.1)/SDSS (DR2) matched sample having SDSS spectroscopy. The diamonds represent galaxies in the DEIMOS sample at $z < 0.25$; crosses represent galaxies in the DEIMOS sample at $z > 0.25$. The IR1.1/DR2 sample is plotted as the shaded contour plot; the darker regions correspond to a higher density of points, and the contours encompass 40% and 80% of the of the objects in the sample.

occupied by blue star-forming galaxies in the IR1.1/DR2 sample, mostly along the blue edge of the distribution, with only one having a $NUV - r$ color greater than 4. The spectrum for this object, at $z = 0.077$, is shown in Figure 1. The spectrum shows absorption features, Na D and Mg 1 b, but also shows weak emission features of $H\alpha$, $[N II] \lambda 6584$, and $[S II] \lambda\lambda 6717, 6731$.

In Figure 7 (*bottom left*) we again plot the CMD for the sample, highlighting the objects with FUV detections (*filled circles*). The other three panels of this figure show the UV color distribution for the 24 galaxies in the DEIMOS sample with FUV detections. Plotted for reference are galaxies from the IR1.1/DR2 sample with FUV detections shown as the shaded contour plots or shaded histograms in the respective panels. Besides the three galaxies in the DEIMOS sample with $FUV - NUV < 0$ the UV color for this sample shows a similar distribution with the IR1.1/DR2 sample. In the FUV CMD the objects with $M_r < 20$ show a range of UV color not seen in the local blue-sequence galaxies of similar optical luminosity. Both Figures 6 and 7 show that this sample probes the type of galaxies that one obtains when looking at objects detected by *GALEX* several magnitudes deeper than the SDSS spectroscopic limit.

5. DERIVED GALAXY PARAMETERS

We derive the following galaxy parameters according to the approach of Salim et al. (2005): the V -band dust attenuations, A_V in magnitudes, stellar metallicity Z , the current SFR averaged over the past 100 Myr in $M_\odot \text{ yr}^{-1}$, the present-day stellar mass, M_* , of the galaxy in M_\odot , the fraction of stellar mass formed in bursts




FIG. 7.—*Top left*: CMD of M_r vs. $FUV - NUV$ plotted for the 24 galaxies in the DEIMOS sample with FUV detections, plotted with objects in the IR1.1/DR2 matched sample with FUV detections, shown as a shaded contour plot. *Top right*: Histogram of $FUV - NUV$ color for the 24 DEIMOS FUV galaxies plotted with distribution of the IR1.1/DR2 FUV sample (*shaded histogram*). *Bottom left*: Same as Fig. 6. The filled (unfilled) circles correspond to objects with (without) FUV detections. *Bottom right*: Color-color diagram of the FUV DEIMOS galaxies again overplotted onto the FUV IR1.1/DEIMOS sample. The symbols used are the same as in Fig. 6, and the contours encompass 40% and 80% of the of the objects in the IR1.1/DR2 sample.

TABLE 2
GALAXIES, REDSHIFTS, COLORS, AND DERIVED PARAMETERS

SDSS Object ID	z	R.A.	Decl.	FUV	NUV	NUV $- r$	$\log M_*$	$\log \text{SFR}$	$\log b$
587725578037166705.....	0.192	17 37 30.681	57 20 8.70	22.53	23.11	1.51	8.69	-0.42	-0.09
587725578037166675.....	0.498	17 37 34.409	57 21 13.71	...	22.62	0.97	9.50	0.50	-0.07
587725578037166672.....	0.189	17 37 39.053	57 21 30.10	...	23.04	3.36	10.05	0.15	-0.55
587725578037166749.....	0.320	17 37 41.177	57 18 46.29	22.87	22.12	1.88	9.91	0.65	-0.14
587725578037100839.....	0.561	17 37 43.542	57 24 5.51	...	22.19	1.19	10.40	0.85	-0.32
587725578037166468.....	0.330	17 37 45.923	57 19 17.19	22.38	21.50	2.15	10.85	0.48	-0.84
587725578037166557.....	0.168	17 37 58.250	57 20 16.80	20.61	20.12	1.18	9.47	0.53	-0.08
587725578037166295.....	0.560	17 37 59.495	57 23 21.80	...	22.16	0.89	9.70	0.79	-0.07
587725578037166628.....	0.652	17 38 2.622	57 23 26.99	...	22.73	1.51	10.51	1.11	-0.24
587725578037166629.....	0.321	17 38 7.756	57 23 34.90	...	22.32	1.04	9.09	0.28	-0.05
587725578037101267.....	0.293	17 38 22.192	57 25 2.82	...	23.36	1.66	9.16	-0.06	-0.18
587725578037100922.....	0.168	17 38 23.628	57 27 56.42	21.48	21.97	2.67	9.97	-0.24	-0.76
587725578037101473.....	0.496	17 38 27.876	57 26 33.90	22.91	22.28	1.22	9.76	0.68	-0.10
587725578037101077.....	0.387	17 38 36.526	57 30 38.59	...	22.32	0.52	9.11	0.51	-0.01
587725578037101221.....	0.518	17 38 37.317	57 26 53.92	...	22.50	1.09	10.07	0.62	-0.26
587725578037100775 ^a	0.176	17 38 37.646	57 29 13.99	21.03	20.34	3.08	11.08	0.69	-0.88
587725578037101176.....	0.281	17 38 41.873	57 28 9.30	...	22.33	0.99	9.24	-0.09	-0.19
587725591458873892.....	0.281	17 38 48.259	57 29 10.40	22.69	22.23	1.80	9.76	0.43	-0.13
587725591458808530.....	0.427	17 38 50.596	57 29 54.31	...	22.88	2.89	10.50	2.12	0.00
587725591458808528.....	0.598	17 38 54.873	57 30 5.80	...	22.58	1.83	10.37	1.33	-0.09
587725578037494286.....	0.134	17 39 45.205	56 40 2.20	20.87	20.47	1.17	9.66	-0.15	-0.28
587725578037494409.....	0.355	17 39 51.035	56 39 13.11	...	23.23	1.78	9.14	0.36	-0.04
587725578037495014.....	0.213	17 39 51.343	56 39 58.89	...	22.73	0.95	8.18	-0.66	-0.02
587725578037494283.....	0.084	17 39 53.130	56 40 18.91	22.24	22.00	2.34	9.17	-0.57	-0.35
587725578037494050.....	0.077	17 39 56.243	56 38 17.20	...	22.11	5.46	10.76	-0.80	-1.91
587725578037494494.....	0.356	17 39 56.287	56 37 21.00	...	22.20	1.96	9.99	1.14	-0.03
587725578037494738.....	0.474	17 39 56.704	56 37 53.80	...	22.72	0.86	9.67	0.38	-0.16
587725591459201436.....	0.066	17 40 1.663	56 44 3.12	20.89	20.74	2.20	9.21	-0.41	-0.35
587725591459201125 ^b	0.066	17 40 3.853	56 41 59.82	20.37	19.99	1.96
587725591459201656.....	0.157	17 40 6.724	56 43 49.51	22.64	22.12	-0.52	8.68	-0.97	-0.24
587725591459201627.....	0.079	17 40 8.196	56 44 57.80	22.18	21.68	1.51	8.48	-0.61	-0.09
587725591459201372.....	0.615	17 40 8.328	56 39 39.82	...	22.68	1.80	10.53	0.81	-0.51
587725591459201929.....	0.436	17 40 15.117	56 46 4.70	22.36	22.16	0.24	9.27	0.22	-0.07
587725591459201726.....	0.439	17 40 17.307	56 41 31.09	...	22.42	0.94	9.98	0.37	-0.22
587725591459136339.....	0.188	17 40 17.468	56 48 5.50	21.23	20.57	0.76	9.98	-0.06	-0.47
587725591459201670.....	0.651	17 40 19.175	56 43 39.18	...	22.46	0.14	8.93	0.61	0.29
587725591459201562.....	0.068	17 40 20.969	56 42 43.30	...	22.77	2.90	8.77	-1.06	-0.53
587725591459201614.....	0.188	17 40 23.225	56 45 42.52	22.37	21.50	0.59	8.72	-0.06	-0.04
587725591459136404.....	0.220	17 40 23.708	56 46 34.10	22.53	21.76	0.79	9.59	-0.47	-0.55
587725591459135913.....	0.293	17 40 29.685	56 50 27.71	21.64	21.93	1.18	9.55	0.12	-0.21
587725591459136024 ^b	0.245	17 40 32.498	56 49 12.51	22.73	20.99	-0.08
587725591459201666.....	0.078	17 40 33.193	56 44 5.32	...	20.82	1.06	9.00	-0.90	-0.42
587725591459136277.....	0.290	17 40 35.691	56 50 29.29	22.66	22.23	3.27	10.99	0.49	-0.98
587725591459136330.....	0.565	17 40 46.362	56 49 13.40	...	22.18	1.33	10.03	1.06	-0.08
587725591459136031.....	0.192	17 40 56.177	56 49 15.38	20.94	20.09	2.45	10.92	0.97	-0.47
587725591459136351.....	0.319	17 40 56.594	56 48 52.20	...	21.35	2.15	10.90	0.75	-0.60
587725578037100834.....	1.515	17 38 9.360	57 25 21.36	...	23.40	2.63
587725591459201473.....	1.387	17 40 9.836	56 40 7.68	...	22.67	1.14
587725591459136108.....	1.632	17 40 43.682	56 48 45.36	...	23.05	1.67
587725591459135981 ^a	1.028	17 40 49.197	56 47 23.65	22.62	21.20	1.94

NOTE.—Units of right ascension are hours, minutes, and seconds, and units of declination are degrees, arcminutes, and arcseconds.

^a Denotes objects with corresponding spectra in SDSS.

^b Denotes objects with failed SED fit.

over the last 100 Myr, F_{burst} , and the Scalo (1986) b -parameter, defined as the ratio of the current SFR to the past time-averaged SFR (averaged over the estimated age, not Hubble time). The galaxy parameters are derived from model libraries of galaxies at redshifts between 0.1 and 1.6 at redshift increments of 0.1 for the DEIMOS sample, and at redshifts of 0.05, 0.1, 0.15, 0.2, and 0.25 for the IR1.1/DR2 sample. Each library consists of up to $\sim 10^5$ models. Each model is parameterized according to galaxy age, optical depth, SFH, and metallicity. The SFH of each model is

parameterized according to Kauffmann et al. (2003) with an underlying, continuous, exponentially declining SFR on which bursts of star formation, random in time and amplitude, are superimposed. Dust attenuation in each model is parameterized using the prescription of Charlot & Fall (2000) using an effective V -band optical depth τ_V and absorption curve, $\tau \propto \lambda^{0.7}$, resulting from both giant molecular clouds and the diffuse interstellar medium (ISM), with the fraction μ of τ_V contributed only by the diffuse ISM. The V -band optical depth from giant molecular clouds is taken to only




FIG. 8.—Derived galaxy parameters, metallicity, and V -band attenuation plotted vs. M_* and $\text{NUV} - r$ color. The DEIMOS spectroscopic sample is plotted over the matched *GALEX* IR1.1/SDSS DR2 spectroscopic sample (shaded contour plot). The crosses correspond to objects with redshifts $z > 0.25$. The diamonds correspond to objects with redshifts $z < 0.25$. The contours for the plots of stellar mass contain 57%, 84%, and 97% of the data. The contours of $\text{NUV} - r$ color enclose 40% and 80% of the sample.

affect stars younger than 10 Myr. A description of the prior distributions of the model parameters is discussed in Salim et al. (2005).

SEDs are created for each galaxy in the library using the population-synthesis code of Bruzual & Charlot (2003). The model SEDs are convolved with the *GALEX* and SDSS filter-response curves. Statistical estimates of physical galaxy parameters are derived by comparing the observed seven-band *GALEX*/SDSS fluxes of each galaxy to all the convolved model SEDs in the nearest redshift library. Probability density functions (PDFs) for each physical parameter are created by assigning weights to the parameters. The χ^2 goodness of fit of the model determines the weight ($\propto \exp[-\chi^2/2]$) that is assigned to the parameters of that model. The median (or most typical parameter value) of the PDF is taken as the estimate of the galaxy parameter.

We perform SED fits for galaxies not identified as QSOs. In Table 2 we list the galaxies, their redshifts, $\text{NUV} - r$ non- k -corrected colors, and their derived parameters. The parameters derived from the SED fits are not used unless the reduced χ^2 fit of at least one model is below 10. Only two of the 46 galaxies do not meet this criterion. Figures 8 and 9 show the derived parameters for the DEIMOS sample of galaxies overplotted on the IR1.1/DR2 sample plotted as shaded contour plots. The contours are labeled and encompass 52%, 84%, and 97% of the data. The diamonds as before represent objects with $z < 0.25$, and the crosses represent objects with $z > 0.25$. The objects span the range of derived parameters of the IR1.1/DR2 sample for Z and A_V .

The derived metallicities, $Z = [\text{Fe}/\text{H}]$, for the sample are not very well constrained. The typical error, 1/4 of the difference between the 97.5 percentile and the 2.5 percentile of the PDF, of the

derived metallicities is $\pm 43\% Z_\odot$. The derived M_* range from 10^8 to $10^{11} M_\odot$ with an average error of $\sim \pm 0.2$ dex, but the majority of the M_* are at or below $10^{10} M_\odot$, about an order of magnitude below the mean of the IR1.1/DR2 sample. The derived SFRs lie between 10^{-1} and $10^2 M_\odot \text{ yr}^{-1}$, with five galaxies having derived SFRs greater than $10 M_\odot \text{ yr}^{-1}$. The average error on the derived SFRs is typically ± 0.3 dex. Comparing the SFRs between galaxies with similar $\text{NUV} - r$ color or M_* , the galaxies at $z > 0.25$ have SFRs about an order of magnitude greater than the galaxies at $z < 0.25$. This is mostly a selection effect since the more distant galaxies are also more luminous on average. A comparison of these SED-derived SFRs with $\text{H}\alpha$ -derived SFRs cannot be performed since the spectra are not flux calibrated, and many lack $\text{H}\alpha$. The range of SFRs we find for this sample is similar to the range of SFRs for local UV galaxies found by Sullivan et al. (2000) derived both from $\text{H}\alpha$ luminosity and from the UV luminosity, but is an order of magnitude larger than the SFRs found by Kobulnicky & Kewley (2004) in their sample of GOODS-North Treasury Keck Redshift Survey galaxies at $0.4 < z < .9$ when compared to galaxies in the DEIMOS sample with similar redshifts.

We find that the galaxies are largely starbursts with $\log b \sim -0.1$. This is shown in Figure 10. The distribution of $\log b$ for objects in the DEIMOS sample with SED fits are plotted with the distribution for the IR1.1/DR2 sample normalized to the number of objects in the DEIMOS sample with SED fits. The DEIMOS sample shows a highly peaked value of $\log b$ compared to the IR1.1/DR2 sample which has two small peaks near $\log b \sim -3$ and ~ -1 , showing that most of these galaxies currently have less star formation now than in the past. The value $\log b \sim -0.1$

