

## Measurements of Charged Current Lepton Universality and $|V_{us}|$ Using Tau Lepton Decays to $e^- \bar{\nu}_e \nu_\tau$ , $\mu^- \bar{\nu}_\mu \nu_\tau$ , $\pi^- \nu_\tau$ , and $K^- \nu_\tau$

B. Aubert,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J. P. Lees,<sup>1</sup> V. Poireau,<sup>1</sup> E. Prencipe,<sup>1</sup> X. Prudent,<sup>1</sup> V. Tisserand,<sup>1</sup> J. Garra Tico,<sup>2</sup> E. Grauges,<sup>2</sup> M. Martinelli,<sup>3a,3b</sup> A. Palano,<sup>3a,3b</sup> M. Pappagallo,<sup>3a,3b</sup> G. Eigen,<sup>4</sup> B. Stugu,<sup>4</sup> L. Sun,<sup>4</sup> M. Battaglia,<sup>5</sup> D. N. Brown,<sup>5</sup> L. T. Kerth,<sup>5</sup> Yu. G. Kolomensky,<sup>5</sup> G. Lynch,<sup>5</sup> I. L. Osipenko,<sup>5</sup> K. Tackmann,<sup>5</sup> T. Tanabe,<sup>5</sup> C. M. Hawkes,<sup>5</sup> N. Soni,<sup>6</sup> A. T. Watson,<sup>6</sup> H. Koch,<sup>7</sup> T. Schroeder,<sup>7</sup> D. J. Asgeirsson,<sup>8</sup> B. G. Fulsom,<sup>8</sup> C. Hearty,<sup>8</sup> T. S. Mattison,<sup>8</sup> J. A. McKenna,<sup>8</sup> M. Barrett,<sup>9</sup> A. Khan,<sup>9</sup> A. Randle-Conde,<sup>9</sup> V. E. Blinov,<sup>10</sup> A. D. Bukin,<sup>10,\*</sup> A. R. Buzykaev,<sup>10</sup> V. P. Druzhinin,<sup>10</sup> V. B. Golubev,<sup>10</sup> A. P. Onuchin,<sup>10</sup> S. I. Serednyakov,<sup>10</sup> Yu. I. Skovpen,<sup>10</sup> E. P. Solodov,<sup>10</sup> K. Yu. Todyshev,<sup>10</sup> M. Bondioli,<sup>11</sup> S. Curry,<sup>11</sup> I. Eschrich,<sup>11</sup> D. Kirkby,<sup>11</sup> A. J. Lankford,<sup>11</sup> P. Lund,<sup>11</sup> M. Mandelkern,<sup>11</sup> E. C. Martin,<sup>11</sup> D. P. Stoker,<sup>11</sup> H. Atmacan,<sup>12</sup> J. W. Gary,<sup>12</sup> F. Liu,<sup>12</sup> O. Long,<sup>12</sup> G. M. Vitug,<sup>12</sup> Z. Yasin,<sup>12</sup> V. Sharma,<sup>13</sup> C. Campagnari,<sup>14</sup> T. M. Hong,<sup>14</sup> D. Kovalskyi,<sup>14</sup> M. A. Mazur,<sup>14</sup> J. D. Richman,<sup>14</sup> T. W. Beck,<sup>15</sup> A. M. Eisner,<sup>15</sup> C. A. Heusch,<sup>15</sup> J. Kroseberg,<sup>15</sup> W. S. Lockman,<sup>15</sup> A. J. Martinez,<sup>15</sup> T. Schalk,<sup>15</sup> B. A. Schumm,<sup>15</sup> A. Seiden,<sup>15</sup> L. Wang,<sup>15</sup> L. O. Winstrom,<sup>15</sup> C. H. Cheng,<sup>15</sup> D. A. Doll,<sup>16</sup> B. Echenard,<sup>16</sup> F. Fang,<sup>16</sup> D. G. Hitlin,<sup>16</sup> I. Narsky,<sup>16</sup> P. Ongmongkolkul,<sup>16</sup> T. Piatenko,<sup>16</sup> F. C. Porter,<sup>16</sup> R. Andreassen,<sup>17</sup> G. Mancinelli,<sup>17</sup> B. T. Meadows,<sup>17</sup> K. Mishra,<sup>17</sup> M. D. Sokoloff,<sup>17</sup> P. C. Bloom,<sup>18</sup> W. T. Ford,<sup>18</sup> A. Gaz,<sup>18</sup> J. F. Hirschauer,<sup>18</sup> M. Nagel,<sup>18</sup> U. Nauenberg,<sup>18</sup> J. G. Smith,<sup>18</sup> S. R. Wagner,<sup>18</sup> R. Ayad,<sup>19,†</sup> W. H. Toki,<sup>19</sup> R. J. Wilson,<sup>19</sup> E. Feltresi,<sup>20</sup> A. Hauke,<sup>20</sup> H. Jasper,<sup>20</sup> T. M. Karbach,<sup>20</sup> J. Merkel,<sup>20</sup> A. Petzold,<sup>20</sup> B. Spaan,<sup>20</sup> K. Wacker,<sup>20</sup> M. J. Kobel,<sup>21</sup> R. Nogowski,<sup>21</sup> K. R. Schubert,<sup>21</sup> R. Schwierz,<sup>21</sup> D. Bernard,<sup>22</sup> E. Latour,<sup>22</sup> M. Verderi,<sup>22</sup> P. J. Clark,<sup>23</sup> S. Playfer,<sup>23</sup> J. E. Watson,<sup>23</sup> M. Andreotti,<sup>24a,24b</sup> D. Bettoni,<sup>24a</sup> C. Bozzi,<sup>24a</sup> R. Calabrese,<sup>24a,24b</sup> A. Cecchi,<sup>24a,24b</sup> G. Cibinetto,<sup>24a,24b</sup> E. Fioravanti,<sup>24a,24b</sup> P. Franchini,<sup>24a,24b</sup> E. Luppi,<sup>24a,24b</sup> M. Munerato,<sup>24a,24b</sup> M. Negrini,<sup>24a,24b</sup> A. Petrella,<sup>24a,24b</sup> L. Piemontese,<sup>24a</sup> V. Santoro,<sup>24a,24b</sup> R. Baldini-Ferroli,<sup>25</sup> A. Calcaterra,<sup>25</sup> R. de Sangro,<sup>25</sup> G. Finocchiaro,<sup>25</sup> S. Pacetti,<sup>25</sup> P. Patteri,<sup>25</sup> I. M. Peruzzi,<sup>25,‡</sup> M. Piccolo,<sup>25</sup> M. Rama,<sup>25</sup> A. Zallo,<sup>25</sup> R. Contri,<sup>26a,26b</sup> E. Guido,<sup>26a</sup> M. Lo Vetere,<sup>26a,26b</sup> M. R. Monge,<sup>26a,26b</sup> S. Passaggio,<sup>26a</sup> C. Patrignani,<sup>26a,26b</sup> E. Robutti,<sup>26a</sup> S. Tosi,<sup>26a,26b</sup> K. S. Chaisanguanthum,<sup>27</sup> M. Morii,<sup>27</sup> A. Adamez,<sup>28</sup> J. Marks,<sup>28</sup> S. Schenk,<sup>28</sup> U. Uwer,<sup>28</sup> F. U. Bernlochner,<sup>29</sup> V. Klohe,<sup>29</sup> H. M. Lacker,<sup>29</sup> T. Lueck,<sup>29</sup> A. Volk,<sup>29</sup> D. J. Bard,<sup>30</sup> P. D. Dauncey,<sup>30</sup> M. Tibbetts,<sup>30</sup> P. K. Behera,<sup>31</sup> M. J. Charles,<sup>31</sup> U. Mallik,<sup>31</sup> J. Cochran,<sup>32</sup> H. B. Crawley,<sup>32</sup> L. Dong,<sup>32</sup> V. Eyges,<sup>32</sup> W. T. Meyer,<sup>32</sup> S. Prell,<sup>32</sup> E. I. Rosenberg,<sup>32</sup> A. E. Rubin,<sup>32</sup> Y. Y. Gao,<sup>33</sup> A. V. Gritsan,<sup>33</sup> Z. J. Guo,<sup>33</sup> N. Arnaud,<sup>34</sup> J. Béquilleux,<sup>34</sup> A. D'Orazio,<sup>34</sup> M. Davier,<sup>34</sup> D. Derkach,<sup>34</sup> J. Firmino da Costa,<sup>34</sup> G. Grosdidier,<sup>34</sup> F. Le Diberder,<sup>34</sup> V. Lepeltier,<sup>34</sup> A. M. Lutz,<sup>34</sup> B. Malaescu,<sup>34</sup> S. Pruvot,<sup>34</sup> P. Roudeau,<sup>34</sup> M. H. Schune,<sup>34</sup> J. Serrano,<sup>34</sup> V. Sordini,<sup>34,§</sup> A. Stocchi,<sup>34</sup> G. Wormser,<sup>34</sup> D. J. Lange,<sup>35</sup> D. M. Wright,<sup>35</sup> I. Bingham,<sup>36</sup> J. P. Burke,<sup>36</sup> C. A. Chavez,<sup>36</sup> J. R. Fry,<sup>36</sup> E. Gabathuler,<sup>36</sup> R. Gamet,<sup>36</sup> D. E. Hutchcroft,<sup>36</sup> D. J. Payne,<sup>36</sup> C. Touramanis,<sup>36</sup> A. J. Bevan,<sup>37</sup> C. K. Clarke,<sup>37</sup> F. Di Lodovico,<sup>37</sup> R. Sacco,<sup>37</sup> M. Sigamani,<sup>37</sup> G. Cowan,<sup>38</sup> S. Paramesvaran,<sup>38</sup> A. C. Wren,<sup>38</sup> D. N. Brown,<sup>39</sup> C. L. Davis,<sup>39</sup> A. G. Denig,<sup>40</sup> M. Fritsch,<sup>40</sup> W. Gradl,<sup>40</sup> A. Hafner,<sup>40</sup> K. E. Alwyn,<sup>41</sup> D. Bailey,<sup>41</sup> R. J. Barlow,<sup>41</sup> G. Jackson,<sup>41</sup> G. D. Lafferty,<sup>41</sup> T. J. West,<sup>41</sup> J. I. Yi,<sup>41</sup> J. Anderson,<sup>42</sup> C. Chen,<sup>42</sup> A. Jawahery,<sup>42</sup> D. A. Roberts,<sup>42</sup> G. Simi,<sup>42</sup> J. M. Tuggle,<sup>42</sup> C. Dallapiccola,<sup>43</sup> E. Salvati,<sup>43</sup> R. Cowan,<sup>44</sup> D. Dujmic,<sup>44</sup> P. H. Fisher,<sup>44</sup> S. W. Henderson,<sup>44</sup> G. Sciolla,<sup>44</sup> M. Spitznagel,<sup>44</sup> R. K. Yamamoto,<sup>44</sup> M. Zhao,<sup>44</sup> P. M. Patel,<sup>45</sup> S. H. Robertson,<sup>45</sup> M. Schram,<sup>45</sup> P. Biassoni,<sup>46a,46b</sup> A. Lazzaro,<sup>46a,46b</sup> V. Lombardo,<sup>46a</sup> F. Palombo,<sup>46a,46b</sup> S. Stracka,<sup>46a,46b</sup> L. Cremaldi,<sup>47</sup> R. Godang,<sup>47,||</sup> R. Kroeger,<sup>47</sup> P. Sonnek,<sup>47</sup> D. J. Summers,<sup>47</sup> H. W. Zhao,<sup>47</sup> M. Simard,<sup>48</sup> P. Taras,<sup>48</sup> H. Nicholson,<sup>49</sup> G. De Nardo,<sup>50a,50b</sup> L. Lista,<sup>50a</sup> D. Monorchio,<sup>50a,50b</sup> G. Onorato,<sup>50a,50b</sup> C. Sciacca,<sup>50a,50b</sup> G. Raven,<sup>51</sup> H. L. Snoek,<sup>51</sup> C. P. Jessop,<sup>52</sup> K. J. Knoepfel,<sup>52</sup> J. M. LoSecco,<sup>52</sup> W. F. Wang,<sup>52</sup> L. A. Corwin,<sup>53</sup> K. Honscheid,<sup>53</sup> H. Kagan,<sup>53</sup> R. Kass,<sup>53</sup> J. P. Morris,<sup>53</sup> A. M. Rahimi,<sup>53</sup> S. J. Sekula,<sup>53</sup> Q. K. Wong,<sup>53</sup> N. L. Blount,<sup>54</sup> J. Brau,<sup>54</sup> R. Frey,<sup>54</sup> O. Igonkina,<sup>54</sup> J. A. Kolb,<sup>54</sup> M. Lu,<sup>54</sup> R. Rahmat,<sup>54</sup> N. B. Sinev,<sup>54</sup> D. Strom,<sup>54</sup> J. Strube,<sup>54</sup> E. Torrence,<sup>54</sup> G. Castelli,<sup>55a,55b</sup> N. Gagliardi,<sup>55a,55b</sup> M. Margoni,<sup>55a,55b</sup> M. Morandin,<sup>55a</sup> M. Posocco,<sup>55a</sup> M. Rotondo,<sup>55a</sup> F. Simonetto,<sup>55a,55b</sup> R. Stroili,<sup>55a,55b</sup> C. Voci,<sup>55a,55b</sup> P. del Amo Sanchez,<sup>56</sup> E. Ben-Haim,<sup>56</sup> G. R. Bonneaud,<sup>56</sup> H. Briand,<sup>56</sup> J. Chauveau,<sup>56</sup> O. Hamon,<sup>56</sup> Ph. Leruste,<sup>56</sup> G. Marchiori,<sup>56</sup> J. Ocariz,<sup>56</sup> A. Perez,<sup>56</sup> J. Prendki,<sup>56</sup> S. Sitt,<sup>56</sup> L. Gladney,<sup>57</sup> M. Biasini,<sup>58a,58b</sup> E. Manoni,<sup>58a,58b</sup> C. Angelini,<sup>59a,59b</sup> G. Batignani,<sup>59a,59b</sup> S. Bettarini,<sup>59a,59b</sup> G. Calderini,<sup>59a,59b,¶</sup> M. Carpinelli,<sup>59a,59b,\*</sup> A. Cervelli,<sup>59a,59b</sup> F. Forti,<sup>59a,59b</sup> M. A. Giorgi,<sup>59a,59b</sup> A. Lusiani,<sup>59a,59c</sup> M. Morganti,<sup>59a,59b</sup> N. Neri,<sup>59a,59b</sup> E. Paoloni,<sup>59a,59b</sup> G. Rizzo,<sup>59a,59b</sup> J. J. Walsh,<sup>59a</sup> D. Lopes Pegna,<sup>60</sup> C. Lu,<sup>60</sup> J. Olsen,<sup>60</sup> A. J. S. Smith,<sup>60</sup> A. V. Telnov,<sup>60</sup> F. Anulli,<sup>61a</sup> E. Baracchini,<sup>61a,61b</sup> G. Cavoto,<sup>61a</sup> R. Faccini,<sup>61a,61b</sup> F. Ferrarotto,<sup>61a</sup> F. Ferroni,<sup>61a,61b</sup> M. Gaspero,<sup>61a,61b</sup> P. D. Jackson,<sup>61a</sup> L. Li Gioi,<sup>61a</sup> M. A. Mazzoni,<sup>61a</sup> S. Morganti,<sup>61a</sup> G. Piredda,<sup>61a</sup> F. Renga,<sup>61a,61b</sup> C. Voena,<sup>61a</sup> M. Ebert,<sup>62</sup> T. Hartmann,<sup>62</sup> H. Schröder,<sup>62</sup> R. Waldi,<sup>62</sup> T. Adye,<sup>63</sup> B. Franek,<sup>63</sup>

E. O. Olaiya,<sup>63</sup> F. F. Wilson,<sup>63</sup> S. Emery,<sup>64</sup> L. Esteve,<sup>64</sup> G. Hamel de Monchenault,<sup>64</sup> W. Kozanecki,<sup>64</sup> G. Vasseur,<sup>64</sup> Ch. Yèche,<sup>64</sup> M. Zito,<sup>64</sup> M. T. Allen,<sup>65</sup> D. Aston,<sup>65</sup> R. Bartoldus,<sup>65</sup> J. F. Benitez,<sup>65</sup> R. Cenci,<sup>65</sup> J. P. Coleman,<sup>65</sup> M. R. Convery,<sup>65</sup> J. C. Dingfelder,<sup>65</sup> J. Dorfan,<sup>65</sup> G. P. Dubois-Felsmann,<sup>65</sup> W. Dunwoodie,<sup>65</sup> R. C. Field,<sup>65</sup> M. Franco Sevilla,<sup>65</sup> A. M. Gabareen,<sup>65</sup> M. T. Graham,<sup>65</sup> P. Grenier,<sup>65</sup> C. Hast,<sup>65</sup> W. R. Innes,<sup>65</sup> J. Kaminski,<sup>65</sup> M. H. Kelsey,<sup>65</sup> H. Kim,<sup>65</sup> P. Kim,<sup>65</sup> M. L. Kocian,<sup>65</sup> D. W. G. S. Leith,<sup>65</sup> S. Li,<sup>65</sup> B. Lindquist,<sup>65</sup> S. Luitz,<sup>65</sup> V. Luth,<sup>65</sup> H. L. Lynch,<sup>65</sup> D. B. MacFarlane,<sup>65</sup> H. Marsiske,<sup>65</sup> R. Messner,<sup>65,\*</sup> D. R. Muller,<sup>65</sup> H. Neal,<sup>65</sup> S. Nelson,<sup>65</sup> C. P. O'Grady,<sup>65</sup> I. Ofte,<sup>65</sup> M. Perl,<sup>65</sup> B. N. Ratcliff,<sup>65</sup> A. Roodman,<sup>65</sup> A. A. Salnikov,<sup>65</sup> R. H. Schindler,<sup>65</sup> J. Schwiening,<sup>65</sup> A. Snyder,<sup>65</sup> D. Su,<sup>65</sup> M. K. Sullivan,<sup>65</sup> K. Suzuki,<sup>65</sup> S. K. Swain,<sup>65</sup> J. M. Thompson,<sup>65</sup> J. Va'vra,<sup>65</sup> A. P. Wagner,<sup>65</sup> M. Weaver,<sup>65</sup> C. A. West,<sup>65</sup> W. J. Wisniewski,<sup>65</sup> M. Wittgen,<sup>65</sup> D. H. Wright,<sup>65</sup> H. W. Wulsin,<sup>65</sup> A. K. Yarritu,<sup>65</sup> C. C. Young,<sup>65</sup> V. Ziegler,<sup>65</sup> X. R. Chen,<sup>66</sup> H. Liu,<sup>66</sup> W. Park,<sup>66</sup> M. V. Purohit,<sup>66</sup> R. M. White,<sup>66</sup> J. R. Wilson,<sup>66</sup> M. Bellis,<sup>67</sup> P. R. Burchat,<sup>67</sup> A. J. Edwards,<sup>67</sup> T. S. Miyashita,<sup>67</sup> S. Ahmed,<sup>68</sup> M. S. Alam,<sup>68</sup> J. A. Ernst,<sup>68</sup> B. Pan,<sup>68</sup> M. A. Saeed,<sup>68</sup> S. B. Zain,<sup>68</sup> A. Soffer,<sup>69</sup> S. M. Spanier,<sup>70</sup> B. J. Wogland,<sup>70</sup> R. Eckmann,<sup>71</sup> J. L. Ritchie,<sup>71</sup> A. M. Ruland,<sup>71</sup> C. J. Schilling,<sup>71</sup> R. F. Schwitters,<sup>71</sup> B. C. Wray,<sup>71</sup> B. W. Drummond,<sup>72</sup> J. M. Izen,<sup>72</sup> X. C. Lou,<sup>72</sup> F. Bianchi,<sup>73a,73b</sup> D. Gamba,<sup>73a,73b</sup> M. Pelliccioni,<sup>73a,73b</sup> M. Bomben,<sup>74a,74b</sup> L. Bosisio,<sup>74a,74b</sup> C. Cartaro,<sup>74a,74b</sup> G. Della Ricca,<sup>74a,74b</sup> L. Lanceri,<sup>74a,74b</sup> L. Vitale,<sup>74a,74b</sup> V. Azzolini,<sup>75</sup> N. Lopez-March,<sup>75</sup> F. Martinez-Vidal,<sup>75</sup> D. A. Milanes,<sup>75</sup> A. Oyanguren,<sup>75</sup> J. Albert,<sup>76</sup> Sw. Banerjee,<sup>76</sup> B. Bhuyan,<sup>76</sup> H. H. F. Choi,<sup>76</sup> K. Hamano,<sup>76</sup> G. J. King,<sup>76</sup> R. Kowalewski,<sup>76</sup> M. J. Lewczuk,<sup>76</sup> I. M. Nugent,<sup>76</sup> J. M. Roney,<sup>76</sup> R. J. Sobie,<sup>76</sup> T. J. Gershon,<sup>77</sup> P. F. Harrison,<sup>77</sup> J. Ilic,<sup>77</sup> T. E. Latham,<sup>77</sup> G. B. Mohanty,<sup>77</sup> E. M. T. Puccio,<sup>77</sup> H. R. Band,<sup>78</sup> X. Chen,<sup>78</sup> S. Dasu,<sup>78</sup> K. T. Flood,<sup>78</sup> Y. Pan,<sup>78</sup> R. Prepost,<sup>78</sup> C. O. Vuosalo,<sup>78</sup> and S. L. Wu<sup>78</sup>

(BABAR Collaboration)

<sup>1</sup>Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP), Université de Savoie, CNRS/IN2P3, F-74941 Annecy-Le-Vieux, France

<sup>2</sup>Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

<sup>3a</sup>INFN Sezione di Bari, I-70126 Bari, Italy

<sup>3b</sup>Dipartimento di Fisica, Università di Bari, I-70126 Bari, Italy

<sup>4</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway

<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

<sup>6</sup>University of Birmingham, Birmingham, B15 2TT, United Kingdom

<sup>7</sup>Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany

<sup>8</sup>University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

<sup>9</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>10</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

<sup>11</sup>University of California at Irvine, Irvine, California 92697, USA

<sup>12</sup>University of California at Riverside, Riverside, California 92521, USA

<sup>13</sup>University of California at San Diego, La Jolla, California 92093, USA

<sup>14</sup>University of California at Santa Barbara, Santa Barbara, California 93106, USA

<sup>15</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

<sup>16</sup>California Institute of Technology, Pasadena, California 91125, USA

<sup>17</sup>University of Cincinnati, Cincinnati, Ohio 45221, USA

<sup>18</sup>University of Colorado, Boulder, Colorado 80309, USA

<sup>19</sup>Colorado State University, Fort Collins, Colorado 80523, USA

<sup>20</sup>Technische Universität Dortmund, Fakultät Physik, D-44221 Dortmund, Germany

<sup>21</sup>Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

<sup>22</sup>Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

<sup>23</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

<sup>24a</sup>INFN Sezione di Ferrara, I-44100 Ferrara, Italy

<sup>24b</sup>Dipartimento di Fisica, Università di Ferrara, I-44100 Ferrara, Italy

<sup>25</sup>INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy

<sup>26a</sup>INFN Sezione di Genova, I-16146 Genova, Italy

<sup>26b</sup>Dipartimento di Fisica, Università di Genova, I-16146 Genova, Italy

<sup>27</sup>Harvard University, Cambridge, Massachusetts 02138, USA

<sup>28</sup>Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany

<sup>29</sup>Humboldt-Universität zu Berlin, Institut für Physik, Newtonstrasse 15, D-12489 Berlin, Germany

<sup>30</sup>Imperial College London, London, SW7 2AZ, United Kingdom

<sup>31</sup>University of Iowa, Iowa City, Iowa 52242, USA

- <sup>32</sup>*Iowa State University, Ames, Iowa 50011-3160, USA*
- <sup>33</sup>*Johns Hopkins University, Baltimore, Maryland 21218, USA*
- <sup>34</sup>*Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, Boîte Postale 34, F-91898 Orsay Cedex, France*
- <sup>35</sup>*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*
- <sup>36</sup>*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- <sup>37</sup>*Queen Mary, University of London, London, E1 4NS, United Kingdom*
- <sup>38</sup>*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*
- <sup>39</sup>*University of Louisville, Louisville, Kentucky 40292, USA*
- <sup>40</sup>*Johannes Gutenberg-Universität Mainz, Institut für Kernphysik, D-55099 Mainz, Germany*
- <sup>41</sup>*University of Manchester, Manchester M13 9PL, United Kingdom*
- <sup>42</sup>*University of Maryland, College Park, Maryland 20742, USA*
- <sup>43</sup>*University of Massachusetts, Amherst, Massachusetts 01003, USA*
- <sup>44</sup>*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA*
- <sup>45</sup>*McGill University, Montréal, Québec, Canada H3A 2T8*
- <sup>46a</sup>*INFN Sezione di Milano, I-20133 Milano, Italy*
- <sup>46b</sup>*Dipartimento di Fisica, Università di Milano, I-20133 Milano, Italy*
- <sup>47</sup>*University of Mississippi, University, Mississippi 38677, USA*
- <sup>48</sup>*Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7*
- <sup>49</sup>*Mount Holyoke College, South Hadley, Massachusetts 01075, USA*
- <sup>50a</sup>*INFN Sezione di Napoli, I-80126 Napoli, Italy*
- <sup>50b</sup>*Dipartimento di Scienze Fisiche, Università di Napoli Federico II, I-80126 Napoli, Italy*
- <sup>51</sup>*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*
- <sup>52</sup>*University of Notre Dame, Notre Dame, Indiana 46556, USA*
- <sup>53</sup>*Ohio State University, Columbus, Ohio 43210, USA*
- <sup>54</sup>*University of Oregon, Eugene, Oregon 97403, USA*
- <sup>55a</sup>*INFN Sezione di Padova, I-35131 Padova, Italy*
- <sup>55b</sup>*Dipartimento di Fisica, Università di Padova, I-35131 Padova, Italy*
- <sup>56</sup>*Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France*
- <sup>57</sup>*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- <sup>58a</sup>*INFN Sezione di Perugia, I-06100 Perugia, Italy*
- <sup>58b</sup>*Dipartimento di Fisica, Università di Perugia, I-06100 Perugia, Italy*
- <sup>59a</sup>*INFN Sezione di Pisa, I-56127 Pisa, Italy*
- <sup>59b</sup>*Dipartimento di Fisica, Università di Pisa, I-56127 Pisa, Italy*
- <sup>59c</sup>*Scuola Normale Superiore di Pisa, I-56127 Pisa, Italy*
- <sup>60</sup>*Princeton University, Princeton, New Jersey 08544, USA*
- <sup>61a</sup>*INFN Sezione di Roma, I-00185 Roma, Italy*
- <sup>61b</sup>*Dipartimento di Fisica, Università di Roma La Sapienza, I-00185 Roma, Italy*
- <sup>62</sup>*Universität Rostock, D-18051 Rostock, Germany*
- <sup>63</sup>*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*
- <sup>64</sup>*CEA, Irfu, SPP, Centre de Saclay, F-91191 Gif-sur-Yvette, France*
- <sup>65</sup>*SLAC National Accelerator Laboratory, Stanford, California 94309 USA*
- <sup>66</sup>*University of South Carolina, Columbia, South Carolina 29208, USA*
- <sup>67</sup>*Stanford University, Stanford, California 94305-4060, USA*
- <sup>68</sup>*State University of New York, Albany, New York 12222, USA*
- <sup>69</sup>*Tel Aviv University, School of Physics and Astronomy, Tel Aviv, 69978, Israel*
- <sup>70</sup>*University of Tennessee, Knoxville, Tennessee 37996, USA*
- <sup>71</sup>*University of Texas at Austin, Austin, Texas 78712, USA*
- <sup>72</sup>*University of Texas at Dallas, Richardson, Texas 75083, USA*
- <sup>73a</sup>*INFN Sezione di Torino, I-10125 Torino, Italy*
- <sup>73b</sup>*Dipartimento di Fisica Sperimentale, Università di Torino, I-10125 Torino, Italy*
- <sup>74a</sup>*INFN Sezione di Trieste, I-34127 Trieste, Italy*
- <sup>74b</sup>*Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy*
- <sup>75</sup>*IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*
- <sup>76</sup>*University of Victoria, Victoria, British Columbia, Canada V8W 3P6*
- <sup>77</sup>*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*
- <sup>78</sup>*University of Wisconsin, Madison, Wisconsin 53706, USA*

(Received 2 December 2009; published 28 July 2010)

Using  $467 \text{ fb}^{-1}$  of  $e^+e^-$  annihilation data collected with the *BABAR* detector, we measure  $\frac{\mathcal{B}(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)}{\mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)} = (0.9796 \pm 0.0016 \pm 0.0036)$ ,  $\frac{\mathcal{B}(\tau^- \rightarrow \pi^- \nu_\tau)}{\mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)} = (0.5945 \pm 0.0014 \pm 0.0061)$ , and  $\frac{\mathcal{B}(\tau^- \rightarrow K^- \nu_\tau)}{\mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)} = (0.03882 \pm 0.00032 \pm 0.00057)$ , where the uncertainties are statistical and systematic, respectively. From these precision  $\tau$  measurements, we test the standard model assumption of  $\mu - e$  and  $\tau - \mu$  charge current lepton universality and provide determinations of  $|V_{us}|$  experimentally independent of the decay of a kaon.

DOI: 10.1103/PhysRevLett.105.051602

PACS numbers: 11.30.Hv, 12.15.Hh, 13.35.Dx, 14.60.Fg

Decays of the  $\tau$  lepton to a single charged particle and neutrino(s) probe the standard model (SM) predictions of charged current lepton universality and the unitarity relation of the first row of the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix [1]. Previous measurements of universality [2,3], expressible in terms of the coupling strength ( $g_\ell$ ) of lepton of flavor  $\ell$  to the charged gauge boson of the electroweak interaction are in agreement with the SM where  $g_\tau/g_\mu = g_\mu/g_e = 1$ . Similarly, kaon decay measurements [3,4] sensitive to  $|V_{us}|$ , the relative weak coupling between up and strange quarks, yield a value consistent with unitarity ( $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$ ), where nuclear beta decays provide  $|V_{ud}|$  [5] and  $|V_{ub}|$  is negligible [3]. However, new physics that couples primarily to the third generation could be revealed through deviations from the SM in precision universality and  $|V_{us}|$  measurements involving the  $\tau$ . Significant deviations of this nature are unambiguous signatures of new physics that provide crucial but complimentary information to the direct searches for Higgs bosons [6] and other new physics models with, e.g., lepto-quarks [7], heavy gauge  $W'$  or  $Z'$  bosons, heavy quarks or leptons, compositeness or extra dimensions [8].

Recent measurements of the sum of strange  $\tau$  branching fractions interpreted in the framework of the operator product expansion (OPE) and finite energy sum rules yield a value of  $|V_{us}|$  that is approximately 3 standard deviations ( $\sigma$ ) lower than expectations from CKM unitarity [9]. This Letter addresses both experimental and theoretical aspects of this question by providing the first precision measurements of  $R_K \equiv \frac{\mathcal{B}(\tau^- \rightarrow K^- \nu_\tau)}{\mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)}$  [10] and  $R_{K/\pi} \equiv \frac{\mathcal{B}(\tau^- \rightarrow K^- \nu_\tau)}{\mathcal{B}(\tau^- \rightarrow \pi^- \nu_\tau)}$  enabled by the unique combination of a very large  $\tau$  sample with particle momenta amenable to particle identification using Cherenkov radiation. By using values of the meson decay constants from lattice QCD [11], we provide two precision determinations of  $|V_{us}|$  from  $\tau$  decays independent of the OPE framework. We also report on new measurements of  $R_\pi \equiv \frac{\mathcal{B}(\tau^- \rightarrow \pi^- \nu_\tau)}{\mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)}$  and  $R_\mu \equiv \frac{\mathcal{B}(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)}{\mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)}$ .  $R_\mu$  provides an improved measurement of  $g_\mu/g_e$  whereas  $R_\pi$  and  $R_K$ , when compared to the muonic branching fractions of the pion and kaon, yield improved measurements of  $g_\tau/g_\mu$  involving pseudoscalar mesons.

The data sample corresponds to an integrated luminosity of  $\mathcal{L} = 467 \text{ fb}^{-1}$  recorded at an  $e^+e^-$  center-of-mass

(CM) energy ( $\sqrt{s}$ ) near 10.58 GeV and was collected with the *BABAR* detector at the SLAC PEP-II  $e^+e^-$  storage rings. With a luminosity-weighted average cross section of  $\sigma_{e^+e^- \rightarrow \tau^+\tau^-} = (0.919 \pm 0.003) \text{ nb}$  [12,13], this corresponds to the production of  $4.29 \times 10^8$   $\tau$ -pair events. The *BABAR* detector [14] is composed of a silicon vertex tracker, drift chamber (DCH), ring-imaging Cherenkov detector (DIRC), and electromagnetic calorimeter (EMC), all contained in a 1.5-T solenoid. The iron flux return for the solenoid is instrumented (IFR) to identify muons.

Tau-pair events are simulated with the KK Monte Carlo (MC) generator [13], which includes higher-order radiative corrections. We simulate  $\tau$  decays with TAUOLA [15] and PHOTOS [16] using measured branching fractions [3]. The detector response is simulated with GEANT4 [17]. Simulated events for signal as well as background processes [13,15,16,18,19] are reconstructed in the same manner as data. The MC samples are used for selection optimization, control sample studies, and systematic error studies. The number of simulated nonsignal events is comparable to the number expected in the data, with the exception of Bhabha and two-photon events, which are not simulated but which data studies show to be negligible.

We study  $e^+e^- \rightarrow \tau^+\tau^-$  events with the  $\tau^-$  decaying via  $\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau$ ,  $\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau$ ,  $\tau^- \rightarrow \pi^- \nu_\tau$  or  $\tau^- \rightarrow K^- \nu_\tau$  modes and the  $\tau^+$  decaying via a  $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \bar{\nu}_\tau$  tagging channel with the selection criteria optimized to minimize the combined statistical and systematic uncertainties [20]. The number of signal events for decay modes  $i = \{e, \mu, \pi, K\} = \{e^- \bar{\nu}_e \nu_\tau, \mu^- \bar{\nu}_\mu \nu_\tau, \pi^- \nu_\tau, K^- \nu_\tau\}$  are  $N_i^S = \mathcal{E}_i^{-1}(N_i^D - N_i^B)$  where  $\mathcal{E}_i$  is the efficiency (including  $\mathcal{B}(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau) = (8.85 \pm 0.13)\%$  [3]),  $N_i^D$  the number of selected data events, and  $N_i^B$  the estimated number of background events for the  $i$ th mode.

We measure the ratios  $R_i = N_i^S/N_e^S$  which normalizes to the most precisely known relevant SM process available, and in which several common sources of systematic uncertainty cancel.  $N_i^D$  are multiplied with reproducible random numbers until all efficiency and uncertainty estimates are finalized. Once unblinded, we use the values of the three branching ratios to update world averages of the branching fractions, which we then use to recalculate the backgrounds for our final results.

Events with a net charge of zero and with four well-reconstructed tracks not originating from the conversion of

a photon in the detector material are selected. For good particle identification, each track is required to be within the acceptance of the DIRC and EMC, and have a transverse momentum greater than 0.25 GeV to ensure that it reaches the DIRC. The plane normal to the thrust axis divides the event into hemispheres in the CMframe. The “signal” hemisphere contains a single track and the “tag” hemisphere the other three tracks.

Each tag hemisphere track is required to be consistent with being a pion and the energy deposited in the EMC unassociated with any tracks in this hemisphere is required to be less than 0.20 GeV. Also, events that contain track pairs consistent with coming from a  $K_S^0$  are vetoed.

The signal track momentum is required to lie between 1 and 4 GeV/ $c$ . Information from the five detector subsystems is combined in likelihood selectors which identify  $e$ ,  $\pi$ , and  $K$  particles and in a neural network which identifies muons. The  $\pi - K$  separation is provided by the DIRC and DCH whereas  $\pi - \mu$  separation is primarily accomplished with the IFR and EMC. The identification efficiencies are given in Table I and cross-contaminations are given below. We suppress di-muon and Bhabha backgrounds by requiring signal tracks identified as a lepton to have CMmomentum less than 80% of  $\sqrt{s}/2c$ . To reduce cross-feed from  $e$  into the  $\pi$  and  $K$  channels, the ratio of deposited electromagnetic energy of a  $\pi$  or  $K$  candidate track to its measured momentum,  $E/pc$ , is required to be less than 0.85. A pion track also passing a loose muon selection is rejected. A similar veto is applied for a kaon track passing the loose muon selection if its measured momentum exceeds 3 GeV/ $c$ . Also, events with an EMC energy  $>\{1.0, 0.5, 0.2, 0.2\}$  GeV in the signal hemisphere unassociated with the  $\{e, \mu, \pi, K\}$  track are removed.

Pion and kaon control samples from  $D^{*+} \rightarrow \pi^+ D^0$ ,  $D^0 \rightarrow \pi^+ K^-$  decays are used to study and correct for small differences between MC and data. We cross-check

these with independent  $\pi^+$  ( $K^-$ ) control samples from  $\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau$  ( $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$ ) decays using particle identification of two of the oppositely charged particles and the fact that the wrong sign  $\tau^- \rightarrow \pi^- \pi^- K^+ \nu_\tau$  decays are heavily suppressed. Samples of radiative Bhabha and radiative  $\mu$ -pair events provide control samples of electrons and muons. The systematic uncertainty associated with charged particle identification is assessed from the control sample statistical errors, consistency between control samples, and the sensitivity of the control sample corrections to the number of particles near the track. The statistical errors in the more limited cross-check control samples dominate these errors. Because we use control samples to correct charge conjugate particles separately, charge-dependent detector responses are accounted for by construction.

To remove two-photon and Bhabha backgrounds, the event must have a missing CMenergy between 10% and 70% of  $\sqrt{s}$ . The angle between the missing momentum and electron beam direction in the CM,  $\theta_{\text{miss}}^{\text{CM}}$ , is constrained to satisfy  $|\cos(\theta_{\text{miss}}^{\text{CM}})| < 0.7$ , the thrust of the event is required to be above 0.9, and the net missing transverse momentum in the CMgreater than  $0.009\sqrt{s}/c$ .

Each of the three tagside tracks has an electron veto applied to further reduce the Bhabha contamination. This results in less than 0.03% contamination from two-photon events and less than 0.1% contamination from Bhabha events in the electron signal sample. These backgrounds were investigated by studying samples enriched in Bhabha and two-photon events by adjusting the requirements on the thrust,  $\cos(\theta_{\text{miss}}^{\text{CM}})$ , and transverse momentum of the event. Potential background from Bhabha events were further probed by studying the number of events having a high signal track momentum as the electron veto was progressively lifted from one, then two, and finally all three tracks in the tag hemisphere.

TABLE I. Number of selected events, purity, total efficiency, component of the efficiency from particle identification, and systematic uncertainties (in %) on  $R_i$  for each decay mode.

	$\mu$	$\pi$	$K$
$N^{\text{D}}$	731 102	369 091	25 123
Purity	97.3%	78.7%	76.6%
Total efficiency	0.485%	0.324%	0.330%
Particle ID efficiency	74.5%	74.6%	84.6%
Systematic uncertainties:			
Particle ID	0.32	0.51	0.94
Detector response	0.08	0.64	0.54
Backgrounds	0.08	0.44	0.85
Trigger	0.10	0.10	0.10
$\pi^- \pi^- \pi^+$ modeling	0.01	0.07	0.27
Radiation	0.04	0.10	0.04
$\mathcal{B}(\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau)$	0.05	0.15	0.40
$\mathcal{L}\sigma_{e^+ e^- \rightarrow \tau^+ \tau^-}$	0.02	0.39	0.20
Total [%]	0.36	1.0	1.5

To suppress backgrounds in the  $\tau^- \rightarrow \pi^- \nu_\tau$  and  $\tau^- \rightarrow K^- \nu_\tau$  channels from  $\tau$  decays with undetected neutral particles other than the  $\nu_\tau$  (e.g.,  $K_L^0$  mesons,  $\nu_\mu$ ), we reconstruct the direction of the back-to-back  $\tau^+ \tau^-$  system in the CMframe. The polar angle of the  $\tau$  momentum with respect to the tagside hadronic system is calculated assuming that the CMenergy of the  $\tau$  is  $\sqrt{s}/2$ , and the azimuthal angle of the  $\tau$  momentum is fixed to a value that has been optimized to minimize the total error on  $\mathcal{B}_{K/\pi}$  [20]. With this estimator for the  $\tau$  momentum, we require the missing mass in the signal hemisphere to be less than  $0.56 \text{ GeV}/c^2$ .

For the selected  $\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau$  events, the dominant backgrounds are  $\tau^- \rightarrow \pi^- \nu_\tau$  ( $1.46 \pm 0.01\%$ ) and  $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$  ( $0.85 \pm 0.01\%$ ). For the  $\tau^- \rightarrow \pi^- \nu_\tau$  channel, the dominant backgrounds are  $\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau$  ( $12.90 \pm 0.07\%$ ),  $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$  ( $5.87 \pm 0.04\%$ ), and non- $\tau$  backgrounds ( $0.34 \pm 0.05\%$ ). The major backgrounds in the  $\tau^- \rightarrow K^- \nu_\tau$  channel are from  $\tau^- \rightarrow \pi^- \nu_\tau$  decays ( $10.06 \pm 0.13\%$ ),  $\tau^- \rightarrow K^- K_L^0 \nu_\tau$  ( $3.87 \pm 0.41\%$ ),  $\tau^- \rightarrow K^- \pi^0 \nu_\tau$  ( $1.97 \pm 0.14\%$ ),  $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$  ( $1.07 \pm 0.06\%$ ), and non- $\tau$  backgrounds ( $2.58 \pm 0.38\%$ ). The uncertainties are from MC statistics, branching fractions and, for non- $\tau$  backgrounds, the systematic uncertainty on background rates. Figure 1 shows the momentum distributions in the




FIG. 1 (color online). Data (points) and MC (histograms) distributions of CM momentum for (a)  $\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau$ , (b)  $\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau$ , (c)  $\tau^- \rightarrow \pi^- \nu_\tau$  and (d)  $\tau^- \rightarrow K^- \nu_\tau$  modes. The small differences between MC and data are accounted for in the systematic errors.

CMframe for each of the four decay modes for data, along with the background MC contributions.

For the  $\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau$  channel, 884426 events are selected with an efficiency and purity of  $(0.589 \pm 0.010)\%$  and  $(99.69 \pm 0.06)\%$ , respectively. The number of selected events, efficiency, purity, and systematic uncertainties on  $R_i$  of the  $\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau$ ,  $\tau^- \rightarrow \pi^- \nu_\tau$ , and  $\tau^- \rightarrow K^- \nu_\tau$  selections are presented in Table I. These uncertainties include contributions from the particle identification, the sensitivity to detector response including the impact of changing the MC momentum scale and DCH resolution, modeling of hadronic and electromagnetic showers in the EMC, the EMC energy scale, and angular measurements made by these detectors within their modelling uncertainties, the backgrounds, initial- and final-state radiation, radiation in  $\tau$  decays, rate and shape of  $\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau$  decays, the trigger, and  $\mathcal{L} \sigma_{e^+ e^- \rightarrow \tau^+ \tau^-}$ . The systematic uncertainty on  $R_\mu$  is dominated by uncertainties in particle identification. The  $R_\pi$  and  $R_K$  measurements have additional dominant contributions from the detector modelling and associated backgrounds, due to stronger cuts on the EMC energy necessary to reduce non- $\tau$  backgrounds. Presence of the  $\sim 20\%$  backgrounds in these channels render them more sensitive to the modelling of the tagside decays. The dominant background uncertainty in the  $R_\pi$  measurement arises from the electron contamination in the  $\pi$  sample investigated by measuring the number of events that fail the  $E/p$  electron veto requirement in data and MC. In the  $R_K$  event sample, the uncertainty arising from  $\tau$  decay branching fractions of background modes is  $0.58\%$ , which is dominated by the uncertainty of the  $\tau^- \rightarrow K_L^0 K^- \nu_\tau$  fraction. There is also a  $0.49\%$  uncertainty assigned for  $q\bar{q}$  backgrounds, which are studied using events with an invariant mass of the tracks in the tag hemisphere above the  $\tau$  mass and cross-checked in regions of thrust and  $\cos(\theta_{\text{miss}}^{\text{CM}})$  enriched with these backgrounds.

The measured branching ratios and fractions are

$$R_\mu = (0.9796 \pm 0.0016 \pm 0.0036),$$

$$R_\pi = (0.5945 \pm 0.0014 \pm 0.0061),$$

$$R_K = (0.03882 \pm 0.00032 \pm 0.00057),$$

$$R_h = R_\pi + R_K = (0.6333 \pm 0.0014 \pm 0.0061),$$

$$\mathcal{B}(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau) = (17.46 \pm 0.03 \pm 0.08)\%$$

$$\mathcal{B}(\tau^- \rightarrow \pi^- \nu_\tau) = (10.59 \pm 0.03 \pm 0.11)\%$$

$$\mathcal{B}(\tau^- \rightarrow K^- \nu_\tau) = (0.692 \pm 0.006 \pm 0.010)\%, \quad (1)$$

where  $h = \pi$  or  $K$  and we use  $\mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau) = (17.82 \pm 0.05)\%$  [3]. The off-diagonal elements of the correlation matrix for the measured ratios (branching fractions) are  $\rho_{\mu\pi} = 0.25$  (0.34),  $\rho_{\mu K} = 0.12$  (0.20), and  $\rho_{\pi K} = 0.33$  (0.36). The  $\mu$  and  $\pi$  measurements are con-

sistent with and of comparable precision as the world averages [3], whereas the  $K$  measurement is consistent with but twice as precise as the world average [3].

Tests of  $\mu - e$  universality can be expressed as

$$\left(\frac{g_\mu}{g_e}\right)_\tau = \frac{\mathcal{B}(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau) f(m_e^2/m_\tau^2)}{\mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau) f(m_e^2/m_\tau^2)},$$

where  $f(x) = 1 - 8x + 8x^3 - x^4 - 12x^2 \log x$ , assuming that the neutrino masses are negligible [21]. This gives  $\left(\frac{g_\mu}{g_e}\right)_\tau = 1.0036 \pm 0.0020$ , yielding a new world average of  $1.0018 \pm 0.0014$ , which is consistent with the SM and the value of  $1.0021 \pm 0.0015$  from pion decays [3,22].

Tau-muon universality is tested with

$$\left(\frac{g_\tau}{g_\mu}\right)_h = \frac{\mathcal{B}(\tau \rightarrow h \nu_\tau)}{\mathcal{B}(h \rightarrow \mu \nu_\mu)} \frac{2m_h m_\mu^2 \tau_h}{(1 + \delta_h) m_\tau^3 \tau_\tau} \left(\frac{1 - m_\mu^2/m_h^2}{1 - m_\tau^2/m_\tau^2}\right)^2,$$

where the radiative corrections are  $\delta_\pi = (0.16 \pm 0.14)\%$  and  $\delta_K = (0.90 \pm 0.22)\%$  [23]. Using the world averaged mass and lifetime values and meson decay rates [3], we determine  $\left(\frac{g_\tau}{g_\mu}\right)_{\pi(K)} = 0.9856 \pm 0.0057$  ( $0.9827 \pm 0.0086$ ) and  $\left(\frac{g_\tau}{g_\mu}\right)_h = 0.9850 \pm 0.0054$  when combining these results; this is  $2.8\sigma$  below the SM expectation and within  $2\sigma$  of the world average.

We use the kaon decay constant  $f_K = 157 \pm 2$  MeV [11], and our value of

$$\mathcal{B}(\tau^- \rightarrow K^- \nu_\tau) = \frac{G_F^2 f_K^2 |V_{us}|^2 m_\tau^3 \tau_\tau}{16\pi\hbar} \left(1 - \frac{m_K^2}{m_\tau^2}\right)^2 S_{EW},$$

where  $S_{EW} = 1.0201 \pm 0.0003$  [24], to determine  $|V_{us}| = 0.2193 \pm 0.0032$ . This measurement is within  $2\sigma$  of the value of  $0.2255 \pm 0.0010$  predicted by CKM unitarity and is also consistent with the value of  $|V_{us}| = 0.2165 \pm 0.0027$  derived from the inclusive sum of strange  $\tau$  decays [9].

Both of our measured  $|V_{us}|$  values depend on absolute strange decay rates. Our value of  $R_{K/\pi} = (0.06531 \pm 0.00056 \pm 0.00093)$ , however, provides a  $|V_{us}|$  value driven by the ratio between strange and nonstrange decays. We use  $f_K/f_\pi = 1.189 \pm 0.007$  [11],  $|V_{ud}|$  [5], and the long-distance correction  $\delta_{LD} = (0.03 \pm 0.44)\%$  estimated [25] using corrections to  $\tau \rightarrow h\nu$  and  $h \rightarrow \mu\nu$  [23,26] in

$$R_{K/\pi} = \frac{f_K^2 |V_{us}|^2}{f_\pi^2 |V_{ud}|^2} \frac{(1 - \frac{m_K^2}{m_\tau^2})^2}{(1 - \frac{m_\pi^2}{m_\tau^2})^2} (1 + \delta_{LD}),$$

to obtain  $|V_{us}| = 0.2255 \pm 0.0024$  where short-distance electro-weak corrections cancel in this ratio. This value is consistent with CKM unitarity [5] and  $2.5\sigma$  higher than  $|V_{us}|$  from the inclusive sum of strange  $\tau$  decays.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organiza-

tions that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MES (Russia), MEC (Spain), and STFC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

\*Deceased.

†Present address: Temple University, Philadelphia, Pennsylvania 19122, USA.

‡Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy.

§Also with Università di Roma La Sapienza, I-00185 Roma, Italy.

||Present address: University of South AL, Mobile, AL 36688, USA.

¶Also with Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France.

\*\*Also with Università di Sassari, Sassari, Italy.

- [1] N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
- [2] A. Pich, *Nucl. Phys. B, Proc. Suppl.* **181–182**, 300 (2008).
- [3] C. Amsler *et al.* (Particle Data Group), *Phys. Lett. B* **667**, 1 (2008); B. Aubert *et al.* (*BABAR* Collaboration), *Phys. Rev. Lett.* **100**, 011801 (2008).
- [4] F. Ambrosino *et al.* (KLOE Collaboration), *J. High Energy Phys.* 04 (2008) 059,  $|V_{us}| = 0.2237 \pm 0.0013$ ; M. Antonelli *et al.* (FlaviaNet Kaon Group), *Nucl. Phys. B, Proc. Suppl.* **181–182**, 83 (2008).
- [5] J. C. Hardy and I. S. Towner, *Phys. Rev. C* **79**, 055502 (2009),  $|V_{ud}| = 0.97425 \pm 0.00022$ .
- [6] H. E. Logan and D. MacLennan, *Phys. Rev. D* **79**, 115022 (2009); M. Krawczyk and D. Temes, *Eur. Phys. J. C* **44**, 435 (2005).
- [7] I. Dorsner, S. Fajfer, J. F. Kamenik, and N. Kosnik, *Phys. Lett. B* **682**, 67 (2009).
- [8] W. Loinaz, N. Okamura, T. Takeuchi, and L. C. R. Wijewardhana, *Phys. Rev. D* **67**, 073012 (2003); A. Czarnecki, W. J. Marciano, and A. Sirlin, *Phys. Rev. D* **70**, 093006 (2004); W. J. Marciano, *Proc. Sci. KAON* (2008) 003.
- [9] E. Gamiz, M. Jamin, A. Pich, J. Prades, and F. Schwab, *Proc. Sci. KAON* (2008) 008; K. Maltman, C. E. Wolfe, S. Banerjee, I. Nugent, and J. M. Roney, *Int. J. Mod. Phys. A* **23**, 3191 (2008).
- [10] Charge conjugate  $\tau$  decays are implied throughout.
- [11] E. Follana, C. T. H. Davies, G. P. Lepage, and J. Shigemitsu, *Phys. Rev. Lett.* **100**, 062002 (2008).
- [12] S. Banerjee, B. Pietrzyk, J. M. Roney, and Z. Was, *Phys. Rev. D* **77**, 054012 (2008).

- [13] S. Jadach, B.F.L. Ward, and Z. Was, *Comput. Phys. Commun.* **130**, 260 (2000).
- [14] B. Aubert *et al.* (BABAR Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 1 (2002).
- [15] S. Jadach, Z. Was, R. Decker, and J.H. Kühn, *Comput. Phys. Commun.* **76**, 361 (1993).
- [16] E. Barberio and Z. Was, *Comput. Phys. Commun.* **79**, 291 (1994).
- [17] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [18] D.J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [19] T. Sjöstrand, *Comput. Phys. Commun.* **82**, 74 (1994).
- [20] I.M. Nugent, Ph.D. thesis, University of Victoria [Institution Report No. SLAC-R-936, 2008].
- [21] Y.S. Tsai, *Phys. Rev. D* **4**, 2821 (1971); **13**, 771(E) (1976).
- [22] V. Cirigliano and I. Rosell, *J. High Energy Phys.* **10** (2007) 005.
- [23] W.J. Marciano and A. Sirlin, *Phys. Rev. Lett.* **71**, 3629 (1993); R. Decker and M. Finkemeier, *Nucl. Phys.* **B438**, 17 (1995); R. Decker and M. Finkemeier, *Phys. Lett. B* **334**, 199 (1994).
- [24] J. Erler, *Rev. Mex. Fis.* **50**, 200 (2004).
- [25] S. Banerjee, *Proc. of ICHEP08*, eConf C080730 (2008).
- [26] W.J. Marciano, *Phys. Rev. Lett.* **93**, 231803 (2004).