

Crystals are not just a sideline at ISOMET. Years of research and development have enabled us to produce consistently the highest quality crystals obtainable and at attractive prices. We go all out to serve you with

HIGHEST QUALITY ATTRACTIVE PRICES • PROMPT DELIVERY --three good reasons why leading spectroscopists everywhere are turning to ISOMET for their crystal needs. Why don't you try us on your next order.

NaCl, KBr, KCl, Kl, CsBr, Csl, BaF₂, MgO Rough blanks, polished windows, prisms, special shapes. Standard liquid and gas cells; special cells. Repair and polishing service for cells and windows.

KBr Pellet Powder.



GOODYEAR ATOMIC CORPORATION

Has Opening For Specialists In:

- · Solid State Physics
- Physical Chemistry

and/or

- Electron Microscopy and
 - anu
- Electron Diffraction

To work on research and development projects in the fields of thin films, crystal perfection and corrosion.

SUBMIT RESUME AND SALARY REQUIREMENTS TO:

EMPLOYMENT DEPARTMENT

GOODYEAR ATOMIC CORPORATION

> P.O. Box 628 Portsmouth, Ohio

Meetings

Quantitative Spectroscopy

TWO-DAY symposium on quantitative spectroscopy with selected military applications was held at the California Institute of Technology on March 28 and 29, 1960. The plans for the meeting were laid in Washington during December 1959 as the result of discussions between S. S. Penner of Caltech and J. W. Bond of the Advanced Research Projects Agency (ARPA). A review of studies connected with long-range missile detection and the physics of missile re-entry suggested that the enormous scientific and technical effort in this field could be planned more effectively and could be executed more efficiently through better utilization of people, facilities, and funds. The basic difficulty appeared to be associated with the tendency to collect essentially uninterpretable data: the measurements were often qualitative and were obtained in such a way that they were not useful either for predicting observable data on missiles under new test conditions or for understanding quantitatively the nature of the physical processes under study. What appeared to be needed was a more deliberate effort to reduce all missile radiation data to quantitative terms, an objective that could perhaps be furthered by a very special type of conference, namely, a symposium on quantitative spectroscopy at which the military applications represented nothing more than a special use of fundamental information.

Discussions with sponsors of military and basic spectroscopic research in various government agencies showed enthusiastic support for this special type of meeting and the final program enjoyed active support from Caltech (S. S. Penner), ARPA (J. W. Bond), the Office of Naval Research (S. Silverman and A. R. Laufer), the Air Force Office of Scientific Research (W. J. Otting), and the Air Force Cambridge Research Center (J. N. Howard). A. R. Laufer of the Pasadena Office of Naval Research did most of the work of arranging the conference; the technical program was planned by S. S. Penner with assistance from the other members of the organizing committee.

Lee A. DuBridge, president of the California Institute of Technology, opened the two-day symposium, expressing his pleasure at noting the participation of former colleagues at a meeting on classical physics that cut clearly across the borders of conventional academic disciplines.

The first session, under the chairmanship of S. S. Penner, was dominated by molecular spectroscopists.

PHYSICS TODAY

W. S. Benedict of Johns Hopkins University discussed absolute intensity and line-width measurements of individual spectral lines in the infrared region; his beautiful experimental data dispelled at once the notion that the widths and contours of spectral lines in vibrationrotation bands could be represented by a single "effective" shape parameter or that the temperature dependence of line widths could be correlated by a simple collision theory. B. L. Crawford, Jr., of the University of Minnesota discussed techniques for the determination of absolute intensity measurements for entire vibration-rotation bands by the method which he chose to denote, "with total disregard for the valencies of tungsten", as the "WaP method" (Wilson-Wells-Penner-Weber method). In this method, the test gas is pressurized with sufficient inert gas to smear out the rotational line structure, and the resulting integrated intensity data are then extrapolated to zero optical depth. The pitfalls involved in data reduction and precautions required in good experimental procedure were clearly defined. Absolute intensity measurements using the same procedure in the visible and ultraviolet regions of the spectrum for NO were considered by G. W. Bethke of the General Electric Company. D. H. Rank of the University of Pennsylvania told of the significant progress made in his laboratory in attaining extremely high-resolution spectra in the infrared. His methods of scanning spectra with etalons and diffraction gratings were covered. The first morning session was concluded by C. E. Treanor of the Cornell Aeronautical Laboratory with a discussion of experimental results of absolute intensity measurements on O2 obtained in shock-tube studies performed in collaboration with W. H. Wurster.

The second session, chaired by A. R. Laufer, opened with a stimulating survey of the theories of line broadening by H. Margenau of Yale University: in 45 minutes the theories of natural, Doppler, pressure, and statistical line broadening were reviewed with admirable clarity and enthusiasm. Illustrative examples of astrophysical applications of quantitative spectroscopy were described by J. L. Greenstein of Caltech who noted a number of unusual features of astronomical spectra. The use of an atomic beam apparatus for the determination of absolute intensities of metallic spectral lines was considered by R. B. King of Caltech who reviewed the results of representative measurements made in his laboratory. In the concluding paper, D. H. Menzel presented an outline of a very successful screening theory developed with D. A. Layzer for application to the calculation of energy levels and transition probabilities of complex atoms.

The early evening was devoted to relaxation, including an industry-sponsored cocktail hour followed by a banquet at the Huntington-Sheraton Hotel. Nobel Laureate George W. Beadle, acting dean of the faculty at Caltech, concluded the educational activities of the first day with an after-dinner talk on "The Nature of Genetic Information". One cannot help but wonder if this choice of topic was designed to put additional

Think Clean

When you think of high vacuums, you have to think clean...and if you think of <u>exceptionally</u> clean vacuums, without fluids or other contaminants, you have to think of UlteVac electronic vacuum pumps — made by Ultek, the only manufacturer devoted exclusively to the technology of fluidless vacuum pumping. UlteVac pumps, using no moving parts, hot filaments, or refrigeration, produce vacuums to 10⁻⁹ mm Hg and below; operate unattended for months, invulnerable to power failure. System vacuum automatically measured.







NOW! a low cost Michelson INTERFEROMETER

This precision instrument is ingeniously designed to provide .2% accuracy at minimum cost. Directly calibrated in inches. Clear, well defined fringes are observable and measurable. Refractive properties of transparent materials can be easily studied. Sturdily constructed for classroom or laboratory research.

Complete with monochromatic mercury light and illustrated experiment manual ... only \$209.95



CENTRAL SCIENTIFIC CO. A Subsidiary of Cenco Instruments Corporation 1718-B trying Park Road • Chicago 13, Illinois Branches and Warehouses-Mountainside, N. J. Beston • Birmingham • Santa Cira • Los Angeles • Julis Houston • Toronta • Montrea) • Vancouver • Ottawa

MANAGER OF Physics research

American-Standard seeks a Manager of Physics Research. This is a newly created position in a newly organized research operation located in suburban New York City. We seek a man with proven creative ability who will be given complete responsibility for the administration and design of new techniques in the field of Physics. Ph.D. preferred. Relocation assistance provided.

> Qualified applicants, please submit, in confidence, a detailed resume, including present salary to Paul Lanier

AMERICAN-STANDARD

40 West 40 Street, New York 18, New York pressure on the participants concerned with early detection of missiles by spectroscopic techniques. Dr. Beadle presented a sober evaluation of genetic degeneration in informal additions to his lecture.

W. J. Otting served as chairman of the third session at which the discussion was opened by J. A. L. Thomson of Convair with a presentation of theoretical considerations on the radiation from rocket exhausts. A statistical treatment for the calculation of infrared emissivities was described. The results of numerical computations of emission from heated solid particles were summarized next by G. N. Plass of Aeronutronic. These data, which required information regarding particle-size distribution and refractive index of the pure solid material, were obtained through application of the Mie theory. Experimental studies on rocket exhaust radiation were outlined by C. C. Ferriso of Convair Astronautics, who conducted double-path experiments for the determination of spectral emissivities and temperatures in the exhausts of small rocket motors. The existence of nonequilibrium vibrational population distributions for CO, appeared to be indicated. D. B. Olfe of Caltech outlined the calculation of emissivities of hydrogen at temperatures up to 10 000°K and pressures up to 100 atmospheres. A. Skumanich noted measurements of gamma-ray-induced air luminescence conducted at the Los Alamos Scientific Laboratory during atomic bomb testing. Experiments by N. Solimene and M. Newstein of TRG, Incorporated, relating to the gas discharge properties of importance in connection with the operation of a LASER (Light Amplification by Stimulated Emission) device were reported by Solimene.

J. N. Howard presided over the final session of the symposium which bore the fetching designation "Hot Air". R. E. Meyerott of Lockheed Missile Systems Division presented an authoritative review of the radiative properties of heated air between 1000 and 200 000°K. Experimental data on the emission of radiation from heated air in shock tubes, based on the pioneering work performed by Kantrowitz's AVCO group, were presented by J. Keck. The meeting concluded with an open discussion on basic problems connected with military applications of quantitative spectroscopy that was both entertaining and informative because of John Howard's ability to make the best of a difficult assignment.

Judging from the comments made by participants at the Caltech Symposium, the technical meeting served as a productive opportunity for an exchange of views. Probably no one will object to the following conclusion: our practical difficulties in military applications of quantitative spectroscopy stem from a lack of knowledge of quantitative data of the type that were collected, in another age, by university professors who never thought of missiles. In quantitative spectroscopy, as in many other fields of science, our technology has simply outpaced our store of basic data. If the Caltech Symposium served no other purpose than to emphasize

PHYSICS TODAY

the overwhelming importance of this fact, and thereby leads to increased effort and support on the accumulation of basic data, then it will have been a worthwhile undertaking.

T. A. Jacobs, M. Lapp, and A. Guttman Daniel and Florence Guggenheim Jet Propulsion Center, California Institute of Technology

Fusion

ELEMENTARY aspects of achieving controllable release of energy in significant amounts from thermonuclear reactions will be covered at a symposium to be held August 22-26 in Gatlinburg, Tenn. Designed for university faculty members who have little familiarity with the subject, the meeting is sponsored by the Oak Ridge National Laboratory and the Oak Ridge Institute of Nuclear Studies, in cooperation with the Atomic Energy Commission. The program will consist of a course of four lectures on basic theory, with supplementary lectures on various separate experimental approaches and aspects of plasma physics of more general interest. Various machines used in thermonuclear studies will be discussed.

Further information on the symposium may be obtained from the University Relations Division, Oak Ridge Institute of Nuclear Studies, P. O. Box 117, Oak Ridge, Tenn.

Automatic Control

THE first annual Joint Automatic Control Conference will take place September 7-9 in Cambridge, Mass., on the campus of the Massachusetts Institute of Technology. The aim of the JACC, which is sponsored by the American Society of Mechanical Engineers, the American Institute of Chemical Engineers, the American Institute of Electrical Engineers, the Institute of Radio Engineers, and the Instrument Society of America, is to establish a single high-level technical conference devoted exclusively to the theory and application of automatic control techniques and equipment.

At this year's meeting which is being organized by the ASME Instrument & Regulators Division, the program will include three sessions on sampled data and adaptive control, three on process dynamics, four on computer control, and three on control components. There will also be a review of the proceedings of the first International Congress of the International Federation of Automatic Control, which was held in Moscow last month. For further information and a copy of the advance program when it becomes available, write to the JACC general chairman, William D. Archibald, Energy Control Company, 5 Beekman Street, New York 38, N. Y.

Nuclear Science

PAPERS on solid-state radiation detectors and lownoise amplifiers and their applications to nuclear science will be featured at the seventh annual meeting of the Professional Group on Nuclear Science of the





Now—a combination galvanometer, four-range voltmeter and four-range ammeter with figures and graduations on a large 17" scale that is plainly seen from both sides over any distance in the classroom. An accurate, versatile lecture meter with a modern, shielded movement. All parts and circuits clearly visible.

Cenco No. 82140 complete with shunts and multipliers \$260.00



CENTRAL SCIENTIFIC CO. A Subsidiary of Cenco Instruments Corporation 1718-B Irving Park Road • Chicago 13, Illinois Branches and Warehouses-Mountainside, N. J. Boston • Birmingham • Santa Clara • Los Angeles • Tulta Houston • Toronto • Montreal • Vancouver • Ottawa

