

transformations are allowed if the *orientation* of M remains undetermined, and (ii) the *dilatations* of undotted spinors by say $\lambda \neq 0$, compensated by the dilatations of the dotted spinors by λ^{-1} . This group component, however, does not act effectively on the fundamental objects g_{AB} .

¹¹"6 - 1 = 5" is to be understood on the basis of the fact that (16c) can be interpreted as $3S^{AB} \wedge S_{CD} = \rho \delta^A_C \delta^B_D$, with ρ to be determined. Then, $S^{AB} \wedge S^{CD} = S^{CD} \wedge S^{AB}$ implies that we end up with $6 - 1 = 5$ of the effective conditions on S_{AB} .

¹²A. Tomimatsu and H. Sato, *Progr. Theor. Phys.* 50, 95 (1973).

¹³M. Yamazaki, Kanazawa Univ. Preprint, 1976.

¹⁴F. Ernst, *J. Math. Phys.* 17, 1091 (1976).

¹⁵R. Penrose, a talk at the "Riddle of Gravity" symposium in honor of P. G. Bergmann's 60th birthday, in Syracuse, March 1975.

¹⁶R. Penrose, "Non-Linear Gravitation," First Award Gravity Foundation Essay (1975) and *Gen. Rel. Grav.* 7, 31 (1976).

¹⁷L. Robinson, King's College Lectures, unpublished (1956).

¹⁸J. F. Plebański and I. Robinson, *Phys. Rev. Lett.* 37, 493 (1976).

¹⁹J. F. Plebański and I. Robinson, in *Proceedings of Symposium on Asymptotic Structures of Space-Time*, Cincinnati, June 1977, Edited by F. Eposito and L. Witten (Plenum, New York, 1977), p. 361 and in *Proceedings of V - e Colloque Int. de Theorie de Groups, Montreal, July 1976*, edited by R. Sharp (Academic, New York, 1977).

²⁰A. García, J. F. Plebański, and I. Robinson, "Null Strings and Complex Einstein-Maxwell Fields with λ ," to be published in *Gen. Rel. Grav.*

²¹J. D. Finley, III and J. F. Plebański, *J. Math. Phys.* 17, 2207 (1976).

²²A. García and J. F. Plebański, "Seven Parameter Type Metric in Canonical Coordinates," to be published in *Nuovo Cimento*.

²³J. D. Finley, III and J. F. Plebański, "Spinorial structures and electromagnetic hyperheavens," *J. Math. Phys.* 18, 1662 (1977).

²⁴For the general theory of the real slices of the complex V_4 's see thesis of K. Różga, IFT, Hoża 69, University of Warsaw, Warsaw, Real Slices of Complex Riemannian Spaces."

²⁵With M real,

\hat{g}_{AB} and $\hat{g}^{\dot{A}\dot{B}}$ can be chosen Hermitian;

thus, the concept of a *real* "time" is meaningful—although uncorrelated—in \int_H and \int^E in the case of a real structure.

²⁶F. J. Ernst, *Phys. Rev.* 168, 1415 (1968), *J. Math. Phys.* 15, 1409 (1974).

²⁷F. Ernst and J. F. Plebański, "Killing Structures and Complex \hat{C} -Potentials" to be published in *Ann. Phys.* (Oct., 1977).

²⁸W. Kinnersley, *J. Math. Phys.* 14, 651 (1973).

²⁹W. Kinnersley, *J. Math. Phys.* 18, 1529 (1977).

³⁰M. Walker and R. Penrose, *Commun. Math. Phys.* 18, 265 (1970).

³¹S. Hacyan and J. F. Plebański, "D(1,0) Killing structures and \hat{C} potentials," *J. Math. Phys.* 18, 1517 (1977).

³²A suggestion in this direction by Dr. Frederick J. Ernst is gratefully appreciated.

ERRATA

Erratum: Lattices of effectively nonintegral dimensionality [*J. Math. Phys.* 18, 577 (1977)]

Deepak Dhar

George W. Downs Laboratory of Physics, California Institute of Technology, Pasadena, California 91125

(Received 29 July 1977)

- The exponent in Eq. (3) is $N_r/2$.
- A right parenthesis is missing after $2J_2^{(r)}$ in the second line of Eq. (14e).
- Page 581, column 2, line 5 should have " $i = 1$ to 4."
- Replace the 35th line in the first column of p. 584 by " $3^{1/2q}$. For the Ising."
- Page 584, column 1, line 38 should have " $q = \exp(-2\beta J)$ and $\kappa = 2$ ".