

ON THE THEORY OF Π_3^1 SETS OF REALS

BY A. S. KECHRIS¹ AND D. A. MARTIN

Communicated by Solomon Feferman, June 2, 1977

1. An ordinal basis theorem. Assuming that $\forall x \in \omega^\omega$ ($x^\#$ exists), let u_α be the α th uniform indiscernible (see [3] or [2]). A canonical coding system for ordinals $< u_\omega$ can be defined by letting $WO_\omega = \{w \in \omega^\omega : w = \langle n, x^\# \rangle, \text{ for some } n \in \omega, x \in \omega^\omega\}$ and for $w = \langle n, x^\# \rangle \in WO_\omega$, $|w| = \tau_n^{L[x]}(u_1, \dots, u_{k_n})$, where τ_n is the n th term in a recursive enumeration of all terms in the language of $ZF + V = L[\dot{x}]$, \dot{x} a constant, taking always ordinal values. Call a relation $P(\xi, x)$, where ξ varies over u_ω and x over ω^ω , Π_k^1 if $P^*(w, x) \Leftrightarrow w \in WO_\omega \wedge P(|w|, x)$ is Π_k^1 . An ordinal $\xi < u_\omega$ is called Δ_k^1 if it has a Δ_k^1 notation i.e. $\exists w \in WO_\omega (w \in \Delta_k^1 \wedge |w| = \xi)$.

THEOREM 1 ($ZF + DC + \text{DETERMINACY } (\Delta_2^1)$). *Every nonempty Π_3^1 subset of u_ω contains a Δ_3^1 ordinal.*

COROLLARY 2 ($ZF + DC + \text{DETERMINACY } (\Delta_2^1)$). *Π_3^1 is closed under quantification over ordinals $< u_\omega$ i.e. if $P(\xi, x)$ is Π_3^1 so are $\exists \xi P(\xi, x)$, $\forall \xi P(\xi, x)$.*

COROLLARY 3 ($ZF + DC + AD$). *The class of Π_3^1 sets of reals is closed under $< \delta_3^1$ intersections and unions.*

Martin [3] has proved the corresponding result for Δ_3^1 .

2. A Kleene theory for Π_3^1 . Kleene has characterized the Π_1^1 relations as those which are inductive (see [7]) on the structure $\langle \omega, < \rangle = Q_1$. Let $j_m : u_\omega \rightarrow u_\omega$, $m \geq 1$, be defined by letting

$$j_m(u_i) = \begin{cases} u_i, & \text{if } i < m, \\ u_{i+1}, & \text{if } i \geq m, \end{cases}$$

and then

$$j_m(\tau_n^{L[x]}(u_1, \dots, u_{k_n})) = \tau_n^{L[x]}(j_m(u_1) \dots j_m(u_{k_n})).$$

Let R be the relation on u_ω coding these embeddings, i.e.

$$R = \{(m, \alpha, \beta) : m \in \omega \wedge \alpha, \beta < u_\omega \wedge j_m(\alpha) = \beta\}.$$

Put $Q_3 = \langle u_\omega, <, R \rangle$.

AMS (MOS) subject classifications (1970). Primary 04A15, 02K30, 28A05, 54H05; Secondary 02F35, 02K25, 02K35, 04A30.

¹Research partially supported by NSF Grant MCS 76-17254.

THEOREM 4. ($ZF + DC + DETERMINACY (\Delta_2^1)$). *A set of reals is Π_3^1 iff it is absolutely inductive on the structure \mathcal{Q}_3 .*

In the second part of the above characterization a relation on reals is viewed as a second order relation on u_ω and absolutely inductive means that only parameters from ω are allowed in the definitions (see [7]).

It should be mentioned here that \mathcal{Q}_3 is up to absolute hyper elementary equivalence the same as $\langle u_\omega, <, T^2 \rangle$, where T^2 is the tree (on $\omega \times u_\omega$) coming from the Martin and Solovay [4] analysis of Π_2^1 sets (see [3] for the definition of T^2).

One also obtains the analog for Π_3^1 of the Souslin-Kleene representation of Π_1^1 sets in terms of well-founded trees.

THEOREM 5 ($ZF + DC + DETERMINACY (\Delta_2^1)$). *A set of reals P is Π_3^1 iff there is a tree T on $\omega \times u_\omega$ which is recursive in the structure \mathcal{Q}_3 and $P(x) \Leftrightarrow T(x)$ is well founded.*

For the notation see [2]. The fact that every Π_3^1 set can be so represented is a well-known result of Martin and Solovay [4], the converse being new here.

Let $\mathcal{Q}_3^- = \langle u_\omega, <, \{u_n\}_{n < \omega} \rangle$. Then we also have the context of full AD , in which case $u_n = \delta_n, \forall n \leq \omega$.

THEOREM 6 ($ZF + DC + AD$). *A set of reals is Π_3^1 iff it is Π_1^1 on the structure \mathcal{Q}_3^- .*

3. Explaining the Q -theory. The results in §2 provide a nice explanation for the Q -theory (see [5], [1]) at level 3, which accounts for the structural differences between Π_3^1 and Π_1^1 sets. For example, a real is Δ_3^1 iff it is absolutely hyper elementary on \mathcal{Q}_3 while it is in \mathcal{Q}_3 iff it is hyper elementary (i.e. parameters $< u_\omega$ are allowed) on \mathcal{Q}_3 . Also if y_0 is the first nontrivial Π_3^1 singleton then y_0 is hyper elementary-in- \mathcal{Q}_3 equivalent to the complete inductive-in- \mathcal{Q}_3 subset of u_ω .

4. Higher level analogs of L . Assuming Projective Determinacy (PD), let T^3 be the tree (on $\omega \times \delta_3^1$) associated with an arbitrary Π_3^1 -scale on a complete Π_3^1 set (see [6] and [2]). Let also C_4 be the largest countable Σ_4^1 set. The next result proves a conjecture of Moschovakis and shows that $L[T^3]$ is a correct higher level analog of L for level 4.

THEOREM 7 ($ZF + DC + DETERMINACY (L[\omega^\omega] \cap \text{power}(\omega^\omega))$). *For any T^3 as above, $L[T^3] \cap \omega^\omega = C_4$. In particular $L[T^3] \cap \omega^\omega$ is independent of the tree T^3 .*

Open problem. Is $L[T^3]$ independent of T^3 ?

Further applications of the methods developed here to the theory of Π_3^1 sets as well as details and proofs of the results announced here will appear elsewhere.

REFERENCES

1. A. S. Kechris, *The theory of countable analytical sets*, Trans. Amer. Math. Soc. **202** (1975), 259–297.
2. ———, *On projective ordinals*, J. Symbolic Logic **39** (1974), 269–282.
3. D. A. Martin, *Projective sets and cardinal numbers; some questions related to the continuum problem*, J. Symbolic Logic (to appear).
4. D. A. Martin and R. M. Solovay, *A basis theorem for Σ_3^1 sets of reals*, Ann. of Math. **89** (1969), 138–160.
5. ———, *Basis theorems for Π_{2k}^1 sets of reals* (to appear).
6. Y. N. Moschovakis, *Uniformization in a playful universe*, Bull. Amer. Math. Soc. **77** (1970), 731–736.
7. ———, *Elementary induction on abstract structures*, North-Holland, Amsterdam, 1974.

DEPARTMENT OF MATHEMATICS, CALIFORNIA INSTITUTE OF TECHNOLOGY,
PASADENA, CALIFORNIA 91125

DEPARTMENT OF MATHEMATICS, ROCKEFELLER UNIVERSITY, NEW YORK,
NEW YORK 10021

Current address (D. A. Martin): Department of Mathematics, University of California,
Los Angeles, California 90024