DENSE GAS TRACERS AND STAR FORMATION LAWS IN ACTIVE GALAXIES: APEX SURVEY OF HCN $J = 4 \rightarrow 3$, HCO⁺ $J = 4 \rightarrow 3$, AND CS $J = 7 \rightarrow 6$

ZHI-YU ZHANG (張智昱)^{1,2,3}, YU GAO (高煜)¹, CHRISTIAN HENKEL^{2,4}, YINGHE ZHAO (趙應和)^{1,5},
 JUNZHI WANG (王均智)⁶, KARL M. MENTEN², AND ROLF GÜSTEN²

¹ Purple Mountain Observatory/Key Lab for Radio Astronomy, 2 West Beijing Road, Nanjing 210008, China; zyzhang@pmo.ac.cn

² Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

³ University of the Chinese Academy of Sciences, 19A Yuquan Road, P.O. Box 3908, Beijing 100039, China

⁴ Astronomy Department, King Abdulaziz University, P.O. Box 80203, Jeddah, Saudi Arabia

⁵ Infrared Processing and Analysis Center, California Institute of Technology, MS 100-22, Pasadena, CA 91125, USA

⁶ Shanghai Astronomical Observatory, 80 Nandan Road, Shanghai 200030, China

Received 2013 November 26; accepted 2014 February 4; published 2014 March 14

ABSTRACT

We report HCN $J = 4 \rightarrow 3$, HCO⁺ $J = 4 \rightarrow 3$, and CS $J = 7 \rightarrow 6$ observations in 20 nearby star-forming galaxies with the Atacama Pathfinder EXperiment 12 m telescope. Combined with four HCN, three HCO⁺, and four CS detections from the literature, we probe the empirical link between the luminosity of molecular gas (L'_{gas}) and that of infrared emission (L_{IR}), up to the highest gas densities ($\sim 10^6 \text{ cm}^{-3}$) that have been probed so far. For nearby galaxies with large radii, we measure the IR luminosity within the submillimeter beam size ($14''\text{--}18''$) to match the molecular emission. We find linear slopes for $L'_{\text{CS } J=7-6} - L_{\text{IR}}$ and $L'_{\text{HCN } J=4-3} - L_{\text{IR}}$, and a slightly super-linear slope for $L'_{\text{HCO}^+ J=4-3} - L_{\text{IR}}$. The correlation of $L'_{\text{CS } J=7-6} - L_{\text{IR}}$ even extends over eight orders of luminosity magnitude down to Galactic dense cores, with a fit of $\log(L_{\text{IR}}) = 1.00(\pm 0.01) \times \log(L'_{\text{CS } J=7-6}) + 4.03(\pm 0.04)$. Such linear correlations appear to hold for all densities $> 10^4 \text{ cm}^{-3}$, and indicate that star formation rate is not related to the free-fall timescale for dense molecular gas.

Key words: galaxies: evolution – galaxies: ISM – infrared: galaxies – ISM: molecules – radio lines: galaxies

Online-only material: color figures

1. INTRODUCTION

The star formation process constantly turns gas into stars. The Kennicutt–Schmidt (K-S) law (Kennicutt 1998) globally correlates the surface densities of star formation rate (Σ_{SFR} traced by H α) with total gas mass (Σ_{gas} traced by CO and H I), with a slope $\alpha \sim 1.4$ ($\Sigma_{\text{SFR}} \propto \Sigma_{\text{gas}}^\alpha$). However, quite a range of deviations from the K-S law have been observed, and no unique slope has been found in the $L_{\text{CO}} - L_{\text{IR}}$ correlations as well (e.g., Liu & Gao 2012; Gao & Solomon 2004a).

Recent studies on star formation indicate that stars, especially the massive stars, are predominantly formed in the dense cores of giant molecular clouds (e.g., Evans 2008). Dense gas directly represents molecular content involved in forming stars (e.g., Lada et al. 2012), traced by the rotational transitions of high dipole moment molecules (e.g., HCN and HCO⁺), because of their high critical densities⁷ (n_{crit}). Gao & Solomon (2004a, 2004b) find a tight linear correlation between the luminosities of IR emission (L_{IR} tracing the star formation rate (SFR)) and HCN $J = 1 \rightarrow 0$ ($L'_{\text{HCN } J=1-0}$ tracing M_{dense}) in galaxies. This correlation even extends to Galactic dense cores undergoing high mass star formation (e.g., Wu et al. 2005). Lada et al. (2012) argue that such a linear correlation is a fundamental relation for SFR and dense gas, and molecular gas with densities above 10^4 cm^{-3} should follow this linearity. Linear correlations have also been found in other dense gas tracers with similar or higher

n_{crit} (e.g., HCO⁺ $J = 1 \rightarrow 0$, HNC $J = 3 \rightarrow 2$, CS $J = 5 \rightarrow 4$), in both Galactic dense cores and galaxies (e.g., Schenck et al. 2011; Ma et al. 2013; Reiter et al. 2011; Wang et al. 2011).

For gas tracers with n_{crit} higher than HCN $J = 1 \rightarrow 0$, the slopes of the $L'_{\text{gas}} - L_{\text{IR}}$ correlations are controversial. Krumholz & Thompson (2007) argue that the deviation of the mean gas density in different types of galaxies and n_{crit} of the gas tracers change the slopes. Numerical simulations predict decreasing slopes against increasing n_{crit} , because of sub-thermal excitation conditions (i.e., Narayanan et al. 2008; Juneau et al. 2009). Observations of HCN, HCO⁺, and CS (typically, $J \leq 3 \rightarrow 2$) have been used to support these and show sub-linear correlations (e.g., Baan et al. 2008; Bussmann et al. 2008; Graciá-Carpio et al. 2008), where HCN $J = 3 \rightarrow 2$ has a slope of $\alpha \sim 0.8$, following the prediction quite well (Bussmann et al. 2008). Up to date, only few detections of higher- J transitions of the above species have been reported (i.e., Knudsen et al. 2007; Jackson et al. 1995; Greve et al. 2009; Bayet et al. 2009; Wilson et al. 2008).

To better probe the densest molecular gas in galaxies, and to test the predictions for higher n_{crit} tracers, we therefore performed a survey of HCN $J = 4 \rightarrow 3$, HCO⁺ $J = 4 \rightarrow 3$, and CS $J = 7 \rightarrow 6$ in 20 nearby actively star-forming galaxies with the Atacama Pathfinder EXperiment (APEX) 12 m telescope.⁸ With the advent of the *Herschel* space telescope, we are able to obtain beam-matched IR luminosities in nearby galaxies, and to compare them with data from single dish (SD) telescopes. In this Letter, we summarize our findings and compare them with the results from Galactic studies. We adopt

⁷ All critical densities (n_{crit}) in this Letter are calculated from $n_{\text{crit}} = \sum_{u>l} A_{ul} / \sum_{u \neq l} C_{ul}(T_{\text{kin}})$ at $T_{\text{kin}} = 100 \text{ K}$ assuming optically thin emission.

A_{ul} and C_{ul} denote the Einstein coefficient for spontaneous emission and the collision rate, respectively. All state-to-state cross sections and rate coefficients are from the LAMDA Web site (<http://home.strw.leidenuniv.nl/~moldata/>) (Schöier et al. 2005).

⁸ This publication is based on data acquired with the Atacama Pathfinder Experiment (APEX). APEX is a collaboration between the Max-Planck-Institut für Radioastronomie, the European Southern Observatory, and the Onsala Space Observatory.

Figure 1. Spectra of the strong detections of $\text{HCO}^+ J = 4 \rightarrow 3$. Filled lines show $\text{CS } J = 7 \rightarrow 6$, thick red lines show $\text{HCO}^+ J = 4 \rightarrow 3$, and dotted blue lines show $\text{HCN } J = 4 \rightarrow 3$. All velocities are labeled relative to the velocity corresponding to their redshifts.

(A color version of this figure is available in the online journal.)

flat cosmology parameters ($\Omega_m = 0.27$, $\Omega_\Lambda = 0.73$, and $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-3}$; Spergel et al. 2007).

2. SAMPLE, OBSERVATIONS AND DATA REDUCTION

We selected 20 galaxies from the *Infrared Astronomical Satellite* (IRAS) Revised Bright Galaxy Sample (Sanders et al. 2003), with $S_\nu(100 \mu\text{m}) > 100 \text{ Jy}$, and declination $< 20^\circ$. In the analysis. We also include data from the literature (Jackson et al. 1995; Knudsen et al. 2007; Papadopoulos 2007; Wilson et al. 2008; Greve et al. 2009; Bayet et al. 2009), which were mostly observed with the James Clerk Maxwell Telescope with a beam-size (full width half power) of $14''$. The sample encompasses galaxies with L_{IR} from $10^{10} L_\odot$ to $10^{12.5} L_\odot$, including nearby normal galaxies, starbursts, and ultra luminous infrared galaxies (ULIRGs; $L_{\text{IR}} \geq 10^{12} L_\odot$), with about half containing active galactic nuclei (AGNs). Table 1 lists the targets with integrated intensity, distance, and luminosities.

Our observations were performed in 2011 April and August with the APEX on the Chajnantor Plateau in Chile, in good ($p_{\text{wv}} < 0.6 \text{ mm}$) to median ($p_{\text{wv}} \sim 1 \text{ mm}$) weather conditions. In total we spent $\sim 30 \text{ hr}$ telescope time on this project. The First Light APEX Submillimeter Heterodyne receiver was employed to observe $\text{CS } J = 7 \rightarrow 6$, $\text{HCN } J = 4 \rightarrow 3$ and $\text{HCO}^+ J = 4 \rightarrow 3$ simultaneously, with dual sidebands. Typical system temperatures were $T_{\text{sys}} \sim 180\text{--}240 \text{ K}$. The Fast Fourier Transform Spectrometer back ends led to a bandwidth of 4 GHz for each sideband, with a channel spacing of 0.2 MHz . The beam size is $\sim 18''$ at 342 GHz .

All observations were performed in a wobbler switching mode. Beam throws range between $2'$ and $4'$, according to the target size. Every 15 minutes we made a chopper wheel calibration. The focus was determined on Saturn or Jupiter every 3–6 hr. Pointing was checked once per hour, resulting in a

typical uncertainty of $2''\text{--}3''$ (rms). Including overhead, we spent $\sim 1.5 \text{ hr}$ on each galaxy. Typical rms noise levels are $1.5\text{--}2 \text{ mK}$ at 20 km s^{-1} velocity resolution. Although the sideband separation is better than 10 dB , there are still some $\text{CO } J = 3 \rightarrow 2$ images present in the upper sideband. Fortunately these CO images do not mix with our HCN spectra.

All data were reduced with the CLASS package in GILDAS.⁹ We checked the line profiles of low- J HCN or CO transitions in the literature and set the baseline ranges accordingly. Linear baselines were subtracted after inspecting each spectrum. We qualified spectra by comparing the measured noise and the theoretical noise before and after a few times of boxcar smoothing. About 5% of the spectra were discarded during the qualification.

We converted the antenna temperatures (T_A) to the main beam brightness temperatures (T_{mb}) using $T_{\text{mb}} = T_A^* F_{\text{eff}} / B_{\text{eff}}$. The adopted forward hemisphere efficiency F_{eff} and beam efficiency B_{eff} are 0.95 and 0.73, respectively. The flux uncertainty was estimated to be $\sim 15\%$. We derived the line luminosity (L'_{gas}) following Solomon et al. (1992).

3. RESULTS AND DISCUSSION

We detected 8, 14, and 16 ($I_{\text{line}} \geq 3\sigma$) galaxies in $\text{CS } J = 7 \rightarrow 6$, $\text{HCN } J = 4 \rightarrow 3$, and $\text{HCO}^+ J = 4 \rightarrow 3$, respectively. Most of them are strong detections in HCN and HCO^+ ($\geq 4\sigma$), while Mrk 331, NGC 7469 and NGC 4039 show marginal detections in $\text{HCO}^+ J = 4 \rightarrow 3$ ($\sim 3\sigma$). Upper limits (3σ) are derived for the non-detections. Figure 1 shows the spectra of strong detections of $\text{HCO}^+ J = 4 \rightarrow 3$, and Figure 2 shows the spectra of weak detections of $\text{HCO}^+ J = 4 \rightarrow 3$. NGC 7771 and NGC 2903 has only non-detections, so we do not show their spectra. Combining data from the literature we

⁹ <http://www.iram.fr/IRAMFR/GILDAS>

Table 1
Observed Properties of the Sample

Source Name	Distance (Mpc)	L_{IR} ($10^{10} L_{\odot}$)	$I_{\text{CS}(7-6)}$ (K km s $^{-1}$)	$I_{\text{HCN}(4-3)}$ (K km s $^{-1}$)	$I_{\text{HCO}^+(4-3)}$ (K km s $^{-1}$)	$L'_{\text{CS}(7-6)}$ (10^6 K km s $^{-1}$ pc 2)	$L'_{\text{HCN}(4-3)}$ (10^6 K km s $^{-1}$ pc 2)	$L'_{\text{HCO}^+(4-3)}$ (10^6 K km s $^{-1}$ pc 2)	R_{SD}	C_{aper}	Band (μm)	Type ^g
IC 1623	82.4	49.0	<0.9	<0.9	0.9 ± 0.2	<66	<62	62 ± 15	0.65	1.29	100	SF
NGC 1068	14.4	20.4	1.0 ± 0.2	5.0 ± 0.2	2.7 ± 0.2	2.5 ± 0.5	10.7 ± 0.4	5.8 ± 0.4	0.16	1.16	70	AGN
NGC 1365	18.3	10.5	0.6 ± 0.2	1.7 ± 0.2	2.9 ± 0.2	2.2 ± 0.7	5.9 ± 0.7	10.0 ± 0.7	0.25	1.16	70	AGN
NGC 1808	12.3	6.92	<0.6	2.5 ± 0.2	2.2 ± 0.2	<1.0	3.9 ± 0.3	3.4 ± 0.3	0.46	1.29	100	AGN
NGC 2903	10.6	2.57	<0.9	<0.9	<0.9	<1.1	<1.0	<1.0	1.0	...	...	SF
NGC 3256	44.3	57.5	0.8 ± 0.2	1.1 ± 0.2	2.8 ± 0.2	45 ± 6	22 ± 4	56 ± 4	0.34	1.29	100	SF
NGC 3627	11.0	2.88	<0.7	0.8 ± 0.2	0.7 ± 0.2	<1.0	1.0 ± 0.3	0.9 ± 0.3	0.08	1.29	100	AGN
NGC 3628	11.0	2.14	0.5 ± 0.2	0.9 ± 0.2	2.5 ± 0.2	<1.2	1.1 ± 0.4	3.1 ± 0.4	0.23	...	850	SF
NGC 4039	25.0	9.33	<1.2	<0.7	0.9 ± 0.3	<8.2	<4.5	5.8 ± 1.8	0.04	1.29	100	SF
NGC 4418	29.5	9.33	1.2 ± 0.2	3.0 ± 0.2	1.2 ± 0.2	11.5 ± 2.0	26.8 ± 2.0	8.9 ± 1.8	1.0	...	...	AGN
NGC 4945	3.8	2.82	7.3 ± 0.3	23.4 ± 0.3	9.9 ± 0.3	1.2 ± 0.05	3.6 ± 0.05	4.5 ± 0.05	0.40	1.29	100	AGN
IRAS 13120-5453	128.3	178	<0.9	2.0 ± 0.3	1.6 ± 0.3	<106	330 ± 30	265 ± 30	1.0	...	...	SF
IRAS 13242-5713	38.9	31.6	1.8 ± 0.3	4.3 ± 0.3	3.5 ± 0.3	30 ± 5	67 ± 5	54 ± 5	0.64	1.29	100	SF
NGC 6240	107	70.8	<0.7	1.3 ± 0.2	2.4 ± 0.2	<99	189 ± 30	260 ± 20	1.0	...	...	SF
IRAS 17578-0400	61.1	24.5	<0.9	1.5 ± 0.3	<0.9	<36.6	57.0 ± 10	<34	1.0	...	...	SF
IRAS 18293-3413	77.4	64.6	<0.9	<0.9	1.6 ± 0.3	<58.6	<55	116 ± 20	1.0	...	...	SF
NGC 7469	71.6	46.8	<0.9	<0.9	1.0 ± 0.2	<33.5	<30	37 ± 10	1.0	...	...	AGN
NGC 7552	19.5	8.91	0.6 ± 0.2	1.3 ± 0.2	2.4 ± 0.2	1.3 ± 0.4	5.1 ± 0.4	9.4 ± 0.4	0.59	1.29	100	SF
NGC 7771	63.0	26.9	<0.9	<0.9	<0.9	<40	<36	<36	0.55	1.29	100	SF
Mrk 331	80.9	33.9	<0.9	1.1 ± 0.2	<0.9	<43	60 ± 13	<40	1.0	...	...	AGN
NGC 253 ^a	3.47	3.45	...	...	...	$0.35 \pm 0.01^{\text{a}}$	$8.9 \pm 1^{\text{b}}$	$7.7 \pm 1.0^{\text{b}}$	0.18	1.33	100	SF
IC 342 ^b	3.93	1.08	...	...	...	$0.05 \pm 0.02^{\text{a}}$	$0.8 \pm 0.2^{\text{c}}$	...	0.05	1.33	100	SF
Arp 220 ^c	79.9	162.2	...	...	...	$190 \pm 40^{\text{d}}$	950 ± 190	$170 \pm 40^{\text{d}}$	1.0	...	...	SF
HE2-10 ^d	16.4	0.6	...	...	...	$0.21 \pm 0.7^{\text{a}}$	...	...	0.29	1.33	100	SF
Mrk 231 ^e	180.8	340	...	...	...	...	$550 \pm 110^{\text{e}}$	$188 \pm 23^{\text{f}}$	1.0	...	...	AGN

Notes.

^a Bayet et al. (2009).

^b Knudsen et al. (2007).

^c Jackson et al. (1995).

^d Greve et al. (2009).

^e Papadopoulos (2007).

^f Wilson et al. (2008).

^g Galaxy types are found in the NASA/IPAC Extragalactic Database (NED).

Figure 2. Spectra of the weak detections of $\text{HCO}^+ J = 4 \rightarrow 3$. The labels are the same as Figure 1.

(A color version of this figure is available in the online journal.)

construct samples containing 14, 17, and 17 detected galaxies for $\text{CS } J = 7 \rightarrow 6$, $\text{HCN } J = 4 \rightarrow 3$, and $\text{HCO}^+ J = 4 \rightarrow 3$, respectively.

Total IR (TIR; 8–1000 μm) luminosity (Sanders et al. 2003) is adopted as a proxy of SFR. However, the molecular lines are obtained in small beam sizes ($\sim 14''$ – $18''$), which only pick up a certain portion of the IR luminosity for a whole galaxy. This is particularly problematic for nearby normal spirals and starbursts because of their large radii (up to a few arcminutes). If this effect was neglected, one would systematically overestimate the IR emission or underestimate the dense gas emission at the low luminosity end of the $L'_{\text{gas}} - L_{\text{IR}}$ correlations, while the ULIRGs are not affected due to the small angular sizes of their gas emission.

We decide to measure the IR emission within the submillimeter beam size rather than to adopt the IR luminosities of entire galaxies. We download either *Herschel*¹⁰ PACS 100 μm or 70 μm (when 100 μm is not available) images from the Herschel Science Archive, and they have been processed to level 2.5 in the pipeline. NGC 3628 does not have *Herschel* data, so we adopt a SCUBA 850 μm archival image instead. Using these IR data, we perform aperture photometry with the submillimeter beam size and with the whole galaxy. The background radii are selected by eye from the outside of galaxies to the edge of the images. Some nearby galaxies are downloaded from the Key Insights on Nearby Galaxies project, and similar to their photometry results (Dale et al. 2012), the impact of using $\sim 10\%$ larger or smaller aperture areas is a median difference of less than 3% on the flux densities for all photometric measurements. The IR luminosity within the beam size is:

$$L_{\text{IR}} = L_{\text{TIR}} \times R_{\text{SD}} \times C_{\text{aper}},$$

where L_{TIR} is the IR luminosity of the entire galaxy, R_{SD} is the ratio of flux densities within the beam size of the SD telescope to that measured in the whole galaxy, and C_{aper} is the aperture correction factor for the beam sizes. The final error comprises the photometric error ($\sim 5\%$), the point source assumption ($\sim 10\%$), the flux calibration error ($\sim 5\%$), and the error of tracing TIR with a chromatic IR band ($\sim 10\%$; e.g., Galametz et al. 2013). In the end we consider 20% to be a conservative uncertainty for L_{IR} .

3.1. The $L'_{\text{gas}} - L_{\text{IR}}$ Correlations

Figure 3 presents the $L'_{\text{gas}} - L_{\text{IR}}$ correlations. All detections are included in the fitting except for NGC 4039 and NGC 3627, because the regions covered by their APEX beams are not active in star formation as indicated by their IR images. In the linear regression we assume Gaussian independent variables and account for the errors in both L_{IR} and L'_{gas} . Upper limits are not adopted in the fitting. The $L'_{\text{gas}} - L_{\text{IR}}$ correlations extend from the nuclear regions of nearby normal galaxies to ULIRGs, covering an L_{IR} range of ~ 2.5 decades. We adopt publicly available IDL routines MPFIT (Markwardt 2009) for the linear least-squares fit and LINMIX_ERR (Kelly 2007) for the Bayesian regression. The best linear least-squares fits with uncertainties are listed below, with r indicating the correlation coefficient:

$$\log(L_{\text{IR}}) = 0.95(\pm 0.04) \times \log(L'_{\text{CS}76}) + 4.33(\pm 0.28); r = 0.99$$

$$\log(L_{\text{IR}}) = 1.00(\pm 0.04) \times \log(L'_{\text{HCN}43}) + 3.67(\pm 0.28); r = 0.97$$

$$\log(L_{\text{IR}}) = 1.10(\pm 0.05) \times \log(L'_{\text{HCO}+43}) + 2.48(\pm 0.38); r = 0.96.$$

In the Bayesian regressions, we present the posterior distributions of possible slopes in the insets of Figure 3, and obtain error estimates by fitting the distributions with Gaussian

¹⁰ *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

Figure 3. Correlations between the gas luminosity $\log(L'_{\text{gas}})$ and the IR luminosity $\log(L_{\text{IR}})$. Upper left: CS $J = 7 \rightarrow 6$. Upper right: HCN $J = 4 \rightarrow 3$. Bottom left: $\text{HCO}^+ J = 4 \rightarrow 3$. Bottom right: CS $J = 7 \rightarrow 6$ for both galaxies and Galactic cores. The horizontal dashed lines indicate $L_{\text{IR}} = 10^{4.5} L_{\odot}$, below where the Galactic cores have less statistical meaning (i.e., Wu et al. 2010). N is the slope of the correlation and r is the correlation coefficient. The thick lines show the fitted results. The oblique dashed lines indicate linear relations. Red empty diamonds are AGNs, open circles are detections for the fitting, and filled circles and arrows are limits. NGC 4039 and NGC 3627 are not adopted for fitting and are plotted in blue. The insets present probability density distributions of the slopes derived from the Bayesian fitting.

(A color version of this figure is available in the online journal.)

profiles. These fittings give slopes of 0.94 ± 0.07 , 1.01 ± 0.07 , and 1.12 ± 0.9 for CS $J = 7 \rightarrow 6$, HCN $J = 4 \rightarrow 3$, and $\text{HCO}^+ J = 4 \rightarrow 3$, respectively. These are consistent with the linear least-squares fit. Both correlations of $L'_{\text{CS } J=7-6} - L_{\text{IR}}$ and $L'_{\text{HCN } J=4-3} - L_{\text{IR}}$ have slopes very close to unity, while the $L'_{\text{HCO}^+ J=4-3} - L_{\text{IR}}$ correlation shows a slightly super-linear slope. The slopes of HCN and HCO^+ are much higher than those predicted by Narayanan et al. (2008) and Juneau et al. (2009). Their model did not predict for CS $J = 7 \rightarrow 6$, but the high n_{crit} of CS $J = 7 \rightarrow 6$ makes it not likely, following a similar trend predicted for HCN and HCO^+ . We also fitted the correlations with our APEX data only, and obtained slopes of 0.98 ± 0.15 , 0.98 ± 0.05 , and 1.08 ± 0.05 , which are very close to the results fitted with the combined data.

Alternatively, we fitted the correlations without the beam-matching correction, and obtained the slopes (N) and correlation coefficients (r) to be: $N = 0.71 \pm 0.08$ and $r = 0.89$ for

CS $J = 7 \rightarrow 6$, $N = 0.77 \pm 0.05$ and $r = 0.94$ for HCN $J = 4 \rightarrow 3$, and $N = 0.94 \pm 0.10$ and $r = 0.92$ for $\text{HCO}^+ J = 4 \rightarrow 3$, respectively. The shallower slopes (sub-linear) and worse correlations indicate that the beam-matching correction has a significant effect in the fitting. On the other hand, it should be emphasized that, with the beam-matching correction we are studying the central $14''$ – $18''$ regions in nearby galaxies (except for the maps of HCN $J = 4 \rightarrow 3$ and $\text{HCO}^+ J = 4 \rightarrow 3$ in NGC 253), rather than whole galaxies.

At $T_{\text{kin}} = 100$ K, HCN $J = 4 \rightarrow 3$, $\text{HCO}^+ J = 4 \rightarrow 3$, and CS $J = 7 \rightarrow 6$ have n_{crit} of $5.6 \times 10^6 \text{ cm}^{-3}$, $1.3 \times 10^6 \text{ cm}^{-3}$, and $2.8 \times 10^6 \text{ cm}^{-3}$ (see Section 1), respectively. These tracers pick up the densest part of the molecular cores, and trace only a small amount of the total dense gas mass involved in star-formation. The linear slopes found in the high- J HCN and CS tend to support the proposed correlations for gas tracers with n_{crit} higher than $\sim 10^4 \text{ cm}^{-3}$ (Lada et al. 2012).

Compared with the neutral HCN and CS molecules, the molecular ion HCO^+ is more sensitive to the average ambient electron abundance $x(e)$, because of the protonation reaction:

The high electron abundance is likely to reduce $[\text{HCO}^+/\text{H}_2]$ efficiently. Considering the impacts of cosmic ray ionization (Papadopoulos 2007), the turbulent diffusion (Xie et al. 1995), and the AGN illumination, HCO^+ is likely deficient when exposed to the extreme physical conditions prevailing at the high luminosity end, i.e., ULIRGs. Such effect could increase the slopes of the L'_{HCO^+} - L_{IR} correlations in galaxies.

3.2. Comparison between Galactic Cores and Galaxies

Wu et al. (2010) measured the $L'_{\text{CS } J=7-6}$ - L_{IR} correlation in individual Galactic dense cores with high mass star formation. Their best fit yields a slope of ~ 0.8 with a large uncertainty because of the significant scatter in the data and the limited range of infrared luminosities they have at their disposal. In the bottom right of Figure 3, we plot their Galactic cores and our galaxies together. Combining both samples, we find a highly linear correlation,

$$\log(L_{\text{IR}}) = 1.00(\pm 0.01) \times \log(L'_{\text{CS } J=7-6}) + 4.03(\pm 0.04),$$

with a correlation coefficient of 0.98. This correlation is remarkably similar to that derived for $\text{HCN } J = 1 \rightarrow 0$ by Wu et al. (2005), though $\text{CS } J = 7 \rightarrow 6$ has much higher excitation requirements. The linear correlation between L'_{gas} and L_{IR} holds over an IR luminosity range of about eight orders of magnitude.

The linearity of the (dense gas)-SFR correlation was interpreted as a “fundamental unit” of star formation on different physical scales, thus both SFR and dense gas mass are simply accumulated by adding in more units (e.g., Wu et al. 2005). Once the studied gas content is denser than a density threshold, the SFR does not depend on the exact value of the gas density, but depends on the total mass of the dense gas (e.g., Lada et al. 2012). In such a context the near-linearity in the Σ_{gas} - Σ_{SFR} correlation in nearby normal spirals (e.g., Bigiel et al. 2008; Schruba et al. 2012) is likely caused by a constant $M_{\text{dense}}/M_{\text{total H}_2}$ fraction.

The gas densities probed by the three lines are beyond the highest average densities in starbursts (Krumholz & Thompson 2007), and their n_{crit} are one to two orders of magnitude higher than the threshold density for star formation ($n_{\text{H}_2} \sim 10^4 \text{ cm}^{-3}$; e.g., Parmentier et al. 2011; Lada et al. 2012). From $\text{HCN } J = 1 \rightarrow 0$, all dense gas tracers with $n_{\text{crit}} > 10^4 \text{ cm}^{-3}$ are linearly correlated with the SFR, which indicates that the SFR in the dense gas is not likely affected by the free-fall time scale (t_{ff}), because $t_{\text{ff}} \propto \rho^{-1/2}$. If $\Sigma_{\text{gas}}/t_{\text{ff}}$ - Σ_{SFR} correlations are linear for all of the dense gas, the shorter t_{ff} for the denser gas would not keep L'_{gas} - L_{IR} linear. Indeed, the gas content traced by $\text{HCN } J = 1 \rightarrow 0$ has 10 times longer t_{ff} than that traced by $\text{HCN } J = 4 \rightarrow 3$, but both transitions have linear correlations with the SFR. The nonlinear correlations found in the K-S law are likely caused by the different fractions of dense gas to total gas in different types of galaxies, rather than the free-fall timescales (Lada et al. 2012). Actually, the K-S law with a slope of ~ 1.4 is likely not a fundamental physical relation, as the nonlinear slope found for $\text{H}_2 + \text{H I}$, seen in very tenuous gas, is related to the relative amounts of dense gas and dust directly

involved in star formation and other gas being too diffuse and diluted to have anything to do with star formation.

3.3. AGN Contamination to the IR Emission

About half of the galaxies in our sample host AGNs, in particular the ULIRGs. Compared to the near- and mid-IR bands, the far- and TIR are well correlated and are thus less contaminated by AGN emission (e.g., U et al. 2012; Juneau et al. 2009). Some AGNs, however, can still contribute significantly to the bolometric luminosity in ULIRGs (e.g., Veilleux et al. 2009). For the most extreme case, Mrk 231, the estimated AGN contribution is $\sim 70\%$, and the SFR indicated from the IR emission should be ~ 0.5 dex lower in the L'_{gas} - L_{IR} correlations (e.g., Veilleux et al. 2009). For the rest of the galaxies, the AGN contamination should be much less ($< 10\%$) for nearby Seyferts, where star formation is expected to dominate over IR emission. Unfortunately, the IR emission for most AGN host galaxies have not been decomposed, so that we could not deduce the IR luminosity from the fraction of the IR luminosity due to star formation. Overall, we do not find significant changes in the correlations if we remove AGNs from our samples. This may be due to the small size of our sample, or the fact that most galaxies hosting AGNs are not strongly contaminated by AGNs.

In conclusion, we present a survey of three dense gas tracers ($\text{CS } J = 7 \rightarrow 6$, $\text{HCN } J = 4 \rightarrow 3$, and $\text{HCO}^+ J = 4 \rightarrow 3$) in 20 nearby galaxies observed with the APEX 12 m telescope. Combined with data from the literature and after a beam-matching correction for nearby galaxies, we find linear L'_{gas} - L_{IR} correlations for $\text{CS } J = 7 \rightarrow 6$ and $\text{HCN } J = 4 \rightarrow 3$, and a slightly super-linear slope for $\text{HCO}^+ J = 4 \rightarrow 3$. These results are consistent with those found in $\text{HCN } J = 1 \rightarrow 0$ (i.e., Gao & Solomon 2004a, 2004b) and $\text{CS } J = 5 \rightarrow 4$ (i.e., Wang et al. 2011; Wu et al. 2010), but contrary to the predictions in Bussmann et al. (2008) and Juneau et al. (2009). We also find that the linear $L'_{\text{CS } J=7-6}$ - L_{IR} correlation holds *universally* across eight orders of luminosity magnitude, down to Galactic cores. If the beam-matching corrections are not applied for nearby galaxies, however, the slopes will be significantly sub-linear. Future ALMA surveys of more dense gas tracers as well as other transitions over large samples of galaxies are necessary to consolidate the above findings and ensure that, apart from the vagaries of HCO^+ chemistry, the simplest (dense gas)-SFR empirical relation is indeed true.

We are grateful to the MPIfR team and the APEX staff for their help and support during and after the observations. We also thank the anonymous referee for very useful comments and suggestions which improved the quality of the paper. Z.Z. thanks Padelis Papadopoulos, Thomas Greve, and Daizhong Liu for their constructive suggestions. This work is partially supported by NSFC grant Nos. 11173059 and 11390373, and CAS No. XDB09000000. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Baan, W. A., Henkel, C., Loenen, A. F., Baudry, A., & Wiklind, T. 2008, *A&A*, **477**, 747
 Bayet, E., Aladro, R., Martín, S., Viti, S., & Martín-Pintado, J. 2009, *ApJ*, **707**, 126

- Bigiel, F., Leroy, A., Walter, F., et al. 2008, *AJ*, **136**, 2846
- Bussmann, R. S., Narayanan, D., Shirley, Y. L., et al. 2008, *ApJL*, **681**, L73
- Dale, D. A., Aniano, G., Engelbracht, C. W., et al. 2012, *ApJ*, **745**, 95
- Evans, N. J., II 2008, in ASP Conf. Ser. 390, Pathways Through an Eclectic Universe, ed. J. H. Knapen, T. J. Mahoney, & A. Vazdekis (San Francisco, CA: ASP), 52
- Galametz, M., Kennicutt, R. C., Calzetti, D., et al. 2013, *MNRAS*, **431**, 1956
- Gao, Y., & Solomon, P. M. 2004a, *ApJS*, **152**, 63
- Gao, Y., & Solomon, P. M. 2004b, *ApJ*, **606**, 271
- Graciá-Carpio, J., García-Burillo, S., Planesas, P., Fuente, A., & Usero, A. 2008, *A&A*, **479**, 703
- Greve, T. R., Papadopoulos, P. P., Gao, Y., & Radford, S. J. E. 2009, *ApJ*, **692**, 1432
- Jackson, J. M., Paglione, T. A. D., Carlstrom, J. E., & Rieu, N.-Q. 1995, *ApJ*, **438**, 695
- Juneau, S., Narayanan, D. T., Moustakas, J., et al. 2009, *ApJ*, **707**, 1217
- Kelly, B. C. 2007, *ApJ*, **665**, 1489
- Kennicutt, R. C., Jr. 1998, *ApJ*, **498**, 541
- Knudsen, K. K., Walter, F., Weiss, A., et al. 2007, *ApJ*, **666**, 156
- Krumholz, M. R., & Thompson, T. A. 2007, *ApJ*, **669**, 289
- Lada, C. J., Forbrich, J., Lombardi, M., & Alves, J. F. 2012, *ApJ*, **745**, 190
- Liu, L., & Gao, Y. 2012, *ScChG*, **55**, 347
- Ma, B., Tan, J. C., & Barnes, P. J. 2013, *ApJ*, **779**, 79
- Markwardt, C. B. 2009, in ASP Conf. Ser. 411, Astronomical Data Analysis Software and Systems XVIII, ed. D. A. Bohlender, D. Durand, & P. Dowler (San Francisco, CA: ASP), 251
- Narayanan, D., Cox, T. J., Shirley, Y., et al. 2008, *ApJ*, **684**, 996
- Papadopoulos, P. P. 2007, *ApJ*, **656**, 792
- Parmentier, G., Kauffmann, J., Pillai, T., & Menten, K. M. 2011, *MNRAS*, **416**, 783
- Reiter, M., Shirley, Y. L., Wu, J., et al. 2011, *ApJS*, **195**, 1
- Sanders, D. B., Mazzarella, J. M., Kim, D.-C., Surace, J. A., & Soifer, B. T. 2003, *AJ*, **126**, 1607
- Schenck, D. E., Shirley, Y. L., Reiter, M., & Juneau, S. 2011, *AJ*, **142**, 94
- Schöier, F. L., van der Tak, F. F. S., van Dishoeck, E. F., & Black, J. H. 2005, *A&A*, **432**, 369
- Schruba, A., Leroy, A. K., Walter, F., et al. 2012, *AJ*, **143**, 138
- Solomon, P. M., Radford, S. J. E., & Downes, D. 1992, *Natur*, **356**, 318
- Spergel, D. N., Bean, R., Doré, O., et al. 2007, *ApJS*, **170**, 377
- U, V., Sanders, D. B., Mazzarella, J. M., et al. 2012, *ApJS*, **203**, 9
- Veilleux, S., Rupke, D. S. N., Kim, D.-C., Genzel, R., et al. 2009, *ApJS*, **182**, 628
- Wang, J., Zhang, Z., & Shi, Y. 2011, *MNRAS*, **416**, L21
- Wilson, C. D., Petitpas, G. R., Iono, D., et al. 2008, *ApJS*, **178**, 189
- Wu, J., Evans, N. J., II, Gao, Y., et al. 2005, *ApJL*, **635**, L173
- Wu, J., Evans, N. J., II, Shirley, Y. L., & Knez, C. 2010, *ApJS*, **188**, 313
- Xie, T., Allen, M., & Langer, W. D. 1995, *ApJ*, **440**, 674