

## On the flux of oxygenated volatile organic compounds from organic aerosol oxidation

Alan J. Kwan,<sup>1</sup> John D. Crouse,<sup>2</sup> Antony D. Clarke,<sup>3</sup> Yohei Shinozuka,<sup>3</sup>  
Bruce E. Anderson,<sup>4</sup> James H. Crawford,<sup>4</sup> Melody A. Avery,<sup>4</sup> Cameron S. McNaughton,<sup>3</sup>  
William H. Brune,<sup>5</sup> Hanwant B. Singh,<sup>6</sup> and Paul O. Wennberg<sup>1,7</sup>

Received 24 February 2006; revised 17 April 2006; accepted 12 June 2006; published 9 August 2006.

[1] Previous laboratory and field studies suggest that oxidation of organic aerosols can be a source of oxygenated volatile organic compounds (OVOC). Using measurements of atmospheric oxidants and aerosol size distributions performed on the NASA DC-8 during the INTEX-NA campaign, we estimate the potential magnitude of the continental summertime OVOC flux from organic aerosol oxidation by OH to be as large as  $\sim 70$  pptv C/day in the free troposphere. Contributions from O<sub>3</sub>, H<sub>2</sub>O<sub>2</sub>, photolysis, and other oxidants may increase this estimate. These processes may provide a large, diffuse source of OVOC that has not been included in current atmospheric models, and thus have a significant impact on our understanding of organic aerosol, OVOC, PAN, and HO<sub>x</sub> chemistry. The potential importance and highly uncertain nature of our estimate highlights the need for more field and laboratory studies on organic aerosol composition and aging.  
**Citation:** Kwan, A. J., et al. (2006), On the flux of oxygenated volatile organic compounds from organic aerosol oxidation, *Geophys. Res. Lett.*, 33, L15815, doi:10.1029/2006GL026144.

### 1. Introduction

[2] Oxygenated volatile organic compounds (OVOC) comprise a large number of the species whose transport to the remote troposphere can impact radical budgets and sequester NO<sub>x</sub> in the form of nitrates [Singh et al., 1995; Wennberg et al., 1998; Muller and Brasseur, 1999]. In addition, they play a role in the formation of organic aerosols (OA) [Kanakidou et al., 2005, and references therein]. Field campaigns have noted large concentrations of OVOC throughout the troposphere, but their budgets are poorly understood [Singh et al., 2001, 2004; Wisthaler et al., 2002].

[3] Ellison et al. [1999] suggest that oxidation of OA may provide an OVOC source to the remote troposphere.

Field campaigns have established the ubiquity of OA throughout the troposphere [Murphy et al., 1998; Kanakidou et al., 2005]. Most aerosols contain both organic and inorganic components, though significantly, the organic fraction tends to be found on aerosol surfaces [Tervahattu et al., 2002a, 2002b, 2005, and references therein].

[4] Laboratory studies have demonstrated that organic surfaces can be oxidized by OH and O<sub>3</sub> [e.g., Rudich, 2003, and references therein; Thornberry and Abbatt, 2004; Molina et al., 2004]. Several of these studies have shown volatilization of OVOC resulting from organic surface oxidation. Molina et al. [2004], for example, report the full volatilization of a C<sub>18</sub> alkane monolayer following heterogeneous loss of 2–3 OH radicals to the surface, with many (though not exclusively) OVOC products. Evidence for such chemistry in the ambient environment include the demonstration by Grannas et al. [2004] that photooxidation of snow phase organic matter may explain the daytime flux of lightweight OVOC from snowpack to the boundary layer, and field observations that atmospheric OA becomes more oxidized with greater oxidant exposure and/or age [Gogou et al., 1996; de Gouw et al., 2005; Quinn et al., 2005; McFiggans et al., 2005; Robinson et al., 2006].

[5] Here, we use data collected on the NASA DC-8 aircraft during the INTEX-NA campaign (H. Singh et al., manuscript in preparation, 2006) to place constraints on OA oxidation's potential contribution to continental summertime OVOC budgets. This campaign was designed to examine the transport and transformation of air masses over continental scales. Most of the flights took place over the North American continent and the North Atlantic in the summer of 2004.

### 2. Method

[6] Aerosol size distributions, surface area, and volume for particles 10 nm to 3 μm in diameter were measured on the DC-8 using a differential mobility analyzer (DMA) and an optical particle counter (OPC) [Clarke et al., 2004]. Comparison of the DMA and OPC data to measurements of larger and ultrafine particles indicates that these instruments generally capture >90% of the total aerosol surface area except in a few select plumes. The aerosol size is quantified in conditions that are often dryer than the ambient atmosphere, so the aerosols may lose water (and thus mass) prior to measurement. Correcting for this effect is non-trivial, particularly for submicron particles, so we neglect it in our calculations. Based on the ratio of ambient to dry aerosol scattering coefficients, we believe the resulting underestimate of ambient surface area is significantly less than a factor

<sup>1</sup>Division of Engineering and Applied Science, California Institute of Technology, Pasadena, California, USA.

<sup>2</sup>Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California, USA.


<sup>3</sup>Department of Oceanography, University of Hawai'i at Manoa, Honolulu, Hawaii, USA.

<sup>4</sup>NASA Langley Research Center, Hampton, Virginia, USA.

<sup>5</sup>Department of Meteorology, Pennsylvania State University, University Park, Pennsylvania, USA.

<sup>6</sup>NASA Ames Research Center, Moffett Field, California, USA.

<sup>7</sup>Also at Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, California, USA.



**Figure 1.** Mean profiles of aerosol collision rates with (a) OH, (b) O<sub>3</sub>, and (c) H<sub>2</sub>O<sub>2</sub>. Horizontal lines show the interquartile range and x's are the elevation bin medians.

of 2. OH measurements were made by laser induced fluorescence [Faloona *et al.*, 2004], O<sub>3</sub> by chemiluminescence (M. Avery *et al.*, FASTOZ: An accurate, fast-response in situ ozone measurement system for aircraft campaigns, submitted to *Journal of Oceanic and Atmospheric Technology*, 2006), and H<sub>2</sub>O<sub>2</sub> by chemical ionization mass spectrometry (J. D. Crounse *et al.*, Measurements of gas-phase hydroperoxides by chemical ionization mass spectrometry, submitted to *Analytical Chemistry*, 2006).

[7] The collision rate of an oxidant with aerosol along the flight track was estimated using 1-minute averages of the measured aerosol surface area and oxidant mixing ratios:

$$\text{collisions} = \frac{1}{4} \times (8RT/\pi M)^{1/2} \times S \times [O_x] \times f(D)$$

where the factor of  $1/4$  converts aerosol surface area to cross sectional area,  $(8RT/\pi M)^{1/2}$  is the thermal speed of the oxidant ( $R$  is the universal gas constant,  $T$  is temperature,  $M$  is the oxidant molar mass),  $S$  the aerosol surface area,  $[O_x]$  the concentration of oxidant (converted to a 24-hour average in the case of OH), and  $f(D)$  the correction applied due to gas-phase diffusion limitations to large particles (for OH only) [Fuchs and Sutugin, 1971].


[8] For each point along the DC-8 flight track, we estimate a 24-hour average [OH] by scaling the observed [OH] by the ratio of the diurnally-averaged to instantaneous [OH] calculated from the highly-constrained NASA LaRC photochemical box model [Crawford *et al.*, 1999]. The main source of uncertainty in determining the diurnal average is that cloud effects, which during INTEX-NA normally altered actinic flux  $\sim 20\%$  relative to clear sky conditions, are assumed to be constant throughout the day, though they are generally transient. We expect that our large data set sufficiently captures the variability of cloud effects such that they will not significantly bias our estimate.

[9] Following the studies of Bertram *et al.* [2001] and Molina *et al.* [2004] we assume that each OH collision is reactive ( $\gamma = 1$ ) and volatilizes 6 organic carbons. Although the alkane used in the Molina *et al.* [2004] study is not representative of all OA surfaces, long chain fatty acids may comprise a significant fraction [Tervahattu *et al.*, 2002a, 2002b, 2005]. These parameters, which also assume full aerosol surface coverage by organic substrates, represent by far the largest uncertainties in our analysis. Because our assumptions imply a unity accommodation coefficient ( $\alpha = 1$ ), we account for diffusion limitations in calculating the OH collision rate.

[10] Estimating the OVOC flux from aerosol collisions with O<sub>3</sub> and H<sub>2</sub>O<sub>2</sub> is significantly more challenging and is not attempted here. While OH is highly reactive with many classes of organic compounds, O<sub>3</sub> reactivity and product yield are very substrate dependent [Rudich, 2003; Thornberry and Abbatt, 2004]; such selectivity also means that  $\gamma_{O_3}$  would be time dependent as reactive sites are depleted [Poschl *et al.*, 2001; Ammann *et al.*, 2003]. Also, unlike for OH [Molina *et al.*, 2004], O<sub>3</sub> reactivity with solid organic surfaces may depend on relative humidity [Poschl *et al.*, 2001]. For H<sub>2</sub>O<sub>2</sub>, we found no experimental studies allowing us to constrain  $\gamma$  or the product yield. For these oxidants, though, estimating the collision rate with aerosols is a first step for assessing their potential contributions to OVOC flux. We expect that the accommodation coefficients for these two oxidants are significantly less than unity – Berkowitz *et al.* [2001], for example, estimate  $\alpha_{O_3} \sim 10^{-3}$  – and thus do not consider diffusion limitations for their collision rates.

### 3. Results and Discussion

[11] Calculated 24-hour average collision rates are plotted as a function of elevation in Figure 1. For OH (Figure 1a) and O<sub>3</sub> (Figure 1b), most of the variability is driven by the (dry) aerosol surface area, which varies from  $\sim 10 \mu\text{m}^2 \text{cm}^{-3}$  in the upper troposphere to  $\sim 150 \mu\text{m}^2 \text{cm}^{-3}$  in the lowest elevation bin. For OH, we estimate the upper tropospheric collision rate to be  $\sim 1 \times 10^3 \text{ collisions cm}^{-3} \text{ s}^{-1}$ . Our assumption of carbon volatilization from Molina *et al.* [2004] thus yields an estimated OVOC source of  $\sim 70 \text{ pptv C/day}$  in the upper troposphere. A significantly larger source



**Figure 2.** Modeled (circle) vs. measured (x) profiles for (a) acetaldehyde and (b) peroxyacetic acid. Marks are bin medians, lines are interquartile range. Model predictions, offset by 0.25 km for clarity, are based on the photochemical box model of Crawford *et al.* [1999], and assume a PAA lifetime of  $\sim 20$  days (upper limit).

would exist in the lower troposphere ( $\sim 500$  pptv C/day in the lowest 2 km), but the contribution of this mechanism to OVOC concentrations will be less significant because of the much larger OVOC sources from gas phase oxidation of larger hydrocarbons in this region.

[12] Assuming a product yield for  $O_3$  similar to that of OH, the ratio of OH to  $O_3$  collisions gives a value of  $\gamma_{O_3} \sim 10^{-6}$  necessary for the OVOC flux from  $O_3$  to equal that from OH. Achieving this value would require only  $\sim 0.1\%$  surface coverage by typical alkene liquids ( $\gamma_{O_3} \sim 10^{-3}$  [de Gouw and Lovejoy, 1998; Rudich, 2003]). Thus, it is plausible that for certain aerosols reaction with  $O_3$  can be a significant oxidation mechanism. This is particularly true in the lower troposphere, where most primary (i.e., more unsaturated) OA is expected to reside. For  $H_2O_2$  (Figure 1c), the equivalent  $\gamma_{H_2O_2}$  ranges from  $\sim 10^{-5}$  in the lower troposphere to  $\sim 10^{-2}$  in the upper troposphere, so  $H_2O_2$ 's contribution to OVOC flux is likely confined to the lower troposphere or in cloud. Because of clouds' high aerosol surface area, increased actinic flux, and potential for aqueous phase chemistry, aerosol oxidation in clouds may warrant particular attention.

[13] Photolysis can, in principle, also lead to the decomposition of OA [Kieber *et al.*, 1990]. To be competitive with the OH chemistry, however, photolysis rates would have to be relatively fast ( $J \sim 10^{-6} s^{-1}$ ), as photolysis is likely less efficient at degrading organic molecules in the condensed

phase than in the gas phase: caging effects can stabilize the intermediates and aid their recombination or polymerization, preventing volatilization.

#### 4. Atmospheric Implications

[14] A dispersed and previously unconsidered source of OVOC from OA oxidation may have important implications for tropospheric photochemistry. For example, measurements of acetaldehyde ( $CH_3CHO$ ) have routinely exceeded model predictions [Singh *et al.*, 2001, 2004; Wisthaler *et al.*, 2002]; this was also true during INTEX-NA (Figure 2a). For each acetaldehyde measurement, we divide the difference between measured and box model-predicted mixing ratios by the photochemical lifetime to estimate the flux necessary to reconcile the difference. We find that a source of  $\sim 90$  pptv/day ( $\sim 180$  pptv C/day) is required to sustain the observed acetaldehyde concentrations in the upper troposphere. INTEX-NA also marked the first extensive airborne measurements of peroxyacetic acid ( $CH_3C(O)OOH$ , PAA), which significantly exceeded model-predicted values in the upper troposphere as well (Figure 2b); a similar analysis for PAA yields an estimated missing source of  $\sim 20$ – $200$  pptv C/day (J. Crouse *et al.*, manuscript in preparation, 2006). Even considering only acetaldehyde, oxidation by OH alone is likely too small to explain the upper tropospheric discrepancies between OVOC measurements and models; other oxidation mechanisms or alternative explanations are needed.

[15] A significant flux of OVOC from aerosol would have consequent impacts on  $HO_x$  and peroxyacyl nitrates. In fact, if the observations and our understanding of the subsequent chemistry of acetaldehyde are correct, peroxyacetyl nitrate ( $CH_3C(O)OONO_2$ , PAN) formation is very fast in the upper troposphere. Observations of PAN are not, however, consistent with those of acetaldehyde, based on current understanding of the chemistry that links these compounds [Staudt *et al.*, 2003].

[16] A large OVOC flux from aerosol is also incompatible with current estimates of OA budgets. The Intergovernmental Panel on Climate Change [2001] estimate of OA production (and loss) is  $\sim 150$  Tg "organic matter"/yr, or  $\sim 100$  Tg C/yr. If our estimated OA oxidation rate from OH were representative of the entire atmosphere, the global flux of organic carbon from aerosol would be as large as  $\sim 150$  Tg C/yr (integrating with bin median collision rates,  $\sim 100$  Tg C/yr; interquartile range,  $\sim 50$ – $200$  Tg C/yr). This is clearly an upper limit due to the assumptions of unity  $\gamma$  and that the continental summertime conditions of INTEX-NA are representative. Nonetheless, even a fraction of this large flux would imply that oxidation may need to be included in OA models, which currently consider only depositional loss. From our carbon flux (and assuming internally mixed aerosols with density  $1 g cm^{-3}$  and organic carbon fraction 0.5), we estimate the median lifetime of aerosol organic carbon to be  $\sim 10$  days, similar to estimates of carbonaceous aerosol lifetime against deposition of  $\sim 5$ – $10$  days [Kanakidou *et al.*, 2005]; thus, oxidation may be an important sink, particularly in the upper troposphere and regions with minimal precipitation.

[17] Consideration of an additional significant sink of OA from oxidation would dramatically increase the required global OA production inferred from top-down analyses

(generally calculated from measured OA burdens and estimates of the depositional loss rate). Bottom-up estimates, deduced from emission inventories and secondary organic aerosol (SOA) yields for precursor gases, may also be too low: *Holzinger et al.* [2005], for example, demonstrate that many biogenic SOA precursors are too short-lived to have been previously measured, and thus have been omitted from emission inventories. Furthermore, because of the diversity of SOA precursor gases, the photochemistry and SOA yield of only a few model compounds have been extensively studied; even for relatively well studied compounds, such as isoprene, estimates of SOA yields are undergoing significant revision upward [*Limbeck et al.*, 2003; *Claeys et al.*, 2004a, 2004b; *Kanakidou et al.*, 2005; *Kroll et al.*, 2005]. In addition, our understanding of other aspects of OA chemistry is poor. For example, *Heald et al.* [2005], considering only wet depositional loss, underpredict OA mass in the free troposphere 1–2 orders of magnitude, which they cannot attribute merely to OA flux underestimates.

## 5. Conclusions and Recommendations

[18] Our estimate of the OVOC source and its atmospheric impact are highly speculative and uncertain due to the complexity of the processes involved and the paucity of laboratory and field data for quantifying key parameters. Continued identification of OA constituents and study of oxidant interactions with a wider range of substrates is necessary to better constrain OVOC flux from atmospheric aerosols. Of particular concern is that much of the OA in the atmosphere may actually consist of highly oxidized humic-like substances (HULIS) [*Limbeck et al.*, 2003; *Claeys et al.*, 2004a]; whether HULIS can be volatilized in a similar fashion as the aliphatic compounds usually studied in the laboratory must be investigated. *De Gouw and Lovejoy* [1998] find that  $O_3$  reacts with liquid aldehydes and ketones ( $\gamma \sim 10^{-4}$ ), presumably with the carbonyl moieties, but do not determine if any gas phase products form. An additional difficulty in applying laboratory studies is that they have utilized reaction parameters, such as low pressures and  $[O_2]$ , and high  $[O_3]$  and  $[OH]$ , which are not representative of the real atmosphere. *Moise and Rudich* [2000] find, for example, that  $\gamma_{O_3}$  drops when  $O_2$  instead of He is used as a carrier gas in their experiments; also, *Molina et al.* [2004] propose that the carbon volatilization in their experiments would be reduced at atmospheric  $[O_2]$ . Other oxidants, such as  $NO_3$  and  $-$  in certain areas such as coastal and polar regions – halogen atoms, may also play an important role in aerosol oxidation. Finally, our analysis assumes that all aerosols are completely covered with organic films; accurate parameterizations of surface coverage require more field studies of aerosol coatings. A full understanding of OA oxidation will only result from continued, multifaceted research endeavors.

[19] **Acknowledgments.** We thank Jonathan Abbatt (University of Toronto) for reviewing a draft of this manuscript, Steven Howell (University of Hawai'i at Manoa) for helpful discussions, Clyde Brown (NASA Langley Research Center) for preparing merged data files, and the NASA Tropospheric Chemistry Program office for support of the INTEX-NA campaign (grant NNG04FA59G). This material is based upon work supported under a National Science Foundation Graduate Research Fellowship (AJK). This article was also developed under a STAR Research Assistance Agreement FP-91633401-2 awarded by the U.S. Environmental Protection Agency

(JDC). It has not been formally reviewed by the EPA. The views expressed in this document are solely those of the authors and the EPA does not endorse any products or commercial services mentioned in this publication.

## References

- Ammann, M., U. Poschl, and Y. Rudich (2003), Effects of reversible adsorption and Langmuir-Hinshelwood surface reactions on gas uptake by atmospheric particles, *Phys. Chem. Chem. Phys.*, *5*, 351–356, doi:10.1039/b208708a.
- Berkowitz, C. M., et al. (2001), Aircraft observations of aerosols,  $O_3$ , and NOy in a nighttime urban plume, *Atmos. Environ.*, *35*, 2395–2404.
- Bertram, A. K., A. V. Ivanov, M. Hunter, L. T. Molina, and M. J. Molina (2001), The reaction probability of OH on organic surfaces of tropospheric interest, *J. Phys. Chem. A*, *105*, 9415–9421.
- Claeys, M., et al. (2004a), Formation of secondary organic aerosols through photooxidation of isoprene, *Science*, *303*, 1173–1176.
- Claeys, M., et al. (2004b), Formation of secondary organic aerosols from isoprene and its gas-phase oxidation products through reaction with hydrogen peroxide, *Atmos. Environ.*, *38*, 4093–4098.
- Clarke, A. D., et al. (2004), Size-distributions and mixtures of black carbon and dust aerosol in Asian outflow: Physico-chemistry, optical properties, *J. Geophys. Res.*, *109*, D15S09, doi:10.1029/2003JD004378.
- Crawford, J., et al. (1999), Assessment of upper tropospheric HOx sources over the tropical Pacific based on NASA GTE/PEM data: Net effect on HOx and other photochemical parameters, *J. Geophys. Res.*, *104*(D13), 16,255–16,273.
- de Gouw, J. A., and E. R. Lovejoy (1998), Reactive uptake of ozone by liquid organic compounds, *Geophys. Res. Lett.*, *25*(6), 931–934.
- de Gouw, J. A., et al. (2005), Budget of organic carbon in a polluted atmosphere: Results from the New England Air Quality Study in 2002, *J. Geophys. Res.*, *110*, D16305, doi:10.1029/2004JD005623.
- Ellison, G. B., A. F. Tuck, and V. Vaida (1999), Atmospheric processing of organic aerosols, *J. Geophys. Res.*, *104*(D9), 11,633–11,641.
- Faloona, I. C., et al. (2004), A laser-induced fluorescence instrument for detecting tropospheric OH and HO<sub>2</sub>: Characteristics and calibration, *J. Atmos. Chem.*, *47*, 139–167.
- Fuchs, N. A., and A. G. Sutugin (1971), High-dispersed aerosols, in *Topics in Current Aerosol Research*, part 2, edited by G. M. Hidy and J. R. Block, pp. 1–60, Elsevier, New York.
- Gogou, A., et al. (1996), Differences in lipid and organic salt constituents of aerosols from eastern and western Mediterranean coastal cities, *Atmos. Environ.*, *28*(7), 1301–1310.
- Grannas, A. M., P. B. Shepson, and T. R. Filley (2004), Photochemistry and nature of organic matter in Arctic and Antarctic snow, *Global Biogeochem. Cycles*, *18*, GB1006, doi:10.1029/2003GB002133.
- Heald, C. L., D. J. Jacob, R. J. Park, L. M. Russell, B. J. Huebert, J. H. Seinfeld, H. Liao, and R. J. Weber (2005), A large organic aerosol source in the free troposphere missing from current models, *Geophys. Res. Lett.*, *32*, L18809, doi:10.1029/2005GL023831.
- Holzinger, R., A. Lee, K. T. Paw U, and A. H. Goldstein (2005), Observations of oxidation products above a forest imply biogenic emissions of very reactive compounds, *Atmos. Chem. Phys.*, *5*, 67–75.
- Intergovernmental Panel on Climate Change (2001), *Climate Change 2001: The Scientific Basis—Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge Univ. Press, New York. (Available at [http://www.grida.no/climate/ipcc\\_tar/wg1/](http://www.grida.no/climate/ipcc_tar/wg1/).)
- Kanakidou, M., et al. (2005), Organic aerosol and global climate modeling: A review, *Atmos. Chem. Phys.*, *5*, 1053–1123.
- Kieber, R. J., X. Zhou, and K. Mopper (1990), Formation of carbonyl compounds from UV-induced photodegradation of humic substances in natural waters: Fate of riverine carbon in the sea, *Limnol. Oceanogr.*, *35*, 1503–1515.
- Kroll, J. H., N. L. Ng, S. M. Murphy, R. C. Flagan, and J. H. Seinfeld (2005), Secondary organic aerosol formation from isoprene photooxidation under high-NOx conditions, *Geophys. Res. Lett.*, *32*, L18808, doi:10.1029/2005GL023637.
- Limbeck, A., M. Kulmala, and H. Puxbaum (2003), Secondary organic aerosol formation in the atmosphere via heterogeneous reaction of gaseous isoprene on acidic particles, *Geophys. Res. Lett.*, *30*(19), 1996, doi:10.1029/2003GL017738.
- McFiggans, G., et al. (2005), Simplification of the representation of the organic component of atmospheric particulates, *Faraday Disc.*, *130*, 341–362, doi:10.1039/b419435.
- Moise, T., and Y. Rudich (2000), Reactive uptake of ozone by proxies for organic aerosols: Surface versus bulk processes, *J. Geophys. Res.*, *105*(D11), 6469–6476.
- Molina, M. J., A. V. Ivanov, S. Trakhtenberg, and L. T. Molina (2004), Atmospheric evolution of organic aerosol, *Geophys. Res. Lett.*, *31*, L22104, doi:10.1029/2004GL020910.

- Muller, J.-F., and G. Brasseur (1999), Sources of upper tropospheric HOx: A three-dimensional study, *J. Geophys. Res.*, *104*(D1), 1705–1715.
- Murphy, D. M., D. S. Thomson, and M. J. Mahoney (1998), In situ measurements of organics, meteoric material, mercury, and other elements in aerosols at 5 to 19 kilometers, *Science*, *282*, 1664–1669.
- Poschl, U., T. Letzel, C. Schauer, and R. Niessner (2001), Interaction of ozone and water vapor with spark discharge soot aerosol particles coated with benzo[a]pyrene: O<sub>3</sub> and H<sub>2</sub>O adsorption, benzo[a]pyrene degradation, and atmospheric implications, *J. Phys. Chem. A*, *105*, 4029–4041.
- Quinn, P. K., et al. (2005), Impact of particulate organic matter on the relative humidity dependence of light scattering: A simplified parameterization, *Geophys. Res. Lett.*, *32*, L22809, doi:10.1029/2005GL024322.
- Robinson, A. L., N. M. Donahue, and W. F. Rogge (2006), Photochemical oxidation and changes in molecular composition of organic aerosol in the regional context, *J. Geophys. Res.*, *111*, D03302, doi:10.1029/2005JD006265.
- Rudich, Y. (2003), Laboratory perspectives on the chemical transformations of organic matter in atmospheric particles, *Chem. Rev.*, *103*, 5097–5124.
- Singh, H. B., M. Kanakidou, P. J. Crutzen, and D. J. Jacob (1995), High concentrations and photochemical fate of oxygenated hydrocarbons in the global troposphere, *Nature*, *378*, 50–54.
- Singh, H., et al. (2001), Evidence from the Pacific troposphere for large global sources of oxygenated organic compounds, *Nature*, *410*, 1078–1081.
- Singh, H. B., et al. (2004), Analysis of the atmospheric distribution, sources, and sinks of oxygenated volatile organic chemicals based on measurements over the Pacific during TRACE-P, *J. Geophys. Res.*, *109*, D15S07, doi:10.1029/2003JD003883.
- Staudt, A. C., D. J. Jacob, F. Ravetta, J. A. Logan, D. Bachiochi, T. N. Krishnamurti, S. Sandholm, B. Ridley, H. B. Singh, and B. Talbot (2003), Sources and chemistry of nitrogen oxides over the tropical Pacific, *J. Geophys. Res.*, *108*(D2), 8239, doi:10.1029/2002JD002139.
- Tervahattu, H., et al. (2002a), New evidence of an organic layer on marine aerosols, *J. Geophys. Res.*, *107*(D7), 4053, doi:10.1029/2000JD000282.
- Tervahattu, H., J. Juhanaja, and K. Kupiainen (2002b), Identification of an organic coating on marine aerosol particles by TOF-SIMS, *J. Geophys. Res.*, *107*(D16), 4319, doi:10.1029/2001JD001403.
- Tervahattu, H., et al. (2005), Fatty acids on continental sulfate aerosol particles, *J. Geophys. Res.*, *110*, D06207, doi:10.1029/2004JD005400.
- Thornberry, T., and J. P. D. Abbatt (2004), Heterogeneous reaction of ozone with liquid unsaturated fatty acids: Detailed kinetics and gas-phase product studies, *Phys. Chem. Chem. Phys.*, *6*, 84–93.
- Wennberg, P. O., et al. (1998), Hydrogen radicals, nitrogen radicals, and the production of ozone in the upper troposphere, *Science*, *279*, 49–53.
- Wisthaler, A., A. Hansel, R. R. Dickerson, and P. J. Crutzen (2002), Organic trace gas measurements by PTR-MS during INDOEX 1999, *J. Geophys. Res.*, *107*(D19), 8024, doi:10.1029/2001JD000576.
- 
- B. E. Anderson, M. A. Avery, and J. H. Crawford, NASA Langley Research Center, Hampton, VA 23681, USA.
- W. H. Brune, Department of Meteorology, Pennsylvania State University, University Park, PA 16802, USA.
- A. D. Clarke, C. S. McNaughton, and Y. Shinozuka, Department of Oceanography, University of Hawai'i at Manoa, Honolulu, HI 96822, USA.
- J. D. Crouse, Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA 91125, USA.
- A. J. Kwan and P. O. Wennberg, Division of Engineering and Applied Science, California Institute of Technology, Pasadena, CA 91125, USA. (kwan@caltech.edu)
- H. B. Singh, NASA Ames Research Center, Moffett Field, CA 94035, USA.