

A CATEGORICAL PROOF OF THE PARSHIN RECIPROCITY LAWS ON ALGEBRAIC SURFACES

DENIS OSIPOV, XINWEN ZHU

ABSTRACT. We define and study the 2-category of torsors over a Picard groupoid, a central extension of a group by a Picard groupoid, and commutator maps in this central extension. Using this in the context of two-dimensional local fields and two-dimensional adèle theory we obtain the two-dimensional tame symbol and a new proof of Parshin reciprocity laws on an algebraic surface.

1. INTRODUCTION

Let C be a projective algebraic curve over a perfect field k . Then there is the following famous Weil reciprocity law:

$$(1.1) \quad \prod_{p \in C} \text{Nm}_{k(p)/k} \{f, g\}_p = 1,$$

where $f, g \in k(C)^\times$, $\{f, g\}_p$ is the one-dimensional tame symbol, which is equal to $(-1)^{\nu_p(f)\nu_p(g)} \frac{f^{\nu_p(g)}}{g^{\nu_p(f)}}(p)$, and $k(p)$ is the residue field of the point p . The product (1.1) contains only finitely many non-equal to 1 terms.

There is the proof of Weil reciprocity law (and the analogous reciprocity law for residues of rational differential forms: sum of residues equals to zero) by reduction to the case of \mathbf{P}_k^1 using the connection between tame symbols (and residues of differentials) in extensions of local fields, see, for example, [S, ch. 2-3].

On the other hand, Tate gave in [T] the definition of local residue of differential form as some trace of an infinite-dimensional matrix. Starting from this definition he gave an intrinsic proof of the residue formula on a projective algebraic curve C using the fact that $\dim_k H^i(C, \mathcal{O}_C) < \infty$, $i = 0, 1$.

The multiplicative analog of Tate's approach, i.e. the case of the tame symbol and the proof of Weil reciprocity law, was done later by Arbarello, De Concini and Kac in [ACK]. They used the central extension of some infinite dimensional group of matrices $\text{GL}(K)$ by the group k^\times such that the group $\text{GL}(K)$ acts on the field $K = k((t))$, and obtained the tame symbol up to sign as the commutator of the lifting of two elements from $K^\times \subset \text{GL}(K)$ to this central extension. Hence, as in Tate's proof mentioned above, they obtained an intrinsic proof of the Weil reciprocity law on an algebraic curve. However, in this proof the exterior algebra of finite-dimensional k -vector spaces was used. Therefore difficult sign conventions were used in this paper to obtain the reciprocity law. To avoid these difficulties, in [BBE] Beilinson, Bloch and Esnault used the category of graded lines instead of the category of lines. The category of graded lines has non-trivial commutativity constraints multipliers $(-1)^{mn}$, where $m, n \in \mathbb{Z}$ are corresponding gradings. In other words, they used the Picard groupoid of graded lines which is a non-strictly commutative instead of

strictly commutative Picard groupoid. It was the first application of this notion of non-strictly commutative Picard groupoid.

Now let X be an algebraic surface over a perfect field k . For any pair $x \in C$, where $C \subset X$ is a curve that $x \in C$ is a closed point, it is possible to define the ring $K_{x,C}$ such that $K_{x,C}$ is isomorphic to the two-dimensional local field $k(x)((t))((s))$ when x is a smooth point on C and X . If x is not a smooth point, then $K_{x,C}$ is a finite direct sum of two-dimensional local fields (see section 5.2 of this paper). For any two-dimensional local field $k'((t))((s))$ one can define the two-dimensional tame symbol of 3 variables with values in k'^{\times} , see section 4.1 and [Pa1], [Pa3, §3]. Parshin formulated and proved the reciprocity laws for two-dimensional tame symbols, but his proof was never published. Contrary to the 1-dimensional case, there are a lot of reciprocity laws for two-dimensional tame symbols, which belong to two types. For the first type we fix a point on the surface and will vary irreducible curves containing this point. For the second type we fix a projective irreducible curve on the surface and will vary points on this curve. Parshin's idea for the proof, for example, of more unexpected first type of reciprocity laws, was to use the chain of successive blowups of points on algebraic surfaces. Later, Kato generalized the reciprocity laws for excellent schemes by using the reduction to the reciprocity law of Bass and Tate for Milnor K-groups of some field $L(t)$, see [K, prop. 1]. He used them to construct an analog of the Gersten-Quillen complex for Milnor K-theory.

In this paper, we give a generalization of Tate's proof of the reciprocity law on an algebraic curve to the case of two-dimensional tame symbols and obtain an intrinsic proof of Parshin reciprocity laws for two-dimensional tame symbols on an algebraic surface.

To fulfill this goal, we first generalize the notion of a central extension of a group by a commutative group and of the commutator map associated to the central extension. More precisely, we define and study in some detail the properties of the category of central extensions of a group G by a (non strictly commutative) Picard groupoid \mathcal{P} . Roughly speaking, an object in this category is a rule to assign every $g \in G$ a \mathcal{P} -torsor, satisfying certain properties. For such a central extension \mathcal{L} we define a map $C_3^{\mathcal{L}}$ which is an analog of the commutator map. In this case when G is abelian, this commutator map is an anti-symmetric and tri-multiplicative map from G^3 to the group $\pi_1(\mathcal{P})$. Let us remark that to obtain some of these properties, we used the results of Breen from [Br3] on group-like monoidal 2-groupoids. We hope these constructions would be of some independent interest.

We then apply this formalism to $\mathcal{P} = \mathcal{P}ic^{\mathbb{Z}}$, where $\mathcal{P}ic^{\mathbb{Z}}$ stands for the Picard groupoid of graded lines. The key ingredient here is Kapranov's graded-determinantal theory from [Kap], which associates a $\mathcal{P}ic^{\mathbb{Z}}$ -torsor to every 1-Tate vector space (a.k.a. locally linearly compact vector space). This allows one to construct the central extension $\mathcal{D}et$ of $GL(\mathbb{K})$ by $\mathcal{P}ic^{\mathbb{Z}}$, where \mathbb{K} is a two-dimensional local field (or more generally, a 2-Tate vector space). It turns out that the two-dimensional tame symbol coincides with the commutator map $C_3^{\mathcal{D}et}$. Finally, using "semilocal" adèle complexes on an algebraic surface we obtain that the corresponding central extension constructed by semilocal fields on the surface is the trivial one. This leads us to a new proof of Parshin's reciprocity laws on an algebraic surface, which is distinct from both Parshin's original approach as well as Kato's.

Our approach to the reciprocity laws on the algebraic surfaces has the following features. First, we use the non-strictly commutative Picard groupoid, which can

be regarded as another application of this notion after [BBE]. However, unlike the one-dimensional case where one can just play with the usual Picard groupoid of lines (though complicated, as done in [ACK]), the use of $\mathcal{P}ic^{\mathbb{Z}}$ is essential here. This indicates that the non-strictly commutative Picard groupoid is an important and fruitful mathematical object that deserves further attention. Also, in order to apply this notion, we develop certain constructions in higher categories (e.g. the commutator map $C_3^{\mathcal{L}}$), which could be potentially useful elsewhere. Second, as in the one-dimensional case, our approach uses certain local-to-global (in other words, factorization) principle. Since the local-to-global (factorization) principle in the one-dimensional story is very important in the Langlands program and conformal field theory, we hope our approach is just a shadow of a whole fascinating yet explored realm of mathematics. Finally, our approach can be generalized by replacing the ground field k by an Artinian ring A (and even more general rings) and we can obtain the reciprocity laws for two-dimensional Contou-Carrère symbols. By choosing appropriate A , this specializes to residue formulas for algebraic surfaces¹. We will carefully discuss this in the next paper.

The paper is organized as follows. In section 2 we describe some categorical constructions, which we need further on. In section 2.1 we recall the definition of a Picard groupoid. In section 2.2 we discuss the difference between strictly commutative and non-strictly commutative Picard groupoids. In section 2.3 we describe the 2-category of \mathcal{P} -torsors, where \mathcal{P} is a Picard groupoid. In section 2.4 we study the Picard groupoid of homomorphisms from a group G to a Picard groupoid \mathcal{P} and describe the "commutator" of two commuting elements from G with values in $\pi_1(\mathcal{P})$. In section 2.5 we define and study the Picard 2-groupoid of central extensions of a group G by a Picard groupoid \mathcal{P} . We define and study properties of the commutator category of such a central extension, and finally study the "commutator" of three commuting elements from G with values in $\pi_1(\mathcal{P})$. This section may be of independent interest.

In section 3 we recall the theory of graded-determinantal theories on Tate vector spaces. We recall the definition and basic properties of the category of n -Tate vector spaces in section 3.1. In section 3.2 we recall the definition of determinant functor from the exact category $(\text{Tate}_0, \text{isom})$ to the Picard groupoid $\mathcal{P}ic^{\mathbb{Z}}$ of graded lines and the definition of graded-determinantal theory on the exact category Tate_1 of 1-Tate vector spaces.

In section 4 we apply the constructions given above to one-dimensional and two-dimensional local fields. In section 4.1 we review one-dimensional and two-dimensional tame symbols. In section 4.2 we obtain a description of the one-dimensional (usual) tame symbol as some commutator. In section 4.3 we obtain the two-dimensional tame symbol as commutator of 3 elements in some central extension of the group $K^{\times} = k((t))((s))^{\times}$ by the Picard groupoid $\mathcal{P}ic^{\mathbb{Z}}$.

In section 5 we obtain the reciprocity laws. In section 5.1 we give the proof of Weil reciprocity law using the constructions given above and adèle complexes on a curve. In section 5.2 we apply the previous results in order to obtain a proof of Parshin's reciprocity laws on an algebraic surface using "semilocal" adèle complexes on an algebraic surface.

¹The generalization of Tate's approach to the n -dimensional residue of differential form was done in [Be1], but this note contains no proofs.

Acknowledgments. The first author is grateful to L. Breen, M. Kapranov and A.N. Parshin for useful discussions on categorical approach to the two-dimensional tame symbol at various periods of time. The first author was financially supported by Russian Foundation for Basic Research (grant no. 08-01-00095) and by the Programme for the Support of Leading Scientific Schools of the Russian Federation (grant no. NSh-4713.2010.1). The second author thanks V. Kac for useful discussions. The research of the second author is supported by NSF grant under DMS-1001280.

2. GENERAL NONSENSE

2.1. Picard groupoid. Let \mathcal{P} be a Picard groupoid, i.e. a symmetric monoidal group-like groupoid. Let us recall that this means that \mathcal{P} is a groupoid, together with a bifunctor

$$+ : \mathcal{P} \times \mathcal{P} \rightarrow \mathcal{P}$$

and natural (functorial) isomorphisms

$$a_{x,y,z} : (x + y) + z \simeq x + (y + z),$$

called the associativity constraints, and natural (functorial) isomorphisms

$$c_{x,y} : x + y \simeq y + x,$$

called the commutativity constraints, such that:

- (i) For each $x \in \mathcal{P}$, the functor $y \mapsto x + y$ is an equivalence.
- (ii) The pentagon axiom holds, i.e. the following diagram is commutative

$$(2.1) \quad \begin{array}{ccc} & (x + y) + (z + w) & \\ & \swarrow \quad \searrow & \\ x + (y + (z + w)) & & ((x + y) + z) + w \\ \downarrow & & \downarrow \\ x + ((y + z) + w) & \xrightarrow{\quad\quad\quad} & (x + (y + z)) + w \end{array}$$

- (iii) The hexagon axiom holds, i.e. the following diagram is commutative

$$(2.2) \quad \begin{array}{ccccc} & & (x + y) + z & & \\ & & \swarrow \quad \searrow & & \\ (y + x) + z & & & & x + (y + z) \\ \downarrow & & & & \downarrow \\ y + (x + z) & & & & x + (z + y) \\ & \swarrow \quad \searrow & & \swarrow \quad \searrow & \\ & (x + z) + y & & & \end{array}$$

- (iv) For any $x, y \in \mathcal{P}$, $c_{y,x}c_{x,y} = id_{x+y}$.

A unit (e, φ) of \mathcal{P} is an object $e \in \mathcal{P}$ together with an isomorphism $\varphi : e + e \simeq e$. It is an exercise to show that (e, φ) exists and is unique up to a unique isomorphism. For any $x \in \mathcal{P}$, there is a unique isomorphism $e + x \simeq x$ such that the following diagram is commutative

$$\begin{array}{ccc} (e + e) + x & \xrightarrow{\quad\quad\quad} & e + (e + x) \\ & \searrow & \swarrow \\ & e + x & \end{array}$$

and therefore $x + e \simeq e + x \simeq x$. For any $x \in \mathcal{P}$, we choose an object, denoted by $-x$, together with an isomorphism $\phi_x : x + (-x) \simeq e$. The pair $(-x, \phi_x)$ is called an inverse of x , and it is unique up to a unique isomorphism. We choose for each x its inverse $(-x, \phi_x)$, then we have a canonical isomorphism

$$(2.3) \quad -(-x) \simeq e + (-(-x)) \simeq (x + (-x)) + (-(-x)) \simeq x + ((-x) + (-(-x))) \simeq x + e \simeq x,$$

and therefore a canonical isomorphism

$$(2.4) \quad (-x) + x \simeq (-x) + (-(-x)) \simeq e.$$

Observe that we have another isomorphism $(-x) + x \simeq x + (-x) \simeq e$ using the commutativity constraint. When the Picard groupoid \mathcal{P} is strictly commutative (cf. §2.2), these two isomorphisms are the same (cf. [Z, lemma 1.6]), but in general they are different.

If $\mathcal{P}_1, \mathcal{P}_2$ are two Picard groupoids, then $\text{Hom}(\mathcal{P}_1, \mathcal{P}_2)$ is defined as follows. Objects are 1-homomorphisms, i.e., functors $F : \mathcal{P}_1 \rightarrow \mathcal{P}_2$ together with isomorphisms $F(x + y) \simeq F(x) + F(y)$ such that the following diagrams are commutative:

$$(2.5) \quad \begin{array}{ccc} F((x + y) + z) & \xrightarrow{\quad\quad\quad} & (F(x) + F(y)) + F(z) \\ \downarrow & & \downarrow \\ F(x + (y + z)) & \xrightarrow{\quad\quad\quad} & F(x) + (F(y) + F(z)), \\ \\ F(x + y) & \xrightarrow{\quad\quad\quad} & F(x) + F(y) \\ \downarrow & & \downarrow \\ F(y + x) & \xrightarrow{\quad\quad\quad} & F(y) + F(x). \end{array}$$

Morphisms in $\text{Hom}(\mathcal{P}_1, \mathcal{P}_2)$ are 2-isomorphisms, i.e., natural transformations

$$\theta : F_1 \rightarrow F_2$$

such that the following diagram is commutative

$$\begin{array}{ccc} F_1(x + y) & \xrightarrow{\quad\quad\quad} & F_1(x) + F_1(y) \\ \theta \downarrow & & \downarrow \theta \\ F_2(x + y) & \xrightarrow{\quad\quad\quad} & F_2(x) + F_2(y). \end{array}$$

It is clear that $\text{Hom}(\mathcal{P}_1, \mathcal{P}_2)$ has a natural structure as a Picard groupoid. Namely,

$$(F_1 + F_2)(x) := F_1(x) + F_2(x),$$

and the isomorphism $(F_1 + F_2)(x + y) \simeq (F_1 + F_2)(x) + (F_1 + F_2)(y)$ is the unique one such that the following diagram is commutative

$$\begin{array}{ccc} F_1(x + y) + F_2(x + y) & \longrightarrow & (F_1(x) + F_1(y)) + (F_2(x) + F_2(y)) \\ \downarrow & & \swarrow \\ (F_1(x) + F_2(x)) + (F_1(y) + F_2(y)) & & \end{array}$$

The associativity constraints and the commutativity constraints for $\text{Hom}(\mathcal{P}_1, \mathcal{P}_2)$ are clear. If $\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3$ are three Picard groupoids, then $\text{Hom}(\mathcal{P}_1, \mathcal{P}_2; \mathcal{P}_3)$ is defined as $\text{Hom}(\mathcal{P}_1, \text{Hom}(\mathcal{P}_2, \mathcal{P}_3))$, called the Picard groupoid of bilinear homomorphisms from $\mathcal{P}_1 \times \mathcal{P}_2$ to \mathcal{P}_3 . The Picard groupoid of trilinear homomorphisms from $\mathcal{P}_1 \times \mathcal{P}_2 \times \mathcal{P}_3$ to \mathcal{P}_4 is defined similarly.

For a (small) monoidal group-like groupoid (or gr-category) C we denote by $\pi_0(C)$ the group² of isomorphism classes of objects. We denote by $\pi_1(C)$ the group $\text{Aut}_C(e)$, where e is the unit object of C . It follows that $\pi_1(C)$ is an abelian group. If C is a Picard groupoid, then $\pi_0(C)$ is also an abelian group.

2.2. Strictly commutative vs. non-strictly commutative Picard groupoids.

If the commutativity constraints c further satisfy $c_{x,x} = \text{id}$, then the Picard groupoid \mathcal{P} is called strictly commutative. It is a theorem of Deligne's (cf. [Del1]) that the 2-category of strictly commutative Picard groupoids is 2-equivalent to the 2-category of 2-term complexes of abelian groups concentrated on degree -1 and 0 , whose degree -1 term abelian groups are injective³.

Example 2.1. The most famous example is $\mathcal{P} = BA$, where A is an abelian group, and BA is the category of A -torsors. The tensor products of A -torsors make BA a strictly commutative Picard groupoid. The 2-term complex of abelian groups that represents BA under Deligne's theorem is any injective resolution of $A[1]$. If $A = k^\times$ is the group of invertible elements in a field k , then BA is also denoted by $\mathcal{P}ic$, which is the symmetric monoidal category of 1-dimensional k -vector spaces.

However, it is also important for us to consider the non strictly commutative Picard groupoids. The following example of a non-strictly commutative Picard groupoid is crucial.

Example 2.2. Let $\mathcal{P}ic^{\mathbb{Z}}$ denote the category of graded lines (i.e. 1-dimensional k -vector spaces with gradings), over a base field k . An object in $\mathcal{P}ic^{\mathbb{Z}}$ is a pair (ℓ, n) where ℓ is a 1-dimensional k -vector space, and n is an integer. The morphism set $\text{Hom}_{\mathcal{P}ic^{\mathbb{Z}}}((\ell_1, n_1), (\ell_2, n_2))$ is empty unless $n_1 = n_2$, and in this case, it is just $\text{Hom}_k(\ell_1, \ell_2) \setminus 0$. Observe that as a groupoid, $\mathcal{P}ic^{\mathbb{Z}}$ is not connected. In fact $\pi_0(\mathcal{P}) \simeq \mathbb{Z}$. The tensor product $\mathcal{P}ic^{\mathbb{Z}} \times \mathcal{P}ic^{\mathbb{Z}} \rightarrow \mathcal{P}ic^{\mathbb{Z}}$ is given as

$$(\ell_1, n_1) \otimes (\ell_2, n_2) \mapsto (\ell_1 \otimes \ell_2, n_1 + n_2).$$

There is a natural associativity constraint that makes $\mathcal{P}ic^{\mathbb{Z}}$ a monoidal groupoid.

Remark 2.1. For the Picard groupoids $\mathcal{P}ic$ and $\mathcal{P}ic^{\mathbb{Z}}$, we will often use in this article the usual notation " \otimes " for monoidal structures in these categories, although for a general Picard groupoid we denoted it as "+".

²The group structure on $\pi_0(C)$ is induced by the monoidal structure of C .

³In fact, Deligne's theorem holds in any topos.

We note that the commutativity constraint in category $\mathcal{P}ic^{\mathbb{Z}}$ is the interesting one. Namely,

$$c_{\ell_1, \ell_2} : (\ell_1 \otimes \ell_2, n_1 + n_2) \simeq (\ell_2 \otimes \ell_1, n_2 + n_1), \quad c_{\ell_1, \ell_2}(v \otimes w) = (-1)^{n_1 n_2} w \otimes v.$$

Of course, there is another commutativity constraint on the category of graded lines given by $c(v \otimes w) = w \otimes v$. Then as a Picard groupoid with this naive commutativity constraints, it is just the strictly commutative Picard groupoid $\mathcal{P}ic \times \mathbb{Z}$. There is a natural monoidal equivalence $\mathcal{P}ic^{\mathbb{Z}} \simeq \mathcal{P}ic \times \mathbb{Z}$, but this equivalence is NOT symmetric monoidal (i.e. a 1-homomorphism of Picard groupoids). We denote by

$$F_{\mathcal{P}ic} : \mathcal{P}ic^{\mathbb{Z}} \rightarrow \mathcal{P}ic$$

the natural monoidal functor.

The importance of $\mathcal{P}ic^{\mathbb{Z}}$ lies in the following observation. Let us make the following convention.

Convention. For any category \mathcal{C} we denote by $(\mathcal{C}, \text{isom})$ a category with the same objects as in the category \mathcal{C} , and morphisms in the category $(\mathcal{C}, \text{isom})$ are the isomorphisms in the category \mathcal{C} .

Now let Tate_0 be the category of finite dimensional vector spaces over a field k . The categories Tate_0 and $(\text{Tate}_0, \text{isom})$ are symmetric monoidal categories under the direct sum. The commutativity constraints in the categories Tate_0 and $(\text{Tate}_0, \text{isom})$ are defined in the natural way. Namely, the map $c_{V, W} : V \oplus W \rightarrow W \oplus V$ is given by $c_{V, W}(v, w) = (w, v)$. Then there is a natural symmetric monoidal functor

$$(2.6) \quad \det : (\text{Tate}_0, \text{isom}) \rightarrow \mathcal{P}ic^{\mathbb{Z}},$$

which assigns to every V its top exterior power and the grading $\dim V$, i.e. the dimension of the vector space V over the field k . Observe, however, that the functor $F_{\mathcal{P}ic} \circ \det : (\text{Tate}_0, \text{isom}) \rightarrow \mathcal{P}ic$ is NOT symmetric monoidal.

It is a folklore theorem that the category of Picard groupoid (not necessarily strict commutative) is equivalent to the category of spectra whose only non-vanishing homotopy groups are π_0 and π_1 ⁴. For example, $\mathcal{P}ic^{\mathbb{Z}}$ should correspond to the truncation $\tau_{\leq 1} \mathcal{K}$, where \mathcal{K} is the spectra of algebraic K -theory of k .

2.3. \mathcal{P} -torsors. Let \mathcal{P} be a Picard groupoid. Recall (see also [BBE, Appendix A6] and [Dr, § 5.1]) that a \mathcal{P} -torsor \mathcal{L} is a module category over \mathcal{P} , i.e., there is a bifunctor

$$+ : \mathcal{P} \times \mathcal{L} \rightarrow \mathcal{L}$$

together with natural isomorphisms

$$a_{x, y, v} : (x + y) + v \simeq x + (y + v), \quad x, y \in \mathcal{P}, v \in \mathcal{L},$$

satisfying

- (i) the pentagon axiom, i.e. a diagram similar to (2.1) holds;
- (ii) for any $x \in \mathcal{P}$, the functor from \mathcal{L} to \mathcal{L} given by $v \mapsto x + v$ is an equivalence;
- (iii) for any $v \in \mathcal{L}$, the functor from \mathcal{P} to \mathcal{L} given by $x \mapsto x + v$ is an equivalence of categories.

⁴Indeed, consider the geometrization of the nerve of \mathcal{P} . Then the Picard structure of \mathcal{P} puts an E_{∞} -structure on this space.

It is clear that we can verify the condition (ii) of this definition only for the unit object e of \mathcal{P} .

For any $v \in \mathcal{L}$, there is a unique isomorphism $e + v \simeq v$ such that the following diagram is commutative

$$\begin{array}{ccc} (e + e) + v & \xrightarrow{\quad} & e + (e + v) \\ & \searrow & \swarrow \\ & e + v & \end{array}$$

If $\mathcal{L}_1, \mathcal{L}_2$ are \mathcal{P} -torsors, then $\text{Hom}_{\mathcal{P}}(\mathcal{L}_1, \mathcal{L}_2)$ is the category defined as follows. Objects are 1-isomorphisms, i.e. equivalences $F : \mathcal{L}_1 \rightarrow \mathcal{L}_2$ together with isomorphisms $\lambda : F(x + v) \simeq x + F(v)$ such that the following diagram is commutative

$$(2.7) \quad \begin{array}{ccc} F((x + y) + v) & \longrightarrow & (x + y) + F(v) \\ \downarrow & & \downarrow \\ F(x + (y + v)) & \longrightarrow & x + (y + F(v)) \end{array}$$

Morphisms are natural transformations $\theta : F_1 \rightarrow F_2$ such that the following diagram is commutative

$$\begin{array}{ccc} F_1(x + v) & \longrightarrow & x + F_1(v) \\ \theta \downarrow & & \downarrow \theta \\ F_2(x + v) & \longrightarrow & x + F_2(v) \end{array}$$

From discussions above it follows that all \mathcal{P} -torsors form a 2-category, denoted by $B\mathcal{P}$. We will choose, once and for all, for any \mathcal{P} -torsors $\mathcal{L}_1, \mathcal{L}_2$ and $F \in \text{Hom}_{\mathcal{P}}(\mathcal{L}_1, \mathcal{L}_2)$, a quasi-inverse F^{-1} of F together with an isomorphism $F^{-1}F \simeq \text{id}$.

Moreover, $B\mathcal{P}$ is a category enriched over itself. That is, for any \mathcal{P} -torsors $\mathcal{L}_1, \mathcal{L}_2$ the category $\text{Hom}_{\mathcal{P}}(\mathcal{L}_1, \mathcal{L}_2)$ is again a \mathcal{P} -torsor, where an action of \mathcal{P} on $\text{Hom}_{\mathcal{P}}(\mathcal{L}_1, \mathcal{L}_2)$ is defined as follows: for any $z \in \mathcal{P}$, $v \in \mathcal{L}_1$, $F \in \text{Hom}_{\mathcal{P}}(\mathcal{L}_1, \mathcal{L}_2)$ we put $z + F \in \text{Hom}_{\mathcal{P}}(\mathcal{L}_1, \mathcal{L}_2)$ as $(z + F)(v) := z + F(v)$. Now the isomorphism λ for the equivalence $z + F$ is defined by means of the braiding maps c in \mathcal{P} (commutativity constraints from section 2.1). Then the diagram (2.7) for the equivalence $z + F$ follows from hexagon diagram (2.2). It is clear that this definition is extended to the definition of a bifunctor

$$(2.8) \quad + : \mathcal{P} \times \text{Hom}_{\mathcal{P}}(\mathcal{L}_1, \mathcal{L}_2) \rightarrow \text{Hom}_{\mathcal{P}}(\mathcal{L}_1, \mathcal{L}_2)$$

such that the axioms of \mathcal{P} -torsor are satisfied (see the beginning of this section).

We note that to prove that the category $B\mathcal{P}$ is enriched over itself we used the commutativity constraints in \mathcal{P} . The commutativity constraints will be important also below to define the sum of two \mathcal{P} -torsors.

The category $B\mathcal{P}$ furthermore forms a Picard 2-groupoid. We will not make the definition of Picard 2-groupoids precise. (However, one refers to [KV, Br2] for details). We will only describe the Picard structure on $B\mathcal{P}$ in the way we need.

First, if $\mathcal{L}_1, \mathcal{L}_2$ are two \mathcal{P} -torsors, then $\mathcal{L}_1 + \mathcal{L}_2$ is defined to be the category whose objects are pairs (v, w) , where $v \in \mathcal{L}_1$ and $w \in \mathcal{L}_2$. The morphisms from (v, w) to (v', w') are defined as the equivalence classes of triples $(x, \varphi_1, \varphi_2)$, where $x \in \mathcal{P}$, $\varphi_1 \in \text{Hom}_{\mathcal{L}_1}(v, x + v')$ and $\varphi_2 \in \text{Hom}_{\mathcal{L}_2}(x + w, w')$, and $(x, \varphi_1, \varphi_2) \sim (y, \phi_1, \phi_2)$ if

there exists a map $f : x \rightarrow y$ such that $\phi_1 = f(\varphi_1)$ and $\varphi_2 = f(\phi_2)$. The identity in $\text{Hom}_{\mathcal{L}_1 + \mathcal{L}_2}((v, w), (v, w))$ and the composition

$$\text{Hom}_{\mathcal{L}_1 + \mathcal{L}_2}((v, w), (v', w')) \times \text{Hom}_{\mathcal{L}_1 + \mathcal{L}_2}((v', w'), (v'', w'')) \rightarrow \text{Hom}_{\mathcal{L}_1 + \mathcal{L}_2}((v, w), (v'', w''))$$

are clear. (To define the composition we have to use the commutativity constraints in $B\mathcal{P}$.) So $\mathcal{L}_1 + \mathcal{L}_2$ is a category. Define the action of \mathcal{P} on $\mathcal{L}_1 + \mathcal{L}_2$ as $x + (v, w) := (x + v, w)$. The natural isomorphism $(x + y) + (v, w) \simeq x + (y + (v, w))$ is the obvious one. It is easy to check that $\mathcal{L}_1 + \mathcal{L}_2$ is a \mathcal{P} -torsor.

There is an obvious 1-isomorphism of \mathcal{P} -torsors

$$A : (\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3 \simeq \mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}_3),$$

which is the associativity constraint. Namely, objects in $(\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3$ and in $\mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}_3)$ are both canonically bijective to triples (v_1, v_2, v_3) where $v_i \in \mathcal{L}_i$. Then A is identity on objects. A morphism from (v_1, v_2, v_3) to (w_1, w_2, w_3) in $(\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3$ is of the form $(x, (y, \varphi_1, \varphi_2), \varphi_3)$, where $x, y \in \mathcal{P}$, $\varphi_1 : v_1 \rightarrow y + (x + w_1)$, $\varphi_2 : y + v_2 \rightarrow w_2$, $\varphi_3 : x + v_3 \rightarrow w_3$. Then A maps $(x, (y, \varphi_1, \varphi_2), \varphi_3)$ to $(x + y, \varphi'_1, (x, \varphi'_2, \varphi'_3))$, where $\varphi'_1 : v_1 \rightarrow (x + y) + w_1$ coming from $v_1 \xrightarrow{\varphi_1} y + (x + w_1) \simeq (y + x) + w_1 \simeq (x + y) + w_1$, $\varphi'_2 : (x + y) + v_2 \rightarrow x + w_2$ coming from $(x + y) + v_2 \simeq x + (y + v_2) \xrightarrow{x + \varphi_2} x + w_2$ and $\varphi'_3 : x + v_3 \rightarrow w_3$ is the same as φ_3 .

To complete the definition of A , we should specify for every $x \in \mathcal{P}$, $(v_1, v_2, v_3) \in (\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3$, an isomorphism $\lambda : A(x + (v_1, v_2, v_3)) \simeq x + A(v_1, v_2, v_3)$ such that the diagram (2.7) is commutative for $F = A$. It is clear that $\lambda = \text{id} : (x + v_1, v_2, v_3) = (x + v_1, v_2, v_3)$ will suffice for this purpose.

It is clear from definition of A that we can similarly construct a 1-morphism A^{-1} of \mathcal{P} -torsors such that the following equalities are satisfied:

$$A^{-1}A = AA^{-1} = \text{id}.$$

From above construction of the associativity constraints (1-morphisms A and A^{-1}) it follows that for any \mathcal{P} -torsors $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3, \mathcal{L}_4$ the following diagram of 1-morphisms (pentagon diagram) is commutative

$$(2.9) \quad \begin{array}{ccc} & (\mathcal{L}_1 + \mathcal{L}_2) + (\mathcal{L}_3 + \mathcal{L}_4) & \\ \swarrow & & \searrow \\ \mathcal{L}_1 + (\mathcal{L}_2 + (\mathcal{L}_3 + \mathcal{L}_4)) & & ((\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3) + \mathcal{L}_4 \\ \downarrow & & \downarrow \\ \mathcal{L}_1 + ((\mathcal{L}_2 + \mathcal{L}_3) + \mathcal{L}_4) & \xrightarrow{\quad\quad\quad} & (\mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}_3)) + \mathcal{L}_4 \end{array}$$

(To prove this diagram we note that this diagram is evident for objects from category $(\mathcal{L}_1 + \mathcal{L}_2) + (\mathcal{L}_3 + \mathcal{L}_4)$. To verify this diagram for morphisms from this category one needs to make some non-complicated routine calculations. The analogous reasonings are also applied to the diagram (2.13) below.)

The following axioms are satisfied in the category $B\mathcal{P}$ and describe the functoriality of the associativity constraints. Let $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3, \mathcal{L}'_1$ be any \mathcal{P} -torsors, and $\mathcal{L}_1 \rightarrow \mathcal{L}'_1$

be any 1-morphism of \mathcal{P} -torsors, then the following diagram of 1-morphisms is commutative

$$(2.10) \quad \begin{array}{ccc} (\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3 & \longrightarrow & (\mathcal{L}'_1 + \mathcal{L}_2) + \mathcal{L}_3 \\ \downarrow & & \downarrow \\ \mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}_3) & \longrightarrow & \mathcal{L}'_1 + (\mathcal{L}_2 + \mathcal{L}_3). \end{array}$$

Let $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3, \mathcal{L}'_2$ be any \mathcal{P} -torsors, and $\mathcal{L}_2 \rightarrow \mathcal{L}'_2$ be any 1-morphism of \mathcal{P} -torsors, then the following diagram of 1-morphisms is commutative

$$(2.11) \quad \begin{array}{ccc} (\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3 & \longrightarrow & (\mathcal{L}_1 + \mathcal{L}'_2) + \mathcal{L}_3 \\ \downarrow & & \downarrow \\ \mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}_3) & \longrightarrow & \mathcal{L}_1 + (\mathcal{L}'_2 + \mathcal{L}_3). \end{array}$$

Let $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3, \mathcal{L}'_3$ be any \mathcal{P} -torsors, and $\mathcal{L}_3 \rightarrow \mathcal{L}'_3$ be any 1-morphism of \mathcal{P} -torsors, then the following diagram of 1-morphisms is commutative

$$(2.12) \quad \begin{array}{ccc} (\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3 & \longrightarrow & (\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}'_3 \\ \downarrow & & \downarrow \\ \mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}_3) & \longrightarrow & \mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}'_3). \end{array}$$

(In diagrams (2.10)-(2.12) the vertical arrows are the associativity constraints.)

Next we define the commutativity constraints. Recall that we have chosen for each $x \in \mathcal{P}$ its inverse $(-x, \phi_x)$, and then obtained the isomorphism (2.4). This gives an obvious 1-isomorphism

$$C : \mathcal{L}_1 + \mathcal{L}_2 \simeq \mathcal{L}_2 + \mathcal{L}_1.$$

Namely, C will map the object (v_1, v_2) to (v_2, v_1) , and $(x, \varphi_1, \varphi_2) : (v_1, v_2) \rightarrow (w_1, w_2)$ to $(-x, \varphi'_1, \varphi'_2) : (v_2, v_1) \rightarrow (w_2, w_1)$, where $\varphi'_1 : v_2 \simeq e + v_2 \simeq (-x + x) + v_2 \simeq -x + (x + v_2) \xrightarrow{-x + \varphi_2} -x + w_2$ and $\varphi'_2 : -x + v_1 \xrightarrow{-x + \varphi_1} -x + (x + w_1) \simeq (-x + x) + w_1 \simeq e + w_1 \simeq w_1$.

We also define for each $x \in \mathcal{P}$, $(v_1, v_2) \in \mathcal{L}_1 + \mathcal{L}_2$, the isomorphism $\lambda : C(x + (v_1, v_2)) = (v_2, x + v_1) \rightarrow x + C(v_1, v_2) = (x + v_2, v_1)$ as $\lambda = (-x, \varphi_1, \varphi_2)$, where $\varphi_1 : v_2 \simeq (-x + x) + v_2 \simeq -x + (x + v_2)$ and $\varphi_2 : -x + (x + v_1) \simeq (-x + x) + v_1 \simeq v_1$.

In addition, by (2.3), there is an equality of 1-morphisms $C^2 = \text{id}$.

The commutativity constrains together with the associativity constrains satisfy the hexagon diagram, i.e. for any \mathcal{P} -torsors $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3$ the following diagram of

1-morphisms is commutative

$$(2.13) \quad \begin{array}{ccc} & (\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3 & \\ & \swarrow \quad \searrow & \\ (\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3 & & \mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}_3) \\ \downarrow & & \downarrow \\ \mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}_3) & & \mathcal{L}_1 + (\mathcal{L}_2 + \mathcal{L}_3) \\ & \swarrow \quad \searrow & \\ & (\mathcal{L}_1 + \mathcal{L}_2) + \mathcal{L}_3 & \end{array}$$

The following axiom is satisfied in the category $B\mathcal{P}$ and describes the functoriality of the commutativity constraints. Let $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}'_1$ be any \mathcal{P} -torsors, and $\mathcal{L}_1 \rightarrow \mathcal{L}'_1$ be any 1-morphism of \mathcal{P} -torsors, then the following diagram of 1-morphisms is commutative

$$(2.14) \quad \begin{array}{ccc} \mathcal{L}_1 + \mathcal{L}_2 & \longrightarrow & \mathcal{L}'_1 + \mathcal{L}_2 \\ \downarrow & & \downarrow \\ \mathcal{L}_2 + \mathcal{L}_1 & \longrightarrow & \mathcal{L}_2 + \mathcal{L}'_1, \end{array}$$

where the vertical arrows are the commutativity constraints.

By regarding \mathcal{P} as a \mathcal{P} -torsor, there is a canonical 1-isomorphism of \mathcal{P} -torsors $\mathcal{P} + \mathcal{L} \rightarrow \mathcal{L}$, $(x, v) \mapsto x + v$ satisfying the associativity and commutativity constraints. This means that \mathcal{P} is the unit in $B\mathcal{P}$. For each $\mathcal{L} \in B\mathcal{P}$, we have an object

$$-\mathcal{L} := \mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{P}),$$

together with a natural 1-isomorphism of \mathcal{P} -torsors $\varphi_{\mathcal{L}} : \mathcal{L} + (-\mathcal{L}) \simeq \mathcal{P}$. This object is called an inverse of \mathcal{L} .

For \mathcal{L} a \mathcal{P} -torsor, $\mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{L})$ is a natural monoidal groupoid (by composition). The natural homomorphism

$$(2.15) \quad \mathcal{Z} : \mathcal{P} \rightarrow \mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{L})$$

given by $\mathcal{Z}(z) = z + \mathrm{id}^5$ is a 1-isomorphism of monoidal groupoids. We will fix once and for all its inverse, i.e., we choose an 1-isomorphism of monoidal groupoids

$$(2.16) \quad \mathcal{Z}^{-1} : \mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{L}) \rightarrow \mathcal{P}$$

together with a 2-isomorphism $\mathcal{Z}^{-1} \circ \mathcal{Z} \simeq \mathrm{id}$.

Remark 2.2. We constructed some "semistrict" version of Picard 2-groupoid, because diagrams (2.9)-(2.14) are true in $B\mathcal{P}$ for 1-morphisms without consideration of additional 2-morphisms which involve higher coherence axioms for braided monoidal 2-categories as in [KV] and [BN]. Besides, from the equality $C^2 = \mathrm{id}$ we obtain at one stroke that our 2-category $B\mathcal{P}$ is strongly braided, i.e, the diagram (8.4.6) in [Br2] pp. 149 holds. Let us mention that in *loc. cit.*, the commutativity constraint C is denoted by R .

⁵Recall that we constructed the bifunctor $+: \mathcal{P} \times \mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{L}) \rightarrow \mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{L})$ in (2.8).

2.4. The case $H^1(BG, \mathcal{P})$. Let \mathcal{P} be a Picard groupoid, and G be a group. Then we define $H^1(BG, \mathcal{P})$ to be the Picard groupoid of homomorphisms from G to \mathcal{P} . That is, the objects are monoidal functors from G to \mathcal{P} , where G is regarded as a discrete monoidal category (i.e. the monoidal groupoid, where objects are elements of the group G , and morphisms in this groupoid are only the unit morphisms of objects), and morphisms between these monoidal functors are monoidal natural transformations. In concrete terms, $f \in H^1(BG, \mathcal{P})$ is a functor $f : G \rightarrow \mathcal{P}$, together with isomorphisms

$$f(gg') \simeq f(g) + f(g')$$

which are compatible with the associativity constraints. The monoidal structure on $H^1(BG, \mathcal{P})$ is given by $(f + f')(g) = f(g) + f'(g)$. The natural isomorphism $(f + f')(gg') \simeq (f + f')(g) + (f + f')(g')$ is the obvious one. The associativity constraints and the commutativity constraints on $H^1(BG, \mathcal{P})$ are clear. Let (e, φ) be a unit of \mathcal{P} , and e is regarded as a discrete Picard groupoid with one object. Then $f : G \rightarrow \mathcal{P}$ is called trivial if it is isomorphic to $G \rightarrow e \rightarrow \mathcal{P}$.

Example 2.3. If $\mathcal{P} = BA$, then $H^1(BG, BA)$ is equivalent to the category of central extensions of G by A as Picard groupoids.

Let $Z_2 \subset G \times G$ be the subset of commuting elements, so that if G itself is an abelian group, then $Z_2 = G \times G$. In general, fix $g \in G$, then $Z_2 \cap (G \times g) \simeq Z_2 \cap (g \times G) \simeq Z_G(g)$, the centralizer of g in G .

Lemma-Definition 2.4. *There is a well defined anti-symmetric bimultiplicative map $\text{Comm}(f) : Z_2 \rightarrow \pi_1(\mathcal{P}) = \text{End}_{\mathcal{P}}(e)$.*

Proof. The definition of $\text{Comm}(f)$ is as follows. For $g_1, g_2 \in Z_2$, we have

$$f(g_1g_2) \simeq f(g_1) + f(g_2) \simeq f(g_2) + f(g_1) \simeq f(g_2g_1) = f(g_1g_2),$$

where the first and the third isomorphisms come from the constraints for the homomorphism f , and the second isomorphism comes from the commutativity constraints of the Picard groupoid \mathcal{P} . We thus obtain an element

$$\text{Comm}(f)(g_1, g_2) \in \text{Aut}_{\mathcal{P}}(f(g_1g_2)) \simeq \pi_1(\mathcal{P}).$$

Since \mathcal{P} is Picard, i.e., the commutativity constraints satisfy $c_{f(g_1), f(g_2)} = c_{f(g_2), f(g_1)}^{-1}$, the map Comm is anti-symmetric. One can check directly by diagram that $\text{Comm}(f)$ is also bimultiplicative (see the analogous diagram (2.29) below).

Here we will give another proof of bimultiplicativity whose higher categorical analogue we will use in the proof of lemma-definition 2.7. We construct the following category H_f , where objects of H_f are all possible expressions

$$f(g_1) + \cdots + f(g_k) := (\cdots (f(g_1) + f(g_2)) + f(g_3)) + \cdots + f(g_k), \text{ where } g_i \in G,$$

and morphisms in H_f are defined as following:

$$\begin{aligned} & \text{Hom}_{H_f}(f(g_{i_1}) + \cdots + f(g_{i_k}), f(g_{j_1}) + \cdots + f(g_{j_l})) = \\ & \begin{cases} \emptyset & \text{if } g_{i_1} \cdots g_{i_k} \neq g_{j_1} \cdots g_{j_l}; \\ \text{Hom}_{\mathcal{P}}(f(g_{i_1}) + \cdots + f(g_{i_k}), f(g_{j_1}) + \cdots + f(g_{j_l})) & \text{if } g_{i_1} \cdots g_{i_k} = g_{j_1} \cdots g_{j_l}. \end{cases} \end{aligned}$$

The category H_f is a monoidal group-like groupoid (or *gr*-category), where the monoidal structure on H_f is given in an obvious way by using the associativity constraints in the category \mathcal{P} . We have $\pi_0(H_f) = G$, and H_f is equivalent to the

trivial gr -category. We consider $\pi_1(\mathcal{P})$ -torsor E over Z_2 which is the commutator of H_f (see [Br3, §3]). The fibre of E over $(g_1, g_2) \in Z_2$ is the set

$$E_{g_1, g_2} = \text{Hom}_{H_f}(f(g_1) + f(g_2), f(g_2) + f(g_1)).$$

The $\pi_1(\mathcal{P})$ -torsor E has a natural structure of a weak biextension of Z_2 by $\pi_1(\mathcal{P})$ (see [Br3, prop. 3.1]), i.e. there are partial composition laws on E which are compatible (see also (2.22)). Now the commutativity constraints $c_{f(g_1), f(g_2)}$ give a section of E over Z_2 which is compatible with partial composition laws on E , i.e. "bimultiplicative". (The compatibility of this section with the composition laws follows at once from the definition of the partial composition laws on E and the hexagon diagram (2.2).) The other section of E which is compatible with partial composition laws on E is obtained as the composition of following two morphisms from definition of f : $f(g_1) + f(g_2) \simeq f(g_1g_2) = f(g_2g_1) \simeq f(g_2) + f(g_1)$. (The compatibility of this section with composition laws follows from diagrams (3.10) and (1.4) of [Br3], because of the compatibility of our homomorphism f with the associativity constraints.) Now the difference between the first section and the second section coincides with $\text{Comm}(f)$, which is, thus, a bimultiplicative function, because both sections are "bimultiplicative". \square

Remark 2.3. In [Br3, §2] the notion of a weak biextension was introduced only for $Z_2 = B \times B$ where B is an abelian group. Here, we generalize this notion by allowing B to be non-commutative and by replacing $B \times B$ by Z_2 . But all the axioms for partial composition laws in *loc. cit.* are still applicable in this setting. The same remark applies when we talk about "(2, 2)-extension" in § 2.5.

Remark 2.4. It is clear that if $f \simeq f'$ in $H^1(BG, \mathcal{P})$, then $\text{Comm}(f) = \text{Comm}(f')$.

Remark 2.5. When $\mathcal{P} = BA$, this construction reduces to the usual construction of inverse to the commutator pairing maps for central extensions.

Corollary 2.5. *One has $\text{Comm}(f + f') = \text{Comm}(f) + \text{Comm}(f')$.*

Proof. It can be easily checked directly by diagrams. See, for example, analogous formulas and diagrams (2.30)-(2.32) below. \square

Corollary 2.6. *Assume that G is abelian so that $Z_2 = G \times G$. Then $\text{Comm}(f)$ is trivial if and only if the 1-homomorphism f is a 1-homomorphism of Picard groupoids. In particular, if the homomorphism f is trivial, then $\text{Comm}(f)$ is trivial.*

Proof. It follows from diagram (2.5). \square

The above two corollaries together can be rephrased as by saying that if G is abelian, then there is an exact sequence of Picard groupoids

$$1 \rightarrow \text{Hom}(G, \mathcal{P}) \rightarrow H^1(BG, \mathcal{P}) \rightarrow \text{Hom}(\wedge^2 G, \pi_1(\mathcal{P})).$$

2.5. The case $H^2(BG, \mathcal{P})$. If \mathcal{P}' is a Picard n -groupoid, and G is a group, one should be able to define $H^1(BG, \mathcal{P}')$ as the Picard n -groupoid of homomorphisms from G to \mathcal{P}' . When $n = 1$, this is what we discussed in the previous subsection. The next step for consideration is $n = 2$. Again, instead of discussing general Picard 2-groupoids, we will focus on the case when $\mathcal{P}' = B\mathcal{P}$, where \mathcal{P} is a Picard groupoid. Then one can interpret $H^1(BG, B\mathcal{P})$ as the Picard groupoid⁶ of central extensions of

⁶As we just mentioned, it is in fact a Picard 2-groupoid.

the group G by the Picard groupoid \mathcal{P} . For this reason, we also denote $H^1(BG, B\mathcal{P})$ by $H^2(BG, \mathcal{P})$.

In concrete terms, an object \mathcal{L} in $H^2(BG, \mathcal{P})$ is a rule to assign to every $g \in G$ a \mathcal{P} -torsor \mathcal{L}_g , and to every g, g' an equivalence $\mathcal{L}_{gg'} \simeq \mathcal{L}_g + \mathcal{L}_{g'}$ of \mathcal{P} -torsors, and to every g, g', g'' an isomorphism between two equivalences

$$(2.17) \quad \begin{array}{ccc} & \mathcal{L}_{gg'g''} & \\ & \swarrow \quad \searrow & \\ \mathcal{L}_{gg'} + \mathcal{L}_{g''} & \Downarrow & \mathcal{L}_g + \mathcal{L}_{g'g''} \\ \downarrow & & \downarrow \\ (\mathcal{L}_g + \mathcal{L}_{g'}) + \mathcal{L}_{g''} & \longrightarrow & \mathcal{L}_g + (\mathcal{L}_{g'} + \mathcal{L}_{g''}) \end{array}$$

such that for every g, g', g'', g''' , the natural compatibility condition holds, which we describe below.

Remark 2.6. Our notation for the 2-arrow in diagram (2.17) is symbolic, and is distinct from the traditional notation of 2-arrows in a 2-category, because this 2-arrow is between a pair of 1-arrows from $\mathcal{L}_{gg'g''}$ to $\mathcal{L}_g + (\mathcal{L}'_{g'} + \mathcal{L}''_{g'})$ and should be written horizontally from left to right rather than vertically. This notation for the 2-arrow will be important for us in diagram (2.28).

We define an isomorphism between two central extensions of G by \mathcal{P} . An isomorphism between two central extensions $\mathcal{L}, \mathcal{L}'$ is a rule which assigns to any g a \mathcal{P} -torsor 1-isomorphism $\mathcal{L}_g \simeq \mathcal{L}'_g$, and to any g, g' the following 2-isomorphism

$$\begin{array}{ccc} \mathcal{L}_{gg'} & \longrightarrow & \mathcal{L}_g + \mathcal{L}_{g'} \\ \downarrow & \nearrow & \downarrow \\ \mathcal{L}'_{gg'} & \longrightarrow & \mathcal{L}'_g + \mathcal{L}'_{g'} \end{array}$$

In addition, these assignments have to be compatible with diagram (2.17) in an obvious way.

Now we describe the compatibility condition which we need after diagram (2.17). If we don't consider the associativity constraints in category $B\mathcal{P}$, then the 2-arrows induced by the one in (2.17) should satisfy the compatibility condition described by

the following cube:

(2.18)

$$\begin{array}{ccccc}
 & & \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''g'''} & \longrightarrow & \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''} + \mathcal{L}_{g'''} \\
 & \nearrow & \uparrow \text{---} & & \nearrow \\
 \mathcal{L}_g + \mathcal{L}_{g'g''g'''} & \longrightarrow & \mathcal{L}_g + \mathcal{L}_{g'g''} + \mathcal{L}_{g'''} & & \\
 \uparrow & & \uparrow & & \uparrow \\
 & & \mathcal{L}_{gg'} + \mathcal{L}_{g''g'''} & \dashrightarrow & \mathcal{L}_{gg'} + \mathcal{L}_{g''} + \mathcal{L}_{g'''} \\
 & \nearrow & & & \nearrow \\
 \mathcal{L}_{gg'g''g'''} & \longrightarrow & \mathcal{L}_{gg'g''} + \mathcal{L}_{g'''} & &
 \end{array}$$

To obtain the correct compatibility diagram for 2-morphisms, we have to replace in diagram (2.18) the arrow (an edge of cube)

$$\mathcal{L}_{gg'} + \mathcal{L}_{g''} + \mathcal{L}_{g'''} \longrightarrow \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''} + \mathcal{L}_{g'''}$$

by the following commutative diagram of 1-morphisms in the category $B\mathcal{P}$

$$\begin{array}{ccc}
 (2.19) & (\mathcal{L}_{gg'} + \mathcal{L}_{g''}) + \mathcal{L}_{g'''} & \longrightarrow & ((\mathcal{L}_g + \mathcal{L}_{g'}) + \mathcal{L}_{g''}) + \mathcal{L}_{g'''} \\
 & \downarrow & & \downarrow \\
 & \mathcal{L}_{gg'} + (\mathcal{L}_{g''} + \mathcal{L}_{g'''}) & \longrightarrow & (\mathcal{L}_g + \mathcal{L}_{g'}) + (\mathcal{L}_{g''} + \mathcal{L}_{g'''})
 \end{array}$$

(where the vertical arrows are associativity constraints); we have to replace in diagram (2.18) the arrow (an edge of the cube)

$$\mathcal{L}_g + \mathcal{L}_{g'g''} + \mathcal{L}_{g'''} \longrightarrow \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''} + \mathcal{L}_{g'''}$$

by the following commutative diagram of 1-morphisms in the category $B\mathcal{P}$

$$\begin{array}{ccc}
 (2.20) & (\mathcal{L}_g + \mathcal{L}_{g'g''}) + \mathcal{L}_{g'''} & \longrightarrow & (\mathcal{L}_g + (\mathcal{L}_{g'} + \mathcal{L}_{g''})) + \mathcal{L}_{g'''} \\
 & \downarrow & & \downarrow \\
 & \mathcal{L}_g + (\mathcal{L}_{g'g''} + \mathcal{L}_{g'''}) & \longrightarrow & \mathcal{L}_g + ((\mathcal{L}_{g'} + \mathcal{L}_{g''}) + \mathcal{L}_{g'''})
 \end{array}$$

(where vertical arrows are associativity constraints); we have to replace in diagram (2.18) the arrow (an edge of the cube)

$$\mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''g'''} \longrightarrow \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''} + \mathcal{L}_{g'''}$$

by the following commutative diagram of 1-morphisms in the category $B\mathcal{P}$

$$\begin{array}{ccc}
 (2.21) & (\mathcal{L}_g + \mathcal{L}_{g'}) + \mathcal{L}_{g''g'''} & \longrightarrow & (\mathcal{L}_g + \mathcal{L}_{g'}) + (\mathcal{L}_{g''} + \mathcal{L}_{g'''}) \\
 & \downarrow & & \downarrow \\
 & \mathcal{L}_g + (\mathcal{L}_{g'} + \mathcal{L}_{g''g'''}) & \longrightarrow & \mathcal{L}_g + (\mathcal{L}_{g'} + (\mathcal{L}_{g''} + \mathcal{L}_{g'''}))
 \end{array}$$

(where vertical arrows are associativity constraints). Besides, instead of the vertex $\mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''} + \mathcal{L}_{g'''}$ in diagram (2.18) we insert the commutative diagram which is the modification of pentagon diagram (2.9) for $\mathcal{L}_g, \mathcal{L}_{g'}, \mathcal{L}_{g''}, \mathcal{L}_{g'''}$, and this diagram is always true in category $B\mathcal{P}$. The correct compatibility diagram for 2-morphisms from diagrams (2.17) has 15 vertices.

We note that diagrams (2.19)-(2.21) are commutative for 1-morphisms, i.e. the corresponding 2-isomorphisms equal identity morphisms. These diagrams express the "functoriality" of associativity constraints in $B\mathcal{P}$ and follow from axioms-diagrams (2.10)-(2.12) in category $B\mathcal{P}$.

The trivial central extension of G by \mathcal{P} is the rule which assign to every $g \in G$ the trivial \mathcal{P} -torsor \mathcal{P} , to every g, g' the natural 1-isomorphism $\mathcal{P} \simeq \mathcal{P} + \mathcal{P}^{\tilde{7}}$, and to every g, g', g'' the corresponding natural 2-isomorphism.

Remark 2.7. A central extension \mathcal{L} of G by \mathcal{P} gives rise to a gr-category, $\tilde{\mathcal{L}}$, together with a short exact sequence of gr-categories in the sense of [Br1, definition 2.1.2]

$$1 \rightarrow \mathcal{P} \xrightarrow{i} \tilde{\mathcal{L}} \xrightarrow{\pi} G \rightarrow 1.$$

Namely, as a category, $\tilde{\mathcal{L}} = \bigcup_{g \in G} \mathcal{L}_g$. Then the natural equivalence $\mathcal{L}_{gg'} \simeq \mathcal{L}_g + \mathcal{L}_{g'}$ together with the compatibility conditions endows $\tilde{\mathcal{L}}$ with a gr-category structure. The natural morphism $\pi : \tilde{\mathcal{L}} \rightarrow G$ is clearly monoidal, and one can show that $\ker \pi = \mathcal{L}_e$ is 1-isomorphic to \mathcal{P} .

As is shown in *loc. cit.*, such a short exact sequence endows every $\tilde{\mathcal{L}}_g := \pi^{-1}(g) = \mathcal{L}_g$ with a \mathcal{P} -bitorsor structure. This \mathcal{P} -bitorsor structure is nothing but the canonical \mathcal{P} -bitorsor structure on \mathcal{L}_g (observe that the morphism $\mathcal{Z} : \mathcal{P} \rightarrow \text{Hom}_{\mathcal{P}}(\mathcal{L}_g, \mathcal{L}_g)$ as in (2.15) induces a canonical \mathcal{P} -bitorsor structure on \mathcal{L}_g).

The upshot is that an object \mathcal{L} in $H^2(BG, \mathcal{P})$ gives rise to a categorical generalization of a central extension of a group by an abelian group. This justifies our terminology. Indeed, one can define a central extension of G by \mathcal{P} as a short exact sequence as above such that the induced \mathcal{P} -bitorsor structure on each $\tilde{\mathcal{L}}_g$ is the canonical one induced from its left \mathcal{P} -torsor structure. Since we do not use this second definition, we will not make it precise.

Finally, let us define the Picard structure on $H^2(BG, \mathcal{P})$. Let \mathcal{L} and \mathcal{L}' be two central extensions of G by \mathcal{P} . Then we define the central extension $\mathcal{L} + \mathcal{L}'$ by the following way:

$$(\mathcal{L} + \mathcal{L}')_g := \mathcal{L}_g + \mathcal{L}'_g,$$

and the equivalence $(\mathcal{L} + \mathcal{L}')_{gg'} \simeq (\mathcal{L} + \mathcal{L}')_g + (\mathcal{L} + \mathcal{L}')_{g'}$ as the composition of the following equivalences

$$\begin{aligned} (\mathcal{L} + \mathcal{L}')_{gg'} &= \mathcal{L}_{gg'} + \mathcal{L}'_{gg'} \simeq (\mathcal{L}_g + \mathcal{L}_{g'}) + (\mathcal{L}'_g + \mathcal{L}'_{g'}) \\ &\simeq (\mathcal{L}_g + \mathcal{L}'_g) + (\mathcal{L}_{g'} + \mathcal{L}'_{g'}) = (\mathcal{L} + \mathcal{L}')_g + (\mathcal{L} + \mathcal{L}')_{g'}. \end{aligned}$$

The corresponding 2-isomorphism for central extension $\mathcal{L} + \mathcal{L}'$ and any elements g, g', g'' of G follows from diagrams (2.17) for central extensions \mathcal{L} and \mathcal{L}' . The further compatibility conditions for these 2-isomorphisms hold as in diagrams (2.18)-(2.21), since they follow at once from the corresponding diagrams for central extensions \mathcal{L} and \mathcal{L}' .

⁷The naturality means that this 1-isomorphism is the chosen quasi-inverse of the natural 1-isomorphism $\mathcal{P} + \mathcal{P} \rightarrow \mathcal{P}$.

Again, let Z_2 denote the subset of $G \times G$ consisting of commuting elements. We will give a categorical analogue of lemma-definition 2.4. For this purpose, let us first explain some terminology. A 1-morphism $f : Z_2 \rightarrow \mathcal{P}$ is called bimultiplicative if for fixed $g \in G$, $(Z_G(g), g) \subset Z_2 \rightarrow \mathcal{P}$ and $(g, Z_G(g)) \subset Z_2 \rightarrow \mathcal{P}$ are homomorphisms, i.e. monoidal functors from discrete monoidal categories $(Z_G(g), g)$ and $(g, Z_G(g))$ to \mathcal{P} . In addition, the following diagram must be commutative (which is the compatibility condition between these two homomorphisms)

$$(2.22) \quad \begin{array}{ccc} f(g_1g_2, g_3) + f(g_1g_2, g_4) & \longrightarrow & (f(g_1, g_3) + f(g_2, g_3)) + (f(g_1, g_4) + f(g_2, g_4)) \\ \uparrow \simeq & & \downarrow \simeq \\ f(g_1g_2, g_3g_4) & & \\ \downarrow \simeq & & \\ f(g_1, g_3g_4) + f(g_2, g_3g_4) & \longrightarrow & (f(g_1, g_3) + f(g_1, g_4)) + (f(g_2, g_3) + f(g_2, g_4)) \end{array}$$

When $\mathcal{P} = BA$, a bimultiplicative 1-morphism from $Z_2 \rightarrow BA$ is the same as a weak biextension of Z_2 by A as defined in [Br3, §2] (see also remark 2.3).

A 1-morphism $f : Z_2 \rightarrow \mathcal{P}$ is called anti-symmetric if there is a 2-isomorphism $\theta : f \simeq -f \circ \sigma$, where σ is the natural flip on Z_2 , such that for any $(g_1, g_2) \in Z_2$, the following diagram is commutative

$$\begin{array}{ccc} f(g_1, g_2) & \xrightarrow{\simeq} & -f(g_2, g_1) \\ \parallel & & \downarrow \simeq \\ f(g_1, g_2) & \xleftarrow{\simeq} & -(-f(g_1, g_2)) \end{array}$$

We need some more terminology. Following [Br3, §7], we *define* a weak $(2, 2)$ -extension of Z_2 by \mathcal{P} as a rule which assigns to every $(g, g') \in Z_2$ a \mathcal{P} -torsor $\mathcal{E}_{(g, g')}$ such that its restrictions to $(g, Z_G(g))$ and $Z_G(g), g)$ are central extensions of $Z_G(g)$ by \mathcal{P} , and that the corresponding diagram (2.22) is 2-commutative (i.e. commutative modulo some 2-isomorphism), and these 2-isomorphisms satisfy further compatibility conditions (see (7.1), (7.3) in *loc. cit.* where these compatibility conditions are carefully spelt out).

Lemma-Definition 2.7. *There is an anti-symmetric bimultiplicative homomorphism $C_2^{\mathcal{L}} : Z_2 \rightarrow \mathcal{P}$.*

Proof. As in the proof of lemma-definition 2.4, using the commutativity constraints $C : \mathcal{L}_g + \mathcal{L}_{g'} \simeq \mathcal{L}_{g'} + \mathcal{L}_g$ in the category $B\mathcal{P}$, one constructs the following composition of 1-isomorphisms:

$$\mathcal{L}_{gg'} \simeq \mathcal{L}_g + \mathcal{L}_{g'} \simeq \mathcal{L}_{g'} + \mathcal{L}_g \simeq \mathcal{L}_{g'g} = \mathcal{L}_{gg'},$$

for $(g, g') \in Z_2$. In this way, we obtain a functor $Z_2 \rightarrow \text{Hom}_{\mathcal{P}}(\mathcal{L}_{gg'}, \mathcal{L}_{gg'})$. Using $\mathcal{Z}^{-1} : \text{Hom}_{\mathcal{P}}(\mathcal{L}_{gg'}, \mathcal{L}_{gg'}) \rightarrow \mathcal{P}$ (see (2.16)), we get a morphism $C_2^{\mathcal{L}} : Z_2 \rightarrow \mathcal{P}$.

We need to construct the following canonical isomorphisms

$$C_2^{\mathcal{L}}(gg', g'') \simeq C_2^{\mathcal{L}}(g, g'') + C_2^{\mathcal{L}}(g', g''), \quad C_2^{\mathcal{L}}(g, g'g'') \simeq C_2^{\mathcal{L}}(g, g') + C_2^{\mathcal{L}}(g, g''),$$

satisfying the natural compatibility conditions. We now construct the first isomorphism. The second is similar. Let $\mathcal{Z} : \mathcal{P} \rightarrow \text{Hom}_{\mathcal{P}}(\mathcal{L}_{gg'g''}, \mathcal{L}_{gg'g''})$ be the canonical

equivalence as monoidal groupoids as in (2.15). It is enough to construct a canonical 2-isomorphism $\mathcal{Z}(C_2^{\mathcal{L}}(gg', g'')) \simeq \mathcal{Z}(C_2^{\mathcal{L}}(g, g'') + C_2^{\mathcal{L}}(g', g''))$.

By the definition of the morphism $C_2^{\mathcal{L}}$, there is a canonical 2-isomorphism from 1-isomorphism $\mathcal{Z}(C_2^{\mathcal{L}}(g, g'') + C_2^{\mathcal{L}}(g', g''))$ to the following composition of 1-isomorphisms:

$$(2.23) \quad \begin{aligned} \mathcal{L}_{gg'g''} &\simeq \mathcal{L}_g + \mathcal{L}_{g'g''} \simeq \mathcal{L}_g + (\mathcal{L}_{g'} + \mathcal{L}_{g''}) \simeq \mathcal{L}_g + (\mathcal{L}_{g''} + \mathcal{L}_{g'}) \simeq \mathcal{L}_g + \mathcal{L}_{g''g'} \simeq \mathcal{L}_{gg''g'} \\ &\simeq \mathcal{L}_{gg''} + \mathcal{L}_{g'} \simeq (\mathcal{L}_g + \mathcal{L}_{g''}) + \mathcal{L}_{g'} \simeq (\mathcal{L}_{g''} + \mathcal{L}_g) + \mathcal{L}_{g'} \simeq \mathcal{L}_{gg''} + \mathcal{L}_{g'} \simeq \mathcal{L}_{gg'g''} \end{aligned}$$

By the definition of the central extension of G by \mathcal{P} (see diagram (2.17)), there is a canonical 2-isomorphism from the above composition of 1-isomorphisms to the following composition of 1-isomorphisms

$$(2.24) \quad \begin{aligned} \mathcal{L}_{gg'g''} &\simeq \mathcal{L}_g + \mathcal{L}_{g'g''} \simeq \mathcal{L}_g + (\mathcal{L}_{g'} + \mathcal{L}_{g''}) \simeq \mathcal{L}_g + (\mathcal{L}_{g''} + \mathcal{L}_{g'}) \\ &\simeq (\mathcal{L}_g + \mathcal{L}_{g''}) + \mathcal{L}_{g'} \simeq (\mathcal{L}_{g''} + \mathcal{L}_g) + \mathcal{L}_{g'} \simeq \mathcal{L}_{gg''} + \mathcal{L}_{g'} \simeq \mathcal{L}_{gg'g''} \end{aligned}$$

From the hexagon axiom for 1-morphisms in the category $B\mathcal{P}$ (see diagram 2.13) we have that the 1-isomorphism which is the composition of the above 1-isomorphisms is equal to the 1-isomorphism which is the composition of the following 1-isomorphisms

$$(2.25) \quad \begin{aligned} \mathcal{L}_{gg'g''} &\simeq \mathcal{L}_g + \mathcal{L}_{g'g''} \simeq \mathcal{L}_g + (\mathcal{L}_{g'} + \mathcal{L}_{g''}) \simeq (\mathcal{L}_g + \mathcal{L}_{g'}) + \mathcal{L}_{g''} \\ &\simeq \mathcal{L}_{g''} + (\mathcal{L}_g + \mathcal{L}_{g'}) \simeq (\mathcal{L}_{g''} + \mathcal{L}_g) + \mathcal{L}_{g'} \simeq \mathcal{L}_{gg''} + \mathcal{L}_{g'} \simeq \mathcal{L}_{gg'g''} \end{aligned}$$

By the "functoriality" of the commutativity constraints in the category $B\mathcal{P}$ (see axiom-diagram (2.14)) we have that the 1-isomorphism which is the composition of the above 1-isomorphisms is equal to the 1-isomorphism which is the composition of the following 1-isomorphisms

$$(2.26) \quad \begin{aligned} \mathcal{L}_{gg'g''} &\simeq \mathcal{L}_g + \mathcal{L}_{g'g''} \simeq \mathcal{L}_g + (\mathcal{L}_{g'} + \mathcal{L}_{g''}) \simeq (\mathcal{L}_g + \mathcal{L}_{g'}) + \mathcal{L}_{g''} \simeq \mathcal{L}_{gg'} + \mathcal{L}_{g''} \\ &\simeq \mathcal{L}_{g''} + \mathcal{L}_{gg'} \simeq \mathcal{L}_{g''} + (\mathcal{L}_g + \mathcal{L}_{g'}) \simeq (\mathcal{L}_{g''} + \mathcal{L}_g) + \mathcal{L}_{g'} \simeq \mathcal{L}_{gg''} + \mathcal{L}_{g'} \simeq \mathcal{L}_{gg'g''} \end{aligned}$$

Again, by the definition of the central extension of G by \mathcal{P} (see diagram (2.17), which we apply twice now), there is a canonical 2-isomorphism from the above composition of 1-isomorphisms to the following composition of 1-isomorphisms

$$(2.27) \quad \mathcal{L}_{gg'g''} \simeq \mathcal{L}_{gg'} + \mathcal{L}_{g''} \simeq \mathcal{L}_{g''} + \mathcal{L}_{gg'} \simeq \mathcal{L}_{gg'g''},$$

which is canonically isomorphic to $\mathcal{Z}(C_2^{\mathcal{L}}(gg', g''))$.

Let us write down a diagram which will represent the above 2-isomorphisms. To simplify the notation, we will denote the 2-commutative diagram (2.17) as

$$(2.28) \quad \begin{array}{c} \mathcal{L}_{gg'g''} \\ \Downarrow \\ \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''} \end{array}$$

Then, the 2-isomorphism $\mathcal{Z}(C_2^{\mathcal{L}}(gg', g'')) \simeq \mathcal{Z}(C_2^{\mathcal{L}}(g, g'')) + \mathcal{Z}(C_2^{\mathcal{L}}(g', g''))$ is represented by the following diagram (2.29)

$$\begin{array}{ccccc}
& & \mathcal{L}_{gg''g'} & & \\
& \nearrow^{\mathcal{Z}(C_2^{\mathcal{L}}(g', g''))} & \Downarrow & \searrow_{\mathcal{Z}(C_2^{\mathcal{L}}(g, g''))} & \\
& & \mathcal{L}_g + \mathcal{L}_{g''} + \mathcal{L}_{g'} & & \\
\mathcal{L}_{gg'g''} & \rightleftarrows & \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''} & \longrightarrow & \mathcal{L}_{g''} + \mathcal{L}_g + \mathcal{L}_{g'} & \rightleftarrows & \mathcal{L}_{g''g'g'} \\
& & \mathcal{Z}(C_2^{\mathcal{L}}(gg', g'')) & & & &
\end{array}$$

To check all the compatibility conditions between these canonical isomorphisms we generalize the proof of lemma-definition 2.4. We construct the following 2-category $H_{\mathcal{L}}$, where objects of $H_{\mathcal{L}}$ are objects from categories given by all expressions

$$\mathcal{L}_{g_1} + \cdots + \mathcal{L}_{g_k} := (\cdots (\mathcal{L}_{g_1} + \mathcal{L}_{g_2}) + \mathcal{L}_{g_3}) + \cdots + \mathcal{L}_{g_k}, \text{ where } g_i \in G,$$

1-morphisms in $H_{\mathcal{L}}$ are defined as following:

$$\text{Hom}_{H_{\mathcal{L}}}(\mathcal{L}_{g_{i_1}} + \cdots + \mathcal{L}_{g_{i_k}}, \mathcal{L}_{g_{j_1}} + \cdots + \mathcal{L}_{g_{j_l}}) = \begin{cases} \emptyset & \text{if } g_{i_1} \cdots g_{i_k} \neq g_{j_1} \cdots g_{j_l}; \\ \text{Hom}_{B\mathcal{P}}(\mathcal{L}_{g_{i_1}} + \cdots + \mathcal{L}_{g_{i_k}}, \mathcal{L}_{g_{j_1}} + \cdots + \mathcal{L}_{g_{j_l}}) & \text{if } g_{i_1} \cdots g_{i_k} = g_{j_1} \cdots g_{j_l}, \end{cases}$$

and 2-morphisms in 2-category $H_{\mathcal{L}}$ come from 2-morphisms of category $B\mathcal{P}$. The category $H_{\mathcal{L}}$ is a monoidal group-like 2-groupoid (or a 2-gr-category), see [Br2, §8], where monoidal structure on $H_{\mathcal{L}}$ is given in an obvious way by using the associativity constraints in the category $B\mathcal{P}$ and pentagon diagram (2.9). We have $\pi_0(H_{\mathcal{L}}) = G$. We consider the \mathcal{P} -torsor $\mathcal{E}^{\mathcal{L}}$ on Z_2 which is the commutator of $H_{\mathcal{L}}$ (see [Br3, §8]⁸). The fibre of $\mathcal{E}^{\mathcal{L}}$ over $(g_1, g_2) \in Z_2$ is the \mathcal{P} -torsor

$$\mathcal{E}_{g_1, g_2}^{\mathcal{L}} = \text{Hom}_{H_{\mathcal{L}}}(\mathcal{L}_{g_1} + \mathcal{L}_{g_2}, \mathcal{L}_{g_2} + \mathcal{L}_{g_1}).$$

The \mathcal{P} -torsor $\mathcal{E}^{\mathcal{L}}$ on Z_2 has a natural structure of a weak (2, 2)-extension (see [Br3, prop. 8.1]), i.e. there are partial composition (group) laws on $\mathcal{E}^{\mathcal{L}}$ which are compatible (see diagrams (7.1), (7.3) in *loc.cit.*). Now the commutativity constraints C from $B\mathcal{P}$ give a trivialization of \mathcal{P} -torsor $\mathcal{E}^{\mathcal{L}}$ on Z_2 which is compatible with partial composition laws on $\mathcal{E}^{\mathcal{L}}$, i.e. "bimultiplicative". (The compatibility of this trivialization with composition laws follows at once from definition of partial composition laws on $\mathcal{E}^{\mathcal{L}}$ and hexagon diagram (2.13). See also the discussion in the end of [Br3, §8] regarding the braiding structure in $H_{\mathcal{L}}$, which gives the "bimultiplicative" trivialization of the \mathcal{P} -torsor $\mathcal{E}^{\mathcal{L}}$ on Z_2 .) The other trivialization of the \mathcal{P} -torsor $\mathcal{E}^{\mathcal{L}}$ on Z_2 which is compatible with partial composition laws on $\mathcal{E}^{\mathcal{L}}$ is obtained as the composition of the following two equivalences from definition of \mathcal{L} :

$$S_{\mathcal{L}_{g_1}, \mathcal{L}_{g_2}} : \mathcal{L}_{g_1} + \mathcal{L}_{g_2} \simeq \mathcal{L}_{g_1g_2} = \mathcal{L}_{g_2g_1} \simeq \mathcal{L}_{g_2} + \mathcal{L}_{g_1}.$$

⁸L. Breen assumed for simplicity in *loc. cit.* that the group π_1 of a 2-gr-category is equal to 0. We have $\pi_1(H_{\mathcal{L}}) \neq 0$, but the constructions and its properties which we need remain true in our situation.

Now the difference between the first trivialization and the second trivialization of the \mathcal{P} -torsor $\mathcal{E}^{\mathcal{L}}$ on Z_2 coincides with $C_2^{\mathcal{L}}$, which is, thus, a bimultiplicative homomorphism, because both trivializations are "bimultiplicative".

We have shown that $C_2^{\mathcal{L}} : Z_2 \rightarrow \mathcal{P}$ is a bimultiplicative 1-morphism. One readily checks from the above constructions that this is anti-symmetric from Z_2 to \mathcal{P} , since $C^2 = \text{id}$. \square

Remark 2.8. If $\mathcal{P} = BA$, then the construction of $C_2^{\mathcal{L}}$ given above is equivalent to the construction of the commutator category of the central extension $-\mathcal{L}$ introduced by Deligne in [Del2].

We also have the following categorical analogue of corollary 2.5. First, let us remark that if $f_1, f_2 : Z_2 \rightarrow \mathcal{P}$ are two bimultiplicative homomorphisms, one can define $f_1 + f_2$, which is again a bimultiplicative homomorphism, in the same way as defining the Picard structure on $H^1(BG, \mathcal{P})$.

Lemma 2.8. *For any two central extensions \mathcal{L} and \mathcal{L}' of G by \mathcal{P} there is a natural bimultiplicative 2-isomorphism (i.e., it respects the bimultiplicative structure) between bimultiplicative 1-morphisms $C_2^{\mathcal{L}+\mathcal{L}'}$ and $C_2^{\mathcal{L}} + C_2^{\mathcal{L}'}$.*

Proof. Recall that we have the following canonical 1-isomorphism

$$\mathcal{Z} : \mathcal{P} \rightarrow \text{Hom}(\mathcal{L}_{gg'} + \mathcal{L}'_{gg'}, \mathcal{L}_{gg'} + \mathcal{L}'_{gg'}).$$

We construct a canonical isomorphism

$$\mathcal{Z}(C_2^{\mathcal{L}+\mathcal{L}'}(g, g')) \simeq \mathcal{Z}(C_2^{\mathcal{L}}(g, g') + C_2^{\mathcal{L}'}(g, g'))$$

for any $(g, g') \in Z_2$ as following. By definition, $\mathcal{Z}(C_2^{\mathcal{L}+\mathcal{L}'}(g, g'))$ is canonically 2-isomorphic to the following composition of 1-morphisms

$$(2.30) \quad \begin{aligned} (\mathcal{L} + \mathcal{L}')_{gg'} &= \mathcal{L}_{gg'} + \mathcal{L}'_{gg'} \simeq (\mathcal{L}_g + \mathcal{L}_{g'}) + (\mathcal{L}'_g + \mathcal{L}'_{g'}) \simeq (\mathcal{L}_g + \mathcal{L}'_g) + (\mathcal{L}_{g'} + \mathcal{L}'_{g'}) \\ &= (\mathcal{L} + \mathcal{L}')_g + (\mathcal{L} + \mathcal{L}')_{g'} \simeq (\mathcal{L} + \mathcal{L}')_{g'} + (\mathcal{L} + \mathcal{L}')_g \simeq (\mathcal{L}_{g'} + \mathcal{L}'_{g'}) + (\mathcal{L}_g + \mathcal{L}'_g) \\ &\simeq (\mathcal{L}_{g'} + \mathcal{L}_g) + (\mathcal{L}'_{g'} + \mathcal{L}'_g) \simeq \mathcal{L}_{g'g} + \mathcal{L}'_{g'g} = (\mathcal{L} + \mathcal{L}')_{g'g} = (\mathcal{L} + \mathcal{L}')_{gg'} \end{aligned}$$

Using the "functoriality" of commutativity constraints, i.e. applying twice diagram (2.14), and using the following commutative diagram (which is written without associativity constraints)

$$(2.31) \quad \begin{array}{ccc} & \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}'_g + \mathcal{L}'_{g'} & \\ & \swarrow & \searrow \\ \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}'_{g'} + \mathcal{L}'_g & \longleftarrow & \mathcal{L}_g + \mathcal{L}'_g + \mathcal{L}_{g'} + \mathcal{L}'_{g'} \\ \downarrow & \searrow & \downarrow \\ \mathcal{L}_{g'} + \mathcal{L}_g + \mathcal{L}'_{g'} + \mathcal{L}'_g & \longleftarrow & \mathcal{L}_{g'} + \mathcal{L}'_{g'} + \mathcal{L}_g + \mathcal{L}'_g \end{array}$$

(to obtain the correct diagram we have to replace every triangle in this diagram by a hexagon coming from (2.13)), we obtain that the composition (2.30) of 1-morphisms

is equal to the following composition of 1-morphisms

$$(2.32) \quad \begin{aligned} \mathcal{L}_{gg'} + \mathcal{L}'_{gg'} &\simeq (\mathcal{L}_g + \mathcal{L}_{g'}) + (\mathcal{L}'_g + \mathcal{L}'_{g'}) \\ &\simeq (\mathcal{L}_{g'} + \mathcal{L}_g) + (\mathcal{L}'_{g'} + \mathcal{L}'_g) \simeq \mathcal{L}_{g'g} + \mathcal{L}'_{g'g} = \mathcal{L}_{gg'} + \mathcal{L}'_{gg'}, \end{aligned}$$

which is, by definition, 2-isomorphic to $\mathcal{Z}(C_2^{\mathcal{L}}(g, g') + C_2^{\mathcal{L}'}(g, g'))$.

To complete the proof, we need to show that the following diagram

$$(2.33) \quad \begin{array}{ccc} C_2^{\mathcal{L}+\mathcal{L}'}(g, g'') + C_2^{\mathcal{L}+\mathcal{L}'}(g', g'') & \longrightarrow & (C_2^{\mathcal{L}}(g, g'') + C_2^{\mathcal{L}'}(g, g'')) + (C_2^{\mathcal{L}}(g', g'') + C_2^{\mathcal{L}'}(g', g'')) \\ \uparrow & & \downarrow \\ C_2^{\mathcal{L}+\mathcal{L}'}(gg', g'') & & \\ \downarrow & & \\ C_2^{\mathcal{L}}(gg', g'') + C_2^{\mathcal{L}'}(gg', g'') & \longrightarrow & (C_2^{\mathcal{L}}(g, g'') + C_2^{\mathcal{L}}(g', g'')) + (C_2^{\mathcal{L}'}(g, g'') + C_2^{\mathcal{L}'}(g', g'')) \end{array}$$

and a similar diagram involving $C_2^{\mathcal{L}+\mathcal{L}'}(g, g'g'')$ are commutative. To prove this, let us recall that the 2-isomorphism $C_2^{\mathcal{L}}(gg', g'') \simeq C_2^{\mathcal{L}}(g, g'') + C_2^{\mathcal{L}}(g', g'')$ is the composition of the following 2-isomorphisms

$$\mathcal{Z}(C_2^{\mathcal{L}}(g, g'') + C_2^{\mathcal{L}}(g', g'')) \rightarrow (2.23) \rightarrow (2.24) \rightarrow \cdots \rightarrow (2.27) \rightarrow \mathcal{Z}(C_2^{\mathcal{L}}(gg', g'')).$$

Let us denote the 1-isomorphism (2.23) for \mathcal{L} (resp. \mathcal{L}' , resp. $\mathcal{L} + \mathcal{L}'$) as $(2.23)_{\mathcal{L}}$ (resp. $(2.23)_{\mathcal{L}'}$, resp. $(2.23)_{\mathcal{L}+\mathcal{L}'}$) and etc. Then it is readily checked that there exists a canonical 2-isomorphism

$$(2.23)_{\mathcal{L}} + (2.23)_{\mathcal{L}'} \simeq (2.23)_{\mathcal{L}+\mathcal{L}'}$$

between corresponding 1-isomorphisms $\mathcal{L}_{gg'g''} + \mathcal{L}'_{gg'g''} \rightarrow \mathcal{L}_{gg'g''} + \mathcal{L}'_{gg'g''}$, and canonical 2-isomorphisms for (2.24)-(2.27) such that the following diagram

$$\begin{array}{ccc} (2.23)_{\mathcal{L}} + (2.23)_{\mathcal{L}'} & \longrightarrow & (2.23)_{\mathcal{L}+\mathcal{L}'} \\ \downarrow & & \downarrow \\ (2.24)_{\mathcal{L}} + (2.24)_{\mathcal{L}'} & \longrightarrow & (2.24)_{\mathcal{L}+\mathcal{L}'} \end{array}$$

and similar diagrams for (2.24)-(2.27) commute. In addition, the following diagrams commute.

$$\begin{array}{ccc} \mathcal{Z}(C_2^{\mathcal{L}+\mathcal{L}'}(g, g'') + C_2^{\mathcal{L}+\mathcal{L}'}(g', g'')) & \longrightarrow & \mathcal{Z}((C_2^{\mathcal{L}}(g, g'') + C_2^{\mathcal{L}}(g', g'')) + (C_2^{\mathcal{L}'}(g, g'') + C_2^{\mathcal{L}'}(g', g''))) \\ \downarrow & & \downarrow \\ (2.23)_{\mathcal{L}+\mathcal{L}'} & \longrightarrow & (2.23)_{\mathcal{L}} + (2.23)_{\mathcal{L}'} \\ & & \\ \mathcal{Z}(C_2^{\mathcal{L}+\mathcal{L}'}(gg', g'')) & \longrightarrow & \mathcal{Z}(C_2^{\mathcal{L}}(gg', g'') + C_2^{\mathcal{L}'}(gg', g'')) \\ \uparrow & & \uparrow \\ (2.27)_{\mathcal{L}+\mathcal{L}'} & \longrightarrow & (2.27)_{\mathcal{L}} + (2.27)_{\mathcal{L}'} \end{array}$$

These facts together imply the commutativity of diagram (2.33). \square

Fix $g \in G$, the induced map $Z_G(g) \rightarrow \mathcal{P}$ given by $g' \mapsto C_2^\mathcal{L}(g, g')$ is denoted by $C_g^\mathcal{L}$. The bimultiplicativity of $C_2^\mathcal{L}$ implies that $C_g^\mathcal{L}$ is an object in $H^1(BZ_G(g), \mathcal{P})$. It is easy to see from the definition the following lemma:

Lemma 2.9. (i) *If two central extensions \mathcal{L} and \mathcal{L}' of G by \mathcal{P} are isomorphic in $H^2(BG, \mathcal{P})$, then for any g the induced two homomorphisms $C_g^\mathcal{L}$ and $C_g^{\mathcal{L}'}$ are isomorphic in $H^1(BZ_G(g), \mathcal{P})$.*

(ii) *One has that $C_g^\mathcal{P}$ is the trivial homomorphism for any $g \in G$.*

Let $Z_3 \subset G \times G \times G$ be the subset of pairwise commuting elements.

Proposition 2.10. *Let a map*

$$C_3^\mathcal{L} : Z_3 \rightarrow \pi_1(\mathcal{P})$$

be defined by

$$C_3^\mathcal{L}(g, g', g'') := \text{Comm}(C_g^\mathcal{L})(g', g'').$$

Then $C_3^\mathcal{L}$ is an anti-symmetric tri-multiplicative homomorphism from Z_3 to $\pi_1(\mathcal{P})$.

Proof. Let us see the tri-multiplicativity of the map $C_3^\mathcal{L}$. The multiplicativity of this map with respect to g' or g'' follows from lemma-definition 2.4. The multiplicativity of this map with respect to g follows from lemma-definition 2.7 and corollary 2.5.

The hard part is to prove now that the map $C_3^\mathcal{L}$ is anti-symmetric. Let us write C_2 instead of $C_2^\mathcal{L}$, and C_3 instead of $C_3^\mathcal{L}$ for simplicity. Let $(g, g', g'') \in Z_3$. First of all, let us observe that by definition, there is a canonical isomorphism

$$(2.34) \quad C_2(g, g'g'') + C_2(g', g'') \simeq C_2(g', g'') + C_2(g, g''g')$$

induced by the following 2-commutative diagram

$$\begin{array}{ccccc}
 & & \xrightarrow{\mathcal{Z}(C_2(g, g'g''))} & & \\
 \mathcal{L}_{gg'g''} & \xrightarrow{\quad} & \mathcal{L}_g + \mathcal{L}_{g'g''} & \xrightarrow{\quad} & \mathcal{L}_{g'g''} + \mathcal{L}_g & \xrightarrow{\quad} & \mathcal{L}_{gg'g''} \\
 & \downarrow & \downarrow & & \downarrow & & \downarrow \\
 & & \mathcal{L}_g + (\mathcal{L}_{g'} + \mathcal{L}_{g''}) & \xrightarrow{\quad} & (\mathcal{L}_{g'} + \mathcal{L}_{g''}) + \mathcal{L}_g & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{L}_g + (\mathcal{L}_{g''} + \mathcal{L}_{g'}) & \xrightarrow{\quad} & (\mathcal{L}_{g''} + \mathcal{L}_{g'}) + \mathcal{L}_g & & \\
 & & \downarrow & & \downarrow & & \\
 \mathcal{L}_{gg''g'} & \xrightarrow{\quad} & \mathcal{L}_g + \mathcal{L}_{g''g'} & \xrightarrow{\quad} & \mathcal{L}_{g''g'} + \mathcal{L}_g & \xrightarrow{\quad} & \mathcal{L}_{gg''g'} \\
 & & \xrightarrow{\mathcal{Z}(C_2(g, g''g'))} & & & & \\
 & & & & & &
 \end{array}$$

$\mathcal{Z}(C_2(g', g''))$ (left vertical arrow), $\mathcal{Z}(C_2(g', g''))$ (right vertical arrow)

The following lemma can be checked using the definition of $B\mathcal{P}$.

Lemma 2.11. *The isomorphism (2.34) is the same as the commutativity constraint in \mathcal{P} .*

Now, there are two isomorphisms between $(C_2(g, g') + C_2(g, g'')) + C_2(g', g'')$ and $C_2(g', g'') + (C_2(g, g'') + C_2(g, g'))$. Namely, the first isomorphism is obtained by the associativity and commutativity constraints in \mathcal{P} . (Recall that such isomorphism is

unique by the "Mac Lane's coherence theorem" for Picard category.) The second isomorphism is

$$(2.35) \quad (C_2(g, g') + C_2(g, g'')) + C_2(g', g'') \simeq C_2(g, g'g'') + C_2(g', g'') \stackrel{(2.34)}{\simeq} \\ C_2(g', g'') + C_2(g, g'g') \simeq C_2(g', g'') + (C_2(g, g'') + C_2(g, g')).$$

By the lemma, the difference between these two isomorphisms is $C_3(g, g', g'')$. If we recall the definition of $C_2(g, g'g'') \simeq C_2(g, g') + C_2(g, g'')$ by (2.29), we see that the isomorphism (2.35) can be represented by the following diagram

$$\begin{array}{ccccc}
& & \mathcal{L}_{g'g'g''} & \xrightarrow{\mathcal{Z}(C_2(g, g''))} & \mathcal{L}_{g'g''g} & & \\
& & \searrow & & \swarrow & & \\
\mathcal{Z}(C_2(g, g')) & & & & & & \mathcal{Z}(C_2(g', g'')) \\
& & \mathcal{L}_{g'} + \mathcal{L}_g + \mathcal{L}_{g''} & \longrightarrow & \mathcal{L}_{g'} + \mathcal{L}_{g''} + \mathcal{L}_g & & \\
& & \swarrow & & \searrow & & \\
\mathcal{L}_{gg'g''} \rightleftharpoons \mathcal{L}_g + \mathcal{L}_{g'} + \mathcal{L}_{g''} & & & & & & \mathcal{L}_{g''} + \mathcal{L}_{g'} + \mathcal{L}_g \rightleftharpoons \mathcal{L}_{g''g'g} \\
& & \swarrow & & \searrow & & \\
\mathcal{Z}(C_2(g', g'')) & & \mathcal{L}_g + \mathcal{L}_{g''} + \mathcal{L}_{g'} & \longrightarrow & \mathcal{L}_{g''} + \mathcal{L}_g + \mathcal{L}_{g'} & & \\
& & \swarrow & & \swarrow & & \\
& & \mathcal{L}_{gg''g'} & \xrightarrow{\mathcal{Z}(C_2(g, g''))} & \mathcal{L}_{g''gg'} & & \\
& & \searrow & & \swarrow & & \\
& & & & & & \mathcal{Z}(C_2(g, g'))
\end{array}$$

This diagram clearly implies that $C_3(g, g', g'') = C_3(g', g'', g)$. This together with the fact that $C_3(g, g', g'') = -C_3(g, g'', g')$ (because the map $\text{Comm}(C_g^{\mathcal{L}})$ is anti-symmetric) implies that C_3 is anti-symmetric. \square

Corollary 2.12. (i) If two central extensions \mathcal{L} and \mathcal{L}' of G by \mathcal{P} are isomorphic in $H^2(BG, \mathcal{P})$, then $C_3^{\mathcal{L}} = C_3^{\mathcal{L}'}$.

(ii) $C_3^{\mathcal{P}}$ is trivial.

Corollary 2.13. For any two central extensions \mathcal{L} and \mathcal{L}' of G by \mathcal{P} we have

$$C_3^{\mathcal{L}+\mathcal{L}'} = C_3^{\mathcal{L}} + C_3^{\mathcal{L}'}$$

Proof. This follows from lemma 2.8, corollary 2.5 and the definition of map C_3 . \square

Remark 2.9. If $\mathcal{P} = BA$, where A is an abelian group, then a central extension \mathcal{L} of a group G by the Picard groupoid \mathcal{P} is a gr -category such that these gr -categories are classified by the group $H^3(G, A)$ with the trivial G -module A . In this case the map $C_3^{\mathcal{L}}$ coincides with the symmetrization of corresponding 3-cocycle, see [Br3, §4]. (It follows from remarks 2.8, 2.5 and [O1, prop. 10].)

3. TATE VECTOR SPACES

3.1. The category of Tate vector spaces. We first review the definition of Tate vector spaces, following [O3, AK]. Let us fix a base field k .

Let us recall that for an exact category \mathcal{E} in the sense of Quillen (cf. [Q]), Beilinson associates another exact category $\lim_{\leftarrow} \mathcal{E}$, which is again an exact category (cf. [Be2]). In nowadays terminology, this is the category of locally compact objects of \mathcal{E} .

For an exact category \mathcal{E} , let $\hat{\mathcal{E}}$ denote the category of left exact additive contravariant functors from the category \mathcal{E} to the category of abelian groups. This is

again an exact category (in fact an abelian category), in which arbitrary small colimits exist. The Yoneda embedding $h : \mathcal{E} \rightarrow \hat{\mathcal{E}}$ is exact. Then the category $\text{Ind}(\mathcal{E})$ of (strict) ind-objects of \mathcal{E} , is the full subcategory of $\hat{\mathcal{E}}$ consisting of objects of the form $\varinjlim_{i \in I} h(X_i)$, where I is a filtered small category, and $X_i \in \mathcal{E}$, such that for $i \rightarrow j$ in I , the map $X_i \rightarrow X_j$ is an admissible monomorphism. This category is a natural exact category. Likewise, one can define $\text{Pro}(\mathcal{E})$ as $\text{Ind}(\mathcal{E}^{op})^{op}$.

Definition 3.1. Let \mathcal{E} be an exact category. Then $\varprojlim \mathcal{E}$ is the full subcategory of $\text{Pro}(\text{Ind}(\mathcal{E}))$ consisting of objects that can be represented as $\varprojlim_{i \in I^{op}} \varinjlim_{j \in I} h(X_{ij})$ such that for any $i \rightarrow i', j \rightarrow j'$, the following diagram is cartesian (which is automatically cocartesian then).

$$\begin{array}{ccc} X_{ij} & \longrightarrow & X_{ij'} \\ \downarrow & & \downarrow \\ X_{i'j} & \longrightarrow & X_{i'j'} \end{array}$$

One can show that $\varprojlim \mathcal{E}$ is an exact category and the following embedding $\varprojlim \mathcal{E} \rightarrow \text{Pro}(\text{Ind}(\mathcal{E}))$ is exact. In addition, there is a natural embedding $\varprojlim \mathcal{E} \rightarrow \text{Ind}(\text{Pro}(\mathcal{E}))$ which is again exact. It is clear that the natural embedding $\text{Ind}(\mathcal{E}) \rightarrow \text{Pro}(\text{Ind}(\mathcal{E}))$ lands in $\varprojlim \mathcal{E}$ and similarly the natural embedding $\text{Pro}(\mathcal{E}) \rightarrow \text{Ind}(\text{Pro}(\mathcal{E}))$ also lands in $\varprojlim \mathcal{E}$.

Definition 3.2. Define Tate_0 to be the category of finite dimensional vector spaces, together with its canonical exact category structure. Define $\text{Tate}_n = \varprojlim \text{Tate}_{n-1}$, together with the canonical exact category structure given by Beilinson.

There is a canonical forgetful functor $F_n : \text{Tate}_n \rightarrow \mathcal{T}op$, where $\mathcal{T}op$ denotes the category of topological vector spaces. As is shown in [O3], the functor is fully faithful when $n = 1$, but this is in general not the case when $n > 1$.

Definition 3.3. Let V be an object of Tate_n . A *lattice* L of V is an object in Tate_n which actually belongs to $\text{Pro}(\text{Tate}_{n-1})$, together with an admissible monomorphism $L \rightarrow V$ such that the object V/L belongs to $\text{Ind}(\text{Tate}_{n-1})$. A *colattice* L^c of V is an object in Tate_n which actually belongs to $\text{Ind}(\text{Tate}_{n-1})$, together with an admissible monomorphism $L^c \rightarrow V$ such that the object V/L^c belongs to $\text{Pro}(\text{Tate}_{n-1})$.

It is clear that if L is a lattice of V and L^c is a colattice, then $L \cap L^c$ belongs to Tate_{n-1} .

The main players of this paper are Tate_1 and Tate_2 . The category Tate_1 is just the category of locally linearly compact k -vector spaces. A typical object in Tate_1 is the field of formal Laurent series $k((t))$, i.e. $k((t))$ is the field of fractions of the ring $k[[t]]$. $k((t))$ is equipped with the standard topology, where the base of neighbourhoods of zero consists of integer powers of the maximal ideal of $k[[t]]$. The subspace $k[[t]]$ is a lattice in $k((t))$ and $k[t^{-1}]$ is a colattice. Observe that $k[t] \subset k((t))$ is neither a lattice nor a colattice, because the subspace $k[t]$ is not closed in the topological space $k((t))$. Therefore the embedding $k[t] \hookrightarrow k((t))$ is not an admissible monomorphism, since any admissible (exact) triple in the category Tate_1 is of the form:

$$0 \longrightarrow V_1 \longrightarrow V_2 \longrightarrow V_3 \longrightarrow 0,$$

where the locally linearly compact vector space V_1 is a closed vector subspace in a locally linearly compact vector space V_2 , and the locally linearly compact vector space V_3 has the quotient topology on the quotient vector space.

A typical object in Tate_2 is $k((t))((s))$, since

$$k((t))((s)) = \varprojlim_{l \in \mathbb{Z}} \varinjlim_{m \leq l} s^m k((t))[[s]] / s^l k((t))[[s]] = \varinjlim_{m \in \mathbb{Z}} \varprojlim_{l \geq m} s^m k((t))[[s]] / s^l k((t))[[s]],$$

and $s^m k((t))[[s]] / s^l k((t))[[s]]$ is a locally linearly compact k -vector space.

The k -space $k((t))[[s]]$ is a lattice, and the k -space $k((t))[s^{-1}]$ is a colattice in the k -space $k((t))((s))$. As just mentioned above, it is not enough to regard them as topological vector spaces. On the other hand $k[[t]]((s))$ is not a lattice in $k((t))((s))$ although the natural map $k[[t]]((s)) \rightarrow k((t))((s))$ is an admissible monomorphism.

Remark 3.1. The category Tate_n coincides with the category of complete C_n -spaces from [O3].

3.2. Determinant theories of Tate vector spaces. We consider Tate_0 as an exact category. Then $\det : (\text{Tate}_0, \text{isom}) \rightarrow \mathcal{P}ic^{\mathbb{Z}}$ (see (2.6)) is a functor satisfying the following additional property: for each injective homomorphism $V_1 \rightarrow V$ in the category Tate_0 , there is a canonical isomorphism

$$(3.1) \quad \det(V_1) \otimes \det(V/V_1) \simeq \det(V),$$

such that:

(i) for $V_1 = 0$ (resp. $V_1 = V$), equality (3.1) is the same as

$$(3.2) \quad \ell_0 \otimes \det(V) \simeq \det(V)$$

resp.

$$(3.3) \quad \det(V) \otimes \ell_0 \simeq \det(V),$$

where ℓ_0 is the trivial k -line of degree zero.

(ii) For any diagram

$$(3.4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & U_1 & \longrightarrow & U & \longrightarrow & U/U_1 \longrightarrow 0 \\ & & \downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\ 0 & \longrightarrow & V_1 & \longrightarrow & V & \longrightarrow & V/V_1 \longrightarrow 0, \end{array}$$

the following diagram is commutative

$$(3.5) \quad \begin{array}{ccc} \det(U_1) \otimes \det(U/U_1) & \longrightarrow & \det(U) \\ \downarrow & & \downarrow \\ \det(V_1) \otimes \det(V/V_1) & \longrightarrow & \det(V). \end{array}$$

(iii) For any diagram

$$(3.6) \quad \begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & U_1 & \longrightarrow & V_1 & \longrightarrow & W_1 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & U & \longrightarrow & V & \longrightarrow & W \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & U/U_1 & \longrightarrow & V/V_1 & \longrightarrow & W/W_1 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array} ,$$

the following diagram is commutative

$$(3.7) \quad \begin{array}{ccc} (\det(U_1) \otimes \det(U/U_1)) \otimes (\det(W_1) \otimes \det(W/W_1)) & \longrightarrow & \det(U) \otimes \det(W) \\ \downarrow \text{ass. and comm. constraints} & & \downarrow \\ (\det(U_1) \otimes \det(W_1)) \otimes (\det(U/U_1) \otimes \det(W/W_1)) & & \\ \downarrow & & \downarrow \\ \det(V_1) \otimes \det(V/V_1) & \longrightarrow & \det(V) \end{array}$$

Definition 3.4. Let \mathcal{P} be a Picard groupoid. A determinant functor from the category $(\text{Tate}_0, \text{isom})$ to \mathcal{P} is a functor $D : (\text{Tate}_0, \text{isom}) \rightarrow \mathcal{P}$ together with isomorphisms (3.1) satisfying equalities and diagrams (3.2)-(3.7), where we have to change notation "det" to notation "D" everywhere in these formulas.

There is the following obvious proposition.

Proposition 3.1. *Let $D : (\text{Tate}_0, \text{isom}) \rightarrow \mathcal{P}$ be a determinant functor. Then there is a 1-homomorphism of Picard groupoids $\tilde{D} : \mathcal{P}\text{ic}^{\mathbb{Z}} \rightarrow \mathcal{P}$ and a monoidal natural transformation $\varepsilon : \tilde{D} \circ \det \simeq D$. Furthermore, the pair (\tilde{D}, ε) is unique up to a unique isomorphism.*

Remark 3.2. All the above discussions are valid when one replaces k by a noetherian commutative ring A , and replaces Tate_0 by the category of finitely generated projective A -modules.

Next we turn to Tate_1 . The following result is fundamental and is due to Kapranov, [Kap], but see also [Dr, §5.1-5.3].

Proposition 3.2. *There is a natural functor*

$$\mathcal{D}et : (\text{Tate}_1, \text{isom}) \rightarrow B\mathcal{P}\text{ic}^{\mathbb{Z}},$$

and for each admissible monomorphism $\mathcal{V}_1 \rightarrow \mathcal{V}$ there is a 1-isomorphism

$$(3.8) \quad \mathcal{D}et(\mathcal{V}_1) + \mathcal{D}et(\mathcal{V}/\mathcal{V}_1) \rightarrow \mathcal{D}et(\mathcal{V})$$

such that if $\mathcal{V}_1 = 0$ (resp. $\mathcal{V}_1 = \mathcal{V}$), this 1-isomorphism is the canonical 1-isomorphism

$$\mathcal{P} + \mathcal{D}et(\mathcal{V}) \simeq \mathcal{D}et(\mathcal{V})$$

resp.

$$\mathcal{D}et(\mathcal{V}) + \mathcal{P} \simeq \mathcal{D}et(\mathcal{V}).$$

For each admissible diagram (3.4) of 1-Tate vector spaces, the corresponding diagram (3.5) is commutative. In addition, for each admissible diagram (3.6) of 1-Tate vector spaces, there is a 2-isomorphism for the corresponding diagram (3.7).

Remark 3.3. Under conditions of proposition 3.2, the 2-isomorphisms which appear from diagram (3.7) satisfy further compatibility conditions.

Proof. For a 1-Tate vector space \mathcal{V} , we recall the definition of a graded-determinantal theory Δ on \mathcal{V} . This is a rule that assign to every lattice $L \subset \mathcal{V}$ an object $\Delta(L)$ from $\mathcal{P}ic^{\mathbb{Z}}$ and to every lattices $L_1 \subset L_2 \subset \mathcal{V}$ an isomorphism

$$\Delta_{L_1, L_2} : \Delta(L_1) \otimes \det(L_2/L_1) \longrightarrow \Delta(L_2)$$

such that for any three lattices $L_1 \subset L_2 \subset L_3 \subset \mathcal{V}$ the following diagram is commutative

$$\begin{array}{ccc} \Delta(L_1) \otimes \det(L_2/L_1) \otimes \det(L_3/L_2) & \longrightarrow & \Delta(L_1) \otimes \det(L_3/L_1) \\ \downarrow & & \downarrow \\ \Delta(L_2) \otimes \det(L_3/L_2) & \longrightarrow & \Delta(L_3). \end{array}$$

Let $\mathcal{D}et(\mathcal{V})$ be the category of graded-determinantal theories on \mathcal{V} . This is a $\mathcal{P}ic^{\mathbb{Z}}$ -torsor, where for any $x \in \mathcal{P}ic^{\mathbb{Z}}$, $\Delta \in \mathcal{D}et(\mathcal{V})$, we have $(x + \Delta)(L) := x \otimes \Delta(L)$.

Now for an admissible (exact) sequence

$$0 \longrightarrow \mathcal{V}_1 \longrightarrow \mathcal{V} \xrightarrow{\epsilon} \mathcal{V}/\mathcal{V}_1 \longrightarrow 0$$

the 1-isomorphism (3.8) is constructed as

$$\Delta(L) := \Delta_1(L \cap \mathcal{V}_1) \otimes \Delta_2(\epsilon(L)),$$

where L is a lattice in \mathcal{V} , $\Delta_1 \in \mathcal{D}et(\mathcal{V}_1)$, $\Delta_2 \in \mathcal{D}et(\mathcal{V}/\mathcal{V}_1)$, $\Delta \in \mathcal{D}et(\mathcal{V})$. (We used that the k -space $L \cap \mathcal{V}_1$ is a lattice in the 1-Tate vector space \mathcal{V}_1 , and the k -space $\epsilon(L)$ is a lattice in the 1-Tate vector space \mathcal{V}_3).

We note that, by construction, $\mathcal{V} \mapsto \mathcal{D}et(\mathcal{V})$ is naturally a contravariant functor from the category $(\text{Tate}_1, \text{isom})$ to the category $B\mathcal{P}ic^{\mathbb{Z}}$. To obtain the covariant functor we have to inverse arrows in the category $(\text{Tate}_1, \text{isom})$. \square

4. APPLICATIONS TO THE CASE $G = \text{GL}(k((t)))$ AND $\text{GL}(k((t))((s)))$

4.1. Tame symbols. Let us first review the tame symbols. Recall that if K is a field with discrete valuation $\nu : K^\times \rightarrow \mathbb{Z}$, and k denote its residue field, then there are so-called boundary maps for any $i \in \mathbb{N}$

$$\partial_i : K_i^M(K) \longrightarrow K_{i-1}^M(k),$$

where $K_i^M(F)$ denotes the i th Milnor K-group of a field F . Let us also recall that for a field F , the i th Milnor K-group $K_i^M(F)$ is the quotient of the abelian group $F^\times \otimes_{\mathbb{Z}} F^\times \otimes_{\mathbb{Z}} \cdots \otimes_{\mathbb{Z}} F^\times$ modulo the so-called Steinberg relations. Then the tame symbol is defined as the composition of the following maps

$$\{\cdot, \cdot\} : K^\times \otimes_{\mathbb{Z}} K^\times \longrightarrow K_2^M(K) \xrightarrow{\partial_2} K_1^M(k) \simeq k^\times.$$

Explicitly, let $\pi \subset K$ be the maximal ideal. Then

$$(4.1) \quad \{f, g\} = (-1)^{\nu(f)\nu(g)} \frac{f^{\nu(g)}}{g^{\nu(f)}} \pmod{\pi}$$

Now, let \mathbb{K} be a two-dimensional local field, whose residue field is denoted by K , whose residue field is k . Then we define the following map as

$$\nu_{\mathbb{K}} : \mathbb{K}^{\times} \otimes_{\mathbb{Z}} \mathbb{K}^{\times} \longrightarrow K_2^M(\mathbb{K}) \xrightarrow{\partial_2} K_1^M(K) \xrightarrow{\partial_1} K_0^M(k) \simeq \mathbb{Z},$$

and define the two-dimensional tame symbol as

$$\{\cdot, \cdot, \cdot\} : \mathbb{K}^{\times} \otimes_{\mathbb{Z}} \mathbb{K}^{\times} \otimes_{\mathbb{Z}} \mathbb{K}^{\times} \longrightarrow K_3^M(\mathbb{K}) \xrightarrow{\partial_3} K_2^M(K) \xrightarrow{\partial_2} K_1^M(k) \simeq k^{\times}.$$

We have the following explicit formulas for $\nu_{\mathbb{K}}$ and $\{\cdot, \cdot, \cdot\}$ (see [O1].) Let $\nu_1 : \mathbb{K} \rightarrow \mathbb{Z}$, and $\nu_2 : K \rightarrow \mathbb{Z}$ be discrete valuations. Let $\pi_{\mathbb{K}}$ be the maximal ideal of \mathbb{K} , π_K be the maximal ideal of K . For an element $f \in \mathcal{O}_{\mathbb{K}}$, let \bar{f} denote its residue class in K . Then

$$(4.2) \quad \nu_{\mathbb{K}}(f, g) = \nu_2\left(\frac{\overline{f^{\nu_1(g)}}}{\overline{g^{\nu_1(f)}}}\right)$$

$$(4.3) \quad \{f, g, h\} = \text{sgn}(f, g, h) f^{\nu_{\mathbb{K}}(g, h)} g^{\nu_{\mathbb{K}}(h, f)} h^{\nu_{\mathbb{K}}(f, g)} \pmod{\pi_{\mathbb{K}}} \pmod{\pi_K}$$

where

$$(4.4) \quad \begin{aligned} \text{sgn}(f, g, h) &= (-1)^A, \\ A &= \nu_{\mathbb{K}}(f, g)\nu_{\mathbb{K}}(f, h) + \nu_{\mathbb{K}}(g, h)\nu_{\mathbb{K}}(g, f) + \nu_{\mathbb{K}}(h, f)\nu_{\mathbb{K}}(h, g) + \nu_{\mathbb{K}}(f, g)\nu_{\mathbb{K}}(g, h)\nu_{\mathbb{K}}(h, f). \end{aligned}$$

Remark 4.1. Originally one used another explicit formula for the sign of the two-dimensional tame symbol. This other formula was introduced in [Pa1].

It is easy to see that tame symbols $\{\cdot, \cdot\}$, $\{\cdot, \cdot, \cdot\}$ and the map $\nu_{\mathbb{K}}$ are anti-symmetric.

4.2. The one-dimensional story. Let \mathcal{V} be a 1-Tate vector space over k . The group of automorphisms of \mathcal{V} in this category is denoted by $\text{GL}(\mathcal{V})$.

Proposition 4.1. *There is a homomorphism $\text{Det}_{\mathcal{V}} : \text{GL}(\mathcal{V}) \rightarrow \text{Pic}^{\mathbb{Z}}$, which is canonical up to a unique isomorphism in $H^1(\text{BGL}(\mathcal{V}), \text{Pic}^{\mathbb{Z}})$.*

Proof. According to proposition 3.2, we have a homomorphism

$$\text{GL}(\mathcal{V}) \rightarrow \text{Hom}_{\text{Pic}^{\mathbb{Z}}}(\text{Det}(\mathcal{V}), \text{Det}(\mathcal{V})) \simeq \text{Pic}^{\mathbb{Z}}$$

via \mathcal{Z}^{-1} , where $\mathcal{Z} : \text{Pic}^{\mathbb{Z}} \rightarrow \text{Hom}_{\text{Pic}^{\mathbb{Z}}}(\text{Det}(\mathcal{V}), \text{Det}(\mathcal{V}))$ is a natural homomorphism from section 2.3. \square

Choose $\mathcal{L} \subset \mathcal{V}$ a lattice. It follows from the proof of proposition 3.2 that in concrete terms, one has to assign to $\text{Det}_{\mathcal{V}}(g)$ the graded line

$$(4.5) \quad \det(\mathcal{L} | g\mathcal{L}) := \det\left(\frac{g\mathcal{L}}{\mathcal{L} \cap g\mathcal{L}}\right) \otimes \det\left(\frac{\mathcal{L}}{\mathcal{L} \cap g\mathcal{L}}\right)^{-1},$$

where $g \in \text{GL}(\mathcal{V})$. Then, it is well-known that there is a canonical isomorphism

$$\det(\mathcal{L} | gg'\mathcal{L}) \simeq \det(\mathcal{L} | g\mathcal{L}) \otimes \det(g\mathcal{L} | g'\mathcal{L}) \simeq \det(\mathcal{L} | g\mathcal{L}) \otimes \det(\mathcal{L} | g'\mathcal{L}),$$

which is compatible with the associativity constraints in the category $\mathcal{P}ic^{\mathbb{Z}}$ (see, for example, [FZ, §1]). For different choice of \mathcal{L} , the resulting objects in $H^1(BGL(\mathcal{V}), \mathcal{P}ic^{\mathbb{Z}})$ are isomorphic.

We also have the following lemma, which easily follows from the construction of homomorphism $\mathcal{D}et_{\mathcal{V}}$ and the discussion in §3.2 (in particular the diagram (3.7)).

Lemma 4.2. *If $0 \rightarrow \mathcal{V}' \rightarrow \mathcal{V} \rightarrow \mathcal{V}'' \rightarrow 0$ is a short exact sequence of 1-Tate vector spaces (recall that Tate_1 is an exact category). Let P be the subgroup of $\text{GL}(\mathcal{V})$ that preserves this sequence, then there is a canonical 1-isomorphism $\mathcal{D}et_{\mathcal{V}'} + \mathcal{D}et_{\mathcal{V}''} \simeq \mathcal{D}et_{\mathcal{V}}$ in $H^1(BP, \mathcal{P}ic^{\mathbb{Z}})$.*

Remark 4.2. The 1-homomorphism $F_{\mathcal{P}ic} \circ \mathcal{D}et_{\mathcal{V}} : \text{GL}(\mathcal{V}) \rightarrow \mathcal{P}ic$ is essentially constructed in [ACK]. However, the above lemma does not hold for this 1-homomorphism. This is the complication of the sign issues in [ACK].

Now let k'/k be a finite extension and $K = k'((t))$ be a local field with residue field k' . Then K has a natural structure as a 1-Tate vector space over k . Let $H = K^{\times}$. The multiplication gives a natural embedding $H \subset \text{GL}(K)$. The following proposition is from [BBE].

Proposition 4.3. *If $f, g \in H$, then*

$$\text{Comm}(\mathcal{D}et_K)(f, g) = \text{Nm}_{k'/k}\{f, g\}^{-1}$$

that is inverse to the tame symbol of f and g .

Remark 4.3. Since the natural functor $F_{\mathcal{P}ic}$ is monoidal, the restriction to H of the functor $F_{\mathcal{P}ic} \circ \mathcal{D}et_K$ determines a homomorphism $H \rightarrow \mathcal{P}ic$. The commutator pairing $\text{Comm}(f, g)$ constructed by this homomorphism is

$$(-1)^{\text{ord}(f)\text{ord}(g)} \text{Nm}_{k'/k}\{f, g\}^{-1}.$$

By definition 3.3, a lattice \mathcal{L} of \mathcal{V} is a linearly compact open k -subspace of \mathcal{V} such that \mathcal{V}/\mathcal{L} is a discrete k -space. A colattice \mathcal{L}^c is a k -subspace of \mathcal{V} such that for any lattice \mathcal{L} , both $\mathcal{L}^c \cap \mathcal{L}$ and $\mathcal{V}/(\mathcal{L}^c + \mathcal{L})$ are finite dimensional.

Lemma 4.4. *Let $P \subset \text{GL}(\mathcal{V})$ be a subgroup of $\text{GL}(\mathcal{V})$ that preserves a lattice (or a colattice) in \mathcal{V} , then the homomorphism $\mathcal{D}et_{\mathcal{V}}$ is trivial on P .*

Proof. Let $\mathcal{L} \subset \mathcal{V}$ be a lattice such that the group P preserves it. We consider an exact sequence of 1-Tate vector spaces

$$0 \rightarrow \mathcal{L} \rightarrow \mathcal{V} \rightarrow \mathcal{V}/\mathcal{L} \rightarrow 0.$$

Then the group P preserves this sequence. Therefore by lemma 4.2, it is enough to prove that the homomorphisms $\mathcal{D}et_{\mathcal{L}}$ and $\mathcal{D}et_{\mathcal{L}/\mathcal{V}}$ are trivial on P . But this is obvious from the proof of proposition 4.1.

For a colattice $\mathcal{L}^c \subset \mathcal{V}$ we have to use the analogous reasonings. □

4.3. The two-dimensional story. If $\mathbb{V} \in \text{Tate}_2$, then we denote by $\text{GL}(\mathbb{V})$ the group of automorphisms of \mathbb{V} in this category.

There should be a determinantal functor from $(\text{Tate}_2, \text{isom})$ to $B^2\mathcal{P}ic^{\mathbb{Z}}$, which assigns to every such \mathbb{V} the graded gerbel theory in the sense of [AK], satisfying properties which generalize properties listed in proposition 3.2 (and further compatibility conditions). We do not make it precise. But we define the corresponding

central extension of $\mathrm{GL}(\mathbb{V})$ as follows. Pick a lattice \mathbb{L} of \mathbb{V} . Then one associates with g the $\mathcal{P}ic^{\mathbb{Z}}$ -torsor

$$(4.6) \quad \mathcal{D}et_{\mathbb{V}}(g) = \mathcal{D}et(\mathbb{L} \mid g\mathbb{L}) := \mathcal{D}et\left(\frac{g\mathbb{L}}{\mathbb{L} \cap g\mathbb{L}}\right) - \mathcal{D}et\left(\frac{\mathbb{L}}{\mathbb{L} \cap g\mathbb{L}}\right).$$

This definition is correct because both k -spaces $\frac{g\mathbb{L}}{\mathbb{L} \cap g\mathbb{L}}$ and $\frac{\mathbb{L}}{\mathbb{L} \cap g\mathbb{L}}$ belong to objects of category Tate_1 . We define the 1-isomorphism as

$$(4.7) \quad \mathcal{D}et(\mathbb{L} \mid gg'\mathbb{L}) \simeq \mathcal{D}et(\mathbb{L} \mid g\mathbb{L}) + \mathcal{D}et(g\mathbb{L} \mid gg'\mathbb{L}) \simeq \mathcal{D}et(\mathbb{L} \mid g\mathbb{L}) + \mathcal{D}et(\mathbb{L} \mid g'\mathbb{L}).$$

One uses proposition 3.2 to check that this defines a central extension of $\mathrm{GL}(\mathbb{V})$ by $\mathcal{P}ic^{\mathbb{Z}}$. This central extension depends on the chosen lattice \mathbb{L} of \mathbb{V} . If we change the lattice, then the central extension constructed by a new lattice will be isomorphic to the previous one.

Remark 4.4. If one replaces $\mathcal{P}ic^{\mathbb{Z}}$ by $\mathcal{P}ic$, such a central extension was constructed in [O1, FZ]. Besides, in [O1] the two-dimensional tame symbol up to sign was obtained as an application of this construction, and the reciprocity laws on algebraic surfaces were proved up to sign.

As generalization of lemma 4.2 and lemma 4.4 it is not difficult to prove the following lemmas.

Lemma 4.5. *If $0 \rightarrow \mathbb{V}' \rightarrow \mathbb{V} \rightarrow \mathbb{V}'' \rightarrow 0$ is a short exact sequence of 2-Tate vector spaces (recall that Tate_2 is an exact category). Let P be the subgroup of $\mathrm{GL}(\mathbb{V})$ that preserves this sequence, then there is a canonical 1-isomorphism $\mathcal{D}et_{\mathbb{V}'} + \mathcal{D}et_{\mathbb{V}''} \simeq \mathcal{D}et_{\mathbb{V}}$ in $H^2(\mathrm{BP}, \mathcal{P}ic^{\mathbb{Z}})$.*

Lemma 4.6. *Let P be subgroup of $\mathrm{GL}(\mathbb{V})$ which preserves a lattice or a colattice in \mathbb{V} , then the central extension restricted to P can be trivialized.*

Let k'/k be a finite field extension, and $\mathbb{K} = k'((t))((s))$ be a two-dimensional local field. Then \mathbb{K} has a natural structure as a 2-Tate vector space over k . The group $H = \mathbb{K}^{\times}$ acts on \mathbb{K} by left multiplications, which gives rise to an embedding $H \rightarrow \mathrm{GL}(\mathbb{K})$.

Theorem 4.7. *For $f, g, h \in H$, one has*

$$C_3^{\mathcal{D}et}(f, g, h) = \mathrm{Nm}_{k'/k}\{f, g, h\},$$

where the map $C_3^{\mathcal{D}et}$ is constructed in lemma 2.10 and $\{\cdot, \cdot, \cdot\}$ is the two-dimensional tame symbol.

In what follows, we will denote the bimultiplicative homomorphism $C_2^{\mathcal{D}et}$ by C_2 , the homomorphism $C_g^{\mathcal{D}et}$ by C_g and the map $C_3^{\mathcal{D}et}$ by C_3 .

Proof. Since both maps C_3 and $\mathrm{Nm}_{k'/k}\{\cdot, \cdot, \cdot\}$ are anti-symmetric and tri-multiplicative, we just need to consider the following cases: (i) $f, g, h \in \mathcal{O}_{\mathbb{K}}^{\times}$; (ii) $f, g \in \mathcal{O}_{\mathbb{K}}^{\times}, h = s$; (iii) $f \in \mathcal{O}_{\mathbb{K}}^{\times}, g = h = s$; (iv) $f = g = h = s$. Here $\mathcal{O}_{\mathbb{K}} = k'((t))[[s]]$ is the ring of integers of the field \mathbb{K} , which is also a lattice in \mathbb{K} . We will fix $\mathbb{L} = \mathcal{O}_{\mathbb{K}}$.

In the first case, we have that both C_3 and $\mathrm{Nm}_{k'/k}\{\cdot, \cdot, \cdot\}$ are trivial (to see that C_3 is trivial, one uses lemma 4.6).

According to formulas (4.2)-(4.4), the second case amounts to proving that

$$C_3(f, g, s) = \mathrm{Nm}_{k'/k}\{\bar{f}, \bar{g}\},$$

where \bar{f}, \bar{g} are the image of elements f, g under the map $\mathcal{O}_{\mathbb{K}}^{\times} \rightarrow K^{\times}$.

Let us consider a little more general situation. Let $f, g \in \mathrm{GL}(\mathbb{K})$ that leave the lattice $\mathcal{O}_{\mathbb{K}}$ invariant, and let $h \in \mathrm{GL}(\mathbb{K})$ such that $h\mathcal{O}_{\mathbb{K}} \subset \mathcal{O}_{\mathbb{K}}$. Let $\mathcal{V} = \mathcal{O}_{\mathbb{K}}/h\mathcal{O}_{\mathbb{K}}$, which is a 1-Tate vector space over the field k . We assume that f, g, h mutually commute with each other. Then $f, g : \mathcal{O}_{\mathbb{K}} \rightarrow \mathcal{O}_{\mathbb{K}}$ induce automorphisms $\pi_h(f), \pi_h(g) : \mathcal{V} \rightarrow \mathcal{V}$. Let $\mathcal{D}et$ be the central extension of $\mathrm{GL}(\mathbb{K})$ by $\mathcal{P}ic^{\mathbb{Z}}$ defined by the lattice $\mathbb{L} = \mathcal{O}_{\mathbb{K}}$. By definition, under the isomorphism $\mathcal{Z} : \mathcal{P}ic^{\mathbb{Z}} \rightarrow \mathrm{Hom}_{\mathcal{P}ic^{\mathbb{Z}}}(\mathcal{D}et(\mathcal{O}_{\mathbb{K}}|hg\mathcal{O}_{\mathbb{K}}), \mathcal{D}et(\mathcal{O}_{\mathbb{K}}|hg\mathcal{O}_{\mathbb{K}}))$, the 1-isomorphism $C_2(h, g)$ corresponds to the following composition of 1-isomorphisms of $\mathcal{P}ic^{\mathbb{Z}}$ -torsors

$$\begin{aligned} & \mathcal{D}et(\mathcal{O}_{\mathbb{K}}|hg\mathcal{O}_{\mathbb{K}}) \rightarrow \mathcal{D}et(\mathcal{O}_{\mathbb{K}}|h\mathcal{O}_{\mathbb{K}}) + \mathcal{D}et(h\mathcal{O}_{\mathbb{K}}|hg\mathcal{O}_{\mathbb{K}}) \rightarrow \mathcal{D}et(\mathcal{O}_{\mathbb{K}}|h\mathcal{O}_{\mathbb{K}}) + \mathcal{D}et(\mathcal{O}_{\mathbb{K}}|g\mathcal{O}_{\mathbb{K}}) \\ & \rightarrow \mathcal{D}et(\mathcal{O}_{\mathbb{K}}|g\mathcal{O}_{\mathbb{K}}) + \mathcal{D}et(\mathcal{O}_{\mathbb{K}}|h\mathcal{O}_{\mathbb{K}}) \rightarrow \mathcal{D}et(\mathcal{O}_{\mathbb{K}}|g\mathcal{O}_{\mathbb{K}}) + \mathcal{D}et(g\mathcal{O}_{\mathbb{K}}|gh\mathcal{O}_{\mathbb{K}}) \rightarrow \mathcal{D}et(\mathcal{O}_{\mathbb{K}}|gh\mathcal{O}_{\mathbb{K}}). \end{aligned}$$

Using the fact that $g\mathcal{O}_{\mathbb{K}} = \mathcal{O}_{\mathbb{K}}$ and proposition 3.2, the above 1-isomorphism is canonically 2-isomorphic to the following 1-isomorphism

$$\mathcal{D}et(\mathcal{O}_{\mathbb{K}} | hg\mathcal{O}_{\mathbb{K}}) \xrightarrow{\mathcal{Z}(\mathcal{D}et_{\mathcal{V}}(\pi_h(g)))} \mathcal{D}et(\mathcal{O}_{\mathbb{K}} | hg\mathcal{O}_{\mathbb{K}}).$$

Therefore, there is a canonical 2-isomorphism $C_2(h, g) \simeq -\mathcal{D}et_{\mathcal{V}}(\pi_h(g))$, because, by definition (see formula (4.6)), $\mathcal{D}et(\mathcal{O}_{\mathbb{K}} | h\mathcal{O}_{\mathbb{K}}) \simeq -\mathcal{D}et(\mathcal{V})$. One readily checks by the construction of lemma-definition 2.7, that these 2-isomorphisms fit into the following commutative diagrams

$$(4.8) \quad \begin{array}{ccc} C_2(h, fg) & \longrightarrow & C_2(h, f) + C_2(h, g) \\ \downarrow & & \downarrow \\ -\mathcal{D}et_{\mathcal{V}}(\pi_h(fg)) & \longrightarrow & -\mathcal{D}et_{\mathcal{V}}(\pi_h(f)) - \mathcal{D}et_{\mathcal{V}}(\pi_h(g)) \end{array}$$

where the natural isomorphism $\mathcal{D}et_{\mathcal{V}}(\pi_h(fg)) \rightarrow \mathcal{D}et_{\mathcal{V}}(\pi_h(f)) + \mathcal{D}et_{\mathcal{V}}(\pi_h(g))$ comes from proposition 4.1. (We have to use that $\mathcal{D}et((0))$ is canonically isomorphic to $\mathcal{P}ic^{\mathbb{Z}}$, and $\mathcal{O}_{\mathbb{K}}/g\mathcal{O}_{\mathbb{K}} = (0)$, where (0) is the zero-space.)

We now return to our proof of case (ii). Let P_s be the subgroup of $\mathrm{GL}(\mathbb{K})$ consisting of elements that preserve the lattice $\mathcal{O}_{\mathbb{K}}$ and commute with the element s . Then elements in the group P_s also preserve the lattice $s\mathcal{O}_{\mathbb{K}}$, and therefore induce a group homomorphism

$$\pi_s : P_s \rightarrow \mathrm{GL}(K),$$

because $\mathbb{K} = \mathcal{O}_{\mathbb{K}}/s\mathcal{O}_{\mathbb{K}}$. Then the commutative diagram (4.8) amounts to the following lemma.

Lemma 4.8. *The homomorphism $C_s : P_s \rightarrow \mathcal{P}ic^{\mathbb{Z}}$ is isomorphic to the minus (or the inverse) of the following homomorphism*

$$\mathcal{D}et_K \circ \pi_s : P_s \rightarrow \mathrm{GL}(K) \rightarrow \mathcal{P}ic^{\mathbb{Z}}.$$

By proposition 4.3, we thus obtain that

$$C_3(f, g, s) = C_3(s, f, g) = \mathrm{Comm}(C_s)(f, g) = \mathrm{Nm}_{k'/k}\{\bar{f}, \bar{g}\}$$

for $f, g \in \mathcal{O}_{\mathbb{K}}^{\times} \subset \mathrm{GL}(\mathbb{K})$. The case (ii) follows.

Case (iii). According to formulas (4.2)-(4.4), one needs to show

$$C_3(f, s, s) = C_f(s, s) = \mathrm{Nm}_{k'/k}(-1)^{\nu_2(\bar{f})} = (-1)^{(\nu_2(\bar{f})[k':k])} = (-1)^{(\nu_2(\bar{f})[k':k])^2}.$$

We have the following exact sequence of 1-Tate vector spaces

$$0 \longrightarrow \frac{s\mathcal{O}_{\mathbb{K}}}{s^2\mathcal{O}_{\mathbb{K}}} \longrightarrow \frac{\mathcal{O}_{\mathbb{K}}}{s^2\mathcal{O}_{\mathbb{K}}} \longrightarrow \frac{\mathcal{O}_{\mathbb{K}}}{s\mathcal{O}_{\mathbb{K}}} \longrightarrow 0.$$

and therefore by lemma 4.2, for any element $p \in P_s$, there is a canonical isomorphism in $\mathcal{P}ic^{\mathbb{Z}}$

$$(4.9) \quad \mathcal{D}et_{\frac{\mathcal{O}_{\mathbb{K}}}{s^2\mathcal{O}_{\mathbb{K}}}}(\pi_{s^2}(p)) \simeq \mathcal{D}et_{\frac{s\mathcal{O}_{\mathbb{K}}}{s^2\mathcal{O}_{\mathbb{K}}}}(\pi_{s^2}(p)) + \mathcal{D}et_{\frac{\mathcal{O}_{\mathbb{K}}}{s\mathcal{O}_{\mathbb{K}}}}(\pi_{s^2}(p)).$$

On the other hand, we have already shown that there are canonical isomorphisms (4.10)

$$C_2(s, p) \simeq -\mathcal{D}et_{\frac{\mathcal{O}_{\mathbb{K}}}{s\mathcal{O}_{\mathbb{K}}}}(\pi_s(p)) = -\mathcal{D}et_{\frac{\mathcal{O}_{\mathbb{K}}}{s\mathcal{O}_{\mathbb{K}}}}(\pi_{s^2}(p)), \quad C_2(s^2, p) \simeq -\mathcal{D}et_{\frac{\mathcal{O}_{\mathbb{K}}}{s^2\mathcal{O}_{\mathbb{K}}}}(\pi_{s^2}(g)).$$

Again, by checking the construction as in lemma-definition 2.7, one obtains that under the isomorphisms (4.10), the canonical isomorphism $C_2(s^2, p) \simeq C_2(s, p) + C_2(s, p)$ corresponds to (4.9).

Now let $p = f$ as in the Case (iii). We know that $\mathcal{D}et_{\frac{\mathcal{O}_{\mathbb{K}}}{s\mathcal{O}_{\mathbb{K}}}}(\pi(f))$ is a graded line of degree $\nu_2(\bar{f})[k' : k]$. Therefore, using $C_2(a, b) \simeq -C_2(b, a)$ for any commuting elements $a, b \in \text{GL}(\mathbb{K})$, we obtain that Case (iii) follows from the definition of the commutativity constraints in $\mathcal{P}ic^{\mathbb{Z}}$.

Case (iv). One needs to show that $C_s(s, s) = 1$. One can easily show that there are canonical isomorphisms $C_2(s, s) \simeq \ell_0$, $C_2(s^2, s) \simeq \ell_0$, and the canonical isomorphism $C_2(s^2, s) \simeq C_2(s, s) + C_2(s, s)$ corresponds to $\ell_0 \simeq \ell_0 + \ell_0$. (We used that for the k' -space $M = k'[[t]]((s))$ we have $sM = M$, and the k' -space M induce a lattice in every 1-Tate vector space $s^n\mathcal{O}_{\mathbb{K}}/s^{n+l}\mathcal{O}_{\mathbb{K}}$, $n \in \mathbb{Z}$, $l \in \mathbb{N}$.) This case also follows. \square

5. RECIPROCITY LAWS

We will use the adèle theory on schemes. Adèles on algebraic surfaces were introduced by Parshin in [Pa2]. On arbitrary noetherian schemes they were considered by Beilinson in [Be1]. See the proof of part of results of [Be1] in [H]. A survey of adèles can be found in [O4].

We fix a perfect field k .

5.1. Weil reciprocity law. To fix the idea, let us first revisit the Weil reciprocity law. Let C be an irreducible projective curve over a field k . Let $k(C)$ be the field of rational functions on the curve C . For a closed point $p \in C$ let $\hat{\mathcal{O}}_p$ be the completion by maximal ideal m_p of the local ring \mathcal{O}_p of point $p \in C$. Let a ring K_p be the localization of the ring $\hat{\mathcal{O}}_p$ with respect to the multiplicative system $\mathcal{O}_p \setminus 0$. (If p is a smooth point, then $K_p = k(C)_p$ is the fraction field of the ring $\hat{\mathcal{O}}_p$, and $K_p = k(p)((t_p))$, $\hat{\mathcal{O}}_p = k(p)[[t_p]]$, where $k(p)$ is the residue field of the point p , t_p is a local parameter at p . For a non-smooth point $p \in C$, the ring K_p is a finite direct product of one-dimensional local fields.)

We have that K_p is a 1-Tate vector space over k , and $\hat{\mathcal{O}}_p$ is a lattice in K_p for any point $p \in C$.

For any coherent subsheaf \mathcal{F} of the constant sheaf $k(C)$ on the curve C we consider the following adèle complex $\mathcal{A}_C(\mathcal{F})$:

$$\mathbb{A}_{C,0}(\mathcal{F}) \oplus \mathbb{A}_{C,1}(\mathcal{F}) \longrightarrow \mathbb{A}_{C,01}(\mathcal{F})$$

whose cohomology groups coincide with the cohomology groups $H^*(C, \mathcal{F})$. Let us recall that

$$\mathbb{A}_{C,0}(\mathcal{F}) = k(C) \otimes_{\mathcal{O}_C} \mathcal{F}, \quad \mathbb{A}_{C,1}(\mathcal{F}) = \prod_{p \in C} \hat{\mathcal{O}}_p \otimes_{\mathcal{O}_C} \mathcal{F},$$

$$\mathbb{A}_{C,01}(\mathcal{F}) = \mathbb{A}_C = \prod'_{p \in C} K_p \otimes_{\mathcal{O}_C} \mathcal{F},$$

where \prod' denotes the restricted (adèle) product with respect to $\prod_{p \in C} \hat{\mathcal{O}}_p$. Observe

that since \mathcal{F} is a subsheaf of $k(C)$, we have

$$k(C) \otimes_{\mathcal{O}_C} \mathcal{F} = k(C), \quad K_p \otimes_{\mathcal{O}_C} \mathcal{F} = K_p.$$

The adèle ring \mathbb{A}_C is a 1-Tate vector space over k . This is because

$$\mathbb{A}_C = \varprojlim_{\mathcal{G} \subset k(C)} \varinjlim_{\mathcal{H} \subset k(C)} \mathbb{A}_{C,1}(\mathcal{H}) / \mathbb{A}_{C,1}(\mathcal{G}),$$

and $\dim_k \mathbb{A}_{C,1}(\mathcal{H}) / \mathbb{A}_{C,1}(\mathcal{G}) < \infty$ for coherent subsheaves $0 \neq \mathcal{G} \subset \mathcal{H}$ of $k(C)$. (We used that $\mathbb{A}_{C,1}(\mathcal{H}) / \mathbb{A}_{C,1}(\mathcal{G}) = \mathbb{A}_{C,1} = \bigoplus_{p \in C} \hat{\mathcal{O}}_p \otimes_{\mathcal{O}_C} (\mathcal{H}/\mathcal{G})$). For any coherent subsheaf

\mathcal{F} of $k(C)$ the space $\mathbb{A}_{C,1}(\mathcal{F})$ is a lattice in the space \mathbb{A}_C . Hence, we have that the k -space $k(C)$ is a colattice in \mathbb{A}_C , since from the adelic complex $\mathcal{A}(\mathcal{F})$ it follows

$$\dim_k k(C) \cap \mathbb{A}_{C,1}(\mathcal{F}) = \dim_k H^0(C, \mathcal{F}) < \infty,$$

$$\dim_k \mathbb{A}_C / (k(C) + \mathbb{A}_{C,1}(\mathcal{F})) = \dim_k H^1(C, \mathcal{F}) < \infty.$$

Let a p be a point of C and f, g a pair of elements of K_p^\times . If $K_p = k(p)((t_p))$, then we denote by $\{f, g\}_p$ the element from $k(p)^\times$ which is the corresponding tame symbol. If the ring K_p is isomorphic to the finite product of fields isomorphic to $k(p)((t))$, then we denote by $\{f, g\}_p$ the element from $k(p)^\times$ which is the same finite product of the corresponding tame symbols. Recall that there is the diagonal embedding $k(C) \hookrightarrow \mathbb{A}_C$.

Proposition 5.1 (Weil reciprocity law). *For any elements $f, g \in k(C)^\times$ the following product contains only finitely many nonequal to 1 terms and*

$$(5.1) \quad \prod_{p \in C} \text{Nm}_{k(p)/k} \{f, g\}_p = 1.$$

Proof. It is clear that, by proposition 4.3, we can change $\text{Nm}_{k(p)/k} \{f, g\}_p$ to $\text{Comm}(\text{Det}_{K_p})(f, g)$ for all $p \in C$ in formula (5.1). There are points $p_1, \dots, p_l \in C$ such that if $p \in C$ and $p \neq p_i$ ($1 \leq i \leq l$), then $f \hat{\mathcal{O}}_p = \hat{\mathcal{O}}_p$, $g \hat{\mathcal{O}}_p = \hat{\mathcal{O}}_p$, and hence, by proposition 4.4, $\text{Comm}(\text{Det}_{K_p})(f, g) = 1$ for points $p \neq p_i$ ($1 \leq i \leq l$).

We define the group H as the subgroup of the group $k(C)^\times$ generated by the elements f and g . We apply lemma 4.2 to the following 1-Tate k -vector spaces:

$$\mathcal{V} = \mathbb{A}_C, \quad \mathcal{V}' = \mathbb{A}_{C \setminus \{p_1, \dots, p_l\}}, \quad \mathcal{V}'' = \prod_{1 \leq i \leq l} K_{p_i}.$$

The group H preserves the lattice $\prod_{p \in C \setminus \{p_1, \dots, p_l\}} \hat{\mathcal{O}}_p$ in the space \mathcal{V}' . Therefore, by proposition 4.4, the homomorphism $\text{Det}_{\mathbb{A}_C}$ is isomorphic to the homomorphism $\text{Det}_{\mathcal{V}''}$, which is (again by lemma 4.2) isomorphic to the sum of homomorphisms

$\mathcal{D}et_{K_{p_1}}, \dots, \mathcal{D}et_{K_{p_l}}$. Since the group H preserves the colattice $k(C)$ in \mathbb{A}_C , the homomorphism $\mathcal{D}et_{\mathbb{A}_C}$ is isomorphic to the trivial one (by proposition 4.4). Now using remark 2.4 and corollaries 2.5 and 2.6 we obtain formula (5.1). \square

Remark 5.1. To obtain the triviality of homomorphism $\mathcal{D}et_{\mathbb{A}_C} : k(C)^\times \rightarrow \mathcal{P}ic^{\mathbb{Z}}$ in an explicit way, one has to use the following canonical isomorphism for any $g \in k(C)^\times$:

$$(5.2) \quad \mathcal{D}et_{\mathbb{A}_C}(g) \simeq \det(H^*(\mathcal{A}_C(g\mathcal{O}_C))) \otimes \det(H^*(\mathcal{A}_C(\mathcal{O}_C)))^{-1},$$

where for any coherent sheaf \mathcal{F} on C

$$\begin{aligned} \det(H^*(\mathcal{A}_C(\mathcal{F}))) &:= \det(H^0(\mathcal{A}_C(\mathcal{F}))) \otimes \det(H^1(\mathcal{A}_C(\mathcal{F})))^{-1} \\ &\simeq \det(H^0(C, \mathcal{F})) \otimes \det(H^1(C, \mathcal{F}))^{-1}. \end{aligned}$$

(Formula (5.2) easily follows from adèle complexes and formula (4.5) if we change in formula (4.5) the lattices \mathcal{L} and $g\mathcal{L}$ in \mathbb{A}_C to any two lattices coming from nonzero coherent subsheaves $\mathcal{G} \subset \mathcal{H}$ of $k(C)$, and change correspondingly in formula (5.2) the sheaves \mathcal{O} and $g\mathcal{O}$ to the sheaves $\mathcal{G} \subset \mathcal{H}$.) Now the homomorphism $\mathcal{D}et_{\mathbb{A}_C}$ is isomorphic to the trivial one by formula (5.2) and the fact that multiplication on an element $g \in k(C)^*$ gives a canonical isomorphism between adèle complexes $\mathcal{A}_C(\mathcal{O}_C)$ and $\mathcal{A}_C(g\mathcal{O}_C)$, which induce the canonical isomorphism between $\det(H^*(\mathcal{A}_C(\mathcal{O}_C)))$ and $\det(H^*(\mathcal{A}_C(g\mathcal{O}_C)))$.

5.2. Parshin reciprocity laws. Let X be an algebraic surface over the field k . We assume, for simplicity, that X is a smooth connected surface.

We consider pairs $x \in C$, where C are irreducible curves on X and x are closed points on C . For every such pair one can define the ring $K_{x,C}$, which will be a finite product of two-dimensional local fields, as follows. Assume that the curve C on X has the following formal branches $\mathbf{C}_1, \dots, \mathbf{C}_n$ at the point $x \in C$, i.e.

$$C|_{\text{Spec}\hat{\mathcal{O}}_x} = \bigcup_{1 \leq i \leq n} \mathbf{C}_i,$$

where $\hat{\mathcal{O}}_x$ is the completion of the local ring \mathcal{O}_x of a point $x \in X$, and \mathbf{C}_i is irreducible in $\text{Spec}\hat{\mathcal{O}}_x$ for any $1 \leq i \leq n$. (Since we assumed X is smooth, $\hat{\mathcal{O}}_x \simeq k(x)[[t_1, t_2]]$.) Now every \mathbf{C}_i defines a discrete valuation on the fraction field $\text{Frac}\hat{\mathcal{O}}_x$. We define a two-dimensional local field K_{x,\mathbf{C}_i} as the completion of the field $\text{Frac}\hat{\mathcal{O}}_x$ with respect to this discrete valuation, and let $\hat{\mathcal{O}}_{x,\mathbf{C}_i}$ be the valuation ring. Then we define

$$K_{x,C} := \bigoplus_{1 \leq i \leq n} K_{x,\mathbf{C}_i}, \quad \hat{\mathcal{O}}_{x,C} := \bigoplus_{1 \leq i \leq n} \hat{\mathcal{O}}_{x,\mathbf{C}_i},$$

Observe that if $x \in C$ is a smooth point, then $\hat{\mathcal{O}}_{x,C} \simeq k(x)((t))[s]$ and $K_{x,C} \simeq k(x)((t))(s)$. It is clear that the ring $\hat{\mathcal{O}}_x$ diagonally embeds into the ring $K_{x,C}$.

Let us also define $B_x \subset K_{x,C}$ as $\varinjlim_{n>0} s_C^{-n} \hat{\mathcal{O}}_x$, where a local equation $s_C = 0$ determines C on some open $X \supset V \ni x$. It is clear that the subring B_x of $K_{x,C}$ does not depend on the choice of such s_C when $V \ni x$. If $x \in C$ is a smooth point, and $K_{x,C} = k(x)((t))(s_C)$, where $s_C = 0$ is a local equation of the curve C on X near the point x and $t = 0$ defines a transversal curve locally on X near x , then $B_x = k(x)[[t]](s_C)$.

Any ring $K_{x,C}$ is a 2-Tate vector space over the field $k(x)$ (and therefore over the field k), and the ring $\hat{\mathcal{O}}_{x,C}$ is a lattice in $K_{x,C}$.

Let f, g, h be from $K_{x,C}^\times$ such that $f = \bigoplus_{1 \leq i \leq n} f_i$, $g = \bigoplus_{1 \leq i \leq n} g_i$, $h = \bigoplus_{1 \leq i \leq n} h_i$. Then we define the following element from $k(x)^\times$ as

$$(5.3) \quad \{f, g, h\}_{x,C} := \prod_{1 \leq i \leq n} \{f_i, g_i, h_i\}_{x,C_i},$$

where $\{f_i, g_i, h_i\}_{x,C_i}$ is the two-dimensional tame symbol associated to the two-dimensional local field K_{x,C_i} (cf. §4.1).

Fix a point $x \in X$. For any free finitely generated $\hat{\mathcal{O}}_x$ -module subsheaf \mathcal{F} of the constant sheaf $\text{Frac}\hat{\mathcal{O}}_x$ on the scheme $\text{Spec}\hat{\mathcal{O}}_x$ we consider the following adèle complex $\mathcal{A}_{X,x}(\mathcal{F})$:

$$\mathbb{A}_{X,x,0}(\mathcal{F}) \oplus \mathbb{A}_{X,x,1}(\mathcal{F}) \longrightarrow \mathbb{A}_{X,x,01}(\mathcal{F}).$$

This is the adèle complex on the 1-dimensional scheme $U_x := \text{Spec}\hat{\mathcal{O}}_x \setminus x$ for the sheaf $\mathcal{F}|_{U_x}$, and, hence, the cohomology groups of this complex coincide with the cohomology groups $H^*(U_x, \mathcal{F}|_{U_x})$. By definition, we have

$$\mathbb{A}_{X,x,0}(\mathcal{F}) = \text{Frac}\hat{\mathcal{O}}_x, \quad \mathbb{A}_{X,x,1}(\mathcal{F}) = \prod_{\mathbf{C} \ni x} \hat{\mathcal{O}}_{x,\mathbf{C}} \otimes_{\hat{\mathcal{O}}_x} \mathcal{F}, \quad \mathbb{A}_{X,x,01}(\mathcal{F}) = \mathbb{A}_{X,x} = \prod'_{\mathbf{C} \ni x} K_{p,\mathbf{C}},$$

where the product is taken over all prime ideals \mathbf{C} of height 1 of the ring $\hat{\mathcal{O}}_x$, and \prod' denotes the restricted (adèle) product with respect to $\prod_{\mathbf{C} \ni x} \hat{\mathcal{O}}_{x,\mathbf{C}}$.

Observe that the adèle ring $\mathbb{A}_{X,x}$ is a 2-Tate vector space over the field $k(x)$. This is because

$$\mathbb{A}_{X,x} = \varprojlim_{\mathcal{G} \subset \text{Frac}\hat{\mathcal{O}}_x} \varinjlim_{\mathcal{H} \subset \text{Frac}\hat{\mathcal{O}}_x} \mathbb{A}_{X,x,1}(\mathcal{H}) / \mathbb{A}_{X,x,1}(\mathcal{G}),$$

and $\mathbb{A}_{X,x,1}(\mathcal{H}) / \mathbb{A}_{X,x,1}(\mathcal{G})$ is a 1-Tate vector space for free $\hat{\mathcal{O}}_x$ -module subsheaves $0 \neq \mathcal{G} \subset \mathcal{H}$ of $\text{Frac}\hat{\mathcal{O}}_x$. (We used that $\mathbb{A}_{X,x,1}(\mathcal{H}) / \mathbb{A}_{X,x,1}(\mathcal{G}) = \bigoplus_{\mathbf{C} \ni x} \hat{\mathcal{O}}_{x,\mathbf{C}} \otimes_{\hat{\mathcal{O}}_x} \mathcal{H}/\mathcal{G}$.)

For any free finitely generated $\hat{\mathcal{O}}_x$ -module subsheaf \mathcal{F} of $\text{Frac}\hat{\mathcal{O}}_x$ the space $\mathbb{A}_{X,x,1}(\mathcal{F})$ is a lattice in the space $\mathbb{A}_{X,x}$.

From [O2, prop. 8] it follows that $k(x)$ -vector spaces $H^0(\mathcal{A}_{X,x}(\mathcal{F}))$ and $H^1(\mathcal{A}_{X,x}(\mathcal{F}))$ are 1-Tate vector spaces. Indeed, since x is a smooth point of X ,

$$H^0(\mathcal{A}_{X,x}(\mathcal{F})) = H^0(U_x, \mathcal{F}|_{U_x}) = \mathcal{F}$$

is a projective limit of finite-dimensional $k(x)$ -vector spaces $\mathcal{F}/m_x^n \mathcal{F}$ (m_x is the maximal ideal of the ring $\hat{\mathcal{O}}_x$), and

$$H^1(\mathcal{A}_{X,x}(\mathcal{F})) = H^1(U_x, \mathcal{F}|_{U_x}) = \varinjlim_{n>0} \text{Ext}_{\hat{\mathcal{O}}_x}^2(\hat{\mathcal{O}}_x/m_x^n, \mathcal{F}),$$

where for any $n > 0$ the space $\text{Ext}_{\hat{\mathcal{O}}_x}^2(\hat{\mathcal{O}}_x/m_x^n, \mathcal{F})$ is a finite-dimensional over the field $k(x)$ vector space (see, for example, [O2, lemma 6]).

Fix an irreducible projective curve C on X . For any invertible \mathcal{O}_X -subsheaf \mathcal{F} of the constant sheaf $k(X)$ on X we consider the following adèle complex $\mathcal{A}_{X,C}(\mathcal{F})$

$$\mathbb{A}_{X,C,0}(\mathcal{F}) \oplus \mathbb{A}_{X,C,1}(\mathcal{F}) \longrightarrow \mathbb{A}_{X,C,01}(\mathcal{F}).$$

where

$$\mathbb{A}_{X,C,0}(\mathcal{F}) := K_C, \quad \mathbb{A}_{X,C,01}(\mathcal{F}) := \mathbb{A}_{X,C} = \mathbb{A}_C((s_C)),$$

$$(5.4) \quad \mathbb{A}_{X,C,1}(\mathcal{F}) := \left(\prod_{x \in C} B_x \otimes_{\mathcal{O}_X} \mathcal{F} \right) \cap \mathbb{A}_{X,C}.$$

Here K_C is the completion of the field $k(X)$ with respect to the discrete valuation given by the curve C on X . (If $s_C = 0$ is a local equation of the curve C on some open subset V of X such that $V \cap C \neq \emptyset$, then $K_C = k(C)((s_C))$.) The ring $\mathbb{A}_{X,C}$ is a subring of $\prod_{x \in C} K_{x,C}$, and does not depend on the choice of s_C . The intersection (5.4) is taken in the ring $\prod_{x \in C} K_{x,C}$.

We note that from [O2, § 5.1] it follows that the complex $\mathcal{A}_{X,C}(\mathcal{F})$ coincides with the following complex

$$\varinjlim_n \varprojlim_{m > n} \mathcal{A}_{(C, \mathcal{O}_X/J_C^{m-n})}(\mathcal{F} \otimes_{\mathcal{O}_X} J_C^m/J_C^n).$$

Here J_C is the ideal sheaf of the curve C on X , $(C, \mathcal{O}_X/J_C^{m-n})$ is a 1-dimensional scheme which has the topological space C and the structure sheaf \mathcal{O}_X/J_C^{m-n} , and $\mathcal{A}_{(C, \mathcal{O}_X/J_C^{m-n})}(\mathcal{F} \otimes_{\mathcal{O}_X} J_C^m/J_C^n)$ is the adèle complex of the coherent sheaf $\mathcal{F} \otimes_{\mathcal{O}_X} J_C^m/J_C^n$ on the scheme $(C, \mathcal{O}_X/J_C^{m-n})$. Hence and from the proof of proposition 12 from [O2] we obtain that

$$H^*(\mathcal{A}_{X,C}(\mathcal{F})) = \varinjlim_n \varprojlim_{m > n} H^*(C, \mathcal{F} \otimes_{\mathcal{O}_X} J_C^m/J_C^n),$$

where for $i = 0$ and $i = 1$ we have $\dim_k H^i(C, \mathcal{F} \otimes_{\mathcal{O}_X} J_C^m/J_C^n) < \infty$. For $i = 0$ and $i = 1$ the k -vector space $H^i(\mathcal{A}_{X,C}(\mathcal{F}))$ has the natural topology of inductive and projective limits. It is not difficult to see that the space $H^0(\mathcal{A}_{X,C}(\mathcal{F}))$ is a locally linearly compact k -vector space, i.e. it is a 1-Tate vector space. But the space $H^1(\mathcal{A}_{X,C}(\mathcal{F}))$ is not a Hausdorff space in this topology. Let $\tilde{H}^1(\mathcal{A}_{X,C}(\mathcal{F}))$ be the quotient space of $H^1(\mathcal{A}_{X,C}(\mathcal{F}))$ by the closure of zero. Then the space $\tilde{H}^1(\mathcal{A}_{X,C}(\mathcal{F}))$ is a locally linearly compact k -vector space, i.e. a 1-Tate vector space.

We note that for any invertible subsheaves $0 \neq \mathcal{G} \subset \mathcal{H}$ of $k(X)$ we have that the space $B_x \otimes_{\hat{\mathcal{O}}_x} (\mathcal{H}/\mathcal{G})$ is a 1-Tate vector space, which is equal to zero for almost all points $x \in C$. Hence, we obtain that the space

$$\mathbb{A}_{X,C,1}(\mathcal{H})/\mathbb{A}_{X,C,1}(\mathcal{G}) = \bigoplus_{x \in C} B_x \otimes_{\hat{\mathcal{O}}_x} (\mathcal{H}/\mathcal{G})$$

is a 1-Tate vector space.

For any point $x \in X$, we define a ring K_x as the localization of the ring $\hat{\mathcal{O}}_x$ with respect to the multiplicative system $\mathcal{O}_x \setminus 0$. (We note that inside of the field $\text{Frac} \hat{\mathcal{O}}_x$ the ring K_x is defined as the product of two subrings: $\hat{\mathcal{O}}_x$ and $k(X)$.)

For any pair $x \in C$ (where C is an irreducible curve on X and $x \in C$ is a closed point), we have the natural embeddings $k(X) \hookrightarrow K_x$, $k(X) \hookrightarrow K_C$ (recall that K_C is the completion of the field $k(X)$ with respect to the discrete valuation given by the curve C). In addition, there are the natural embeddings $K_x, K_C \hookrightarrow K_{x,C}$. Therefore, we obtain

$$k(X) \hookrightarrow K_x \hookrightarrow \mathbb{A}_{X,x} \quad , \quad k(X) \hookrightarrow K_C \hookrightarrow \mathbb{A}_{X,C}.$$

Theorem 5.2 (Parshin reciprocity laws). (1) Fix a point $x \in X$. Consider elements f, g, h of the group K_x^\times of invertible elements of the ring K_x . Then

the following product in $k(x)^\times$ contains only finitely many non-equal to 1 terms and

$$(5.5) \quad \prod_{C \ni x} \{f, g, h\}_{x, C} = 1.$$

- (2) Fix a projective irreducible curve C on X . Let elements f, g, h be from the group K_C^\times . Then the following product in k^\times contains only finitely many non-equal to 1 terms and

$$(5.6) \quad \prod_{x \in C} \text{Nm}_{k(x)/k} \{f, g, h\}_{x, C} = 1.$$

Proof. We first prove formula (5.5). By theorem 4.7, for any $f, g, h \in K_{x, \mathbf{C}}^\times$ we have

$$(5.7) \quad \{f, g, h\}_{x, \mathbf{C}} = C_3^{\mathcal{D}et_{x, \mathbf{C}}} (f, g, h)$$

for all prime ideals \mathbf{C} of height 1 of the ring $\hat{\mathcal{O}}_x$, where the central extension $\mathcal{D}et_{x, \mathbf{C}}$ of the group $K_{x, \mathbf{C}}^\times$ by the Picard groupoid $\mathcal{P}ic^{\mathbb{Z}}$ is constructed by formula (4.6) from the 2-Tate vector space $K_{x, \mathbf{C}}$ over the field $k(x)$ and the lattice $\hat{\mathcal{O}}_{x, \mathbf{C}}$ as in section 4.3. We note that for almost all prime ideals \mathbf{C} of height 1 of ring $\hat{\mathcal{O}}_x$, and for any elements f, g, h from the group $\text{Frac} \hat{\mathcal{O}}_x^\times$, we have $f \mathcal{O}_{x, \mathbf{C}} = \mathcal{O}_{x, \mathbf{C}}$, $g \mathcal{O}_{x, \mathbf{C}} = \mathcal{O}_{x, \mathbf{C}}$, and $h \mathcal{O}_{x, \mathbf{C}} = \mathcal{O}_{x, \mathbf{C}}$. Then by lemma 4.6 and corollary 2.12, for almost all prime ideals \mathbf{C} of height 1 of ring $\hat{\mathcal{O}}_x$ we have $C_3^{\mathcal{D}et_{x, \mathbf{C}}} (f, g, h) = 1$.

We will prove that the central extension $\mathcal{D}et_x$ of $\text{Frac} \hat{\mathcal{O}}_x^\times (\subset \text{GL}(\mathbb{A}_{X, x}))$ by $\mathcal{P}ic^{\mathbb{Z}}$ constructed by the 2-Tate vector space $\mathbb{A}_{X, x}$ and the lattice $\mathbb{A}_{X, x, 1}(\hat{\mathcal{O}}_x)$ using formula (4.6) can be trivialized in an explicit way. Observe that for any $d \in \text{Frac} \hat{\mathcal{O}}_x^\times$, there is a canonical isomorphism of $\mathcal{P}ic^{\mathbb{Z}}$ -torsors:

$$(5.8) \quad \mathcal{D}et(\mathbb{A}_{X, x, 1}(\hat{\mathcal{O}}_x) \mid \mathbb{A}_{X, x, 1}(d\hat{\mathcal{O}}_x)) \simeq \mathcal{D}et(H^*(\mathcal{A}_{X, x}(d\hat{\mathcal{O}}_x))) - \mathcal{D}et(H^*(\mathcal{A}_{X, x}(\hat{\mathcal{O}}_x))),$$

where for any free subsheaf \mathcal{F} of $\text{Frac} \hat{\mathcal{O}}_x$ on the scheme $\text{Spec} \hat{\mathcal{O}}_x$

$$\mathcal{D}et(H^*(\mathcal{A}_{X, x}(\mathcal{F}))) := \mathcal{D}et(H^0(\mathcal{A}_{X, x}(\mathcal{F}))) - \mathcal{D}et(H^1(\mathcal{A}_{X, x}(\mathcal{F}))).$$

Indeed, isomorphism (5.8) follows from proposition 3.2 applied to the long exact sequence (decomposed into the short exact sequences) associated with the following exact sequence of complexes of length 2 for any nonzero free subsheaves $\mathcal{G} \subset \mathcal{H}$ of $\text{Frac} \hat{\mathcal{O}}_x$ on the scheme $\text{Spec} \hat{\mathcal{O}}_x$:

$$0 \longrightarrow \mathcal{A}_{X, x}(\mathcal{G}) \longrightarrow \mathcal{A}_{X, x}(\mathcal{H}) \longrightarrow \mathcal{A}_{X, x, 1}(\mathcal{H}) / \mathbb{A}_{X, x, 1}(\mathcal{G}) \longrightarrow 0,$$

where the last complex consists only of the group placed in degree zero. Now we have

$$(5.9) \quad \mathcal{D}et(H^*(\mathcal{A}_{X, x}(d\hat{\mathcal{O}}_x))) - \mathcal{D}et(H^*(\mathcal{A}_{X, x}(\hat{\mathcal{O}}_x))) \\ \simeq \text{Hom}_{\mathcal{P}ic^{\mathbb{Z}}}(\mathcal{D}et(H^*(\mathcal{A}_{X, x}(\hat{\mathcal{O}}_x))), \mathcal{D}et(H^*(\mathcal{A}_{X, x}(d\hat{\mathcal{O}}_x))).$$

The multiplication by the element $d \in \text{Frac} \hat{\mathcal{O}}_x^\times$ between adèle complexes $\mathcal{A}_{X, x}(\hat{\mathcal{O}}_x)$ and $\mathcal{A}_{X, x}(d\hat{\mathcal{O}}_x)$ gives a natural isomorphism of $\mathcal{P}ic^{\mathbb{Z}}$ -torsor from formula (5.9) to the trivial torsor $\mathcal{P}ic^{\mathbb{Z}}$.

Let H be the subgroup of $\text{Frac} \hat{\mathcal{O}}_x^\times$ generated by the elements $f, g, h \in \text{Frac} \hat{\mathcal{O}}_x^\times$. Now we proceed as the proof of Weil reciprocity law (see proposition 5.1), with the

help of lemma 4.5, lemma 4.6, and corollary 2.13. Then we obtain the following equality:

$$\prod_{\mathbf{C} \ni x} \{f, g, h\}_{x, \mathbf{C}} = 1.$$

Formula (5.5) follows from the last formula, since if a prime ideal \mathbf{C} of height 1 in $\hat{\mathcal{O}}_x$ is not a formal branch at x of some irreducible curve C on X , then for any element $d \in K_x^\times$ we have $d\mathcal{O}_{x, \mathbf{C}} = \mathcal{O}_{x, \mathbf{C}}$. Hence, by formula (5.7), $\{f, g, h\}_{x, \mathbf{C}} = 1$ for such \mathbf{C} and any $f, g, h \in K_x^\times$.

Next we will prove formula (5.6). We construct the central extension $\mathcal{D}et'_{x, C}$ of the group $k(X)^\times$ by the Picard groupoid $\mathcal{P}ic^{\mathbb{Z}}$ in the following way. We fix a point $x \in C$, and associate with the rings $B_x \subset K_{x, C}$ and with an element $d \in k(X)^\times$ the following $\mathcal{P}ic^{\mathbb{Z}}$ -torsor:

$$(5.10) \quad \mathcal{D}et(B_x \mid dB_x) := \mathcal{D}et\left(\frac{dB_x}{B_x \cap dB_x}\right) - \mathcal{D}et\left(\frac{B_x}{B_x \cap dB_x}\right).$$

(We used that $B_x/B_x \cap dB_x$ is a 1-Tate vector space over the field k .) By the formula which is analogous to formula (4.7) we obtain that the central extension $\mathcal{D}et'_{x, C}$ is well defined. In a similar way we define the central extensions $\mathcal{D}et'_C$ and $\mathcal{D}et'_{C \setminus \{x_1, \dots, x_l\}}$ starting from the rings $\mathbb{A}_{X, C, 1}(\mathcal{O}_X) \subset \mathbb{A}_{X, C}$ and $\mathbb{A}_{X, C \setminus \{x_1, \dots, x_l\}, 1}(\mathcal{O}_X) \subset \mathbb{A}_{X, C \setminus \{x_1, \dots, x_l\}}$, where x_1, \dots, x_l are some points on the curve C .

Let the group H be generated in the group $k(X)^\times$ by the elements $f, g, h \in k(X)^\times$. For almost all points x of the curve C we have that the group H preserves the subring B_x . Therefore from formula (5.10) we obtain that the central extension $\mathcal{D}et'_{x, C}$ is isomorphic to the trivial one for almost all points x of the curve C . Therefore for almost all points x of the curve C we have $C_3^{\mathcal{D}et'_{x, C}}(f, g, h) = 1$.

We will prove that the central extension $\mathcal{D}et'_{x, C}$ is inverse (or dual) to the central extension $\mathcal{D}et_{x, C}$, where the last central extension is constructed by formula (4.6) from the lattice $\mathcal{O}_{x, C}$ in the 2-Tate vector space $K_{x, C}$. For any free subsheaf \mathcal{F} of the constant sheaf $\text{Frac}\hat{\mathcal{O}}_x$ on the scheme $\text{Spec}\hat{\mathcal{O}}_x$ there is the following complex $\mathcal{A}_{X, C, x}(\mathcal{F})$:

$$(B_x \otimes_{\hat{\mathcal{O}}_x} \mathcal{F}) \oplus (\hat{\mathcal{O}}_{x, C} \otimes_{\hat{\mathcal{O}}_x} \mathcal{F}) \longrightarrow K_{x, C}.$$

We have canonically that $H^*(\mathcal{A}_{X, C, x}(\mathcal{F})) = H^*(U_x, \mathcal{F} \mid_{U_x})$, where we recall $U_x = \text{Spec}\hat{\mathcal{O}}_x \setminus x$ (see the proof of proposition 13 from [O2]). Therefore the cohomology groups of complex $\mathcal{A}_{X, C, x}(\mathcal{F})$ are 1-Tate vector spaces. Hence, there is a canonical isomorphism between the following $\mathcal{P}ic^{\mathbb{Z}}$ -torsors for any $d \in k(X)^\times$:

$$\begin{aligned} & \mathcal{D}et(B_x \mid dB_x) + \mathcal{D}et(\hat{\mathcal{O}}_{x, C} \mid d\hat{\mathcal{O}}_{x, C}) \quad \text{and} \\ & \text{Hom}_{\mathcal{P}ic^{\mathbb{Z}}}(\mathcal{D}et(H^*(\mathcal{A}_{X, C, x}(\hat{\mathcal{O}}_x))), \mathcal{D}et(H^*(\mathcal{A}_{X, C, x}(d\hat{\mathcal{O}}_x)))). \end{aligned}$$

Now the multiplication by the element d of adèle complexes gives a natural isomorphism from the last $\mathcal{P}ic^{\mathbb{Z}}$ -torsor to the trivial one. Hence from corollary 2.13 we have that

$$C_3^{\mathcal{D}et'_{x, C}}(f, g, h) = C_3^{\mathcal{D}et_{x, C}}(f, g, h)^{-1} = \text{Nm}_{k(x)/k}\{f, g, h\}_{x, C}^{-1}$$

for $f, g, h \in k(X)^\times$.

Now the proof of formula (5.6) for elements $f, g, h \in k(X)^\times$ follows by the same method as in the proof of formula (5.5), but we have to use the adèle ring $\mathbb{A}_{X, C}$

instead of the ring $\mathbb{A}_{X,x}$, and to use the central extension $\mathcal{D}et'_C$ instead of the central extension $\mathcal{D}et_x$. We need only to prove that the central extension $\mathcal{D}et'_C$ constructed by the analog of formula (5.10) from the rings $\mathbb{A}_{X,C,1}(\mathcal{O}_X) \subset \mathbb{A}_{X,C}$ is isomorphic to the trivial central extension. This follows if we consider the following $\mathcal{P}ic^{\mathbb{Z}}$ -torsors for $d \in k(X)^\times$

$$(5.11) \quad \text{Hom}_{\mathcal{P}ic^{\mathbb{Z}}}(\mathcal{D}et(H^*(\mathcal{A}_{X,C}(\mathcal{O}_X))), \mathcal{D}et(H^*(\mathcal{A}_{X,C}(d\mathcal{O}_X)))),$$

where

$$\mathcal{D}et(H^*(\mathcal{A}_{X,C}(d\mathcal{O}_X))) := \mathcal{D}et(H^0(\mathcal{A}_{X,C}(d\mathcal{O}_X)) - \mathcal{D}et(\tilde{H}^1(\mathcal{A}_{X,C}(d\mathcal{O}_X))).$$

Multiplication by $d \in k(X)^\times$ of adèle complexes gives the triviality of the $\mathcal{P}ic^{\mathbb{Z}}$ -torsor (5.11). (See analogous reasonings earlier in the proof of this theorem.)

To obtain formula (5.6) for elements $f, g, h \in K_C^\times$ we have to use that the field $k(X)$ is dense in the field K_C . Therefore for any element $f \in K_C^\times$ there is an element $\tilde{f} \in k(X)^\times$ such that $f = \tilde{f}m$, where the element m is from the subgroup $1 + m_C^n$ of the group K_C^\times for some $n \geq 1$, and m_C is the maximal ideal of the valuation ring of discrete valuation field K_C . Then from formula (4.3) we have that $\{m, g, h\}_{x,C} = 1$ for any point $x \in C$, and any formal branch \mathbf{C} of the curve C at point x . Hence, from the tri-multiplicativity of the two-dimensional tame symbol we obtain that

$$\{f, g, h\}_{x,C} = \{\tilde{f}, g, h\}_{x,C}.$$

Applying successively the same procedure to elements $g, h \in k(C)^\times$ we obtain

$$\{f, g, h\}_{x,C} = \{\tilde{f}, \tilde{g}, \tilde{h}\}_{x,C},$$

where $\tilde{f}, \tilde{g}, \tilde{h} \in k(X)^\times$, and any point $x \in C$, and \mathbf{C} is any formal branch of the curve C at point x . □

Remark 5.2. For the proof of Parshin reciprocity laws we used "semilocal" adèle complexes of length 2 connected with either points or irreducible curves on an algebraic surface. But for the formulation of these reciprocity laws we used the rings K_x and K_C which appear from the "global" adèle complex of length 3 on an algebraic surface. It would be interesting to find direct connections between the "global" adèle complex and "semilocal" adèle complexes of an algebraic surface.

Remark 5.3. We have a symmetric monoidal functor from the Picard torsor $\mathcal{P}ic^{\mathbb{Z}}$ to the Picard groupoid \mathbb{Z} which sends every graded line to its grading element from \mathbb{Z} , where \mathbb{Z} is considered as the groupoid with objects equal to \mathbb{Z} and morphisms equal to identities morphisms. Under this functor a central extension of a group G by a $\mathcal{P}ic^{\mathbb{Z}}$ -torsor goes to the usual central of the group G by the group \mathbb{Z} . In this way the map $\nu_{\mathbb{K}}$ for a two-dimensional local field \mathbb{K} was obtained as the commutator of elements in this central extension in [O2]. Also in [O2] the reciprocity laws for the map $\nu_{\mathbb{K}}$ were proved by the adèle complexes on an algebraic surface.

REFERENCES

- [ACK] Arbarello E., De Concini C., Kac V. G. *The infinite wedge representation and the reciprocity law for algebraic curves*, Comm. Math. Phys., **117** (1988), 1-36.
- [AK] Arkhipov S. and Kremnizer K. *2-gerbes and 2-Tate spaces*, e-print arXiv:0708.4401.
- [BN] Baez J., Neuchl M. *Higher-dimensional algebra I: Braided monoidal categories*, Adv. Math. **121** (1996), 196-244.

- [Be1] Beilinson A.A. *Residues and adeles*, Funct. Anal. Pril., **14** (1980), no. 1, 44-45; English transl. in Func. Anal. Appl., **14**, no. 1 (1980), 34-35.
- [Be2] Beilinson A. A. *How to glue perverse sheaves*, in K-theory, Arithmetic, Geometry, Semin., Moscow Univ. 1984-86, Lecture Notes in Mathematics 1289, 42-51 (1987).
- [BBE] Beilinson A., Bloch S., Esnault H. *ε -factors for Gauss-Manin determinants.*, Mosc. Math. J. **2**, No. 3, 477-532 (2002).
- [Br1] Breen, L. *Théorie de Schreier supérieure*, Annales scientifiques de l'école Normale Supérieure, Sér. 4, 25 no. 5, 1992.
- [Br2] Breen L. *On the classification of 2-gerbes and 2-stacks*, Astérisque. 225. Paris: Société mathématique de France (1994).
- [Br3] Breen L. *Monoidal categories and multiextensions*, Compositio Mathematica **117**: 295-335, 1999.
- [Del1] Deligne P. *La formule de dualité globale*, in SGA 4, tome 3, Exposé XVIII, Lecture Notes in Mathematics, vol. 305, 481-587. Springer-Verlag, 1973.
- [Del2] Deligne P. *Le symbole modéré*, Publ. Math. IHES **73**(1991), 147-181.
- [Dr] Drinfeld V. *Infinite-dimensional vector bundles in algebraic geometry: an introduction*, The unity of mathematics, In honor of the ninetieth birthday of I. M. Gelfand, Papers from the conference held in Cambridge, MA, USA, August 31-September 4, 2003, Progr. Math. 244 (2006), 263-304.
- [FZ] Frenkel, E., Zhu X. *Gerbal representation of double loop groups*, e-print arXiv:0810.1487.
- [H] Huber A. *On the Parshin-Beilinson Adeles for Schemes*, Abh. Math. Sem. Univ. Hamburg, 61 (1991), 249-273.
- [Kap] Kapranov M. *Semiinfinite symmetric powers*, preprint arXiv:math.QA/0107089.
- [KV] Kapranov M. M., Voevodsky V. A. *2-categories and Zamolodchikov tetrahedra equations*, Proc. Symp. Pure Math., v. 56 (1994), pt.2, Amer. Math. Soc., Providence RI 1994, 177-259.
- [K] Kato K., *Milnor K-theory and the Chow group of zero cycles*, Applications of algebraic K-theory to algebraic geometry and number theory, Proc. AMS-IMS-SIAM Joint Summer Res. Conf., Boulder/Colo. 1983, Part I, Contemp. Math. 55, 241-253 (1986).
- [O1] Osipov D. *To the multidimensional tame symbol*, preprint 03-13 of the Humboldt University of Berlin (2003), available at <http://edoc.hu-berlin.de/docviews/abstract.php?id=26204>
- [O2] Osipov D.V. *Central extensions and reciprocity laws on algebraic surfaces*, (Russian) Mat. Sb. 196:10 (2005), 111-136; translation in Sb. Math. 196:10 (2005), 1503-1527; e-print arXiv: math.NT/0501155.
- [O3] Osipov D. *Adeles on n -dimensional schemes and categories C_n* , International Journal of Mathematics, vol. 18, no. 3 (2007), 269-279; e-print arXiv:math.AG/0509189.
- [O4] Osipov D.V. *n -dimensional local fields and adeles on n -dimensional schemes*, Surveys in Contemporary Mathematics, Edited by N. Young, Y. Choi; London Mathematical Society Lecture Note Series, No. 347 (2007), pp. 131-164, Cambridge University Press; e-print math.AG/0508205.
- [Pa1] Parshin A. N. *Class fields and algebraic K-theory*. Uspekhi Mat. Nauk **30** (1975), 253-254 (in russian), English translation in Russian Math. Surveys.
- [Pa2] Parshin A. N. *On the arithmetic of two-dimensional schemes, I. Repartitions and residues*, Izv. Akad. Nauk SSSR Ser. Mat. **40** (1976), no. 4, 736-773; English transl. in Math. USSR Izv. **10** (1976), no. 4, 695-729.
- [Pa3] Parshin A. N. *Local class field theory*, Trudy Matem. Inst. Steklov. 165 (1984) (in russian), English translation: Proceedings of the Steklov Institute of Mathematics 1985, Issue 3, 157-185.
- [Q] Quillen D. *Higher algebraic K-theory, I*, Lecture Notes in Mathematics **341**, pp. 85-147, Springer 1973.
- [S] Serre J.-P. *Algebraic groups and class fields*, Transl. of the French edition. (English) Graduate Texts in Mathematics, 117. New York etc.: Springer-Verlag, (1988).
- [T] Tate J. *Residues of differentials on curves*, Ann. Sci. Éc. Norm. Supér., 4 serie, t.1, 1968, p.149-159.
- [Z] Zhu X. *The 2-group of linear auto-equivalences of an abelian category and its Lie 2-algebra*, e-print arXiv:0910.5699.

Denis Osipov,
Steklov Mathematical Institute, Gubkina str. 8, Moscow, 119991, Russia
E-mail address: *d_osipov@mi.ras.ru*

Xinwen Zhu,
Department of Mathematics, Harvard University, one Oxford Street, Cambridge,
MA 02138, USA
E-mail address: *xinwenz@math.harvard.edu*