

# Evidence for the baryonic decay $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$

J. P. Lees, V. Poireau, and V. Tisserand

*Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP),  
Université de Savoie, CNRS/IN2P3, F-74941 Annecy-Le-Vieux, France*

E. Grauges

*Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain*

A. Palano<sup>ab</sup>

*INFN Sezione di Bari<sup>a</sup>; Dipartimento di Fisica, Università di Bari<sup>b</sup>, I-70126 Bari, Italy*

G. Eigen and B. Stugu

*University of Bergen, Institute of Physics, N-5007 Bergen, Norway*

D. N. Brown, L. T. Kerth, Yu. G. Kolomensky, M. J. Lee, and G. Lynch

*Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA*

H. Koch and T. Schroeder

*Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany*

C. Hearty, T. S. Mattison, J. A. McKenna, and R. Y. So

*University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1*

A. Khan

*Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom*

V. E. Blinov<sup>ac</sup>, A. R. Buzykaev<sup>a</sup>, V. P. Druzhinin<sup>ab</sup>, V. B. Golubev<sup>ab</sup>, E. A. Kravchenko<sup>ab</sup>, A. P. Onuchin<sup>ac</sup>,  
S. I. Serednyakov<sup>ab</sup>, Yu. I. Skovpen<sup>ab</sup>, E. P. Solodov<sup>ab</sup>, K. Yu. Todyshev<sup>ab</sup>, and A. N. Yushkov<sup>a</sup>

*Budker Institute of Nuclear Physics SB RAS, Novosibirsk 630090<sup>a</sup>,*

*Novosibirsk State University, Novosibirsk 630090<sup>b</sup>,*

*Novosibirsk State Technical University, Novosibirsk 630092<sup>c</sup>, Russia*

D. Kirkby, A. J. Lankford, and M. Mandelkern

*University of California at Irvine, Irvine, California 92697, USA*

B. Dey, J. W. Gary, O. Long, and G. M. Vitug

*University of California at Riverside, Riverside, California 92521, USA*

C. Campagnari, M. Franco Sevilla, T. M. Hong, D. Kovalskyi, J. D. Richman, and C. A. West

*University of California at Santa Barbara, Santa Barbara, California 93106, USA*

A. M. Eisner, W. S. Lockman, B. A. Schumm, and A. Seiden

*University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA*

D. S. Chao, C. H. Cheng, B. Echenard, K. T. Flood, D. G. Hitlin, P. Ongmongkolkul, and F. C. Porter

*California Institute of Technology, Pasadena, California 91125, USA*

R. Andreassen, Z. Huard, B. T. Meadows, B. G. Pushpawela, M. D. Sokoloff, and L. Sun

*University of Cincinnati, Cincinnati, Ohio 45221, USA*

P. C. Bloom, W. T. Ford, A. Gaz, U. Nauenberg, J. G. Smith, and S. R. Wagner

*University of Colorado, Boulder, Colorado 80309, USA*

R. Ayad\* and W. H. Toki

*Colorado State University, Fort Collins, Colorado 80523, USA*

B. Spaan

*Technische Universität Dortmund, Fakultät Physik, D-44221 Dortmund, Germany*

R. Schwierz

*Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany*

D. Bernard and M. Verderi

*Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS/IN2P3, F-91128 Palaiseau, France*

S. Playfer

*University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*

D. Bettoni<sup>a</sup>, C. Bozzi<sup>a</sup>, R. Calabrese<sup>ab</sup>, G. Cibinetto<sup>ab</sup>, E. Fioravanti<sup>ab</sup>,

I. Garzia<sup>ab</sup>, E. Luppi<sup>ab</sup>, L. Piemontese<sup>a</sup>, and V. Santoro<sup>a</sup>

*INFN Sezione di Ferrara<sup>a</sup>; Dipartimento di Fisica e Scienze della Terra, Università di Ferrara<sup>b</sup>, I-44122 Ferrara, Italy*

R. Baldini-Ferrolì, A. Calcaterra, R. de Sangro, G. Finocchiaro,

S. Martellotti, P. Patteri, I. M. Peruzzi,<sup>†</sup> M. Piccolo, M. Rama, and A. Zallo

*INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy*

R. Contri<sup>ab</sup>, E. Guido<sup>ab</sup>, M. Lo Vetere<sup>ab</sup>, M. R. Monge<sup>ab</sup>, S. Passaggio<sup>a</sup>, C. Patrignani<sup>ab</sup>, and E. Robutti<sup>a</sup>

*INFN Sezione di Genova<sup>a</sup>; Dipartimento di Fisica, Università di Genova<sup>b</sup>, I-16146 Genova, Italy*

B. Bhuyan and V. Prasad

*Indian Institute of Technology Guwahati, Guwahati, Assam, 781 039, India*

M. Morii

*Harvard University, Cambridge, Massachusetts 02138, USA*

A. Adametz and U. Uwer

*Universität Heidelberg, Physikalisches Institut, D-69120 Heidelberg, Germany*

H. M. Lacker

*Humboldt-Universität zu Berlin, Institut für Physik, D-12489 Berlin, Germany*

P. D. Dauncey

*Imperial College London, London, SW7 2AZ, United Kingdom*

U. Mallik

*University of Iowa, Iowa City, Iowa 52242, USA*

C. Chen, J. Cochran, W. T. Meyer, and S. Prell

*Iowa State University, Ames, Iowa 50011-3160, USA*

A. V. Gritsan

*Johns Hopkins University, Baltimore, Maryland 21218, USA*

N. Arnaud, M. Davier, D. Derkach, G. Grosdidier, F. Le Diberder,

A. M. Lutz, B. Malaescu,<sup>†</sup> P. Roudeau, A. Stocchi, and G. Wormser

*Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11,  
Centre Scientifique d'Orsay, F-91898 Orsay Cedex, France*

D. J. Lange and D. M. Wright

*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*

J. P. Coleman, J. R. Fry, E. Gabathuler, D. E. Hutchcroft, D. J. Payne, and C. Touramanis

*University of Liverpool, Liverpool L69 7ZE, United Kingdom*

A. J. Bevan, F. Di Lodovico, and R. Sacco  
*Queen Mary, University of London, London, E1 4NS, United Kingdom*

G. Cowan  
*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*

J. Bougher, D. N. Brown, and C. L. Davis  
*University of Louisville, Louisville, Kentucky 40292, USA*

A. G. Denig, M. Fritsch, W. Gradl, K. Griessinger, A. Hafner, E. Prencipe, and K. R. Schubert  
*Johannes Gutenberg-Universität Mainz, Institut für Kernphysik, D-55099 Mainz, Germany*

R. J. Barlow<sup>§</sup> and G. D. Lafferty  
*University of Manchester, Manchester M13 9PL, United Kingdom*

E. Behn, R. Cenci, B. Hamilton, A. Jawahery, and D. A. Roberts  
*University of Maryland, College Park, Maryland 20742, USA*

R. Cowan, D. Dujmic, and G. Sciolla  
*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA*

R. Cheaib, P. M. Patel,<sup>¶</sup> and S. H. Robertson  
*McGill University, Montréal, Québec, Canada H3A 2T8*

P. Biassoni<sup>ab</sup>, N. Neri<sup>a</sup>, and F. Palombo<sup>ab</sup>  
*INFN Sezione di Milano<sup>a</sup>; Dipartimento di Fisica, Università di Milano<sup>b</sup>, I-20133 Milano, Italy*

L. Cremaldi, R. Godang,<sup>\*\*</sup> P. Sonnek, and D. J. Summers  
*University of Mississippi, University, Mississippi 38677, USA*

M. Simard and P. Taras  
*Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7*

G. De Nardo<sup>ab</sup>, D. Monorchio<sup>ab</sup>, G. Onorato<sup>ab</sup>, and C. Sciacca<sup>ab</sup>  
*INFN Sezione di Napoli<sup>a</sup>; Dipartimento di Scienze Fisiche,  
 Università di Napoli Federico II<sup>b</sup>, I-80126 Napoli, Italy*

M. Martinelli and G. Raven  
*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*

C. P. Jessop and J. M. LoSecco  
*University of Notre Dame, Notre Dame, Indiana 46556, USA*

K. Honscheid and R. Kass  
*Ohio State University, Columbus, Ohio 43210, USA*

J. Brau, R. Frey, N. B. Sinev, D. Strom, and E. Torrence  
*University of Oregon, Eugene, Oregon 97403, USA*

E. Feltresi<sup>ab</sup>, M. Margoni<sup>ab</sup>, M. Morandin<sup>a</sup>, M. Posocco<sup>a</sup>, M. Rotondo<sup>a</sup>, G. Simi<sup>a</sup>, F. Simonetto<sup>ab</sup>, and R. Stroili<sup>ab</sup>  
*INFN Sezione di Padova<sup>a</sup>; Dipartimento di Fisica, Università di Padova<sup>b</sup>, I-35131 Padova, Italy*

S. Akar, E. Ben-Haim, M. Bomben, G. R. Bonneaud, H. Briand,  
 G. Calderini, J. Chauveau, Ph. Leruste, G. Marchiori, J. Ocariz, and S. Sitt  
*Laboratoire de Physique Nucléaire et de Hautes Energies,  
 IN2P3/CNRS, Université Pierre et Marie Curie-Paris6,*

*Université Denis Diderot-Paris7, F-75252 Paris, France*

M. Biasini<sup>ab</sup>, E. Manoni<sup>a</sup>, S. Pacetti<sup>ab</sup>, and A. Rossi<sup>a</sup>  
*INFN Sezione di Perugia<sup>a</sup>; Dipartimento di Fisica, Università di Perugia<sup>b</sup>, I-06123 Perugia, Italy*

C. Angelini<sup>ab</sup>, G. Batignani<sup>ab</sup>, S. Bettarini<sup>ab</sup>, M. Carpinelli<sup>ab,††</sup>, G. Casarosa<sup>ab</sup>, A. Cervelli<sup>ab</sup>, F. Forti<sup>ab</sup>,  
 M. A. Giorgi<sup>ab</sup>, A. Lusiani<sup>ac</sup>, B. Oberhof<sup>ab</sup>, E. Paoloni<sup>ab</sup>, A. Perez<sup>a</sup>, G. Rizzo<sup>ab</sup>, and J. J. Walsh<sup>a</sup>  
*INFN Sezione di Pisa<sup>a</sup>; Dipartimento di Fisica, Università di Pisa<sup>b</sup>; Scuola Normale Superiore di Pisa<sup>c</sup>, I-56127 Pisa, Italy*

D. Lopes Pegna, J. Olsen, and A. J. S. Smith  
*Princeton University, Princeton, New Jersey 08544, USA*

R. Faccini<sup>ab</sup>, F. Ferrarotto<sup>a</sup>, F. Ferroni<sup>ab</sup>, M. Gaspero<sup>ab</sup>, L. Li Gioi<sup>a</sup>, and G. Piredda<sup>a</sup>  
*INFN Sezione di Roma<sup>a</sup>; Dipartimento di Fisica,  
 Università di Roma La Sapienza<sup>b</sup>, I-00185 Roma, Italy*

C. Büniger, O. Grünberg, T. Hartmann, T. Leddig, C. Voß, and R. Waldi  
*Universität Rostock, D-18051 Rostock, Germany*

T. Adye, E. O. Olaiya, and F. F. Wilson  
*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*

S. Emery, G. Hamel de Monchenault, G. Vasseur, and Ch. Yèche  
*CEA, Irfu, SPP, Centre de Saclay, F-91191 Gif-sur-Yvette, France*

F. Anulli,<sup>‡‡</sup> D. Aston, D. J. Bard, J. F. Benitez, C. Cartaro, M. R. Convery, J. Dorfan, G. P. Dubois-Felsmann,  
 W. Dunwoodie, M. Ebert, R. C. Field, B. G. Fulsom, A. M. Gabareen, M. T. Graham, C. Hast,  
 W. R. Innes, P. Kim, M. L. Kocian, D. W. G. S. Leith, P. Lewis, D. Lindemann, B. Lindquist, S. Luitz,  
 V. Luth, H. L. Lynch, D. B. MacFarlane, D. R. Muller, H. Neal, S. Nelson, M. Perl, T. Pulliam,  
 B. N. Ratcliff, A. Roodman, A. A. Salnikov, R. H. Schindler, A. Snyder, D. Su, M. K. Sullivan, J. Va'vra,  
 A. P. Wagner, W. F. Wang, W. J. Wisniewski, M. Wittgen, D. H. Wright, H. W. Wulsin, and V. Ziegler  
*SLAC National Accelerator Laboratory, Stanford, California 94309 USA*

W. Park, M. V. Purohit, R. M. White,<sup>§§</sup> and J. R. Wilson  
*University of South Carolina, Columbia, South Carolina 29208, USA*

A. Randle-Conde and S. J. Sekula  
*Southern Methodist University, Dallas, Texas 75275, USA*

M. Bellis, P. R. Burchat, T. S. Miyashita, and E. M. T. Puccio  
*Stanford University, Stanford, California 94305-4060, USA*

M. S. Alam and J. A. Ernst  
*State University of New York, Albany, New York 12222, USA*

R. Gorodeisky, N. Guttman, D. R. Peimer, and A. Soffer  
*Tel Aviv University, School of Physics and Astronomy, Tel Aviv, 69978, Israel*

S. M. Spanier  
*University of Tennessee, Knoxville, Tennessee 37996, USA*

J. L. Ritchie, A. M. Ruland, R. F. Schwitters, and B. C. Wray  
*University of Texas at Austin, Austin, Texas 78712, USA*

J. M. Izen and X. C. Lou  
*University of Texas at Dallas, Richardson, Texas 75083, USA*

F. Bianchi<sup>ab</sup>, F. De Mori<sup>ab</sup>, A. Filippi<sup>a</sup>, D. Gamba<sup>ab</sup>, and S. Zambito<sup>ab</sup>

INFN Sezione di Torino<sup>a</sup>; Dipartimento di Fisica, Università di Torino<sup>b</sup>, I-10125 Torino, Italy

L. Lanceri<sup>ab</sup> and L. Vitale<sup>ab</sup>

INFN Sezione di Trieste<sup>a</sup>; Dipartimento di Fisica, Università di Trieste<sup>b</sup>, I-34127 Trieste, Italy

F. Martinez-Vidal, A. Oyanguren, and P. Villanueva-Perez

IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain

H. Ahmed, J. Albert, Sw. Banerjee, F. U. Bernlochner, H. H. F. Choi, G. J. King, R. Kowalewski,

M. J. Lewczuk, T. Lueck, I. M. Nugent, J. M. Roney, R. J. Sobie, and N. Tasneem

University of Victoria, Victoria, British Columbia, Canada V8W 3P6

T. J. Gershon, P. F. Harrison, and T. E. Latham

Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

H. R. Band, S. Dasu, Y. Pan, R. Prepost, and S. L. Wu

University of Wisconsin, Madison, Wisconsin 53706, USA

Evidence is presented for the baryonic  $B$  meson decay  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  based on a data sample of  $471 \times 10^6$   $B\bar{B}$  pairs collected with the BABAR detector at the PEP-II asymmetric  $e^+e^-$  collider located at the SLAC National Accelerator Laboratory. The branching fraction is determined to be  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}) = (9.8^{+2.9}_{-2.6} \pm 1.9) \times 10^{-6}$ , corresponding to a significance of 3.4 standard deviations including systematic uncertainties. A search for the related baryonic  $B$  meson decay  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  with  $\Sigma^0 \rightarrow \Lambda \gamma$  is performed and an upper limit  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda} + \bar{B}^0 \rightarrow D^0 \Lambda \bar{\Sigma}^0) < 3.1 \times 10^{-5}$  is determined at 90% confidence level.

PACS numbers: 13.25.Hw, 13.60.Rj, 14.20.Lq

## I. INTRODUCTION

Little is known about the mechanism of baryon production in weak decays or in the hadronization process. Baryons are produced in  $(6.8 \pm 0.6)\%$  of all  $B$  meson decays [1]. Due to this large rate,  $B$  meson decays can provide important information about baryon production. Due to the low energy scale, perturbative quantum chromodynamics (QCD) cannot be applied to this process. Furthermore, lattice QCD calculations are not available. The description of baryonic  $B$  decays thus relies on phenomenological models.

Pole models [2] are a common tool used in theoretical studies of hadronic decays. Meson pole models predict an enhancement at low baryon-antibaryon masses. In many three-body decays into a baryon, an antibaryon and a meson, the baryon-antibaryon pair can be described by a meson pole, i.e., the decay of a virtual meson with a mass below threshold. This leads to a steeply falling amplitude at the threshold of the baryon-antibaryon mass, and explains the enhancement observed in decays such as  $B^- \rightarrow \Lambda_c \bar{p} \pi^-$  [3, 4],  $B^- \rightarrow p \bar{p} K^-$  [5–7], and  $\bar{B}^0 \rightarrow D^0 p \bar{p}$  [8, 9].

In addition to the meson pole models described above, there are baryon pole models in which the initial state decays through the strong interaction into a pair of baryons. Then, one of these baryons decays via the weak interaction into a baryon and a meson. For such baryon pole models no enhancement at threshold in the dibaryon invariant mass is expected.

The decay of a  $B$  meson into a  $D^0$  meson and a pair of baryons has been the object of several theoretical investigations [10, 11]. Ref. [11] predicts the branching fractions for  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  decays and for the sum of the  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Sigma}^0$  and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  decays to be

$$\begin{aligned} \mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}) &= (2 \pm 1) \times 10^{-6}, \\ \mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Sigma}^0 + \bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}) &= (1.8 \pm 0.5) \times 10^{-5}. \end{aligned} \quad (1)$$

It is impractical to separate the  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Sigma}^0$  and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  decays since each leads to the final state  $\Lambda \bar{\Lambda} \gamma$ .

\*Now at the University of Tabuk, Tabuk 71491, Saudi Arabia

†Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

‡Now at Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Paris, France

§Now at the University of Huddersfield, Huddersfield HD1 3DH, UK

¶Deceased

\*\*Now at University of South Alabama, Mobile, Alabama 36688, USA

††Also with Università di Sassari, Sassari, Italy

‡‡Also with INFN Sezione di Roma, Roma, Italy

§§Now at Universidad Técnica Federico Santa María, Valparaíso, Chile 2390123




FIG. 1: Leading-order Feynman diagrams for the decays  $\bar{B}^0 \rightarrow D^0 N \bar{N}$ . Setting  $q = u$  leads to the  $D^0 p \bar{p}$  final state and setting  $q = s$  to the  $D^0 \Lambda \bar{\Lambda}$ ,  $D^0 \Sigma^0 \bar{\Lambda}$ ,  $D^0 \Lambda \bar{\Sigma}^0$ , and  $D^0 \Sigma^0 \bar{\Sigma}^0$  final states.

As can be seen from the Feynman diagrams shown in Fig. 1, the only difference between the  $\bar{B}^0 \rightarrow D^0 p \bar{p}$  decay on the one hand and the  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  decays on the other hand is the replacement of a  $u\bar{u}$  pair with an  $s\bar{s}$  pair. In the hadronization process,  $s\bar{s}$ -pair production is suppressed by about a factor of three compared to  $u\bar{u}$ - or  $d\bar{d}$ -pair production [12]. Furthermore, since both  $\Lambda$  and  $\Sigma^0$  baryons can be produced, there are four possible final states with an  $s\bar{s}$  pair ( $\Lambda \bar{\Lambda}$ ,  $\Lambda \bar{\Sigma}^0$ ,  $\Sigma^0 \bar{\Lambda}$ , and  $\Sigma^0 \bar{\Sigma}^0$ ) compared to only one for a  $u\bar{u}$  pair ( $p \bar{p}$ ), neglecting the production of excited baryons. Assuming equal production rates for these four modes and that the spin-1/2 states dominate, a suppression of a factor of  $\sim 12$  is expected for  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  decays compared to  $\bar{B}^0 \rightarrow D^0 p \bar{p}$  decays, where the branching fraction of the latter process is  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 p \bar{p}) = (1.04 \pm 0.04) \times 10^{-4}$  [1].

The branching fraction for  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  has been measured by the Belle Collaboration to be  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}) = (10.5^{+5.7}_{-4.4} \pm 1.4) \times 10^{-6}$  [13]. There are no previous results for the  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  decay mode.

## II. THE BABAR EXPERIMENT

This analysis is based on a data sample of  $429 \text{ fb}^{-1}$  [14], corresponding to  $471 \times 10^6 B \bar{B}$  pairs, collected with the BABAR detector at the PEP-II asymmetric-energy  $e^+e^-$  collider at the SLAC National Accelerator Laboratory at center-of-mass energies near and equal to the  $\Upsilon(4S)$  mass. The reconstruction efficiency is determined through use of Monte Carlo (MC) simulation, based on the EvtGen [15] program for the event generation and the GEANT4 [16] package for modeling of the detector response. The MC events are generated uniformly in the  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  phase space.

The BABAR detector is described in detail elsewhere [17, 18]. Charged particle trajectories are measured with a five-layer double-sided silicon vertex tracker and a 40-layer drift chamber immersed in a 1.5 T axial magnetic field. Charged particle identification is provided by ionization energy measurements in the tracking chambers and by Cherenkov-radiation photons recorded with an internally reflecting ring-imaging detector. Electrons

and photons are reconstructed with an electromagnetic calorimeter.

## III. RECONSTRUCTION OF $\Lambda$ BARYON, $D^0$ MESON, AND $\bar{B}^0$ MESON CANDIDATES

We reconstruct  $\Lambda$  baryons through the decay mode  $\Lambda \rightarrow p \pi^-$  and  $D^0$  mesons through the modes  $D^0 \rightarrow K^- \pi^+$ ,  $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ , and  $D^0 \rightarrow K^- \pi^+ \pi^0$  [19]. Charged kaon and proton candidates are required to satisfy particle identification criteria. Charged pions are selected as charged tracks that are not identified as a kaon or proton.

Candidate  $\pi^0$  mesons are reconstructed from two separated energy deposits in the electromagnetic calorimeter not associated with charged tracks. To discriminate against neutral hadrons, the shower shape of each deposit is required to be consistent with that of a photon [20]. Furthermore, we require  $E(\gamma_1) > 0.125 \text{ GeV}$  and  $E(\gamma_2) > 0.04 \text{ GeV}$ , where  $E(\gamma_1)$  and  $E(\gamma_2)$  are the energies of the photon candidates, with  $E(\gamma_1) > E(\gamma_2)$ . The photon-photon invariant mass is required to lie in the range  $m(\gamma\gamma) \in [0.116, 0.145] \text{ GeV}/c^2$ .

The  $\Lambda$  daughters are fit to a common vertex and the reconstructed mass is required to lie within three standard deviations of the nominal value [1], where the standard deviation is the mass resolution. We select  $\Lambda$  candidates by requiring the flight significance  $L_t/\sigma_{L_t}$  to exceed 4, where  $L_t$  is the  $\Lambda$  flight length in the transverse plane and  $\sigma_{L_t}$  its uncertainty. The  $\Sigma^0$  baryons are produced in the decay  $\Sigma^0 \rightarrow \Lambda \gamma$ , and the photon is not reconstructed.

The  $D^0$  daughter candidates are fit to a common vertex and the reconstructed mass is required to lie within three times the mass resolution from their nominal values [1]. The signal-to-background ratio for  $D^0 \rightarrow K^- \pi^+ \pi^0$  is improved by making use of the resonant substructure of this decay, which is well known. Using results from the E691 Collaboration [21], we calculate the probability  $w_{\text{Dalitz}}$  for a  $D^0$  candidate to be located at a certain position in the Dalitz plane. We require  $w_{\text{Dalitz}} > 0.02$ . Figure 2 shows the Dalitz plot distributions, based on simulation, for candidates selected with and without the  $w_{\text{Dalitz}}$  re-




FIG. 2: Dalitz plot for simulated  $D^0 \rightarrow K^- \pi^+ \pi^0$  events before (gray stars) and after (black crosses) the  $w_{\text{Dalitz}} > 0.02$  requirement. Resonant decays are indicated.

quirement.

The  $D^0$  and  $\Lambda$  candidates are constrained to their nominal masses in the reconstruction of the  $\bar{B}^0$  candidates. We apply a fit to the entire decay chain and require the probability for the vertex fit to be larger than 0.001.

To reduce background from  $e^+e^- \rightarrow q\bar{q}$  events with  $q = u, d, s, c$ , we apply a selection on a Fisher discriminant  $\mathcal{F}$  that combines the values of  $|\cos \theta_{\text{Thr}}|$ , where  $\theta_{\text{Thr}}$  is the angle between the thrust axis of the  $B$  candidate and the thrust axis formed from the remaining tracks and clusters in the event;  $|\cos \theta_z|$ , where  $\theta_z$  is the angle between the  $B$  thrust axis and the beam axis;  $|\cos \phi|$ , where  $\phi$  is the angle between the  $B$  momentum and the beam axis; and the normalized second Fox Wolfram moment [22]. All these quantities are defined in the center-of-mass frame. All selection criteria are summarized in Table I.

TABLE I: Summary of selection criteria.

Selection criterion	Selected candidates
$\Lambda/\bar{\Lambda}$ mass	$m_{p\pi} \in [1.112, 1.120] \text{ GeV}/c^2$
Flight significance	$L_t/\sigma_{L_t} > 4$
$D^0 \rightarrow K^- \pi$ mass	$m_{K\pi} \in [1.846, 1.882] \text{ GeV}/c^2$
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ mass	$m_{K\pi\pi\pi} \in [1.852, 1.876] \text{ GeV}/c^2$
Lateral parameter $\gamma_1$	$0.05 < \text{LAT}(\gamma_1) < 0.55$
Lateral parameter $\gamma_2$	$\text{LAT}(\gamma_2) > 0.075$
Calorimeter energy $\gamma_1$	$E(\gamma_1) > 0.125 \text{ GeV}$
Calorimeter energy $\gamma_2$	$E(\gamma_2) > 0.04 \text{ GeV}$
$\pi^0$ mass	$m_{\gamma\gamma} \in [0.116, 0.145] \text{ GeV}/c^2$
$D^0 \rightarrow K^- \pi^+ \pi^0$ mass	$m_{K\pi\pi^0} \in [1.81, 1.89] \text{ GeV}/c^2$
Dalitz weight	$w_{\text{Dalitz}} > 0.02$
$B$ vertex probability	$p(B) > 0.001$
Fisher discriminant	$\mathcal{F} > 0.1$

#### IV. FIT STRATEGY

We determine the number of signal candidates with a two-dimensional unbinned extended maximum likelihood fit to the invariant mass  $m(D^0 \Lambda \bar{\Lambda})$  and the energy substituted mass  $m_{\text{ES}}$ . The latter is defined as

$$m_{\text{ES}} = \sqrt{(s/2 + \mathbf{p}_0 \cdot \mathbf{p}_B)^2/E_0^2 - |\mathbf{p}|_B^2}, \quad (2)$$

where  $\sqrt{s}$  is the center-of-mass energy,  $\mathbf{p}_B$  the  $B$  candidate's momentum, and  $(E_0, \mathbf{p}_0)$  the four-momentum vector of the  $e^+e^-$  system, each given in the laboratory frame. Both  $m(D^0 \Lambda \bar{\Lambda})$  and  $m_{\text{ES}}$  are centered at the  $B$  mass for well reconstructed  $B$  decays.

Due to the small mass difference of  $76.9 \text{ MeV}/c^2$  [1] between the  $\Lambda$  and  $\Sigma^0$  baryons,  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  decays, where the  $\Sigma^0$  decays radiatively as  $\Sigma^0 \rightarrow \Lambda \gamma$ , are a source of background. Such events peak at the  $B$  mass in  $m_{\text{ES}}$  and are slightly shifted in  $m(D^0 \Lambda \bar{\Lambda})$  with respect to  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  (Fig. 3). We account for this decay by including an explicit term in the likelihood function (see below), whose yield is determined in the fit.

We divide the data sample into three subsamples corresponding to the  $D^0$  decay modes. Given their different signal-to-background ratios, we determine the number of signal candidates in a simultaneous fit to the three independent subsamples. We describe each  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  signal sample with the product of a Novosibirsk function in  $m_{\text{ES}}$  and a sum of two Gaussian functions  $f^{\mathcal{G}\mathcal{G}}$  in  $m(D^0 \Lambda \bar{\Lambda})$  assuming that  $m_{\text{ES}}$  and  $m(D^0 \Lambda \bar{\Lambda})$  are not correlated. We study simulated samples of signal and background events and find no significant correlation between  $m_{\text{ES}}$  and  $m(D^0 \Lambda \bar{\Lambda})$ . The Novosibirsk function is defined as

$$f^{\text{Novo}}(m_{\text{ES}}) = \exp \left[ -\frac{1}{2} \left( \frac{\ln^2[1 + \lambda \alpha(m_{\text{ES}} - \mu)]}{\alpha^2} + \alpha^2 \right) \right],$$

$$\lambda = \sinh(\alpha \sqrt{\ln 4}) / (\sigma \alpha \sqrt{\ln 4}), \quad (3)$$

with  $\mu$  the mean value,  $\sigma$  the width, and  $\alpha$  the tail parameter. The decay  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  is described by the product of a Novosibirsk  $f^{\text{Novo}1, \Sigma^0}$  function in  $m_{\text{ES}}$  and a sum of another Novosibirsk function  $f^{\text{Novo}2, \Sigma^0}$  and a Gaussian  $\mathcal{G}^{\Sigma^0}$  in  $m(D^0 \Lambda \bar{\Lambda})$ . All parameters are determined using Monte Carlo simulated events and are fixed in the final fit. Background from  $e^+e^- \rightarrow q\bar{q}$  events and other  $B$  meson decays is modeled by the product of an ARGUS function [23] in  $m_{\text{ES}}$  and a first order polynomial in  $m(D^0 \Lambda \bar{\Lambda})$ .

The full fit function is defined as




FIG. 3: Distributions for  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  (left) and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  reconstructed as  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  (right) for the  $D^0 \rightarrow K^- \pi^+$  mode in simulated events.

$$\begin{aligned}
 f_j^{\text{Fit}} &= f_j^{\Lambda} + f_j^{\Sigma^0} + f_j^{\text{Bkg}} \\
 &= f_j^{\text{Novo}, \Lambda}(m_{\text{ES}}) \times f_j^{\mathcal{G}\mathcal{G}}(m(D^0 \Lambda \bar{\Lambda})) + f_j^{\text{Novo}, \Sigma^0}(m_{\text{ES}}) \times \left[ f_j^{\text{Novo}, \Sigma^0}(m(D^0 \Lambda \bar{\Lambda})) + \mathcal{G}_j^{\Sigma^0}(m(D^0 \Lambda \bar{\Lambda})) \right] \\
 &\quad + f_j^{\text{ARGUS}}(m_{\text{ES}}) \times f_j^{\text{Poly}}(m(D^0 \Lambda \bar{\Lambda})),
 \end{aligned} \tag{4}$$

where the index  $j$  corresponds to the three  $D^0$  decay modes.

The branching fraction is determined from

$$\begin{aligned}
 \mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}) &= \frac{N(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda})}{2N_{B^0 \bar{B}^0} \times \varepsilon^{\Lambda}} \\
 &\quad \times \frac{1}{\mathcal{B}(\Lambda \rightarrow p\pi)^2 \mathcal{B}(D^0 \rightarrow X)},
 \end{aligned} \tag{5}$$

where  $N(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda})$  is the fitted signal yield,  $N_{B^0 \bar{B}^0}$  the number of the  $B^0 \bar{B}^0$  pairs assuming  $\mathcal{B}(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.5$ ,  $\varepsilon^{\Lambda}$  the total reconstruction efficiency, and  $\mathcal{B}(\Lambda \rightarrow p\pi)$

and  $\mathcal{B}(D^0 \rightarrow X)$  the branching fractions for the daughter decays of  $\Lambda$  and  $D^0$ , respectively. An analogous expression holds for  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda})$ . We perform a simultaneous fit of the three  $D^0$  decay channels to obtain:

$$\begin{aligned}
 N_{\Lambda} &= \frac{N(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda})}{\varepsilon^{\Lambda} \mathcal{B}(D^0 \rightarrow X)}, \\
 N_{\Sigma^0} &= \frac{N(\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda})}{\varepsilon^{\Sigma^0} \mathcal{B}(D^0 \rightarrow X)}.
 \end{aligned} \tag{6}$$

The likelihood function is given by

$$\begin{aligned}
 L &= \prod_j \frac{e^{-(\varepsilon_j^{\Lambda} \mathcal{B}_j N_{\Lambda} + N_j^{\text{Bkg}} + \varepsilon_j^{\Sigma^0} \mathcal{B}_j N_{\Sigma^0})}}{N(j)!} \prod_k^{N(j)} \left[ \varepsilon_j^{\Lambda} \mathcal{B}_j N_{\Lambda} f_j^{\Lambda}(m_{\text{ES}k}, m(D^0 \Lambda \bar{\Lambda})_k) + N_j^{\text{Bkg}} f_j^{\text{Bkg}}(m_{\text{ES}k}, m(D^0 \Lambda \bar{\Lambda})_k) \right. \\
 &\quad \left. + \varepsilon_j^{\Sigma^0} \mathcal{B}_j N_{\Sigma^0} f_j^{\Sigma^0}(m_{\text{ES}k}, m(D^0 \Lambda \bar{\Lambda})_k) \right],
 \end{aligned} \tag{7}$$

where  $\mathcal{B}_j$  is the branching fraction for the  $j$ th  $D^0$  decay,  $N_j^{\text{Bkg}}$  the number of combinatorial background events in the  $j$ th subsample,  $N_{\Lambda}$  and  $N_{\Sigma^0}$  the yields of  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$ , and  $\varepsilon_j^{\Lambda}$  and  $\varepsilon_j^{\Sigma^0}$  the efficiencies for the

$j$ th  $D^0$  decay.



## V. SYSTEMATIC UNCERTAINTIES

We consider the following systematic uncertainties: the uncertainties associated with the number of  $B\bar{B}$  events, the particle identification (PID) algorithm, the tracking algorithm, the  $\pi^0$  reconstruction, the  $D^0$  and  $\Lambda$  branching fractions, the efficiency correction, and the fitting algorithm.

The uncertainty associated with the number of  $B\bar{B}$  pairs is 0.6%. We determine the systematic uncertainty associated with the PID by applying different PID selections and comparing the result with the nominal selection. The difference is 0.8%, which is assigned as the PID uncertainty. The systematic uncertainty associated with the tracking algorithm depends on the number of charged tracks in the decay. We assign a systematic uncertainty of 0.9% for the  $D^0 \rightarrow K^- \pi^+$  and  $D^0 \rightarrow K^- \pi^+ \pi^0$  decays and 1.2% for the  $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$  decay. A 3% uncertainty is assigned to account for the  $\pi^0$  reconstruction in  $D^0 \rightarrow K^- \pi^+ \pi^0$  decays. A detailed description of these detector-related systematic uncertainties is given in Ref. [18].

We rely on the known  $D^0$  branching fractions in our fit. To estimate the associated systematic uncertainty we vary each branching fraction by one standard deviation of its uncertainty [1] and define the systematic uncertainty to be the maximum deviation observed with respect to the nominal analysis. We divide  $m(\Lambda\bar{\Lambda})$  into six bins and determine the total reconstruction efficiency  $\varepsilon_i$  in each bin. We determine the uncertainty due to the use of the average efficiency  $\bar{\varepsilon}$  by studying  $|\varepsilon_i - \bar{\varepsilon}|/\bar{\varepsilon}$  as a function of  $m(\Lambda\bar{\Lambda})$ . We average these values and take the result of 16.3% ( $D^0 \rightarrow K^- \pi^+$ ), 19.6% ( $D^0 \rightarrow K^- \pi^+ \pi^0$ ), and 16.8% ( $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ ) as our estimate of the systematic uncertainty for the efficiency. We estimate the systematic uncertainty due to the fit procedure by independently varying the fit ranges of  $m_{ES}$  and  $m(D^0 \Lambda\bar{\Lambda})$ . The largest differences in the signal yield are 3.9% for the change of the  $m_{ES}$  fit range and 2.1% for the change of the  $m(D^0 \Lambda\bar{\Lambda})$  fit range. To check our background model, we use a second-order polynomial in  $m(D^0 \Lambda\bar{\Lambda})$  instead of a first-order polynomial. The signal yield changes by 1.1%. We use an ensemble of simulated data samples reflecting our fit results to verify the stability of the fit. We generate 1000 such samples with shapes and yields fixed to our results and repeat the final fit. We find no bias in the signal-yield results. All systematic uncertainties are summarized in Table II.

The total systematic uncertainty, obtained by adding all sources in quadrature, is 20.1%.

TABLE II: Summary of the systematic uncertainties for  $\bar{B}^0 \rightarrow D^0 \Lambda\bar{\Lambda}$ .

Source	Relative uncertainty
$B\bar{B}$ counting	0.6%
Particle identification	0.8%
Tracking	
$D^0 \rightarrow K^- \pi^+$	0.9%
$D^0 \rightarrow K^- \pi^+ \pi^0$	0.9%
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	1.2%
$\pi^0$ systematics	
$D^0 \rightarrow K^- \pi^+ \pi^0$	3.0%
$D^0$ and $\Lambda$ branching fractions	2.9%
Efficiency correction	
$D^0 \rightarrow K^- \pi^+$	16.3%
$D^0 \rightarrow K^- \pi^+ \pi^0$	19.6%
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	16.8%
Fit procedure	4.6%
Total uncertainty	20.1%

## VI. RESULTS

The one-dimensional projections of the fit are shown in Fig. 4. We find

$$\begin{aligned} N_{\Lambda} &= 1880^{+560}_{-500}, \\ N_{\Sigma^0} &= 2870^{+1680}_{-1560}. \end{aligned} \quad (8)$$

The statistical significance is calculated as  $\sqrt{-2 \log L_0/L_S}$ , where  $L_0$  is the likelihood value for a fit without a signal component and  $L_S$  is the likelihood value for the nominal fit. The statistical significance of the combined  $\bar{B}^0 \rightarrow D^0 \Lambda\bar{\Lambda}$  and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  yields is 3.9 standard deviations ( $\sigma$ ), while those of the individual  $\bar{B}^0 \rightarrow D^0 \Lambda\bar{\Lambda}$  and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  results are  $3.4\sigma$  and  $1.2\sigma$ , respectively. Compared to the statistical uncertainty the additive systematic uncertainties are negligible. We therefore quote the statistical significance as the global significance.

The branching fractions are

$$\begin{aligned} \mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda\bar{\Lambda}) &= (9.8^{+2.9}_{-2.6} \pm 1.9) \times 10^{-6}, \\ \mathcal{B}(\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda} + \bar{B}^0 \rightarrow D^0 \Lambda\bar{\Sigma}^0) &= \\ &= (15^{+9}_{-8} \pm 3) \times 10^{-6}, \end{aligned} \quad (9)$$

where the first uncertainties represent the statistical uncertainties and the second the systematic uncertainties. As a cross-check of the method, independent fits to the three sub-samples are performed. The results of each of these fits are consistent with each other and with the nominal combined fit.

Since the statistical significance for  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda} + \bar{B}^0 \rightarrow D^0 \Lambda\bar{\Sigma}^0)$  is low, a Bayesian upper limit at the 90% confidence level is calculated by integrating the likelihood function:

$$\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda} + \bar{B}^0 \rightarrow D^0 \Lambda\bar{\Sigma}^0) < 3.1 \times 10^{-5}. \quad (10)$$




FIG. 4: Results of the combined fit. The  $m_{ES}$  projection is shown for  $m(D^0 \Lambda \bar{\Lambda}) \in [5.15, 5.31] \text{ GeV}/c^2$  and the  $m(D^0 \Lambda \bar{\Lambda})$  projection for  $m_{ES} \in [5.272, 5.286] \text{ GeV}/c^2$ . The solid line shows the result of the fit, the dashed curve indicates the  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  contribution, and the shaded histogram the combinatorial background. From top to bottom:  $D^0 \rightarrow K^- \pi^+$ ,  $D^0 \rightarrow K^- \pi^+ \pi^0$ , and  $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$  subsamples.

To investigate the threshold dependence, we perform the fit in bins of  $m(\Lambda \bar{\Lambda})$  and examine the resulting distribution after accounting for the reconstruction efficiency and  $D^0$  branching fractions. The results are shown in Fig. 5. No enhancement in the  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  event rate is observed at the baryon-antibaryon mass threshold within the uncertainties, in contrast to  $\bar{B}^0 \rightarrow D^0 p \bar{p}$  decays, which do exhibit such an enhancement [8].

We compare our results for the  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  branching fractions to theoretical predictions. The result we obtain for the  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  branching fraction is consistent with the prediction of  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda} + \bar{B}^0 \rightarrow D^0 \Lambda \bar{\Sigma}^0) = (18 \pm 5) \times 10^{-6}$  from Ref. [11]. However, the obtained result for the  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  branching fraction is larger than the prediction of  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}) = (2 \pm 1) \times 10^{-6}$  [11] by a factor of

$$\frac{\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda})_{\text{exp}}}{\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda})_{\text{theo}}} = 4.9 \pm 3.0. \quad (11)$$

We further determine

$$\frac{\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda} + \bar{B}^0 \rightarrow D^0 \Lambda \bar{\Sigma}^0)}{\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda})} = 1.5 \pm 0.9, \quad (12)$$

which is in agreement with our assumption that all four modes  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$ ,  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$ ,  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Sigma}^0$ , and  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Sigma}^0$  are produced at equal rates. For the ratio of branching fractions we find

$$\frac{\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda})}{\mathcal{B}(\bar{B}^0 \rightarrow D^0 p \bar{p})} = \frac{1}{10.6 \pm 3.7}, \quad (13)$$

using  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 p \bar{p}) = (1.04 \pm 0.04) \times 10^{-4}$  [1]. This is in agreement with the expected suppression of 1/12 discussed in the introduction.




FIG. 5: Distribution of the invariant baryon-antibaryon mass for  $D^0$ -branching-fraction and efficiency-corrected  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$  signal candidates. The data points represent the BABAR data and the shaded histogram indicates phase-space-distributed simulated events, scaled to match the area under the data.

## VII. SUMMARY

We find evidence for the baryonic  $B$  decay  $\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}$ . We determine the branching fraction to be  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Lambda \bar{\Lambda}) = (9.8^{+2.9}_{-2.6} \pm 1.9) \times 10^{-6}$  with a significance of  $3.4\sigma$  including systematic uncertainties. This is in agreement with the Belle measurement [13]. We find no evidence for an enhancement in the invariant baryon-antibaryon mass distribution near threshold. Our re-

sult for the branching fraction deviates from theoretical predictions based on measurements of  $\bar{B}^0 \rightarrow D^0 p \bar{p}$  but agrees with simple models of hadronization. We find no evidence for the decay  $\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda}$  and calculate a Bayesian upper limit at 90% confidence level of  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \Sigma^0 \bar{\Lambda} + \bar{B}^0 \rightarrow D^0 \Lambda \bar{\Sigma}^0) < 3.1 \times 10^{-5}$ . This result is in agreement with the theoretical expectation.

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support BABAR. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), the Commissariat à l’Énergie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Education and Science of the Russian Federation, Ministerio de Ciencia e Innovación (Spain), and the Science and Technology Facilities Council (United Kingdom). Individuals have received support from the Marie-Curie IEF program (European Union), the A. P. Sloan Foundation (USA) and the Binational Science Foundation (USA-Israel).

- 
- [1] J. Beringer *et al.* (Particle Data Group), Phys. Rev. D **86**, 010001 (2012) and 2013 partial update for the 2014 edition.
  - [2] D. S. Carlstone, S. P. Rosen, and S. Pakvasa, Phys. Lett. **174**, 18771881 (1968).
  - [3] N. Gabychev *et al.* (Belle Collaboration), Phys. Rev. Lett. **97**, 242001 (2006).
  - [4] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **78**, 112003 (2008).
  - [5] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **72**, 051101 (2005).
  - [6] J.T. Wei *et al.* (Belle Collaboration), Phys. Lett. B **659**, 80 (2008).
  - [7] R. Aaij *et al.* (LHCb Collaboration), Phys. Rev. D **88**, 052015 (2013).
  - [8] P. del Amo Sanchez *et al.* (BABAR Collaboration), Phys. Rev. D **85**, 092017 (2012).
  - [9] K. Abe *et al.* (Belle Collaboration), Phys. Rev. Lett. **89**, 151802 (2002).
  - [10] C.H. Chen *et al.*, Phys. Rev. D **78**, 054016 (2008).
  - [11] Y.K. Hsiao, Int. J. Mod. Phys. A **24**, 3638 (2009).
  - [12] G. Lafferty, J. Phys. G **23**, 731 (1997).
  - [13] Y.W. Chang *et al.* (Belle Collaboration), Phys. Rev. D **79**, 052006 (2009).
  - [14] J. P. Lees *et al.* (BABAR collaboration), Nucl. Instrum. Meth. A **726**, 203 (2013).
  - [15] D. J. Lange, Nucl. Instrum. Meth. A **462**, 152 (2001).
  - [16] S. Agostinelli *et al.* (GEANT4 Collaboration), Nucl. Instrum. Meth. A **506**, 250 (2003).
  - [17] B. Aubert *et al.* (BABAR Collaboration), Nucl. Instrum. Meth. A **479**, 1 (2002).
  - [18] B. Aubert *et al.* (BABAR Collaboration), Nucl. Instrum. Meth. A **729**, 2013 (615).
  - [19] Throughout this paper, all decay modes include the charge conjugated process.
  - [20] A. Drescher *et al.*, Nucl. Instrum. Meth. A **237**, 464 (1985).
  - [21] J.C. Anjos *et al.* (E691 Collaboration), Phys. Rev. D **48**, 56 (1993).
  - [22] G.C. Fox, S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
  - [23] H. Albrecht *et al.* (ARGUS Collaboration), Phys. Lett. B **241**, 278 (1990).