Robust estimates of hydrogen-impact ionization cross sections over a wide energy range

A. F. Barghouty

California Institute of Technology, MC 220-47, Pasadena, California 91125

(Received 26 October 1999; published 29 March 2000)

We describe a simple and effective procedure to estimate hydrogen-impact ionization cross sections over the energy range of tens of keV/nucleon to tens of MeV/nucleon. The procedure is valid in the first Born approximation using known or estimated electron-impact cross sections. The original Bates-Griffing relation between the two sets of cross sections is reformulated and a correction factor due to multiple transitions is introduced. Sample calculated ionization cross sections for helium and carbon collisions with hydrogen are presented.

PACS number(s): 34.10.+x, 34.70.+e, 52.20.-j

I. INTRODUCTION

Hydrogen-impact ionization cross sections for light ions in the energy range of $\sim$ keV/nucleon to a few hundreds of MeV/nucleon are poorly known for many reactions of interest to astrophysics and plasma physics. Such cross sections and the corresponding ionization rates are critically needed in, e.g., transport models of energetic ions in space and laboratory plasmas.

For the energy range relevant to these studies the first Born approximation can be used to estimate the cross section σ_p for the ion-hydrogen electron-loss process (e.g., [1]):

$$X^q + H \rightarrow X^{q+1} + H^* + e', \quad (1)$$

if the ion's kinetic energy is above a certain threshold energy E_B , which is ionization-potential dependent. In process (1), H^* denotes all bound and continuum states of H , but without specifying the states, while e' denotes the ejected electron, which can emerge with some kinetic energy imparted to it. The same approximation can also be applied for σ_e from the electron-impact process

$$X^q + e \rightarrow X^{q+1} + e + e'. \quad (2)$$

The original Bates-Griffing relation [2] is a general, but open, functional between the two sets of cross sections for the above two processes. For the same incident relative velocity and at high enough energies (i.e., the incident kinetic energy $\gg E_B$), the two cross sections approach a single value. For energies below this asymptotic region, which, for example, are more relevant to space plasma transport studies, the Bates-Griffing relation gives σ_e in terms of σ_p as a functional since the incident kinetic energies do not directly correspond to each other. The functional is not closed in the sense that σ_p is evaluated at two different energy points in order to estimate σ_e at a single energy point that corresponds to the same incident velocity.

The availability of E_B -dependent, parametric estimates for σ_e (e.g., [3–5]) makes it possible to estimate σ_p using a relation like the Bates-Griffing. To that end, we will express σ_p as a closed functional of σ_e using the Bates-Griffing relation in the first Born approximation. In addition, a correction factor associated with multiple transitions, i.e., sum over shells, is introduced. The procedure is only meant to give a

crude, but robust, first-order approximation to needed cross sections that otherwise either have not been measured or are difficult to calculate theoretically. Sample applications to the direct-impact ionization of helium and carbon in collisions with hydrogen along with some available data are presented.

An implementation of this formalism in F77/90 and C/C++ portable routines along with the needed shells' data files (for elements up to Fe) are available via anonymous FTP from thor.srl.caltech.edu/pub/ace/atomic. Implementation for other elements where estimates (or data) for σ_e are available is straightforward.

II. THE BATES-GRIFFING RELATION

In the first Born approximation the collision is approximated as a transition from continuum to continuum states. In the center-of-mass frame the initial and final wave functions become pure momentum states of a single particle, and the transition matrix is essentially determined by a Coulombic potential and the amount of momentum transfer in the collision. For processes (1) and (2), ignoring any energy imparted to the ejected electron, the Born cross section is written as

$$\sigma_B(E_e) = \frac{8}{\pi E_e} \int_{k_{\min}}^{k_{\max}} dk k^{-3} |\mathcal{M}_{fi}|^2, \quad (3)$$

where E_e is the electron's incident kinetic energy, $k = k_i - k_f$ is the relative momentum, and k_i and k_f are the initial and final relative momenta. The limits $k_{\min}$ and $k_{\max}$ are determined from energy conservation, and are dependent on the incident particle. Kinematically, the two cross sections σ_e and σ_p are different because these limits are different. For a single transition with ionization potential V_j , for protons¹ the formal limits [6] are

$$k_{\min} = \frac{V_j}{2k_i} \left(1 + \frac{1}{4} \frac{m_e}{\mu} \frac{V_j}{E_e} \right), \quad k_{\max} \rightarrow \infty, \quad (4)$$

where m_e is the electron's rest mass and μ the reduced mass of the ion-hydrogen system. For electrons the limits become

¹We treat atomic hydrogen as a proton because the correction factor (in $\mathcal{M}_{fi}$) due to the bound electron behaves like k^{-4} in Eq. (3), which is negligible for the energy range of interest here.

$$k_{\min} = k_i - k_i \left(1 - \frac{V_j}{E_e}\right)^{1/2}, \quad k_{\max} = k_i + k_i \left(1 + \frac{V_j}{E_e}\right)^{1/2}. \quad (5)$$

The matrix elements $\mathcal{M}_{fi}$ need not be evaluated explicitly. Instead, Eq. (3) for protons can be written as

$$\sigma_p(E_p; k) = \frac{\mu}{m_e} \frac{\mathcal{W}(k)}{E_p}, \quad (6)$$

where E_p is the proton's incident kinetic energy such that $k_{\min}$ is the same in both the ion-electron and ion-proton systems. The function $\mathcal{W}$, whose exact form is not required, is defined as

$$\frac{d\mathcal{W}(k)}{dk} \propto -k^{-3} |\mathcal{M}_{fi}|^2, \quad (7)$$

with the property that in the limit $k \rightarrow \infty$, $\mathcal{W} \rightarrow 0$. Similarly, for electrons the Born cross section becomes

$$\sigma_e(E_e; k) = \frac{\mathcal{W}(k) - \mathcal{W}(k')}{E_e}, \quad (8)$$

where $k' = V_j/k$.

Now, defining E_B as $\mu V_j/4m_e$ and γ as $(1 + E_B/E_p)^2$, and realizing that $E_e(k) = (\gamma m_e/\mu) E_p(k)$, the Bates-Griffing relation for σ_e in terms of σ_p can be written as

$$\sigma_e(E_e; k) = \gamma^{-1} \left(\sigma_p(E_p; k) - \frac{E'_p}{E_p} \sigma_p(E'_p; k') \right), \quad (9)$$

with E'_p defined as E_B^2/E_p . In Eq. (9) σ_p needs to be evaluated at two different energy points, E_p and E'_p , corresponding to two different relative momenta, k and k' , in order to estimate σ_e at the energy point E_e . As such the Bates-Griffing relation for σ_e in terms of σ_p is not a closed one. Next, we show how one can, to first order in $(k - k')$, reexpress the Bates-Griffing relation for σ_p in terms of σ_e at the same k .

III. EXPRESSING σ_p IN TERMS OF σ_e

To express σ_p as a closed functional of σ_e , we Taylor expand $\sigma_p(E_p; k)$ about $\sigma_p(E'_p; k')$ to first order in $(k - k')$ as

$$\sigma_p(E_p; k) = \sigma_p(E'_p; k') + (k - k') \frac{d\sigma_p(E_p; k)}{dk}. \quad (10)$$

From Eqs. (6) and (8), to first order in $(k - k')$ with $\mu' = \mu/m_e$, we have

$$\frac{d\sigma_p(E_p; k)}{dk} = \frac{\mu'}{E_p} \frac{d\mathcal{W}}{dk} + \mu' \mathcal{W} \frac{dE_p^{-1}}{dk}, \quad (11)$$

$$\frac{d\mathcal{W}}{dk} = \frac{\mathcal{W}(k) - \mathcal{W}(k')}{k - k'} = \frac{E_e}{k - k'} \sigma_e(E_e; k). \quad (12)$$

FIG. 1. Estimated hydrogen-impact ionization cross sections for helium and carbon. Curves depict the cross sections $\sigma_{0 \rightarrow 1}$ (i.e., the differential cross section for the reaction $X^0 + H \rightarrow X^1 + H^* + e'$), $\sigma_{1 \rightarrow 2}$, etc., and terminate as the kinetic energy $E_p \rightarrow E_B^+$. Data for helium are from Refs. [7–10], and for carbon from Refs. [10,11].

Upon substituting Eqs. (11) and (12) in Eq. (10) and equating terms with Eq. (9), we get

$$\sigma_p(E'_p; k') = -\frac{k - k'}{1 - \epsilon^2} \mu' \mathcal{W} \frac{dE_p^{-1}}{dk} = \frac{2}{1 + \epsilon} \sigma_p(E_p; k), \quad (13)$$

where $\epsilon = E_B/E_p$ and $E_p = 1/4\mu'k^2$. Eliminating $\sigma_p(E'_p; k')$ in Eq. (9) and solving for $\sigma_p(E_p; k)$ gives

$$\sigma_p(E_p; k) = \gamma \left(1 - \frac{2\epsilon^2}{1 + \epsilon} \right)^{-1} \sigma_e(E_e; k), \quad (14)$$

which is a closed relation for σ_p in terms of σ_e at the same k .

Finally, since the Arnaud and Rothenflug [4,5] estimates we use for σ_e are shell dependent and the above relation for σ_p depends on each shell parametrically, the total, i.e., sum over shells, cross section becomes

$$\sigma_p(E_p; k) = \frac{1}{2} (1 + N^2) \sum_{j=1}^N \sigma_p^j(E_p; k) / j^2, \quad (15)$$

where N is the number of shells and $\sigma_p^j(E_p; k)$ is given by Eq. (14) for each ionization potential V_j . The factor $1/2j^2$ assumes hydrogenlike degeneracy and $(1 + N^2)$ is a normalization factor.

To illustrate, in Fig. 1 we show calculated ionization cross sections for helium and carbon where some available data are also shown.

ACKNOWLEDGMENTS

This work has been supported by NSF Grant No. 9810653 and NASA-JOVE Grant No. NAG8-1208.

- [1] M. R. C. McDowell and J. P. Coleman, *Introduction to the Theory of Ion-Atom Collisions* (North-Holland, Amsterdam, 1970).
- [2] D. R. Bates and G. Griffing, Proc. Phys. Soc. London **66**, 961 (1953).
- [3] V. P. Shevelko *et al.*, Mon. Not. R. Astron. Soc. **203**, 45P (1983).
- [4] M. Arnaud and R. Rothenflug, Astron. Astrophys. Suppl. Ser. **60**, 425 (1985).
- [5] M. Arnaud and J. Raymond, Astrophys. J. **398**, 394 (1992).
- [6] G. Peach, Proc. Phys. Soc. London **85**, 709 (1965).
- [7] J. M. Peart, K. Rinn, and K. T. Dolder, J. Phys. B **16**, 1461 (1983).
- [8] M. B. Shah and H. B. Gilbody, J. Phys. B **18**, 899 (1985).
- [9] M. F. Watts, K. F. Dunn, and H. B. Gilbody, J. Phys. B **19**, L355 (1986).
- [10] M. M. Sant'Anna *et al.*, Phys. Rev. A **58**, 1204 (1998).
- [11] T. V. Goffe, M. B. Shah, and H. B. Gilbody, J. Phys. B **12**, 3763 (1979).