Magnetic axis safety factor of finite β spheromaks and transition from spheromaks to toroidal magnetic bubbles

Paul M. Bellan and Roberto Paccagnella

Citation: *Physics of Plasmas* (1994-present) **22**, 022513 (2015); doi: 10.1063/1.4908553

View online: <http://dx.doi.org/10.1063/1.4908553>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/pop/22/2?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Determination of the minimum value of the safety factor from geodesic Alfvén eigenmodes in Joint European Torus](#)

Phys. Plasmas **17**, 110705 (2010); 10.1063/1.3494212

[Ideal magnetohydrodynamic simulation of magnetic bubble expansion as a model for extragalactic radio lobes](#)

Phys. Plasmas **15**, 072905 (2008); 10.1063/1.2948347

[Transition from spherical cap to toroidal bubbles](#)

Phys. Fluids **18**, 052102 (2006); 10.1063/1.2196451

[Deceleration mechanism of spheromak-like compact toroid penetrating into magnetized plasmas](#)

Phys. Plasmas **7**, 5033 (2000); 10.1063/1.1322061

[A new formulation of the resistive magnetohydrodynamics stability problem for finite \$\beta\$ toroidal plasmas](#)

Phys. Plasmas **7**, 4070 (2000); 10.1063/1.1289688

PFEIFFER VACUUM

VACUUM SOLUTIONS FROM A SINGLE SOURCE

Pfeiffer Vacuum stands for innovative and custom vacuum solutions worldwide, technological perfection, competent advice and reliable service.

Magnetic axis safety factor of finite β spheromaks and transition from spheromaks to toroidal magnetic bubbles

Paul M. Bellan¹ and Roberto Paccagnella²

¹Applied Physics and Materials Science, Caltech, Pasadena, California 91107, USA

²Consorzio RFX and Istituto Gas Ionizzati del C.N.R., Corso Stati Uniti, 4 35127 Padova, Italy

(Received 10 November 2014; accepted 3 February 2015; published online 23 February 2015)

The value of the safety factor on the magnetic axis of a finite-beta spheromak is shown to be a function of beta in contrast to what was used in Bellan, *Phys. Plasmas* **9**, 3050 (2002); this dependence on beta substantially reduces the gradient of the safety factor compared to the previous calculation. The method for generating finite-beta spheromak equilibria is extended to generate equilibria describing toroidal magnetic “bubbles,” where the hydrodynamic pressure on the magnetic axis is less than on the toroid surface. This “anti-confinement” configuration can be considered an equilibrium with an inverted beta profile and is relevant to interplanetary magnetic clouds as these clouds have lower hydrodynamic pressure in their interior than on their surface.

© 2015 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4908553>]

I. INTRODUCTION

In Ref. 1, one of the authors (PMB) examined analytic forms of finite β spheromak equilibria and used a well-known expression for the value of the safety factor q on the magnetic axis, denoted as q_{axis} , to argue that finite β causes the beneficial effect of a much larger q gradient than when $\beta = 0$. However, co-author (RP) numerically calculated q_{axis} for these finite β analytic equilibria and found numerical results substantially different from the q_{axis} given in Ref. 1. The reasons for this difference are identified as resulting from a subtle misuse of an expression for q_{axis} . Resolution of this issue revealed that the analytic equilibria presented in Ref. 1 could be extended to give an interesting toroidal equilibria where the pressure on the magnetic axis of a toroid is lower than the pressure at the surface (edge) of the toroid rather than higher as in a tokamak; i.e., the beta profile is inverted and the configuration is bubble-like. Increase of a parameter $\bar{\gamma}$ (to be defined below) results in solutions to a Grad-Shafranov equation evolving from characterizing finite β spheromak equilibria, to a conventional zero β spheromak, to magnetic “bubbles” which are tokamak-like configurations having inverted β profiles, and then to a tokamak with conventional β profile. This evolution is characterized by the ratio of two Bessel functions changing sign as their argument $\bar{\gamma}$ is progressively increased. Interplanetary magnetic clouds are an example of the magnetic bubble situation because on the magnetic axis these clouds have lower hydrodynamic pressure than at their edge. Magnetic clouds have been previously modeled using numerical solutions to Grad-Shafranov equations^{2,3} in a slab approximation (i.e., equations are solved in Cartesian geometry in the xy plane with the z direction ignorable); the model presented here differs by being analytic and axisymmetric (i.e., equations are solved in cylindrical geometry in the rz plane with the ϕ direction ignorable) so that, in contrast to a slab approximation, toroidal geometry effects are inherently included. The analytic model has only a few parameters and so has less

freedom than a numerical model but nevertheless has the useful feature of revealing parametric dependence and scaling. The analytic model also offers the possibility of providing a useful framework for other calculations, for example, calculating particle orbits in an axisymmetric cloud; the virtues of developing a repertoire of analytic solutions to the Grad-Shafranov equation has been discussed in Ref. 4.

II. BASIC RELATIONS

We use a cylindrical coordinate system $\{r, \phi, z\}$ and consider the general axisymmetric magnetic field

$$\mathbf{B} = \frac{1}{2\pi} \nabla\psi \times \nabla\phi + B_\phi r \nabla\phi, \quad (1)$$

where ψ is the poloidal flux function and B_ϕ is the toroidal field. The ϕ direction is called the toroidal direction and any direction lying in the poloidal plane (rz plane) is called a poloidal direction. From Ampere’s law the associated current density is

$$\mathbf{J} = \frac{1}{2\pi\mu_0} \nabla(B_\phi r) \times \nabla\phi - \frac{r^2 \nabla\phi}{2\pi\mu_0} \nabla \cdot \left(\frac{1}{r^2} \nabla\psi \right). \quad (2)$$

We are interested in configurations where the poloidal flux function has a local extremum in the r, z plane; both spheromaks and tokamaks are this type of configuration. The location of this extremum is called the magnetic axis and its vertical location defines the z origin while its radial location is defined to be r_{axis} ; ψ is thus at a maximum or a minimum at $r = r_{axis}, z = 0$. If ψ is at a maximum on the magnetic axis then J_ϕ is positive at the axis whereas if ψ is at a minimum on the magnetic axis then J_ϕ is negative at the axis.

Spheromaks and tokamaks are characterized by the safety factor q which is the number of times a field line goes around toroidally for each time it goes poloidally around the magnetic axis. Tokamaks typically have near-unity q on the magnetic axis with q increasing with increasing distance

from the magnetic axis whereas spheromaks have near-unity q on the magnetic axis and q decreasing on moving away from the magnetic axis. The gradient of q , denoted as q' , provides stability properties and detailed calculations show that a zero β spheromak has small q' .

The safety factor at the magnetic axis is given by⁵

$$q_{axis} = \frac{e^{1/2} + e^{-1/2}}{r_{axis}} \frac{B_{\phi,axis}}{\mu_0 J_{\phi,axis}}, \quad (3)$$

where

$$e = \left(\frac{\psi_{rr}}{\psi_{zz}} \right)_{axis} \quad (4)$$

is a measure of the ellipticity of $\psi(r, z)$ in the vicinity of the magnetic axis such that $e > 1$ corresponds to vertically elongated equilibria (prolate) while $e < 1$ corresponds to vertically shortened equilibria (oblate). The force-free relation $\mu_0 J_{\phi,axis}/B_{\phi,axis} = \lambda$ was invoked in Ref. 5 to give $q_{axis} = (e^{1/2} + e^{-1/2})/(\lambda r_{axis})$, but this result is valid only if the plasma is indeed force-free (i.e., has zero β and equilibrium given by $\nabla \times \mathbf{B} = \lambda \mathbf{B}$). If β is finite, then $\mu_0 J_{\phi,axis} \neq \lambda B_{\phi,axis}$ and it is necessary to calculate the actual value of $\mu_0 J_{\phi,axis}/B_{\phi,axis}$ by consideration of the details of the finite β equilibrium.

To do this, we start by defining β_{rel}

$$\beta_{rel} = \mu_0 \frac{P_{axis} - P_{lc}}{B_{axis}^2} \quad (5)$$

where P_{axis} and P_{lc} are, respectively, the hydrodynamic pressures on the magnetic axis and on the last closed flux surface. Positive β_{rel} thus corresponds to a conventional β profile whereas negative β_{rel} corresponds to an inverted β profile. This definition differs from that used in Ref. 1 because (i) here B_{axis}^2 is used and (ii) a relative rather than absolute pressure is used. The definition in Ref. 1 used, in contrast, the *average poloidal* field linking the circular surface lying in the $z=0$ plane between the geometric axis and the magnetic axis. Because the definition of β_{rel} uses the relative hydrodynamic pressure, it is seen that β_{rel} can be positive or negative. In particular, if P_{axis} is smaller than P_{lc} , then β_{rel} will be negative. The definition of β_{rel} is useful because it provides a simple mathematical way to distinguish toroidal equilibria with inverted β profiles from those with normal β profiles. The former are toroidal magnetic bubbles while the latter are toroidal confinement configurations such as spheromaks and tokamaks.

On expressing the magnetic field as

$$\mathbf{B} = \frac{1}{2\pi} (\nabla\psi(r, z) \times \nabla\phi + \mu_0 I(r, z) \nabla\phi), \quad (6)$$

where $I = 2\pi r B_{\phi}/\mu_0$ is the poloidal current, MHD equilibrium $\mathbf{J} \times \mathbf{B} = \nabla P$ can be expressed as the Grad-Shafranov equation^{6,7}

$$r \frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial \psi}{\partial r} \right) + \frac{\partial^2 \psi}{\partial z^2} + 4\pi^2 \mu_0 r^2 \frac{dP}{d\psi} + \mu_0^2 I \frac{dI}{d\psi} = 0. \quad (7)$$

We assume that P is a linear function of the poloidal flux ψ and so can be expressed as

$$P = \frac{P_{axis} - P_{lc}}{(\psi_{axis} - \psi_{lc})} \psi - \frac{P_{axis}\psi_{lc} - P_{lc}\psi_{axis}}{(\psi_{axis} - \psi_{lc})}, \quad (8)$$

where ψ_{lc} is the last closed flux surface of the configuration.

The poloidal current is similarly assumed to be a linear function of the poloidal flux and can be expressed as

$$\mu_0 I = \lambda \psi. \quad (9)$$

We note that the assumed linear dependence in Eq. (9) differs from the assumption used in Solov'ev-type solutions such as in Ref. 4, where it is assumed that $I^2 \sim \psi + \text{const.}$ For the linear dependence assumed here, $dI/d\psi$ is linear in ψ whereas for the Solov'ev-type assumption, $dI/d\psi$ is a constant.

Using Eq. (9), the toroidal component of Eq. (6) gives

$$B_{\phi,axis} = \frac{\lambda \psi_{axis}}{2\pi r_{axis}}. \quad (10)$$

The gradient of P with respect to ψ can then be expressed in terms of β_{rel} as

$$\frac{dP}{d\psi} = \beta_{rel} \frac{B_{axis}^2}{\mu_0 \psi_{axis} (1 - \psi_{lc}/\psi_{axis})}. \quad (11)$$

Since B_{pol} vanishes at the magnetic axis, $B_{axis}^2 = B_{\phi,axis}^2$ and so

$$\frac{dP}{d\psi} = \beta_{rel} \frac{\lambda^2 \psi_{axis}}{4\pi^2 r_{axis}^2 \mu_0 (1 - \psi_{lc}/\psi_{axis})}. \quad (12)$$

III. CYLINDRICAL SOLUTIONS TO FINITE β_{rel} GRAD-SHAFRANOV EQUATION

We now introduce dimensionless quantities

$$\bar{\psi} = \frac{\psi}{\psi_{axis}}, \quad \bar{r} = \frac{r}{r_{axis}}, \quad \bar{z} = \frac{z}{r_{axis}}, \quad \bar{\lambda} = \lambda r_{axis}, \quad (13)$$

so Eq. (7) can be expressed as

$$\bar{r} \frac{\partial}{\partial \bar{r}} \left(\frac{1}{\bar{r}} \frac{\partial \bar{\psi}}{\partial \bar{r}} \right) + \frac{\partial^2 \bar{\psi}}{\partial \bar{z}^2} + \bar{\lambda}^2 \left(\frac{\beta_{rel} \bar{r}^2}{1 - \bar{\psi}_{lc}} + \bar{\psi} \right) = 0. \quad (14)$$

We define

$$\bar{\chi} = \frac{\beta_{rel} \bar{r}^2}{(1 - \bar{\psi}_{lc})} + \bar{\psi}, \quad (15)$$

so Eq. (14) becomes

$$\bar{r} \frac{\partial}{\partial \bar{r}} \left(\frac{1}{\bar{r}} \frac{\partial \bar{\chi}}{\partial \bar{r}} \right) + \frac{\partial^2 \bar{\chi}}{\partial \bar{z}^2} + \bar{\lambda}^2 \bar{\chi} = 0. \quad (16)$$

We assume a solution of the form

$$\bar{\chi} = \bar{r} g(\bar{r}) \cos(\bar{k} \bar{z}), \quad (17)$$

so Eq. (16) becomes

$$\frac{\partial^2 g}{\partial \bar{r}^2} + \frac{1}{\bar{r}} \frac{\partial g}{\partial \bar{r}} + \left(\bar{\gamma}^2 - \frac{1}{\bar{r}^2} \right) g = 0, \quad (18)$$

where

$$\bar{\gamma}^2 = \bar{\lambda}^2 - \bar{k}^2. \quad (19)$$

Equation (18) is Bessel's equation with general solution for real $\bar{\gamma}$

$$g(\bar{r}) = \sigma_J J_1(\bar{\gamma} \bar{r}) + \sigma_Y Y_1(\bar{\gamma} \bar{r}), \quad (20)$$

where σ_J and σ_Y are constant coefficients to be determined by boundary conditions.

From Eqs. (15) and (17), the solution to the normalized Grad-Shafranov equation is

$$\bar{\psi} = \bar{r}(\sigma_J J_1(\bar{\gamma} \bar{r}) + \sigma_Y Y_1(\bar{\gamma} \bar{r})) \cos(\bar{k} \bar{z}) - \bar{r}^2 Q, \quad (21)$$

where

$$Q = \frac{\beta_{rel}}{1 - \bar{\psi}_{lc}}. \quad (22)$$

However, $\bar{\psi} = 1$ is required at $\bar{r} = 1$, $\bar{z} = 0$ (i.e., at the magnetic axis) so

$$\sigma_J J_1(\bar{\gamma}) + \sigma_Y Y_1(\bar{\gamma}) = 1 + Q. \quad (23)$$

The following three Bessel identities where $C_n = J_n$ or Y_n will now be used repeatedly in the rest of the discussion:

$$\frac{dC_0(s)}{ds} = -C_1(s), \quad (24a)$$

$$\frac{d}{ds}(sC_1(s)) = sC_0(s), \quad (24b)$$

$$sC_2(s) = 2C_1(s) - sC_0(s). \quad (24c)$$

The magnetic axis is also where $\partial \bar{\psi} / \partial \bar{r}$ vanishes and so taking the derivative of Eq. (21) with respect to $\bar{r}$, using Eq. (24b), and then setting $\bar{r} = 1$ and $\bar{z} = 0$ gives

$$\sigma_J J_0(\bar{\gamma}) + \sigma_Y Y_0(\bar{\gamma}) = \frac{2}{\bar{\gamma}} Q. \quad (25)$$

Equations (23) and (25) constitute two linear inhomogeneous algebraic equations for the coefficients σ_J and σ_Y . Solving these equations for σ_J and σ_Y and using the Wronskian

$$J_1(s)Y_0(s) - J_0(s)Y_1(s) = \frac{2}{\pi s}, \quad (26)$$

and Eq. (24c) gives

$$\sigma_J = \frac{\pi \bar{\gamma}}{2} (Y_0(\bar{\gamma}) - Q Y_2(\bar{\gamma})), \quad (27a)$$

$$\sigma_Y = \frac{\pi \bar{\gamma}}{2} (-J_0(\bar{\gamma}) + Q J_2(\bar{\gamma})). \quad (27b)$$

IV. SPHEROMAK-TYPE SOLUTIONS

Spheromaks are *singly connected* Grad-Shafranov equilibria (i.e., there is no “hole” in the “doughnut”) and so the domain includes $\bar{r} = 0$. A spheromak therefore cannot contain a $Y_1(\bar{\gamma} \bar{r})$ component because $Y_1(\bar{\gamma} \bar{r})$ diverges at $\bar{r} = 0$. It is thus necessary to impose $\sigma_Y = 0$ for a spheromak in which case Eq. (27b) yields the relation

$$Q = \frac{J_0(\bar{\gamma})}{J_2(\bar{\gamma})}. \quad (28)$$

Substituting for Q in Eq. (27a) and using Eqs. (24c) and (26) gives

$$\sigma_J = \frac{2}{\bar{\gamma} J_2(\bar{\gamma})}. \quad (29)$$

Using Eq. (24c) to substitute for $J_2(\bar{\gamma})$ in Eq. (28) shows that Eq. (28) can alternately be written as

$$Q = \frac{2J_1(\bar{\gamma})}{\bar{\gamma} J_2(\bar{\gamma})} - 1, \quad (30)$$

so one can also write σ_J as

$$\sigma_J = \frac{1 + Q}{J_1(\bar{\gamma})}. \quad (31)$$

Because $\sigma_Y = 0$ for a spheromak Eqs. (22) and (28) show that a spheromak has

$$\beta_{rel} = \frac{J_0(\bar{\gamma})}{J_2(\bar{\gamma})} (1 - \bar{\psi}_{lc}) \quad (32)$$

and

$$\sigma_J = \frac{1}{J_1(\bar{\gamma})} \left(1 + \frac{\beta_{rel}}{1 - \bar{\psi}_{lc}} \right). \quad (33)$$

On substituting for σ_J and Q in Eq. (21) the solution to the normalized Grad-Shafranov equation becomes

$$\bar{\psi} = \bar{r} \frac{J_1(\bar{\gamma} \bar{r})}{J_1(\bar{\gamma})} \left(1 + \frac{\beta_{rel}}{1 - \bar{\psi}_{lc}} \right) \cos(\bar{k} \bar{z}) - \bar{r}^2 \frac{\beta_{rel}}{1 - \bar{\psi}_{lc}}. \quad (34)$$

If $\beta_{rel} = 0$ and $\bar{\psi}_{lc} = 0$ are additionally assumed, the standard result for a zero-beta spheromak in a cylindrical flux conserver of radius a is retrieved, namely, $\bar{\gamma} = x_{01} = 2.405$ where x_{01} is the first root of J_0 . Since $\bar{\gamma} = \gamma r_{axis}$, and the last closed flux surface is at the cylinder radius, then the assumption $\bar{\psi}_{lc} = 0$ and $\beta_{rel} = 0$ in Eq. (34) implies $J_1(\gamma a) = 0$ in which case $\gamma a = x_{11} = 3.83$ where x_{11} is the first root of J_1 . Thus, for a $\beta_{rel} = 0$ spheromak, $r_{axis}/a = \bar{\gamma}/(\gamma a) = x_{01}/x_{11} = 0.63$ as is well known. Equation (32) shows that spheromaks with finite positive β_{rel} are restricted to the range $0 < \bar{\gamma} < 2.405$ but, as will be discussed in Sec. VI, physically relevant non-spheromak configurations with negative β_{rel} exist when $\bar{\gamma} > 2.405$.

Substitution of Eq. (32) into Eq. (34) gives

$$\bar{\psi} = \frac{1}{J_2(\bar{\gamma})} \left(\frac{2\bar{r}}{\bar{\gamma}} J_1(\bar{\gamma}\bar{r}) \cos(\bar{k}\bar{z}) - \bar{r}^2 J_0(\bar{\gamma}) \right), \quad (35)$$

which reverts to the $\beta_{rel}=0$ solution when $\bar{\gamma} = 2.405$ as can be seen using Eq. (24c) to give $\bar{\gamma} J_2(\bar{\gamma}) = 2J_1(\bar{\gamma})$ if $J_0(\bar{\gamma}) = 0$.

V. SAFETY FACTOR OF SPHEROMAKS WITH FINITE β_{rel}

The last closed flux surface of a spheromak has $\bar{\psi}_{lc} = 0$ and $P_{lc} = 0$ in which case Eqs. (5) and (10) give

$$\beta_{rel} = 4\pi^2 r_{axis}^2 \frac{\mu_0 P_{axis}}{\lambda^2 \psi_{axis}^2} \quad (36)$$

and Eq. (34) becomes

$$\bar{\psi} = \bar{r} \frac{J_1(\bar{\gamma}\bar{r})}{J_1(\bar{\gamma})} (1 + \beta_{rel}) \cos(\bar{k}\bar{z}) - \beta_{rel} \bar{r}^2, \quad (37)$$

which is the same as Eq. (2) of Ref. 1 except for the different definition of β_{rel} .

In order to determine q_{axis} , Eq. (3) shows that it is necessary to calculate $\mu_0 J_{\phi,axis} / B_{\phi,axis}$. Equation (2) shows that

$$\mu_0 J_{\phi} = -\frac{\psi_{axis}}{2\pi \bar{r} r_{axis}^3} \left[\bar{r} \frac{\partial}{\partial \bar{r}} \left(\frac{1}{\bar{r}} \frac{\partial \bar{\psi}}{\partial \bar{r}} \right) + \frac{\partial^2 \bar{\psi}}{\partial \bar{z}^2} \right] \quad (38)$$

so, using Eqs. (10) and (14), it is seen that

$$\frac{\mu_0 J_{\phi,axis}}{B_{\phi,axis}} = (1 + \beta_{rel}) \lambda. \quad (39)$$

Thus, $\mu_0 J_{\phi,axis} / B_{\phi,axis} = \lambda$ only if $\beta_{rel} = 0$. Inserting Eq. (39) in Eq. (3) gives

$$q_{axis} = \frac{e^{1/2} + e^{-1/2}}{\bar{\lambda}(1 + \beta_{rel})}, \quad (40)$$

which differs from Eq. (30) of Ref. 1 by having an extra and important factor of $(1 + \beta_{rel})$ in the denominator.

From Eq. (35) and use of the Bessel identities, it is seen that

$$(\bar{\psi}_{zz})_{axis} = -\frac{2\bar{k}^2 J_1(\bar{\gamma})}{\bar{\gamma} J_2(\bar{\gamma})}, \quad (41a)$$

$$(\bar{\psi}_{rr})_{axis} = \frac{-2\bar{\gamma} J_1(\bar{\gamma})}{J_2(\bar{\gamma})}, \quad (41b)$$

so the ellipticity is

$$e = \frac{\bar{\gamma}^2}{\bar{k}^2}. \quad (42)$$

This indicates that the poloidal flux surfaces will be circular near the magnetic axis (i.e., have $e=1$) if $\bar{\gamma} = \bar{k}$ in which case $\bar{\lambda} = \sqrt{2}\bar{\gamma}$. Combination of Eqs. (32), (40), and (42) gives

$$q_{axis} = \frac{\bar{\lambda}}{\bar{\gamma}\bar{k}} \frac{1}{1 + \beta_{rel}} = \frac{\bar{\lambda}}{2\bar{k}} \frac{J_2(\bar{\gamma})}{J_1(\bar{\gamma})}. \quad (43)$$

Equation (43) has been validated by direct numerical integration of field lines in the vicinity of the magnetic axis of a magnetic configuration characterized by Eq. (6) with $\bar{\psi}$ given by Eq. (35). In the $\beta_{rel}=0$ limit, $J_0(\bar{\gamma}) = 0$ and $q_{axis} \rightarrow \bar{\lambda}/(\bar{\gamma}\bar{k})$ which is Eq. (33) of Ref. 1, but for finite positive β_{rel} , Eq. (43) shows that q_{axis} is reduced from its $\beta_{rel}=0$ value.

The safety factor at the wall is¹

$$q_{wall} = \frac{\bar{\lambda}}{2\pi\bar{k}} \cos^{-1}(J_0(\bar{\gamma})), \quad (44)$$

and so the ratio of safety factor at the wall to that at the axis is

$$\frac{q_{wall}}{q_{axis}} = \frac{J_1(\bar{\gamma})}{J_2(\bar{\gamma})} \frac{\cos^{-1}(J_0(\bar{\gamma}))}{\pi}, \quad (45)$$

which is plotted in Fig. 1. Contrary to Ref. 1, it is seen that the shear (difference between q_{wall} and q_{axis}) decreases with increasing β_{rel} (i.e., with $\bar{\gamma}$ decreasing below 2.405). Using $i\theta = \ln(\cos \theta + i \sin \theta)$ to write

$$\cos^{-1}(J_0(\bar{\gamma})) = -i \ln \left(J_0(\bar{\gamma}) + i \sqrt{1 - (J_0(\bar{\gamma}))^2} \right) \quad (46)$$

and then using $J_0(\bar{\gamma}) = 1 - \bar{\gamma}^2/4$ for $\bar{\gamma} \ll 1$, it is seen that for $\bar{\gamma} \ll 1$

$$\cos^{-1}(J_0(\bar{\gamma})) \simeq -i \ln \left(1 - \frac{\bar{\gamma}^2}{4} + i \frac{\bar{\gamma}}{\sqrt{2}} \right) \simeq \frac{\bar{\gamma}}{\sqrt{2}}. \quad (47)$$

Since $J_1(\bar{\gamma}) \simeq \bar{\gamma}/2$ and $J_2(\bar{\gamma}) \simeq \bar{\gamma}^2/8$ for $\bar{\gamma} \ll 1$, Eq. (45) has the limiting behavior

$$\frac{q_{wall}}{q_{axis}} \rightarrow \frac{4}{\sqrt{2}\pi} = 0.900 \text{ for } \bar{\gamma} \ll 1 \quad (48)$$

which is seen in Fig. 1. Furthermore, Eq. (32) has the limiting behavior

$$\beta_{rel} \rightarrow \frac{8}{\bar{\gamma}^2} \text{ for } \bar{\gamma} \ll 1; \quad (49)$$

i.e., β_{rel} diverges at small $\bar{\gamma}$ which is also seen in Fig. 1.

FIG. 1. q_{wall}/q_{axis} plotted as black solid line v. $\bar{\gamma}$ from Eq. (45) and β_{rel} plotted as red dashed line.

We note that numerical calculations reported in Ref. 8 assumed $I^2 \sim \psi^2(1 + 2\alpha\psi/3)$ and $dP/d\psi \sim \psi - \psi_0$ in a spherical geometry and found that the gradient of the shear had a strong dependence on α . The analytic solution given here would correspond approximately to the $\alpha = 0$ numerical solution reported in Ref. 8; the correspondence is not exact because of the different assumptions for the dependence of P on ψ , the shape of the boundary (cylinder v. sphere), and the assumption of a central hole in Ref. 8.

VI. TOROIDAL MAGNETIC BUBBLE: NEGATIVE β_{rel}

We now consider the situation where $\beta_{rel} < 0$ and $\bar{\psi}_{lc} \neq 0$. We consider the $\sigma_Y = 0$ case first as was assumed for spheromaks and then later consider the more general case where both σ_J and σ_Y are finite.

A. $\sigma_Y = 0$ case

In the $\sigma_Y = 0$ case $\bar{\psi}(\bar{r}, \bar{z})$ is mathematically identical to the spheromak solution considered in Sec. IV, i.e., Eq. (35) provides the relevant flux function. The difference here is that $\bar{\psi}_{lc}$ is no longer assumed to be zero. Plots of $\bar{\psi}(\bar{r}, \bar{z})$ using $\bar{\gamma} > 2.405$ show that $\bar{\psi}(\bar{r}, 0)$ has periodic maxima and minima because of its $J_1(\bar{\gamma}\bar{r})$ dependence. Equation (13) defined $\bar{\psi}$ to be unity on the magnetic axis, i.e., $\bar{\psi}(\bar{r}, 0) = 1$ at $\bar{r} = 1$ and the magnetic axis was defined to be where $\bar{\psi}$ was a maximum or minimum. Because of the oscillatory behavior of Bessel functions, maxima or minima of $\bar{\psi}$ occur not only at $r = 1$ but also for $\bar{r} > 1$. However, the maxima and minima occurring where $\bar{r} > 1$ do not have $\bar{\psi} = 1$ and so do not satisfy the $\bar{\psi} = 1$ condition given in Eq. (13). Thus, only the maximum of $\bar{\psi}(\bar{r}, 0)$ at $\bar{r} = 1$ will be considered since maxima or minima at larger $\bar{r}$ do not satisfy the $\bar{\psi} = 1$ requirement stipulated in Eq. (13).

Examination of Eq. (35) shows that $\bar{\psi}$ is independent of $\bar{z}$ if $J_1(\bar{\gamma}\bar{r}) = 0$; at this radius $\bar{r} = x_{11}/\bar{\gamma}$ where $x_{11} = 3.832$ is the first root of J_1 . We now show that this radius $\bar{r} = x_{11}/\bar{\gamma}$ is infinitesimally larger than the radius of the last closed flux surface. Since $\bar{\psi}$ is independent of $\bar{z}$ when $J_1(\bar{\gamma}\bar{r}) = 0$, the flux surface passing through $\bar{r} = x_{11}/\bar{\gamma}$, $\bar{z} = 0$ must be a straight vertical line, i.e., $\bar{\psi}(x_{11}/\bar{\gamma}, \bar{z}) = \bar{\psi}(x_{11}/\bar{\gamma}, 0)$ for all $\bar{z}$. Because a straight vertical line goes to $\bar{z} = \pm\infty$, the flux surface passing through $\bar{r} = x_{11}/\bar{\gamma}$, $\bar{z} = 0$ is open. Immediately to the left of this line, the flux surfaces are closed and so the last closed flux surface is at the radius $\bar{r}_{lc}$ where

$$\bar{r}_{lc} = \lim_{\delta \rightarrow 0} \left(\frac{x_{11}}{\bar{\gamma}} - \delta \right) = \frac{x_{11}}{\bar{\gamma}}. \quad (50)$$

This can also be seen graphically from the flux surface contours shown in Fig. 2 (to be discussed in more detail later) where it is seen that a straight vertical line separatrix lies between the blue-purple closed flux surfaces having magnetic axis at $\bar{r} = 1$, $\bar{z} = 0$ and the green-orange flux surfaces to the right. Equation (50) gives the radial location of this vertical line.

A toroidal inverse aspect ratio (ratio of torus minor to major radius) can be defined as

FIG. 2. Contour plot of $\bar{\psi}(\bar{r}, \bar{z})$ given by Eq. (35) for $\bar{\gamma} = \bar{k} = 2.5$. The radius of the last closed flux surface is at $\bar{r} = 1.53$. The hydrodynamic pressure is lower in the pink region than in the green region so the configuration is a magnetic bubble.

$$\varepsilon = \frac{r_{lc} - r_{axis}}{r_{axis}} = \bar{r}_{lc} - 1 = \frac{x_{11} - \bar{\gamma}}{\bar{\gamma}}. \quad (51)$$

Using $J_1(\bar{\gamma}\bar{r}_{lc}) = 0$ at the last closed flux surface, Eq. (35) may be evaluated at $\bar{r} = \bar{r}_{lc}$, $\bar{z} = 0$ to give

$$\bar{\psi}_{lc} = -\frac{x_{11}^2 J_0(\bar{\gamma})}{\bar{\gamma}^2 J_2(\bar{\gamma})}. \quad (52)$$

Inserting $\bar{\psi}_{lc}$ in Eq. (32) gives

$$\beta_{rel} = \left(1 + \frac{x_{11}^2 J_0(\bar{\gamma})}{\bar{\gamma}^2 J_2(\bar{\gamma})} \right) \frac{J_0(\bar{\gamma})}{J_2(\bar{\gamma})}. \quad (53)$$

In order to have $r_{lc} > r_{axis}$ Eq. (51) shows that it is necessary to have $\bar{\gamma} < x_{11} = 3.832$. A plot of Eq. (53) shows that β_{rel} is negative if $2.405 < \bar{\gamma} < 3.736$; β_{rel} changes sign at $\bar{\gamma} = 3.736$ because the quantity in parenthesis in Eq. (53) changes sign at $\bar{\gamma} = 3.736$. Thus if $2.405 < \bar{\gamma} < 3.736$, β_{rel} is negative and also $r_{lc} > r_{axis}$.

Because the minimum of $(1 + s)s$ occurs when $s = -1/2$, identifying $s = x_{11}^2 J_0(\bar{\gamma})/(\bar{\gamma}^2 J_2(\bar{\gamma}))$ it is seen that β_{rel} is at a minimum when $x_{11}^2 J_0(\bar{\gamma})/(\bar{\gamma}^2 J_2(\bar{\gamma})) = -1/2$ in which case

$$\min[\beta_{rel}] = -\frac{\bar{\gamma}^2}{4x_{11}^2}. \quad (54)$$

Using the Bessel identities, the magnetic field components are

$$B_r = -\frac{\psi_{axis}}{2\pi r_{axis}^2 \bar{r}} \frac{\partial \bar{\psi}}{\partial \bar{z}} = \frac{\psi_{axis}}{2\pi r_{axis}^2} \frac{2\bar{k}}{\bar{\gamma}} \frac{J_1(\bar{\gamma}\bar{r})}{J_2(\bar{\gamma})} \sin(\bar{k}\bar{z}), \quad (55a)$$

$$B_\phi = \frac{\lambda\psi}{2\pi r} = \frac{\psi_{axis}}{2\pi r_{axis}^2} \frac{\bar{\lambda}}{J_2(\bar{\gamma})} \left(\frac{2}{\bar{\gamma}} J_1(\bar{\gamma}\bar{r}) \cos(\bar{k}\bar{z}) - \bar{r} J_0(\bar{\gamma}) \right), \quad (55b)$$

$$B_z = \frac{\psi_{axis}}{2\pi r_{axis}^2 \bar{r}} \frac{\partial \bar{\psi}}{\partial \bar{r}} = \frac{\psi_{axis}}{2\pi r_{axis}^2} \frac{2}{J_2(\bar{\gamma})} \left(J_0(\bar{\gamma}\bar{r}) \cos(\bar{k}\bar{z}) - J_0(\bar{\gamma}) \right). \quad (55c)$$

Using Eqs. (24c) and (55b), it is seen that

$$B_{\phi,axis} = \frac{\psi_{axis}}{2\pi r_{axis}^2} \bar{\lambda}. \quad (56)$$

A normalized magnetic field can be defined as $\bar{\mathbf{B}} = \mathbf{B}(\bar{r}, \bar{z})/B_{\phi,axis}$ with components

$$\bar{B}_r(\bar{r}, \bar{z}) = \frac{2\bar{k} J_1(\bar{\gamma}\bar{r})}{\bar{\gamma}\bar{\lambda} J_2(\bar{\gamma})} \sin(\bar{k}\bar{z}), \quad (57a)$$

$$\bar{B}_\phi(\bar{r}, \bar{z}) = \frac{\frac{2}{\bar{\gamma}} J_1(\bar{\gamma}\bar{r}) \cos(\bar{k}\bar{z}) - \bar{r} J_0(\bar{\gamma})}{J_2(\bar{\gamma})}, \quad (57b)$$

$$\bar{B}_z(\bar{r}, \bar{z}) = \frac{2}{\bar{\lambda}} \frac{(J_0(\bar{\gamma}\bar{r}) \cos(\bar{k}\bar{z}) - J_0(\bar{\gamma}))}{J_2(\bar{\gamma})}. \quad (57c)$$

As required, both $\bar{B}_r$ and $\bar{B}_z$ vanish on the magnetic axis (i.e., at $\bar{r} = 1, \bar{z} = 0$) and $\bar{B}_\phi = 1$ on the magnetic axis.

Equation (35) with $2.405 < \bar{\gamma} < 3.736$ thus gives the flux surface for a magnetic bubble, i.e., a toroidal configuration with closed field lines, where the pressure on the magnetic axis is *lower* than the pressure at the surface of the toroid. The direction of the $\mathbf{J} \times \mathbf{B}$ force is thus outwards rather than inwards in contrast to a tokamak. This configuration is relevant to axisymmetric interplanetary magnetic clouds ejected from the sun by coronal mass ejections. Spacecraft measurements indicate that P is smaller in the interior of these clouds than outside so these clouds have negative β_{rel} . Another possible situation would be in the solar interior where a toroidal bubble configuration as described here would be a toroidal region of stronger magnetic field but reduced hydrodynamic pressure compared to the surroundings.

As a concrete example of such a configuration, consider the situation where $\bar{\gamma} = \bar{k} = 2.5$ and $\bar{\lambda} = \sqrt{2}\bar{\gamma}$. In this case $e = 1$ so the poloidal flux surfaces are circular near the magnetic axis, the last closed flux surface is at $\bar{\psi}_{lc} = 0.25$ and from Eq. (53) $\beta_{rel} = -0.081$. From Eq. (51), it is seen that the inverse aspect ratio is $\varepsilon = 0.53$. Figure 2 plots contours of $\psi(\bar{r}, \bar{z})$ and it is seen that the last closed flux surface intersects $\bar{z} = 0$ to the right of the magnetic axis at indeed $\bar{r}_{lc} = x_{11}/\bar{\gamma} = 1.53$. Figures 3–6 plot $\bar{\psi}(\bar{r}, 0)$, $\bar{B}_\phi(\bar{r}, 0)$, $\bar{B}_z(\bar{r}, 0)$, and $\bar{B}^2(\bar{r}, 0)$, respectively.

From Eq. (5), it is seen that

$$\frac{\mu_0}{B_{axis}^2} P_{axis} = \frac{\mu_0}{B_{axis}^2} P_{lc} + \beta_{rel}, \quad (58)$$

FIG. 3. Plot of $\bar{\psi}(\bar{r}, 0)$ as given by Eq. (35) for $\bar{\gamma} = \bar{k} = 2.5$ and $\bar{\lambda} = 2\sqrt{\bar{\gamma}}$. The radius of the last closed flux surface is at $\bar{r} = 1.53$ and $\bar{\psi}_{lc} = 0.25$.

FIG. 4. $\bar{B}_\phi(\bar{r}, 0)$; Note that magnetic axis is at $\bar{r} = 1.0$ and that maximum of $\bar{B}_\phi$ occurs to left of magnetic axis.

so the hydrodynamic pressure on the magnetic axis is lower than on the last closed flux surface. If P_{axis} is set to zero, then the external pressure would be

$$\frac{\mu_0}{B_{axis}^2} P_{lc} = -\beta_{rel}, \quad (59)$$

in which case the configuration would be a vacuum at the magnetic axis (zero plasma pressure) with increasing pressure going away from the magnetic axis toward the last closed flux surface.

If $\bar{\gamma}$ is further increased, the sign of β_{rel} can become positive again in which case the equilibrium will become tokamak-like (higher pressure on magnetic axis). Additional increase of $\bar{\gamma}$ will cause β_{rel} to oscillate in sign giving a sequence of bubble-like and tokamak-like configurations. Also, for a given configuration one could elect to truncate the flux at some value larger than ψ_{lc} and so obtain a smaller aspect ratio equilibrium. In accordance with the Shafranov virial theorem, any one of these configurations will involve a jump in the magnetic field at the surface of the toroid if it is assumed that at the surface the external magnetic field differs from the internal field. This jump corresponds to the existence of surface currents. In a tokamak, these surface currents are provided by a set of coils immediately external to the toroidal volume and these coils are called the vertical field coils. The field produced by these coils is mainly in the z direction and will be referred to here as B_z^{ext} . This field B_z^{ext} constitutes a portion of the total field inside the toroidal volume and provides equilibrium in the major radius direction. This takes place via a radial force $\sim J_\phi B_z^{ext}$ directed towards $\bar{r} = 0$ that balances the radially outward hoop force as well as some hydrodynamic pressure forces. The hoop force is a property of any toroidal current system and occurs because a

FIG. 5. $\bar{B}_z(\bar{r}, 0)$; note that $\bar{B}_z$ changes sign at magnetic axis.

FIG. 6. $\bar{B}_r^2 + \bar{B}_\phi^2 + \bar{B}_z^2$ as a function of $\bar{r}$ for $\bar{z} = 0$.

toroidal current produces a stronger poloidal field near $\bar{r} = 0$, $\bar{z} = 0$ (inside) than at $\bar{r} = \bar{r}_{lc}$, $\bar{z} = 0$ (outside). This stronger poloidal field on the inside compared to the outside corresponds to greater magnetic pressure on the inside than on the outside; for low β the force resulting from magnetic pressure imbalance dominates any hydrodynamic pressure imbalance. Without the offsetting force provided by B_z^{ext} , the hoop force would act to expand the torus major radius.

At first sight, it might appear that the flux contours in Fig. 2 are such that the magnetic pressure is higher on the outside than on the inside because the midplane poloidal flux surfaces in Fig. 2 are more tightly packed outside the magnetic axis (e.g., at $\bar{r} \simeq 1.5$) than inside the magnetic axis (e.g., at $\bar{r} \simeq 0.25$). However, the density of field lines and hence the poloidal field is nevertheless stronger inside the magnetic axis than outside because of the inverse $\bar{r}$ dependence in $B_z = (2\pi r)^{-1} \partial\psi/\partial r$. The twice as tight midplane flux surface packing in Fig. 2 at $\bar{r} \simeq 1.5$ compared to at $\bar{r} \simeq 0.25$ gives a twice as large $\partial\psi/\partial r$ on the outside compared to the inside. However, this twice as tight radial packing is overcome by the $(2\pi r)^{-1}$ factor, a toroidal geometry effect that produces an approximately six-fold inside-to-outside enhancement with the net result that $|B_z|$ is about three times larger at $\bar{r} \simeq 0.25$ than at $\bar{r} \simeq 1.5$. This three-fold inside-to-outside ratio of $|B_z|$ is evident in Fig. 5.

In order to have the B_z^{ext} required for equilibrium, it would be necessary to have surface currents flowing on the surface of the toroid. Since there are no powered coils to sustain surface currents exterior to a magnetic cloud, it is unlikely that such surface currents would exist in the magnetic cloud context. Without the B_z^{ext} provided by surface currents (and intrinsic to the equilibrium given here), the hoop force resulting from the imbalance between B_z^2 on the inside and B_z^2 on the outside will cause the major radius of magnetic clouds to increase with time. The difference between poloidal flux surfaces with and without incorporation of B_z^{ext} is of the order of the inverse aspect ratio ε because B_z^{ext} is a toroidal effect and so scales as ε .

B. Finite σ_J and σ_Y case

The spheromak solution required σ_Y to be zero to avoid singularity at $\bar{r} = 0$. The magnetic bubble solution discussed above used the same functional form as the spheromak solution (i.e., had $\sigma_Y = 0$ and used Eq. (34)) and found that a tokamak-like solution with $\beta_{rel} < 0$ (i.e., inverted beta profile) occurred if $2.405 < \bar{\gamma} < 3.736$. If $\bar{r} = 0$ is excluded from the

domain so the configuration is *doubly connected*, the singular nature of $Y_1(\bar{\gamma}\bar{r})$ at $\bar{r} = 0$ is no longer a constraint and the more general solution given by Eqs. (21), (27a), and (27b) can be used. The consequence of imposing $\sigma_Y = 0$ was for Eq. (27b) to force the relationship between Q and $\bar{\gamma}$ given by Eq. (28). If σ_Y is not forced to be zero, then this relationship between Q and $\bar{\gamma}$ is no longer imposed and the only remaining condition is that the domain must exclude $\bar{r} = 0$.

Consideration of Eq. (21) and recalling the discussion that led to Eq. (50) shows that $\bar{\psi}(\bar{r}_{lc}, \bar{z})$ is independent of $\bar{z}$ at $\bar{r}_{lc}$ where $\bar{r}_{lc}$ is now defined by

$$\sigma_J J_1(\bar{\gamma}\bar{r}_{lc}) + \sigma_Y Y_1(\bar{\gamma}\bar{r}_{lc}) = 0. \quad (60)$$

Thus Eq. (60) provides a radial shift of the location of the last closed flux surface and generalizes the discussion that led to Eq. (50). Because σ_J and σ_Y depend on $\bar{\gamma}$ and on Q (hence on β_{rel}), Eq. (60) shows that $\bar{r}_{lc}$ depends on both $\bar{\gamma}$ and β_{rel} . However, by assumption $\bar{r}_{lc} > 1$ (last closed flux surface radius is to the right of the magnetic axis) which restricts the allowed values of $\bar{\gamma}$ and β_{rel} . Introduction of the $Y_1(\bar{\gamma}\bar{r})$ solution and the coefficients σ_J and σ_Y is thus analogous to generalizing the solution of some harmonic equation from being $\sin(kx)$ to being $\sin(kx + \Delta) = \sin(kx)\cos(k\Delta) + \cos(kx)\sin(k\Delta)$ where $\sin(kx)$, $\cos(kx)$ are the analogs of $J_1(\bar{\gamma}\bar{r})$, $Y_1(\bar{\gamma}\bar{r})$ and σ_J , σ_Y are the analogs of $\cos(k\Delta)$, $\sin(k\Delta)$. Introducing finite Δ changes the phase of the solution and shifts the location of the solution.

Substitution for σ_J and σ_Y in Eq. (60) using Eqs. (27a) and (27b) gives

$$Q(\bar{\gamma}, \bar{r}_{lc}) = \frac{Y_0(\bar{\gamma})J_1(\bar{\gamma}\bar{r}_{lc}) - J_0(\bar{\gamma})Y_1(\bar{\gamma}\bar{r}_{lc})}{Y_2(\bar{\gamma})J_1(\bar{\gamma}\bar{r}_{lc}) - J_2(\bar{\gamma})Y_1(\bar{\gamma}\bar{r}_{lc})}. \quad (61)$$

It is seen that Eq. (61) reduces to Eq. (28) if $J_1(\bar{\gamma}\bar{r}_{lc}) = 0$, i.e., the situation considered in Sec. VIA and that Q becomes infinite when $\bar{r}_{lc}$ is such that the denominator in the right hand side of Eq. (61) vanishes.

Using Eqs. (60) in Eq. (21), it is seen that the last closed flux surface is given by

$$\bar{\psi}_{lc} = -\bar{r}_{lc}^2 Q \quad (62)$$

and inserting this in Eq. (22) gives

$$\beta_{rel} = Q + \bar{r}_{lc}^2 Q^2. \quad (63)$$

The derivative of Eq. (63) shows that the minimum possible β_{rel} is $\beta_{rel} = -1/(4\bar{r}_{lc}^2)$ which occurs when $Q = -1/(2\bar{r}_{lc}^2)$; this generalizes Eq. (54).

Figure 7 plots the dependence of Q and β_{rel} on $\bar{r}_{lc}$ for $1 < \bar{r}_{lc} < 2$ with $\bar{\gamma} = 2.5$; it is seen that, as predicted, β_{rel} has a minimum at $\beta_{rel} = -1/(4\bar{r}_{lc}^2)$ which occurs when $Q = -1/(2\bar{r}_{lc}^2)$. It is also seen from this figure that when $\bar{r}_{lc} = x_{11}/\bar{\gamma} = 1.5328$ the Sec. VIA result $\beta_{rel} = -0.081$ and $Q = J_0(\bar{\gamma})/J_2(\bar{\gamma}) = -0.108$ is recovered. For this $\bar{\gamma} = 2.5$ value, the denominator in Eq. (61) vanishes when $\bar{r}_{lc} \rightarrow 1.885$.

The following chain of dependence thus exists for doubly connected configurations:

FIG. 7. Black solid line is plot of $Q(\bar{\gamma}, \bar{r}_{lc})$ as function of $\bar{r}_{lc}$ with $\bar{\gamma} = 2.5$; red dashed line is plot of β_{rel} . For this $\bar{\gamma}$, Q diverges when $\bar{r}_{lc} = 1.885$. The equilibrium changes from bubble-like (negative β_{rel}) to conventional (positive β_{rel}) at $r_{lc} = 1.573$.

- (1) Independent values for $\bar{r}_{lc}$ and $\bar{\gamma}$ can be selected which then determine Q via Eq. (61),
- (2) Using Eqs. (27a) and (27b) in Eq. (21) the flux function $\bar{\psi}(\bar{r}, \bar{z})$ is given by

$$\bar{\psi}(\bar{r}, \bar{z}) = \frac{\pi}{2} \bar{\gamma} \bar{r} \left\{ \begin{array}{l} [Y_0(\bar{\gamma}) - QY_2(\bar{\gamma})]J_1(\bar{\gamma}\bar{r}) \\ + [-J_0(\bar{\gamma}) + QJ_2(\bar{\gamma})]Y_1(\bar{\gamma}\bar{r}) \end{array} \right\} \times \cos(\bar{k}\bar{z}) - \bar{r}^2 Q, \quad (64)$$

- (3) β_{rel} is given by Eq. (63),
- (4) ψ_{lc} is given by Eq. (62).

This chain of dependence for doubly connected configurations differs from that of a finite β spheromak. Specifically the chain of dependence for a finite β spheromak is: $\bar{\psi}_{lc} = 0$ is imposed because of the singly connected topology, $\bar{r}_{lc}$ is determined from setting the left hand side of Eq. (35) to zero on the midplane, and Eq. (32) gives $\beta_{rel} = J_0(\bar{\gamma})/J_2(\bar{\gamma})$.

Another and equivalent point of view differentiating singly and doubly connected configurations from each other is the following:

- (i) because the midplane of a singly connected configuration contains $\bar{r} = 0$ and because $\bar{\psi} = 0$ at $\bar{r} = 0$, the last closed flux surface for a singly connected configuration must always have $\bar{\psi} = 0$, whereas in contrast,

- (ii) because the midplane of a doubly connected configuration excludes $\bar{r} = 0$, the last closed flux surface of a doubly connected configuration cannot be $\bar{\psi} = 0$ as such a flux surface would have to pass through $\bar{r} = 0$.

The magnetic field components associated with Eq. (64) normalized to $B_{\phi,axis} = \bar{\lambda}\psi_{axis}/(2\pi r_{axis}^2)$ are

$$\bar{B}_r(\bar{r}, \bar{z}) = \frac{\bar{k}\pi\bar{\gamma}}{\bar{\lambda}} \left\{ \begin{array}{l} [Y_0(\bar{\gamma}) - QY_2(\bar{\gamma})]J_1(\bar{\gamma}\bar{r}) \\ + [-J_0(\bar{\gamma}) + QJ_2(\bar{\gamma})]Y_1(\bar{\gamma}\bar{r}) \end{array} \right\} \sin(\bar{k}\bar{z}), \quad (65a)$$

$$\bar{B}_\phi(\bar{r}, \bar{z}) = \frac{\pi\bar{\gamma}}{2} \left\{ \begin{array}{l} [Y_0(\bar{\gamma}) - QY_2(\bar{\gamma})]J_1(\bar{\gamma}\bar{r}) \\ + [-J_0(\bar{\gamma}) + QJ_2(\bar{\gamma})]Y_1(\bar{\gamma}\bar{r}) \end{array} \right\} \cos(\bar{k}\bar{z}) - \bar{r}Q, \quad (65b)$$

$$\bar{B}_z(\bar{r}, \bar{z}) = \frac{1}{\bar{\lambda}} \left(\frac{\pi\bar{\gamma}^2}{2} \left\{ \begin{array}{l} [Y_0(\bar{\gamma}) - QY_2(\bar{\gamma})]J_0(\bar{\gamma}\bar{r}) \\ + [-J_0(\bar{\gamma}) + QJ_2(\bar{\gamma})]Y_0(\bar{\gamma}\bar{r}) \end{array} \right\} \cos(\bar{k}\bar{z}) - 2Q \right). \quad (65c)$$

Using Eqs. (24c) and (26), it is seen that Eq. (65) reverts to Eq. (57) when $Q = J_0(\bar{\gamma})/J_2(\bar{\gamma})$.

ACKNOWLEDGMENTS

This material was based upon work supported by the U.S. Department of Energy Office of Science, Office of Fusion Energy Sciences under Award Nos. DE-FG02-04ER54755 and DE-SC0010471, by the National Science Foundation under Award No. 1059519, and by the Air Force Office of Scientific Research under Award No. FA9550-11-1-0184. The authors wish to thank an anonymous reviewer for making the valuable suggestion that Y_n Bessel solutions be allowed when $\bar{r} = 0$ is excluded from the domain.

¹P. M. Bellan, *Phys. Plasmas* **9**, 3050 (2002).

²B. U. O. Sonnerup, H. Hasegawa, W. L. Teh, and L. N. Hau, *J. Geophys. Res.-Space Phys.* **111**, A09204, doi:10.1029/2006JA011717 (2006).

³Q. Hu, C. J. Farrugia, V. A. Osherovich, C. Mostl, A. Szabo, K. W. Ogilvie, and R. P. Lepping, *Sol. Phys.* **284**, 275 (2013).

⁴A. J. Cerfon and J. P. Freidberg, *Phys. Plasmas* **17**, 032502 (2010).

⁵P. M. Bellan, *Spheromaks: A Practical Application of Magnetohydrodynamic Dynamos and Plasma Self-Organization* (Imperial College Press, London, 2000).

⁶H. Grad and H. Rubin, in *Proceedings of the 2nd U.N. Conference on the Peaceful uses of Atomic Energy* (IAEA, Vienna, 1958), vol. 31, p. 190.

⁷V. D. Shafranov, in *Reviews of Plasma Physics* (Consultants Bureau, New York, 1966), Vol. 2, p. 103.

⁸P. Gautier, R. Gruber, and F. Troyon, *Nucl. Fusion* **21**, 1399 (1981).