

# Twenty *Drosophila* visual system cDNA clones: One is a homolog of human arrestin

(eye/retina/photoreceptor/phototransduction/*Arr* gene)

DAVID R. HYDE\*<sup>†</sup>, KIRK L. MECKLENBURG\*<sup>‡</sup>, JOHN A. POLLOCK\*<sup>§</sup>, THOMAS S. VIHTELIC\*,  
AND SEYMOUR BENZER\*<sup>¶</sup>

\*Department of Biological Sciences, University of Notre Dame, Notre Dame, IN 46556; and <sup>‡</sup>Division of Biology, 156-29, California Institute of Technology, Pasadena, CA 91125

Contributed by Seymour Benzer, November 6, 1989

**ABSTRACT** From a group of 436 *Drosophila melanogaster* cDNA clones, we selected 39 that are expressed exclusively or predominantly in the adult visual system. By sequence analysis, 20 of the clones appear to represent previously unreported distinct cDNAs. The corresponding transcripts are detected in the retina and optic lobes. The genes are scattered throughout the genome, some near mutations known to affect eye function. One of these clones has been identified, by sequence analysis, as the structural gene (*Arr*) for a *Drosophila* homolog of human arrestin. Vertebrate arrestin interacts with rhodopsin in phototransduction and has been associated with an autoimmune form of uveitis in primates. The presence of an arrestin homolog in *Drosophila* suggests that both the vertebrate and invertebrate phototransduction cascades are regulated in a similar manner.

Previous molecular screens have identified important *Drosophila* eye-specific molecules that are required for proper eye development and phototransduction. Levy *et al.* (1) isolated 20 genomic clones that hybridized to transcripts more highly expressed in head than in body. Seven of these were shown to be eye-specific, including the genes for opsin Rh3, transient receptor potential (*trp*), and neither inactivation nor afterpotential C (*ninaC*); (2–4). Shieh *et al.* (5) identified eye-specific genes using, as a probe, total wild-type head cDNA that was subtracted with poly(A)<sup>+</sup> RNA from bodies and heads of the mutant eyes absent (*eya*). The *eya* mutation removes the eyes and reduces the optic lobes but leaves the light-detecting ocelli intact (6). Their screen identified many eye-specific genes, including *ninaA*, a cyclophilin homolog (5), and an eye-specific protein kinase C (7).

Recently, 436 *Drosophila* cDNA clones corresponding to genes expressed predominantly in the head but not in the early embryo were isolated, and their expression patterns described (8). Comparison of Northern (RNA) blots of RNA from wild-type and from *eya* heads led to the identification of 39 clones, described here, that correspond to mRNAs expressed in wild-type but reduced in *eya* heads. By DNA sequencing, we have identified 20 of the clones as being cDNAs that are distinct from each other and do not correspond to previously reported cDNAs. In addition, four previously sequenced genes expressed in the retina were also found in our collection, some more than once, confirming the validity of the approach. One of the new clones<sup>||</sup> corresponds to a *Drosophila* gene with strong sequence similarity to the vertebrate phototransduction molecule arrestin. The *Drosophila* arrestin gene has been independently cloned by Smith *et al.* (9).

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

## MATERIALS AND METHODS

**DNA Purification, Isolation of RNA, Northern Blots, and Southern Blots.** These were performed as described (8).

**DNA Sequence Analysis.** cDNA inserts were isolated by digesting phage DNA with *EcoRI* and *Xba I* and subcloning into either SK<sup>+</sup> or KS<sup>+</sup> Bluescript vectors (Stratagene). Either single-stranded phage DNA or double-stranded plasmid templates were sequenced with the United States Biochemical sequencing kit.

**Isolation of Longer cDNA and Genomic Clones.** Longer cDNAs were isolated by screening the unsubtracted *Drosophila* λ SWAJ2 head cDNA library (8) with radiolabeled probes (10). Over 50,000 plaques were screened for each clone tested. Positives were picked, replated, reprobated, plaque purified, and confirmed by Southern blots. Genomic clones were isolated from the Maniatis *Drosophila* genomic library (11).

**Tissue *in Situ* Hybridization.** Third instar larvae, prepupae, or adult flies were frozen in Tissuetek OCT embedding compound on dry ice and equilibrated at –20°C. Frozen sections were cut at 6 μm and processed for hybridization by the method of Hafen and Levine (12), omitting the Pronase step. Nick-translated, <sup>3</sup>H-labeled DNA probes were prepared with all four nucleotides labeled to a specific activity of about 5 × 10<sup>7</sup> cpm/μg.

**Chromosomal *in Situ* Mapping.** Biotinylated DNA probes were hybridized to larval salivary chromosomes as in ref. 13, except that the acetic anhydride treatment of the slides was omitted. The probes were λ SWAJ2 or SWAJ3 (8) cDNAs subcloned into Bluescript. Plasmids were randomly primed to incorporate Bio-16-dUTP (Enzo Biochemicals), or <sup>35</sup>S-labeled dATP and ethanol precipitated with 1/10th vol of 3M ammonium acetate. The hybridization buffer contained dextran sulfate at a final concentration of 10% (wt/vol).

**Materials.** T4 DNA polymerase, *EcoRI*, and *Xba I* were purchased from New England Biolabs. RNasin, T7 RNA polymerase, SP6 RNA polymerase, and cRNA transcription systems were from Promega Biotec. *Spe I* restriction enzyme and helper phage were from Stratagene. Radioactive nucleotide triphosphates, Hybond-N paper, and cDNA cloning kits were from Amersham. All other enzymes and oligo(dT)-cellulose were from Boehringer Mannheim.

## RESULTS

**Identification of Visual System-Specific Clones.** We identified genes that are expressed predominantly in the eye by

<sup>†</sup>These authors contributed equally to this work.

<sup>§</sup>Present address: Department of Biological Sciences, Carnegie Mellon University, 4400 Fifth Avenue, Pittsburgh, PA 15213.

<sup>¶</sup>To whom reprint requests should be addressed.

<sup>||</sup>The sequence reported in this paper has been deposited in the GenBank data base (accession no. M30140).




FIG. 1. RNA gel blots of poly(A)<sup>+</sup> RNA probed with different radioactive cRNAs to identify eye-specific transcripts. In each case, RNA was from wild-type heads (a), heads of *eya* (b), or wild-type bodies (c). Each lane contained 0.5  $\mu$ g of poly(A)<sup>+</sup> RNA. The cRNA probes in these examples recognized transcripts of various sizes, reduced in the *eya* mutant, and not detected in wild-type bodies.

determining which wild-type head cDNAs hybridized to transcripts that were reduced or undetectable in an eyeless mutant. Antisense cRNA was synthesized *in vitro* from each

of 436 head-not-early embryo *Drosophila* cDNA clones and used to probe Northern blots of poly(A)<sup>+</sup> RNA extracted separately from wild-type heads, *eya* mutant heads, and wild-type bodies (8). Body RNA was included as a negative control because the adult visual system is restricted to the head. Thirty-nine cDNAs detected transcripts expressed in wild-type heads that were reduced in *eya* heads and not detected in bodies. Examples of the various expression patterns are illustrated in Fig. 1; transcript sizes in the examples shown ranged from 0.4 to 2.9 kilobases (kb). Clone 7F10 is an example of a cDNA with a signal that is still detectable in *eya* but at a reduced level, suggesting that the expression of the corresponding gene is not restricted to the retina.

**The Corresponding Genes Are Located at Many Sites.** The cDNA clones in the original collection were rather short; therefore, homologous, longer cDNAs and genomic clones were isolated for hybridization *in situ* to third instar larval polytene chromosomes. The genes are scattered throughout the genome (Table 1), with some near to, or coincident with, previously identified mutant eye loci. Sixteen of the original 39 clones mapped to single bands; two mapped to multiple bands, suggesting that they correspond to repeated elements in the genome.

**Identification of Previously Reported Genes.** The 39 putative visual system cDNAs were tested for homology with previously cloned eye-specific genes. Eleven hybridized with the *Drosophila* gene *ninaE* (structural gene for opsin *Rh1*) (13, 14), two hybridized to opsin *Rh4* (15), and one each with clones 549 and 557 of Levy *et al.* (1). The fact that our screen for eye clones detected these previously identified eye-specific molecules demonstrated the effectiveness of the screening procedure. Our failure to detect opsin *Rh2* was consistent with the fact that it is expressed only in the ocelli (16–18), which are intact in the *eya* mutant.

**Sequence Analysis Identifies Other Distinct cDNAs.** We sequenced 25 clones that did not cross-hybridize with *ninaE*. Each contained a poly(A)<sup>+</sup> stretch corresponding to the 3' end of a directionally cloned transcript and a consensus

Table 1. Twenty distinct eye cDNA clones

Clone	Expression onset	Transcript size, kb	Approximate gene location	Tissue <i>in situ</i>
7F10	24-hr embryo	2.4	66BC	Retina, cortex
6H8	Late 2nd instar	0.4	46F–47A	Negative
9C5	25-hr pupa	9.2	93AB	ND
10A2	25-hr pupa	1.3	95AB	ND
2D3	50-hr pupa	1.8	2D–F	Retina
2G1	50-hr pupa	3.6	46E	ND
9C4	50-hr pupa	0.9	83AB	Retina
1C2	After 75-hr pupa	Smear	Multiple sites	ND
2G8	After 75-hr pupa	0.8	ND	ND
9H9	After 75-hr pupa	2.9	10E	Retina
<b>3H10</b>	<b>After 75-hr pupa (arrestin)</b>	<b>1.4</b>	<b>36D</b>	<b>Retina, ocellus, larval photoreceptor</b>
5B3	After 75-hr pupa	2.9	72D	Retina, cortex
4A11	After 75-hr pupa	0.8	72E	Retina, cortex
3C12	After 75-hr pupa	0.8	62DE	ND
8H7	After 75 hr pupa	0.7	67EF	Negative
1D10	After 75 hr pupa	0.9	17A, 86F	ND
6A7	After 75 hr pupa	1.6	ND	Retina, cortex
6C5	After 75 hr pupa	1.6	ND	Negative
6F7	After 75 hr pupa	0.8	ND	Negative
7H1	After 75 hr pupa	0.8	ND	Retina

ND, not determined. Transcript size and expression onset were determined by Northern blots. Onset is listed as the earliest time a transcript could be detected by this method. Gene localization was determined by hybridization to salivary gland polytene chromosomes. Some clones hybridized to more than one site on the chromosomes, possibly due to repeated sequences in the genome. Tissue *in situ* expression was determined in cryostat sections of heads by using <sup>3</sup>H-labeled cDNA probes. A negative result in the tissue *in situ* experiment indicates that the transcript was difficult to detect with that particular probe.




FIG. 2. Expression of the arrestin-homologous RNA in the three distinct visual organs of the fly. The endogenous transcript pattern is revealed autoradiographically in 6- $\mu$ m cryostat sections after *in situ* hybridization with  $^3$ H-labeled probe made from 3H10 cDNA. Phase illumination shows the tissue morphology; darkfield illumination of the same field of view shows the silver grain signal. (A and B) Adult compound eye in radial section of 1-day-old adult. (A) Phase; R, retina; L, lamina; M, medulla. (B) Darkfield, indicating abundant RNA expression in the retina. (Bar = 50  $\mu$ m.) (C and D) Three adult ocelli, in a grazing tangential section of the dorsal head of a 1-day-old adult. (C) Phase shows the retinal layers (OR1, OR2, and OR3) of the three ocelli, surrounded by head cuticle (CU). (D) In darkfield, white arrows indicate the positions of the ocelli. The arrestin RNA is expressed, but at a reduced level compared with the compound eye. (Bar = 25  $\mu$ m.) (E and F) Bolwig's organs in horizontal section of a third instar larva. E reveals the two clusters of larval photoreceptor neurons (BO). They lie within the cephalopharyngeal skeleton, just anterior to the indicated parts of the mouth hooks (MH) at the base of the protocerebral bridge. In the darkfield view, white arrows indicate the photoreceptor organs, which show robust expression of the arrestin RNA. (Note: The apparent signals in CU and MH are artifactual; the probe tends to stick to cuticular material.) (Bar = 25  $\mu$ m.)

poly(A)<sup>+</sup> addition sequence. Each sequence was compared with all the others and with the GenBank nucleic acid data base. Two cDNAs corresponded to *norPA* (no receptor potential) (19) and one to opsin *Rh3* (2, 20). Three clones were identical but not previously reported. This yielded 20 distinct, not previously reported *Drosophila* cDNAs (Table 1).

1	AATCGATTTC	TCGAGTATT	TCGTGACATT	CACCTGTCTT	GCATTGCTAA	TACGCCGTGT
61	TGCTCGCGGC	TTATTTAATG	TTTGACCGAT	CGATGCGCAT	CTCGTCTGTC	AATGTCARAT
121	TCAAAGACAC	TGCCACACG	AAGGTTCCCG	AGCCCGTAGT	AGTTTACCCG	CGTGCAATTG
181	CTGAATTTCCT	GCTGTGCGGC	TAAATGAATT	TAGAGGGGGG	ACAGGTCCCA	CAATCCAGG
11 bp conserved sequence						
241	TATAAATGCC	GGATTGCCAA	AGAGCGCTAA	TTAATAGCCT	AGTGGACACC	GCAACCGCGC
TATA box      TATA box      TATA box      TATA box      TATA box      TATA box						
301	GTATACCATC	GAGACGAGC	CGGAAACGAT	AAAGGCACAT	CCAAGCTTTA	AACTATTTC
361	GCAGAGATT	TGATAAACAG	CTCCAAAATG	CTGGTCAATT	TCAAGGTGTT	CAGAAGTGT
			Met	ValValAsnP	heLysValIph	eLysLysCys
421	TCCCGAACA	ACATGATCAC	GCTCTACATG	AACAGCGCTG	ATTTGTAGA	TCCGTGTAG
	SerProAsnA	snMetIleTh	rLeuTyrMet	AsnArgArgA	spPheValAs	pSerValThr
481	CAGGTGGAC	CCATTGGTAG	GTGTACAGA	CCGAAACCTT	TGAGCAATGG	GATTTACGAA
	GlnValGluP	rolIleA				
541	TGAGGAAATC	CATCAAAAA	TAAATTCGTG	TAGGAATGG	TACCCATATT	CGATTGAAGT
601	ATCTTATAGT	TTGAAAATA	CTTCAGTGTA	ACTTTTGT	CAACTTAA	CAITGGAATT
661	TTTAAATGCT	TCCTTGAAA	GTGATATCAA	ATCAAAATTA	TATTATAAAA	ACTCCATTTC
721	GAATCTGACT	ATGCCGACCA	GCAAAATATA	GCAATATAAA	TATTCAAAA	TTCGACTGTA
781	ATTCAATAGC	CTAGGGAGTT	AAGTAAAT	GGCTGCGCTA	CGAGTAACT	AACTTCATG
841	ATTACGGCAG	CAAGCAAAAC	AAGAGCGTGG	CGCTTGTAA	ATAATATTCT	CCAAATGTA
901	AGCGTCTCT	TGCAGATGGA	ATCATTGTGC	TGGACGATGA	GTACGTGGC	CAGAACCGCA
		spGly	IleIleValI	euAspAspG	uTyrValArg	GlnAsnArgL
961	AGATCTTGGT	GCAGTTGGT	TGCAATTTC	GATATGGGG	CGAGGAGAC	GAGATGATG
	ysIlePheVa	IgInLeuVal	CysAsnPheA	rgTyrGlyAr	GluuAspAsp	GluMetIleG
1021	GTCGCGGTT	CCAGAAGGAA	CTGACCCTGG	TCTCGCAGCA	GGTGTGCCA	CCCCAAGC
	lyLeuArgPh	eGlnLysGlu	LeuThrLeuV	alSerGlnG	nValCysPro	ProGlnLysG
1081	AGGACATCCA	GTTGACCAAG	ATGCAGGAGC	GTCTGCTGAA	GAAGCTGGC	TCCATGCTT
	InAspIleG	nLeuThrLys	MetGlnGluA	rgLeuLeuL	sLysLeuGly	AsnAsnIleA
1141	ATCCCTTGGT	GATGCAGATG	CCACCAAGCT	CGCCGGCTC	GGTGTCTCT	CAGCAGAGG
	yrProPheVa	IMetGlnMet	ProProSerS	erProAlaSe	rValValLeu	GlnGlnLysA
1201	CCAGTGAOGA	GAGCCAGCCC	TGCGGAGTCC	AGTACTTGGT	AAAGATCTTT	ACCGGAGACA
	laSerAspGl	uSerGlnPro	CysGlyValG	InTyrPheVa	ILysIlePhe	ThrGlyAspS
1261	GCGACTGCGA	TCGATGCGAT	CGCAGGAGCA	CCATTAACTT	GGGCATCGC	AAGGTGCGAT
	erAspCysAs	pArgSerHis	ArgArgSerT	hrIleAsnLe	uGlyIleArg	LysValGlnT
1321	AGCACCAGCA	CAAGCAGGCG	ATCCAGCCCT	GCACCGTGT	TGCAGGAGC	TTCCTTCTGT
	ryAlaPProTh	rLysGlnGly	IleGlnProC	ysThrValVa	lArgLysAsp	PheLeuLeuS
1381	CGCCGGAGA	GCTCGAAGTC	GAGTCCACC	TCGACAAGCA	GCTGTACCAT	CACGGCGAGA
	erProGlyGl	uLeuGluLeu	GluValThrL	euAspLysGl	nLeuTyrHis	HisGlyGluL
1441	AGATCTCGGT	GAACATCTGC	GTAOCGAACA	ACTCCAACA	GGTGGTGAAG	ANGATCAAGG
	IleSerVaI	lAsnIleCys	ValArgAsnA	snSerAsnLy	sValValLys	LysIleLysA
1501	CCATGGTGCA	GCAGGGCGTC	GATGTGGTCC	TTTTCCAGAA	CGGTCACTTC	CGCAACAAGA
	laMetValGl	nGlnGlyVal	AspValValL	euPheGlnAs	nGlyGlnPhe	ArgAsnThrI
1561	TGCGCTTCAT	GGAGCAGGAG	GAGGGATGTC	CCCTGAACCC	GGGATCCAGC	CTGCAGAAGG
	IleAlaPheTh	tGluThrSer	GluGlyCysP	roLeuAsnPr	oGlySerSer	LeuGlnLysV
1621	TCATGTATCT	GGTGCCCAAC	CTGGTGGCCA	ATTGCGACCG	CGCAGGCATC	GCCGTGAGG
	alMetTyrLe	uValProThr	LeuValAlaA	snCysAspAr	gAlaGlyIle	AlaValGluG
1681	GTGATATCAA	GCGCAAGAC	ACAGCTCTGG	CCTCGACCAC	ACTGTGAGTA	AAATTTATTC
	lyAspIleL	sArgLysAsp	ThrAlaLeuA	laSerThrTh	rIle	
1741	ACATCATAGC	TTAGCAGATG	AAACATTAAT	ATTATACTCT	ATTAAGTATC	AACTTAAAT
1801	CATACCATAA	AATCAATCAA	ATTTTAAAGT	TAGGAACCTT	TTTAAAAATC	GTATTTTCCC
1861	GGTGACTAAC	AGTCTTTTAC	CTAAATGTGT	TTACAAAATG	GCATMAAACG	CATACTAATA
1921	CTAATGTAAA	AATGCATATT	TAAAAATCTA	TTTCAGTATT	GCCAGTCAGG	ATGCGAGGGA
				uile	AlaSerGlnA	spAlaArgAs
1981	TGCCCTTGGC	ATAATTGTTT	CATATGCTGT	GAAGGTCAAG	CTTTTCTTGG	GAGCCCTGGG
	pAlaPheGly	IleIleValS	erTyrAlaVa	ILysValLys	LeuPheLeuG	lyAlaLeuGl
2041	CGCGGAGCTC	TGCGCTGAGC	TACCATTAT	TCTGATGCAC	CCGAAGGTTA	TAAAAGGTGT
	gYlGluLeu	CysAlaGluL	euProPheIl	eLeuMetHis	ProLys	
2101	GCCCAATAT	TTGAATAGTT	ATTGAAAGGC	AATCAATTAT	TTACAGCCAA	GTCCGAAGGC
					ProS	erArgLysAl
2161	CCAACTGGAA	GCCGAGGGCT	CCATTGAGGC	CTAAACTGAA	AGGGCTACCT	CAACCAACGA
	eGlnLeuGlu	AlaGluGlyS	eIleGluAl	a		
2221	AAAAAATGCC	GTATTCTAC	AAGTCAACCC	GATTTTGTGA	GATCCTAAAA	AATGCTGATG
2281	TTGCTGAAAT	GTCTGAACT	GCAGTCTGCT	TACTTTTCTT	ATATAGCAAA	TCCAATATCC
2341	ATATATTGTA	TGTTGTATG	TGTTATATAT	TTATAACTCA	ACTAAACAATT	AAATATGAC
					poly(A) <sup>+</sup> add site	
2401	AGAGTTTATG	T				

FIG. 3. Nucleotide sequence of the *Drosophila* arrestin gene. The 11-bp conserved sequence found in the promoters of genes expressed specifically in *Drosophila* photoreceptor cells (22), TATA box, transcription initiation site, and putative polyadenylation sites are shown. The transcription start site indicates the major primer-extension product that we detect. Introns are indicated as interruptions in the deduced amino acid sequence. A poly(A)<sup>+</sup> tract after nucleotide 2411 was found in the cDNA.

**Temporal Expression.** From each of the 20 distinct clones, antisense cRNA was synthesized and used to probe Northern blots of poly(A)<sup>+</sup> RNA isolated from various developmental stages. Several different patterns were observed. For example, the transcript detected by clone 6H8 was seen as early as the late second larval instar. Clone 2D3 first detected a transcript in the 50-hr pupa. Clone 7F10 identified an RNA expressed at low levels in all the stages tested. Two cDNAs



less (C. Zuker, personal communication). Moreover, many of the genes are represented by a single clone. Thus, it is clear that many more eye-specific genes remain to be found.

In certain cases, tissue *in situ* hybridization analysis proved more sensitive than Northern blots. This can be true when the gene is expressed in a relatively small number of cells, as is the case for the cDNAs expressed in the larval photoreceptor organ (Fig. 2). The organ consists of only two dozen photoreceptor cells in the anterior of the animal, and RNA extracted from whole larvae will contain only a small contribution from these cells.

Based upon their sequence similarity and tissue-specific expression, the *Drosophila Arr* gene product and vertebrate arrestin may be functional homologs, both molecules negatively regulating the transduction process. Data suggest that bovine arrestin binds to phosphorylated metarhodopsin II, thus inhibiting the transduction cascade (26, 27). Rhodopsin in flies is also phosphorylated upon light activation (28, 29), which further suggests that *Drosophila* arrestin and rhodopsin binding may be similar to vertebrates. It is of interest to note that the *Drosophila* homolog lacks the 31 C-terminal amino acids of human arrestin, suggesting that this segment does not have a necessary function, or that the truncated *Drosophila* molecule functions by a slightly different mechanism than the vertebrate molecule.

Several *Drosophila* visual system-specific genes show sequence similarity with genes from vertebrate species, reflecting possible similarities in function. The phototransduction cascade is initiated by the absorption of photons by chromophores bound to opsin proteins. The opsin molecules have both *Drosophila* and vertebrate counterparts (13, 14, 30). The identification of a *Drosophila* arrestin molecule makes it possible to examine the regulation of the invertebrate phototransduction cascade, which may improve our understanding of the process in vertebrates. The availability of many distinct eye-specific *Drosophila* cDNAs will allow us to investigate the extent to which these molecules are conserved between species.

Recently, a technique for accessing the function of the gene corresponding to a cDNA clone has been developed for the *Drosophila* system by Ballinger and Benzer (31) and Kaiser and Goodwin (Kaiser, K. and Goodwin, S. F., personal communication). The technique consists of creating large numbers of mutant flies with *P* element insertions. Individual flies with transposon inserts in or near the cloned gene of interest are identified by polymerase chain reaction amplification by using oligonucleotide primers derived from the sequences of the *P* element and the cDNA in question. This method can be applied to the clones described here; this may allow use of the fly system to identify the functions of conserved genes through specific mutations.

We thank C. Zuker and coworkers for communicating results before publication and for critical review of the manuscript. We thank Seema Shah for isolating the arrestin genomic clones and characterizing the intron/exon relationship. We also thank Christine Pham for excellent technical assistance, Fintan Steele for confirming the map positions of clones 4A11, 5B3, and 9H9, and G. Guttman for the sequence analysis program. We also thank our colleagues for their constant stimulation. This work was supported by National Science Foundation Grant DCB8409366 and National Institutes of Health Grant GM40499 to S.B., National Eye Institute Grant

EY08058 and the University of Notre Dame Jesse H. Jones Faculty Research and Equipment Fund to D.R.H., a Developmental Biology Grant from the Lucille P. Markey Charitable Trust, and National Institutes of Health Postdoctoral Fellowships from the National Eye Institute to D.R.H., K.L.M., and J.A.P.

1. Levy, L. S., Ganguly, R., Ganguly, N. & Manning, J. (1982) *Dev. Biol.* **94**, 451–464.
2. Fryxell, K. J. & Meyerowitz, E. M. (1987) *EMBO J.* **6**, 443–451.
3. Montell, C., Jones, K., Hafen, E. & Rubin, G. (1985) *Science* **230**, 1040–1043.
4. Montell, C. & Rubin, G. (1988) *Cell* **52**, 757–772.
5. Shieh, B.-H., Stamnes, M. A., Seavello, S., Harris, G. L. & Zuker, C. S. (1989) *Nature (London)* **338**, 67–70.
6. Sved, J. (1986) *Drosoph. Inf. Serv.* **63**, 169.
7. Schaeffer, E., Smith, D., Mardon, G., Quinn, W. & Zuker, C. (1989) *Cell* **57**, 403–412.
8. Palazzolo, M. J., Hyde, D. R., VijayRaghavan, K., Mecklenburg, K., Benzer, S. & Meyerowitz, E. (1989) *Neuron* **3**, 527–539.
9. Smith, D. P., Shieh, B.-H. & Zuker, C. (1990) *Proc. Natl. Acad. Sci. USA* **87**, 1003–1007.
10. Maniatis, T., Fritsch, E. F. & Sambrook, J. (1982) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab., Cold Spring Harbor, NY).
11. Maniatis, T., Hardison, R. C., Lacy, E., Lauer, J., O'Connell, C., Quon, D., Sim, G. K. & Efstratiadis, A. (1978) *Cell* **15**, 687–701.
12. Hafen, E. & Levine, M. (1986) in *Drosophila: A Practical Approach*, ed. Roberts, D. B. (IRL, Washington, DC), pp. 139–157.
13. Zuker, C. S., Cowman, A. F. & Rubin, G. M. (1985) *Cell* **40**, 851–858.
14. O'Tousa, J. E., Baehr, W., Martin, R. L., Hirsh, J., Pak, W. L. & Applebury, M. L. (1985) *Cell* **40**, 839–850.
15. Montell, C., Jones, K., Zuker, C. & Rubin, G. (1987) *J. Neurosci.* **7**, 1558–1566.
16. Pollock, J. A. & Benzer, S. (1988) *Nature (London)* **333**, 779–782.
17. Feiler, R., Harris, W. A., Kirschfeld, K., Wehrhahn, C. & Zuker, C. (1988) *Nature (London)* **333**, 737–741.
18. Mismer, D., Michael, W. M., Laverty, T. R. & Rubin, G. M. (1988) *Genetics* **120**, 173–180.
19. Bloomquist, B., Shortridge, R., Schneuwly, S., Perdew, M., Montell, C., Steller, H., Rubin, G. & Pak, W. L. (1988) *Cell* **54**, 723–733.
20. Zuker, C. S., Montell, C., Jones, K., Laverty, T. & Rubin, G. M. (1987) *J. Neurosci.* **7**, 1550–1557.
21. Waddington, C. H. & Perry, M. M. (1960) *Proc. R. Soc. London Ser. B* **153**, 155–178.
22. Misner, D. & Rubin, G. M. (1989) *Genetics* **121**, 77–87.
23. Kozak, M. (1984) *Nucleic Acids Res.* **12**, 857–872.
24. Shinohara, T., Dietzschold, B., Craft, C. M., Wistow, G., Early, J. J., Donoso, L. A., Horwitz, J. & Tao, R. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 6975–6979.
25. Yamaki, K., Tsuda, M. & Shinohara, T. (1988) *FEBS Lett.* **234**, 39–43.
26. Kühn, H., Hall, S. W. & Wilden, U. (1984) *FEBS Lett.* **176**, 473–478.
27. Wilden, U., Hall, S. W. & Kühn, H. (1986) *Proc. Natl. Acad. Sci. USA* **83**, 1174–1178.
28. Paulsen, R. & Bontrop, J. (1983) *Nature (London)* **302**, 417–419.
29. Matsumoto, H. & Pak, W. L. (1984) *Science* **223**, 184–186.
30. Nathans, J. & Hogness, D. S. (1984) *Proc. Natl. Acad. Sci. USA* **81**, 4851–4855.
31. Ballinger, D. B. & Benzer, S. (1989) *Proc. Natl. Acad. Sci. USA* **86**, 9402–9406.