

Purification and characterization of sortase, the transpeptidase that cleaves surface proteins of *Staphylococcus aureus* at the LPXTG motif

Hung Ton-That[†], Gwen Liu[†], Sarkis K. Mazmanian[†], Kym F. Faull[‡], and Olaf Schneewind^{†§}

[†]Department of Microbiology and Immunology, and [‡]The Pasarow Mass Spectrometry Laboratory, Departments of Psychiatry and Biobehavioral Sciences, Chemistry and Biochemistry, and the Neuropsychiatric Institute, University of California, Los Angeles School of Medicine, 10833 Le Conte Avenue, Los Angeles, CA 90095

Edited by Christopher T. Walsh, Harvard Medical School, Boston, MA, and approved September 1, 1999 (received for review June 23, 1999)

Surface proteins of *Staphylococcus aureus* are linked to the bacterial cell wall by sortase, an enzyme that cleaves polypeptides at the threonine of the LPXTG motif. Surface proteins can be released from staphylococci by treatment with hydroxylamine, resulting in the formation of threonine hydroxamate. Staphylococcal extracts, as well as purified sortase, catalyze the hydroxylaminolysis of peptides bearing an LPXTG motif, a reaction that can be inhibited with sulfhydryl-modifying reagents. Replacement of the single conserved cysteine at position 184 of sortase with alanine abolishes enzyme activity. Thus, sortase appears to catalyze surface-protein anchoring by means of a transpeptidation reaction that captures cleaved polypeptides as thioester enzyme intermediates.

Surface proteins of *Staphylococcus aureus* are anchored to the bacterial cell wall by a mechanism requiring a C-terminal sorting signal with a conserved LPXTG motif (1). Cleavage between the threonine and the glycine of the LPXTG motif liberates the carboxyl of threonine to form an amide bond with the amino group of the pentaglycine crossbridge (2, 3), thereby tethering the C terminus end of surface proteins to the bacterial peptidoglycan (4–7). The LPXTG motif is conserved in more than 100 surface proteins of Gram-positive pathogens, suggesting that anchoring of these polypeptides occurs by a universal mechanism (1). Although surface protein anchoring can be followed by pulse-labeling polypeptides *in vivo* (8, 9), the transpeptidation reaction has thus far not been measured *in vitro*. Furthermore, sortase, the enzyme that is thought to catalyze this reaction, has not yet been purified and characterized.

To identify the sortase gene, we recently screened a collection of temperature-sensitive staphylococcal mutants for a defect in surface-protein anchoring (cell wall sorting) (10). A mutant *S. aureus* strain that displayed a severe sorting defect was transformed with a plasmid library of staphylococcal genomic DNA, and individual clones were screened for complementation of surface-protein anchoring. One gene, named *srtA* (surface protein sorting A), complemented the sorting defect of the temperature-sensitive variant (10). Overexpression of *srtA* in wild-type staphylococci was sufficient to increase the rate of surface-protein anchoring, suggesting that SrtA is involved in the cell wall sorting reaction (10). Nevertheless, this work left unresolved whether *srtA* encodes sortase, the enzyme that has been proposed to catalyze the transpeptidation reaction that links surface proteins to the staphylococcal cell wall.

Surface-protein anchoring in *S. aureus* can be inhibited with sulfhydryl-modifying reagents such as methanethiosulfonate and organic mercurial, indicating that sortase must be a sulfhydryl (cysteine)-containing enzyme (11). We wondered whether staphylococcal sortase captures surface proteins after their cleavage at the LPXTG motif as acyl-enzyme intermediates. In this model, the sulfhydryl of sortase may function as a nucleophile at the peptide bond between threonine and glycine, thereby forming a thioester with the carboxyl of threonine and releasing the amino group of the cleaved C-terminal-sorting signal (Fig. 1).

If so, an acyl-enzyme intermediate between surface proteins and sortase should be sensitive to hydroxylaminolysis (12, 13). Lipmann first used hydroxylamine (NH₂OH) to demonstrate the existence of acyl-enzyme intermediates, as this strong nucleophile attacks thioester bonds to form hydroxamate with carboxyl groups, thereby regenerating the enzyme active-site sulfhydryl (14). We show here that hydroxylamine treatment causes the formation of a C-terminal threonine hydroxamate of surface proteins, which are thereby released into the culture medium. Staphylococcal extracts, as well as purified SrtA, catalyze the hydroxylaminolysis of peptides bearing an LPXTG motif. Replacement of the single conserved cysteine with alanine abolished enzymatic activity, suggesting that sortase catalyzes a transpeptidation reaction by means of the formation of a thioester acyl-enzyme intermediate.

Experimental Procedures

Hydroxylaminolysis of Surface Proteins *in Vivo*. *S. aureus* OS2 (*spa*⁻:*ermC*) or *S. aureus* BB270 (wild type) carrying a plasmid encoding staphylococcal enterotoxin B (SEB) fused to a fragment of staphylococcal protein A (SPA)_{490–524} (5, 9) were grown in minimal medium until OD₆₀₀ reached 0.5. A 1-ml aliquot of the staphylococci culture (10⁹ cells) was pulse-labeled with 100 μCi of Pro-Mix for 1 min, chase solution (50 μl of 100 mg/ml casamino acids, 20 mg/ml methionine and cysteine) was added, and cells were incubated at 37°C for 5 min. A 0.5-ml aliquot was centrifuged at 15,000 × *g* for 5 min, and the supernatant was precipitated with 0.5 ml of 10% trichloroacetic acid (TCA). Another 0.5-ml culture aliquot was directly precipitated with 0.5 ml of 10% TCA and suspended in 1 ml of 0.5 M Tris-HCl (pH 7.0), and peptidoglycan was digested with 100 μg of lysostaphin (15) for 1 hr at 37°C. Proteins were precipitated with TCA, washed in acetone, dried and then boiled in SDS. Aliquots were subjected to immunoprecipitation with anti-SEB and analyzed by PAGE and PhosphorImager (Molecular Dynamics).

Purification of Surface Proteins. Staphylococci [10¹³ BB270(pSEB-MH₆-CWS) cells] (5) were incubated in 200 ml of 50 mM Tris-HCl (pH 7.0), with or without 0.1 M NH₂OH, for 60 min. Samples were centrifuged at 10,000 × *g* for 15 min, and the supernatants were subjected to affinity chromatography on a

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: TCA, trichloroacetic acid; SEB, staphylococcal enterotoxin B; Ni-NTA, nickel nitrilotriacetic acid; TFA, trifluoroacetic acid; MALDI-MS, matrix-assisted laser desorption ionization MS; ESI-MS, electrospray ionization MS; DabcyI, 4-(4-dimethylaminophenylazo)benzoic acid; Edans, 5-[(2-aminoethyl)amino]naphthalene-1-sulfonic acid; MTSET, [2-(trimethylammonium)ethyl]methanethiosulfonate; pHMB, *p*-hydroxymecuribenzoic acid; SPA, staphylococcal protein A; T*, threonine modified to an unknown structure (presumably threonine hydroxamate – 17 Da); T, threonine hydroxamate.

§To whom reprint requests should be addressed. E-mail: olafs@ucla.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.




Fig. 1. Sortase catalyzes the anchoring of surface proteins to the staphylococcal cell wall. Surface proteins bearing sorting signals are cleaved between the threonine (T) and the glycine (G) of their LPXTG motif, and the carboxyl of threonine is amide-linked to the amino of the pentaglycine crossbridge within the staphylococcal cell wall. GN-MN, *N*-acetylglucosamine-*N*-acetylmuramic acid.

nickel nitrilotriacetic acid (Ni-NTA) column and eluted with 0.5 M imidazole. SEB-MH₆-CWS was precipitated with trifluoroacetic acid (TFA), washed in acetone, dried, and cleaved with cyanogen bromide (CNBr) in 70% formic acid (5). C-terminal peptides were suspended in 6 M guanidine-hydrochloride (pH 8.0), purified by affinity chromatography on Ni-NTA, eluted in 4 ml of 6 M guanidine-hydrochloride and 0.2 M acetic acid, desalted over a C18 cartridge, and dried. Pellets were solubilized in 50 μ l of buffer B [8 M urea/0.1 M NaH₂PO₄/0.01 M Tris-HCl (pH 8.0)] and subjected to RP-HPLC on a C18 column with a linear gradient from 1% to 90% acetonitrile (CH₃CN) in 0.1% TFA in 90 min. Matrix-assisted laser desorption ionization MS (MALDI-MS), electrospray ionization MS (ESI-MS), and affinity chromatography were performed as previously described (4).

Hydroxylaminolysis of LPXTG Peptide in Staphylococcal Extracts. Staphylococci [10^{13} *S. aureus* OS2 cells in 50 ml 50 mM Tris-HCl (pH 7.5)] were disrupted in a bead-beater instrument, and the crude extract was centrifuged at $1,500 \times g$ for 15 min to remove beads and unbroken cells. A 10- μ l aliquot of the supernatant (10 mg/ml protein) was used as enzyme preparation. Reactions were assembled in a volume of 260 μ l containing 50 mM Tris-HCl and 150 mM NaCl (pH 7.5). The concentration of LPXTG peptide substrate 4-(4-dimethylaminophenylazo)benzoic acid (Dabcy)-QALPETGEE-5-[(2-aminoethyl)amino]naphthalene-1-sulfonic acid (Edans) was 10 μ M; of *p*-hydroxymecuribenzoic acid (pHMB) was 5 mM; and of NH₂OH was 0.2 M. Incubations were carried out for 1 hr at 37°C, followed by centrifugation at $15,000 \times g$ for 5 min. Supernatant was analyzed in a fluorometer using 350 nm for excitation and 495 nm for recordings.

Purification of SrtA_{ΔN} Protein. The primers orf6N-ds-B (5'-AAAGGATCCAAACCACATATCGATAATTATC-3') and orf6C-B (5'-AAGGATCCTTATTTGACTTCTGTAGCTACAA-3') were used to PCR amplify the *srtA* sequence from the chromosome of *S. aureus* OS2. The DNA fragment was digested with *Bam*HI, inserted into pQE30 (Qiagen, Chatsworth, CA)-cut *Bam*HI to generate pHTT14, transformed into *Escherichia coli* XL-1 Blue, and selected on Luria agar with ampicillin (100 μ g/ml). *E. coli* XL-1 Blue (pHTT14) (10^{12} cells) in 30 ml of buffer C [50 mM Tris-HCl/150 mM NaCl/10% glycerol (pH 7.2)] were lysed in a French pressure cell at 14,000 psi (1 psi = 6.89 kPa). The extract was centrifuged at $29,000 \times g$ for 30 min, and the supernatant was applied to 1 ml of Ni-NTA resin, preequilibrated with buffer C. The column was washed with 40 ml of buffer C, and SrtA_{ΔN} protein was eluted in 4 ml of buffer

C with 0.5 M imidazole at a concentration of 3 μ g/ μ l. The mutant protein SrtA_{ΔN}, C184A was generated by PCR amplification of pHTT14 with the primers C184A-1 (5'-CAATTAACA-TTAATTACTGCTGATGATTACAATGAAAAG-3') and C184A-2 (5'-CTTTTCATTGTAATCATCAGCAGTAATTA-ATGTTAATTG-3'). The PCR product was digested with *Dpn*I and transformed into *E. coli* XL-1 Blue, and the mutant plasmid pHTT16 was isolated and analyzed. When purified proteins were subjected to ESI-MS, we observed an average mass of 22139.02 Da (± 2.37 Da) (SrtA_{ΔN}) and 22103.22 Da (± 3.05 Da) (SrtA_{ΔN}, C184A), respectively. These measurements are consistent with the predicted average mass of SrtA_{ΔN} (22133.93 Da) and SrtA_{ΔN}, C184A (22101.86 Da).

Hydroxylaminolysis of LPXTG Peptide by Purified SrtA_{ΔN}. Reactions were assembled in a volume of 260 μ l containing 50 mM Tris-HCl buffer, 150 mM NaCl (pH 7.5), and, as indicated, 5 μ M SrtA_{ΔN} in 50 mM Tris-HCl (pH 7.5), 10 μ M LPXTG peptide (Dabcy)-QALPETGEE-Edans, 10 μ M GLXTP peptide (Dabcy)-QAGLETPEE-Edans, 5 mM [2-(trimethylammonium)ethyl]methanethiosulfonate (MTSET), 0.2 M NH₂OH, 5 mM pHMB, or 10 mM DTT. Incubations were carried out for 1 hr at 37°C. Samples were analyzed in a fluorometer using 350 nm for excitation and 495 nm for recordings.

Results

Hydroxylamine Release of Surface Proteins into the Medium of *S. aureus* Cultures. To examine whether hydroxylamine can release surface proteins from cell-bound acyl-enzyme intermediate complexes, staphylococci were pulse-labeled with [³⁵S]methionine in either the presence or absence of 0.2 M NH₂OH (Fig. 2). Labeled cultures were divided into two aliquots. One sample was centrifuged, and proteins released into the culture medium were recovered in the supernatant (S). The other culture aliquot was treated with lysostaphin (L) (15), an endopeptidase that cuts the pentaglycine crossbridges of peptidoglycan and releases all cell wall-anchored surface protein of staphylococci (9). Surface proteins (SEB-SPA₄₉₀₋₅₂₄) of mock-treated staphylococci were cell wall anchored; however, treatment with 0.2 M NH₂OH caused 25.3% of all labeled SEB-SPA₄₉₀₋₅₂₄ to be released into the extracellular medium (Fig. 2A). This observation was not strain-specific, as *S. aureus* OS2 (*spa*:*ermC*) and *S. aureus* BB270 (wild type) displayed similar amounts of surface protein release.

Hydroxylaminolysis of acyl-enzyme intermediates should occur only during or shortly after the pulse-labeling of staphylococci, as SEB-SPA₄₉₀₋₅₂₄ is rapidly anchored to the cell wall (9). Addition of hydroxylamine either before or during the pulse with [³⁵S]methionine released surface proteins into the extracellular medium (16.9% and 12.7%, respectively) (Fig. 2B). Very little SEB-SPA₄₉₀₋₅₂₄ was detected in the culture medium when hydroxylamine was added after the pulse (4%), indicating that hydroxylaminolysis occurs during surface protein sorting to the cell wall, but does not affect mature, anchored polypeptides. Increasing the amount of hydroxylamine prior to pulse-labeling caused increased amounts of surface protein to be released into the culture medium (Fig. 2C). Addition of 0.5 M or greater amounts of NH₂OH prevented metabolic labeling of staphylococci with [³⁵S]methionine (data not shown).

Hydroxylaminolysis Causes the Formation of C-Terminal Threonine Hydroxamate. If sortase acyl-enzyme intermediates can be resolved with NH₂OH, the released surface proteins should contain hydroxamate at the threonine of the LPXTG motif. To test this prediction, 10^{13} staphylococci expressing the surface protein SEB-MH₆-CWS (4) were incubated in the presence or absence of 0.1 M NH₂OH. The cells were sedimented by centrifugation, and SEB-MH₆-CWS (Fig. 3A) was purified from the culture




Fig. 2. Release of surface protein from *S. aureus* by treatment with hydroxylamine. (A) Surface-protein (SEB-SPA₄₉₀₋₅₂₄) anchoring to the bacterial cell wall was measured by pulse-labeling staphylococci with [³⁵S]methionine in duplicate. One sample was precipitated with TCA, and anchored surface protein was released by lysostaphin-digestion of the cell wall envelope (L). The other sample was centrifuged, and surface protein released into the culture medium (S, supernatant) was precipitated with TCA. Surface protein was quantified by immunoprecipitation with anti-SEB (α-Seb), followed by SDS/PAGE and PhosphorImager analysis. Surface protein release was measured as the percent of SEB-SPA₄₉₀₋₅₂₄ in the culture medium, as compared with the total amount after lysostaphin solubilization. Hydroxylamine (0.2 M NH₂OH) was added 15 sec prior to labeling of *S. aureus* strains OS2 (*spa*:*ermC*) or BB270 (wild type). (B) Hydroxylamine was added 15 sec prior to labeling (prior), during pulse-labeling (pulse), or 5 min after the addition of the chase to *S. aureus* OS2 cultures (chase). (C) Increasing amounts of hydroxylamine added 15 sec prior to labeling of *S. aureus* OS2 cultures caused increasing amounts of surface-protein release.

medium by affinity chromatography on Ni-NTA. Coomassie-stained SDS/PAGE revealed the presence of released surface protein (Fig. 3B). Polypeptides were cleaved at methionine with CNBr, and C-terminal peptides containing the H₆ tag were purified by a second affinity chromatography step on Ni-NTA (4). RP-HPLC of peptides recovered from mock-treated bacteria revealed an absorbance peak at 30 min (25% CH₃CN) (Fig. 3C), which, when examined by ESI-MS, contained a compound of 2236 Da. This measurement is consistent with the structure of NH₂-L-Ala-D-iGln-L-Lys(NH₂-H₆AQALPET-Gly₅)-D-Ala-

COOH (predicted mass 2235), a C-terminal peptide that is likely the product of cell wall degradation and, together with other peptidoglycan fragments, released into the culture medium (1).

RP-HPLC of peptides released with 0.1 M NH₂OH generated an absorbance peak at 32 min (27% CH₃CN) (Fig. 3D) containing a compound with average mass of 1548 Da. Edman degradation revealed the peptide sequence NH₂-H₆AQALPET*, in which the thirteenth cleavage cycle released a phenylthiohydantoin moiety of unknown structure. The predicted mass of NH₂-H₆AQALPET (T indicates threonine hydroxamate) is 1565 Da, 17 Da more than the observed mass of 1548 Da. RP-HPLC fractions were scanned for the presence of ion signals with an average mass of 1548, 1565, or 2236. Mock-treated staphylococci released only the C-terminal peptide with mass 2236, whereas hydroxylamine-treated bacteria released peptides of 1548 Da, 1565 Da, and 2236 Da (Fig. 3D). Thus, NH₂OH treatment of staphylococci released surface protein hydroxamate, suggesting that sortase forms an acyl-enzyme intermediate with cleaved surface protein.

The amounts of purified C-terminal peptide with *m/z* 1565 were insufficient for Edman degradation. Therefore, we compared the structure of the peptides with *m/z* 1548 and 1565 by tandem mass spectrometry using the triply charged parent ions at *m/z* 517.2 and 522.9, respectively. Collisionally induced dissociation produced daughter ion spectra consistent with the peptide sequences NH₂-H₆AQALPET) and NH₂-H₆AQALPET* (the structure of T* is unknown) (data not shown).

Staphylococcal Extracts Catalyze Hydroxylaminolysis at the LPXTG Motif. If NH₂OH releases surface proteins from staphylococci *in vivo*, sortase may catalyze the cleavage of LPXTG motif-bearing peptides in the presence of NH₂OH *in vitro*. Fluorescence of the Edans fluorophore within the peptide Dabcyl-QALPETGEE-Edans is quenched by the close proximity of Dabcyl (16). When the peptide is cleaved and the fluorophore is separated from Dabcyl, an increase in fluorescence is observed (17). Incubation of the LPXTG peptide with crude staphylococcal extracts caused only a small increase in fluorescence (Fig. 4). However, the addition of 0.1 M NH₂OH to staphylococcal extracts resulted in an increase in fluorescence intensity (Fig. 4). This activity appears to be specific for sortase, as it can, at least in part, be inhibited by preincubation of staphylococcal extracts with pHMB or MTSET (18), known inhibitors of the sorting reaction (11). These results suggest that sortase catalyzes the hydroxylaminolysis of LPXTG peptide *in vitro*.

Purified Soluble SrtA_{ΔN} Catalyzes Hydrolysis and Hydroxylaminolysis at the LPXTG Motif. When expressed in *E. coli* and analyzed by centrifugation of crude lysates, the staphylococcal SrtA protein sedimented with membranes (data not shown). Crude *E. coli* extracts containing recombinant SrtA catalyzed cleavage of LPXTG peptide *in vitro* (data not shown). To obtain a soluble enzyme and examine its properties, the NH₂-terminal membrane anchor segment of SrtA (residues 2–25) was replaced with a six-histidine tag (SrtA_{ΔN}). SrtA_{ΔN} was expressed in *E. coli* XL-1 Blue and purified by affinity chromatography from cleared lysates (Fig. 5A). When incubated with the LPXTG peptide and measured as an increase in fluorescence, SrtA_{ΔN} catalyzed cleavage of the substrate (Fig. 5B). Addition of 0.2 M NH₂OH to this reaction resulted in an increase in fluorescence, indicating that cleavage of the LPXTG peptide occurred more efficiently. Hydroxylaminolysis of LPXTG peptide depended on the sulfhydryl of SrtA_{ΔN}, as preincubation with either MTSET or pHMB abolished all enzymatic activity.

Methanethiosulfonate forms disulfide with sulfhydryl groups that can be reversed by reducing agents such as DTT (19). MTSET-inactivated SrtA_{ΔN} was incubated in the presence of 10




Fig. 3. Characterization of surface protein released from staphylococci by hydroxylamine treatment. (A) The drawing shows the primary structure of SEB-MH₆-CWS and the cleavage sites for signal peptidase, cyanogen bromide (CnBr), and sortase. (B) Staphylococci (10^{13} cells of strains BB270 or OS2) expressing the surface protein SEB-MH₆-CWS were incubated in either presence (+) or absence (-) of 0.1 M NH₂OH. Cells were sedimented by centrifugation, and the supernatant was subjected to affinity purification on Ni-NTA resin. Surface protein was eluted with imidazole and analyzed on Coomassie-stained SDS/PAGE. The position of molecular mass standards (M) is indicated in kDa. (C) Purified SEB-MH₆-CWS surface protein released from mock-treated *S. aureus* BB270 cells was cleaved with CNBr at methionine residues. C-terminal peptides were purified by affinity chromatography and analyzed on RP-HPLC. The absorption peak at 30 min was analyzed by ESI-MS to contain a compound with an average mass of 2236 Da, consistent with the structure of surface protein linked to murein tetrapeptide. (D) RP-HPLC chromatogram of C-terminal anchor peptides released from *S. aureus* BB270 cells by means of treatment with 0.1 M NH₂OH. The absorption peak at 32 min was analyzed by ESI-MS to harbor a compound with average mass 1548 Da. Edman degradation revealed the peptide sequence NH₂-H₆AQALPET* (T* is an unknown structure, see text for detail). Arrows point to the elution of compounds with mass 1548 (NH₂-H₆AQALPET*), 1565 (NH₂-H₆AQALPET); T is threonine hydroxamate, and 2236 Da.




Fig. 4. Hydroxylaminolysis of LPXTG peptide by staphylococcal extracts *in vitro*. Staphylococcal extracts were incubated with the sorting substrate Dabcyl-QALPETGEE-Edans (LPXTG), and peptide cleavage was monitored as an increase in fluorescence. The addition of 0.2 M NH₂OH increased peptide cleavage, whereas the addition of pHMB, a known inhibitor of sortase, inhibited cleavage.

mM DTT, which restored 89% of its activity (Fig. 5B). If hydroxylaminolysis of LPXTG peptide by SrtA_{ΔN} is a measure of the transpeptidation reaction that links surface protein to the staphylococcal cell wall, alterations of the LPXTG motif should prevent peptide cleavage. Dabcyl-QAGLETPEE-Edans, in which the LPXTG motif sequence has been changed without altering the overall amino acid composition, was incubated with SrtA_{ΔN} in either the presence or absence of 0.2 M NH₂OH (Fig. 5B). No increase in fluorescence was observed, indicating that this peptide did not serve as a substrate for hydroxylaminolysis by SrtA_{ΔN}.

Cysteine-184 of SrtA Is Necessary for Sortase Activity. If sortase forms a thioester linkage with cleaved surface protein, replacement of the single conserved cysteine at position 184 with alanine should abolish all enzymatic activity. We purified such a mutant enzyme (SrtA_{ΔN, C184A}) and, when it was incubated with LPXTG peptide, observed only a very small increase in fluorescence. This increase may be caused by the binding of the LPXTG substrate to SrtA_{ΔN, C184A}. However, addition of 0.2 M NH₂OH to this reaction did not cause a further increase in fluorescence, indicating that the LPXTG peptide was not cleaved. Thus, cysteine-184 of SrtA is necessary for both efficient hydrolysis and hydroxylaminolysis of LPXTG-bearing peptides (Fig. 6).

Discussion

Hydroxylamine has been used to characterize several transpeptidation reactions (13, 20, 21). The best studied example is bacterial transpeptidases, which catalyze the crosslinking of cell wall peptides in the peptidoglycan, a reaction that is inhibited by penicillin (22–24). Transpeptidases or penicillin-binding proteins cleave the cell wall precursor at D-Ala-D-Ala and capture the carboxyl of the wall peptide as an ester with the active site hydroxyl of serine (25, 26). The acyl-enzyme intermediate is resolved by the nucleophilic attack of the amino within the cell wall crossbridge (pentaglycine in staphylococci), thereby allowing the formation of crosslinked wall peptides and regenerating the enzyme active-site hydroxyl (26, 27). Hydroxylamine can release the physiological acyl-enzyme intermediate as well as ester-linked penicilloyl, as this compound is a much stronger nucleophile than either the amino of the pentaglycine crossbridge or water (12). Although hydroxylamine can attack both ester and thioester bonds, the thioester enzyme intermediates appear to be more sensitive to hydroxylaminolysis (13).

The observation that surface-protein anchoring is sensitive to sulfhydryl-modifying reagents (11) prompted us to examine




Fig. 5. Purification and characterization of sortase (SrtA). (A) *E. coli* XL-1 Blue(pHTT14) expressing SrtA_{ΔN}, in which the NH₂-terminal membrane anchor of sortase (SrtA) has been replaced with a six-histidine tag, was lysed by French press. SrtA_{ΔN} was purified by affinity chromatography on Ni-NTA and analyzed on Coomassie-stained SDS/PAGE. Lanes: 1, uninduced culture; 2, 1 mM isopropyl β-D-thiogalactoside-induced culture; 3, French press extract; 4, the supernatant of centrifuged French press extracts; 5, the sediment of French press extracts; 6, flow-through of affinity chromatography on Ni-NTA; 7, column wash; and 8–10, 1-ml fractions eluted with 0.5 M imidazole. (B) Purified SrtA_{ΔN} was incubated with the LPXTG peptide, and cleavage was monitored as an increase in fluorescence. The reaction was inhibited by the addition of MTSET or organic mercurial (pHMB), whereas the addition of 0.2 M NH₂OH accelerated cleavage. MTSET-treated SrtA_{ΔN} could be rescued by incubation with 10 mM DTT. The peptide Dabcyl-QAGLEPPEE-Edans (GLXTP) served as a control for cleavage specificity of purified SrtA_{ΔN}.

hydroxylaminolysis of sortase acyl-enzyme intermediates. We show here that hydroxylamine releases surface proteins from staphylococci. Two polypeptide species were purified from the culture medium, surface protein threonine hydroxamate (T) and the more abundant form with a loss of 17 Da (T*). Although the structure of T* is unknown, we think it is likely that this compound is generated from T) during our purification scheme, which includes treatment with TFA, formic acid, guanidine-hydrochloride, and urea. *In vitro*, hydroxylaminolysis and hydrolysis of LPXTG peptide were sensitive to inhibition with the sulfhydryl reagents. Protein sequence comparison revealed the striking conservation of cysteine-184, as well as flanking amino acid residues, within sortase enzymes from several different Gram-positive bacteria (Fig. 6A). Replacement of cysteine-184 with alanine abolished sortase activity, suggesting that the cysteine sulfhydryl may function as an active-site nucleophile.

We propose a model in which surface proteins of *S. aureus* are linked to the cell wall by a transpeptidation reaction. Sortase (SrtA) cleaves polypeptides between the threonine and the





Fig. 6. Cysteine-184 of SrtA is necessary for sortase activity. (A) Alignment of *S. aureus* SrtA (Saur), i.e., residues 179–186 comprising cysteine-184, with corresponding amino acids of sortase homologs from *Streptococcus pyogenes* (Spyo), *Enterococcus faecalis* (Efea), *Actinomyces naeslundii* (Anei), *Streptococcus mutans* (Smut), *Bacillus subtilis* (Bsub), and *Streptococcus pneumoniae* (SpnA, SpnB, and SpnC). (B) Purified SrtA_{ΔN,C184A} contains a replacement of cysteine-184 with alanine. When incubated with LPXTG peptide, SrtA_{ΔN,C184A} caused only a small increase in fluorescence, which may be because of the binding of substrate to the mutant enzyme. The addition of NH₂OH to SrtA_{ΔN,C184A} did not generate an increase in fluorescence, indicating that no cleavage of LPXTG peptide occurred.

glycine of the LPXTG motif, resulting in the formation of a hydroxylamine-sensitive thioester linkage between the carboxyl of threonine and the enzyme sulfhydryl (Eq. 1). *In vivo*, the acyl-enzyme intermediate is resolved by the nucleophilic attack of the amino group within the pentaglycine crossbridge (Eq. 2). Recent observations suggest that the pentaglycine crossbridge of the lipid II precursor may function as a nucleophile for the sorting reaction (11); however, hydroxylamine can substitute for pentaglycine both *in vivo* and *in vitro*.



Purified sortase (SrtA) catalyzed hydrolysis of LPXTG-bearing peptide, suggesting that *in vitro* even water can function as a nucleophile to resolve the acyl-enzyme intermediate. As a spontaneous release of surface protein into the culture medium cannot be measured during pulse-labeling experiments, it appears that the availability of pentaglycine amino groups at the anchoring site of surface proteins prevents hydrolysis from occurring *in vivo*. The availability of purified, soluble sortase (SrtA_{ΔN}) and an *in vitro* assay for the hydroxylaminolysis of LPXTG peptide should allow the screening for compounds that inhibit anchoring of surface proteins in Gram-positive bacteria. As surface proteins are believed to be essential for the pathogenesis of bacterial disease (1), such compounds may be useful for the therapy of human infections caused by Gram-positive bacteria that have gained resistance to all known antibiotics (28).

We thank Drs. Dominique Missiakas [University of California, Los Angeles (UCLA)], Peter Model, Marjorie Russel (Rockefeller University), and members of our laboratory for discussion and critical reading of this manuscript. H.T.T. was supported by the Predoctoral Training Program in Microbial Pathogenesis at UCLA (AI07323). S.K.M. was supported by the

Predoctoral Training Program in Genetic Mechanisms at UCLA (T32GM07104). The W. M. Keck Foundation provided support toward instrument purchase. Work in the laboratory of O.S. is supported by Grant AI33987 from the National Institutes of Health-National Institute of Allergy and Infectious Diseases, Infectious Disease Branch.

1. Navarre, W. W. & Schneewind, O. (1999) *Microbiol. Mol. Biol. Rev.* **63**, 174–229.
2. Navarre, W. W. & Schneewind, O. (1994) *Mol. Microbiol.* **14**, 115–121.
3. Schneewind, O., Fowler, A. & Faull, K. F. (1995) *Science* **268**, 103–106.
4. Ton-That, H., Faull, K. F. & Schneewind, O. (1997) *J. Biol. Chem.* **272**, 22285–22292.
5. Ton-That, H., Labischinski, H., Berger-Bächi, B. & Schneewind, O. (1998) *J. Biol. Chem.* **273**, 29143–29149.
6. Navarre, W. W., Ton-That, H., Faull, K. F. & Schneewind, O. (1998) *J. Biol. Chem.* **273**, 29135–29142.
7. Navarre, W. W., Ton-That, H., Faull, K. F. & Schneewind, O. (1999) *J. Biol. Chem.* **274**, 15847–15856.
8. Schneewind, O., Model, P. & Fischetti, V. A. (1992) *Cell* **70**, 267–281.
9. Schneewind, O., Mihaylova-Petkov, D. & Model, P. (1993) *EMBO J.* **12**, 4803–4811.
10. Mazmanian, S., Liu, G., Ton-That, H. & Schneewind, O. (1999) *Science* **285**, 760–765.
11. Ton-That, H. & Schneewind, O. (1999) *J. Biol. Chem.* **274**, 24316–24320.
12. Lawrence, P. & Strominger, J. L. (1970) *J. Biol. Chem.* **245**, 3653–3659.
13. Kozarich, J. W., Tokuzo, N., Willoughby, E. & Strominger, J. L. (1977) *J. Biol. Chem.* **252**, 7525–7529.
14. Lipmann, F. & Tuttle, L. C. (1945) *J. Biol. Chem.* **161**, 415–416.
15. Schindler, C. A. & Schurhard, V. T. (1964) *Proc. Natl. Acad. Sci. USA* **51**, 414–421.
16. Wang, G. T., Matayoshi, E., Huffaker, H. J. & Krafft, G. A. (1990) *Tetrahedron Lett.* **31**, 6493–6496.
17. Matayoshi, E. D., Wang, G. T., Krafft, G. A. & Erickson, J. (1989) *Science* **247**, 954–958.
18. Smith, D. J., Maggio, E. T. & Kenyon, G. L. (1975) *Biochemistry* **14**, 764–771.
19. Pathak, R., Hendrickson, T. L. & Imperiali, B. (1995) *Biochemistry* **34**, 4179–4185.
20. Shao, Y., Xu, M.-Q. & Paulus, H. (1996) *Biochemistry* **35**, 3810–3815.
21. Ramalingam, S., Maxwell, S. E., Medof, M. E., Chen, R., Gerber, L. D. & Udenfriend, S. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 7528–7533.
22. Strominger, J. L., Izaki, K., Matsuhashi, M. & Tipper, D. J. (1967) *Fed. Proc.* **26**, 9–18.
23. Ghuyssen, J.-M. (1991) *Annu. Rev. Microbiol.* **45**, 37–67.
24. Waxman, D. J. & Strominger, J. L. (1983) *Annu. Rev. Biochem.* **52**, 825–869.
25. Rasmussen, J. R. & Strominger, J. L. (1978) *Proc. Natl. Acad. Sci. USA* **75**, 84–88.
26. Yocum, R. R., Waxman, D. J., Rasmussen, J. R. & Strominger, J. L. (1979) *Proc. Natl. Acad. Sci. USA* **76**, 2730–2734.
27. Tipper, D. J. & Strominger, J. L. (1965) *Proc. Natl. Acad. Sci. USA* **54**, 1133–1141.
28. Sieradzki, K., Roberts, R. B., Haber, S. W. & Tomasz, A. (1999) *N. Engl. J. Med.* **340**, 517–523.