

# Draft Genome Sequence of *Herpetosiphon geysericola* GC-42, a Nonphototrophic Member of the *Chloroflexi* Class *Chloroflexia*

Lewis M. Ward,<sup>a</sup> James Hemp,<sup>a</sup> Laura A. Pace,<sup>b</sup> Woodward W. Fischer<sup>a</sup>

Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, California, USA<sup>a</sup>; Department of Medicine, University of California San Diego, La Jolla, California, USA<sup>b</sup>

**We report here the draft genome sequence of *Herpetosiphon geysericola* GC-42, a predatory nonphototrophic member of the class *Chloroflexia* in the phylum *Chloroflexi*. This genome provides insight into the evolution of phototrophy and aerobic respiration within the *Chloroflexi*.**

Received 29 September 2015 Accepted 5 October 2015 Published 19 November 2015

**Citation** Ward LM, Hemp J, Pace LA, Fischer WW. 2015. Draft genome sequence of *Herpetosiphon geysericola* GC-42, a nonphototrophic member of the *Chloroflexi* class *Chloroflexia*. *Genome Announc* 3(6):e01352-15 doi:10.1128/genomeA.01352-15.

**Copyright** © 2015 Ward et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to James Hemp, jimhemp@caltech.edu.

The majority of cultured members of the bacterial phylum *Chloroflexi* belong to the class *Chloroflexia*, which are prominently anoxygenic phototrophs (1). However, the most basal *Chloroflexia*, members of the orders *Herpetosiphonales* and *Kallotenuales*, are nonphototrophic (2). This makes these clades of central importance to understanding the evolution of phototrophy within the *Chloroflexi*. The *Herpetosiphonales* order currently contains only two species, *Herpetosiphon geysericola* and *Herpetosiphon aurantiacus*, which are pigmented filamentous organoheterotrophs that exhibit gliding motility (3). In culture, *Herpetosiphon* strains are obligate aerobes that prefer microaerobic conditions (3). The physiology and ecology of these organisms are not well understood, but it has been suggested that members of the *Herpetosiphon* genus might be capable of predation via a “wolf pack” strategy (4). *H. geysericola* was isolated from a biofilm at a hot spring in Baja California, Mexico (5).

The genome of *H. geysericola* GC-42 (DSM 7119) was sequenced as part of a project to expand the phylogenetic breadth of *Chloroflexi* genomes. Genome sequencing was performed at SeqMatic using the Illumina MiSeq sequencing platform. SPAdes 3.1.1 (6) was used to assemble the genome. The genome was screened for contaminants based on sequence coverage, G+C composition, and BLAST hits of conserved single-copy genes. Genome annotation was performed using the NCBI Prokaryotic Genome Annotation Pipeline. The draft genome is 6.24 Mb in size, assembled into 46 contigs. It contains 5,335 genes, 4,688 coding sequences (CDSs), 2 16S RNAs, 47 tRNAs, and 6 clustered regularly interspaced short palindromic repeat (CRISPR) arrays. It is estimated to be ~99% (111/111) complete based on conserved single-copy genes.

Analysis of the *H. geysericola* genome revealed genes for a branched aerobic respiratory chain, including two different complex I (NADH dehydrogenase), complex II (succinate dehydrogenase), complex III (cytochrome *bc* complex), an A-family heme-copper oxygen reductase, and a quinol *bd* oxidase. No genes for phototrophy were found. In addition, no genes associated with the 3-hydroxypropionate cycle, a CO<sub>2</sub> fixation pathway present in

photosynthetic members of *Chloroflexi* (7), were found in *H. geysericola*; this is consistent with its predicted heterotrophic physiology.

The sequencing of *H. geysericola*, along with *Herpetosiphon aurantiacus* (4), completes the genomic knowledge of the cultured members of the genus *Herpetosiphon*. These data provide important constraints that help ordinate the acquisition of phototrophy, aerobic respiration, and carbon fixation within the *Chloroflexi*.

**Nucleotide sequence accession number.** This whole-genome shotgun project has been deposited in DDBJ/EMBL/GenBank under the accession no. [LGKP00000000](https://www.ncbi.nlm.nih.gov/nuccore/LGKP00000000).

## ACKNOWLEDGMENTS

Genomic DNA was obtained from the Leibniz-Institut DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH. Sequencing was performed at SeqMatic, Fremont, CA.

This work was funded in part by the Center for Environmental Microbial Interactions (CEMI) at Caltech, the Packard Foundation (to W.W.F.), the Agouron Institute (to J.H. and W.W.F.), and NSF GRFP (to L.M.W.).

## REFERENCES

- Overmann J. 2008. Green nonsulfur bacteria. In *Encyclopedia of Life Sciences (ELS)*. John Wiley & Sons, Chichester, United Kingdom.
- Cole JK, Gieler BA, Heisler DL, Palisoc MM, Williams AJ, Dohnalkova AC, Ming H, Yu TT, Dodsworth JA, Li W-J, Hedlund BP. 2013. *Kallotenua papyrolyticum* gen. nov., sp. nov., a cellulolytic and filamentous thermophile that represents a novel lineage (*Kallotenuales* ord. nov., *Kallotenuaceae* fam. nov.) within the class *Chloroflexia*. *Int J Syst Evol Microbiol* 63:4675–4682. <http://dx.doi.org/10.1099/ijso.0.053348-0>.
- Lee N, Reichenbach H. 2006. The genus *Herpetosiphon*, p 854–877. In Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E (ed), *The prokaryotes: vol 7: Proteobacteria: Delta, Epsilon subclass*. Springer Science + Business Media, New York, NY.
- Kiss H, Nett M, Martin K, Maresca JA, Copeland A, Lapidus A, Lucas S, Berry KW, Glavina Del Rio T, Dalin E, Tice H, Pitluck S, Richardson P, Bruce D, Goodwin L, Han C, Detter JC, Schmutz J, Brettin T, Land M, Hauser L, Kyrpidis NC, Ivanova N, Göker M, Woyke T, Klenk H-P, Bryant DA. 2011. Complete genome sequence of the filamentous gliding predatory bacterium *Herpetosiphon aurantiacus* type

- strain (114-95(T)). *Stand Genomic Sci* 5:356–370. <http://dx.doi.org/10.4056/sigs.2194987>.
5. Lewin RA. 1970. New *Herpetosiphon* species (*Flexibacterales*). *Can J Microbiol* 16:517–520. <http://dx.doi.org/10.1139/m70-087>.
  6. Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Prjibelski AD, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, Clingenpeel SR, Woyke T, McLean JS, Lasken R, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J Comput Biol* 20:714–737. <http://dx.doi.org/10.1089/cmb.2013.0084>.
  7. Strauss G, Fuchs G. 1993. Enzymes of a novel autotrophic CO<sub>2</sub> fixation pathway in the phototrophic bacterium *Chloroflexus aurantiacus*, the 3-hydroxypropionate cycle. *Eur J Biochem* 215:633–643. <http://dx.doi.org/10.1111/j.1432-1033.1993.tb18074.x>.