Non-commutative Nash inequalities

Michael Kastoryano and Kristan Temme

Citation: [Journal of Mathematical Physics](#) **57**, 015217 (2016); doi: 10.1063/1.4937382

View online: <http://dx.doi.org/10.1063/1.4937382>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jmp/57/1?ver=pdfcov>

Published by the [AIP Publishing](#)

Non-commutative Nash inequalities

Michael Kastoryano¹ and Kristan Temme²

¹NBIA, Niels Bohr Institute, University of Copenhagen, 2100 Copenhagen, Denmark

²Institute for Quantum Information and Matter, California Institute of Technology, Pasadena California 91125, USA

(Received 11 August 2015; accepted 24 November 2015; published online 16 December 2015)

A set of functional inequalities—called *Nash inequalities*—are introduced and analyzed in the context of quantum Markov process mixing. The basic theory of Nash inequalities is extended to the setting of non-commutative $\mathbb{L}_p$ spaces, where their relationship to Poincaré and log-Sobolev inequalities is fleshed out. We prove Nash inequalities for a number of unital reversible semigroups. © 2015 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4937382>]

I. INTRODUCTION

Bounding the mixing times of quantum channels is becoming an increasingly important topic in quantum information sciences, with applications ranging from self-correcting quantum memories²¹ to Gibbs sampling¹⁰ to quantum Shannon theory^{18,12} and dissipative many-body dynamics.^{4,11} The main task in the mixing time analysis of quantum dynamical semigroups is to find the least time after which the following bound holds: $\|e^{t\mathcal{L}^*}(\sigma) - \rho\|_1 \leq \epsilon$, for any initial state σ and some desired precision ϵ . Here, ρ is the stationary state of the semigroup $S_t^* \equiv e^{t\mathcal{L}^*}$.²⁷ Operationally, the mixing time corresponds to the time after which all information in σ is erased by the noise process S_t . Upon optimization over all input states, the trace norm bound can be rewritten as

$$\sup_{\|g\|_\infty \leq 1} \|S_t(g) - \rho(g)\|_\infty \equiv \xi(t) \leq \epsilon, \quad (1)$$

where $\rho(f) \equiv \text{tr}[\rho f] \mathbb{1}$. In general, assuming that the generator of the semigroup satisfies detailed balance (see Eq. (10)), a generic bound available on $\xi(t)$ is $\xi(t) \leq \sqrt{\|\rho^{-1}\|} e^{-t\lambda}$, where λ is the spectral gap of $\mathcal{L}$ (see Eq. (11)). However, $\|\rho^{-1}\|$ scales as the size of the phase space (dimension of the operator algebra) and often overshoots by quite a bit on the actual mixing time. We show that if the generator $\mathcal{L}$ of the semigroup satisfies a *non-commutative Nash inequality*

$$\|f - \rho(f)\|_{2,\rho}^{2+4/\nu} \leq C \langle f, -\mathcal{L}(f) \rangle_\rho \|f\|_{1,\rho}^{4/\nu}, \quad (2)$$

with some positive constants $C, \nu \in \mathbb{R}^+$ for all observables f , then the following improvements on the mixing time can be obtained:

$$\|S_t - \rho\|_{\infty \rightarrow \infty} \leq \|S_{t_1} - \rho\|_{2 \rightarrow \infty, \rho} \|S_{t_2} - \rho\|_{2 \rightarrow 2, \rho} \quad (3)$$

$$= \|S_{t_1} - \rho\|_{2 \rightarrow \infty, \rho} e^{-t_2 \lambda}, \quad (4)$$

where $t_1 + t_2 = t$, and the last line follows from the definition of the spectral gap (Eq. (11)). The norms appearing in Eq. (26) are non-commutative $\mathbb{L}_p$ norms^{19,26,13} with respect to the stationary state ρ of S_t^* . If the semigroup satisfies a non-commutative Nash inequality, by Theorem 1, we get

$$\|S_t - \rho\|_{2 \rightarrow \infty, \rho} \leq \left(\frac{\nu C}{4t} \right)^{\nu/4}. \quad (5)$$

Then, taking the supremum over $t_1 + t_2 = t$ yields

$$\|S_t - \rho\|_{\infty \rightarrow \infty} \leq 2e^{-\lambda(t - \nu C/4)}, \quad (6)$$

whenever $t \geq \nu C/4$. Thus if a Nash inequality can be established with good constants ν and C , then extremely tight bounds on the mixing time of the semigroup can be obtained, which are often

optimal when the underlying graph structure of the phase space is highly constrained but regular. To the knowledge of the authors, Nash inequalities only appear to be very powerful for single particle problems on local regular graphs. In general, obtaining good estimates on the constants ν, C can be very challenging.

In this paper, we extend the theory of Nash inequalities to the non-commutative setting. We show that two different types of Nash inequalities lead to close to optimal mixing time bounds. A lower bound on the Log-Sobolev constant is proved in terms of the constants ν, C in the Nash inequality, as well as strong bounds on the entire spectrum of the generator $\mathcal{L}$. Finally, as applications, we prove Nash inequalities for the depolarizing semigroup, for unital qubit semigroups, and for a random walk on the ring.

II. PRELIMINARIES

We denote the set of $d \times d$ complex matrices $\mathcal{M}_d$ and the subset of Hermitian matrices $\mathcal{A}_d = \{X \in \mathcal{M}_d, X = X^\dagger\}$, as well as the subset of positive definite matrices $\mathcal{A}_d^+ = \{X \in \mathcal{A}_d, X > 0\}$. The set of states will be denoted $\mathcal{S}_d = \{X \in \mathcal{A}_d, X \geq 0, \text{tr}[X] = 1\}$, and the full rank states will be analogously denoted $\mathcal{S}_d^+$. Observables will always be represented by lower case Latin letters ($f, g \in \mathcal{A}_d$) and states by Greek letters ($\rho, \sigma \in \mathcal{S}_d$).

Non-commutative Nash inequalities are defined with respect to finite dimensional non-commutative $\mathbb{L}_p$ spaces.¹⁹ The non-commutative $\mathbb{L}_p$ spaces are characterized by a norm and an inner product, which for any $f, g \in \mathcal{A}_d$ and some $\rho \in \mathcal{S}_d^+$ are defined as

$$\|f\|_{p,\rho}^p = \text{tr}[\rho^{1/2p} f \rho^{1/2p}] \quad \text{and} \quad \langle f, g \rangle_\rho = \text{tr}[\rho^{1/2} f \rho^{1/2} g]. \quad (7)$$

We will also make extensive use of the $\mathbb{L}_p$ variance which is defined as

$$\text{Var}_\rho(g) = \langle f - \rho(f), f - \rho(f) \rangle_\rho, \quad (8)$$

where $\rho(f) = \text{tr}[\rho f] \mathbb{1}$ is the uniform projection onto the expectation value of f with respect to ρ .

Throughout this paper, the time evolution of an observable ($f_t \in \mathcal{A}_d$) will be described by one-parameter semigroups of completely positive trace preserving maps (cpt-maps), whose generator (Liouvillian) can always be written in standard *Lindblad form*

$$\partial_t f_t = \mathcal{L}(f_t) \equiv i[H, f_t] + \sum_i L_i^\dagger f_t L_i - \frac{1}{2} \{L_i^\dagger L_i, f_t\}_+, \quad (9)$$

where $H \in \mathcal{A}_d$ is the system Hamiltonian and accounts for coherent dynamics, and the $L_i \in \mathcal{M}_d$ are Lindblad operators describing dissipation on the system. Note that we have written the dynamics on operators (Heisenberg picture) for convenience. We denote the dual of $\mathcal{L}$, with respect to the Hilbert-Schmidt inner product, by $\mathcal{L}^*$ which amounts to the evolution of states, i.e., the Schrödinger picture. The trace preserving condition ensures that $\mathcal{L}(\mathbb{1}) = 0$. If in addition $\mathcal{L}^*(\mathbb{1}) = 0$, then the dynamics are said to be unital. A Liouvillian $\mathcal{L}$ is said to be *primitive* if it has a unique full-rank stationary state. As the framework of non-commutative $\mathbb{L}_p$ spaces depends on a full rank reference state, which will most often be the stationary state of a some dissipative dynamics, we will almost exclusively consider primitive Liouvillians. The reference state ρ should always be clear from the context and will almost always be the unique full rank stationary state of $S_t^* \equiv e^{t\mathcal{L}^*}$.

We say a Liouvillian $\mathcal{L} : \mathcal{M}_d \rightarrow \mathcal{M}_d$ satisfies *detailed balance* (or is *reversible*) with respect to the state $\rho \in \mathcal{S}_d^+$, if for any $f, g \in \mathcal{A}_d$,

$$\langle f, \mathcal{L}(g) \rangle_\rho = \langle \mathcal{L}(f), g \rangle_\rho. \quad (10)$$

The class of reversible generators has a number of particularly nice properties. The one most often exploited is that if $\mathcal{L}$ satisfies detailed balance with respect to some $\rho \in \mathcal{S}_d^+$, then ρ is a stationary state of $\mathcal{L}$. Furthermore, the detailed balance condition ensures that the generator is Hermitian with respect to the weighted inner product $\langle f, g \rangle_\rho$, which ensures that $\mathcal{L}$ has a real spectrum.

We will frequently need the *Dirichlet form* of $\mathcal{L}$: $\mathcal{E}(f) = -\langle f, \mathcal{L}(f) \rangle_\rho$, which can be used in the following Poincaré inequality, to characterize the *spectral gap* of a primitive reversible

generator $\mathcal{L}$ as:²²

$$\lambda = \min_{g \in \mathcal{A}_d} \frac{\mathcal{E}(g)}{\text{Var}_\rho(g)}. \quad (11)$$

III. NASH INEQUALITIES

The results presented in this paper aim to develop a useful theory of Nash inequalities for quantum dynamical semigroups. Most of our arguments follow closely the classical works of Diaconis and Saloff-Coste,^{7,20} which in turn are based on the seminal work in Refs. 6, 5, 3, and 24. In some cases, the classical and quantum proofs are almost identical. We try as much as possible to point out what aspects of the quantum theory differ from the classical theory.

The cornerstone of the theory of Nash inequalities for mixing time analysis is a partial equivalence between so-called *ultracontractivity* of the semigroup and a Nash inequality. This connection is reminiscent of the relation between hypercontractivity and the Log-Sobolev inequalities.^{7,26,2,13} The following theorem sketches out one direction of the implication.

Theorem 1 (Nash I). *Let $\mathcal{L} : \mathcal{M}_d \rightarrow \mathcal{M}_d$ be a primitive reversible Liouvillian. If for all $f \in \mathcal{A}_d$, $\mathcal{L}$ satisfies a type I Nash inequality*

$$\text{Var}_\rho(f)^{1+2/\nu} \leq C \mathcal{E}(f) \|f\|_{1,\rho}^{4/\nu} \quad (12)$$

for some positive constants $\nu, C \in \mathbb{R}^+$, then

$$\|S_t - \rho\|_{1 \rightarrow 2} \leq \left(\frac{\nu C}{4t} \right)^{\nu/4}, \quad (13)$$

for all $t \geq 0$.

Proof. Set $u(t) := \|S_t(f) - \rho(f)\|_{2,\rho}^2 \equiv \text{Var}(S_t(f))$, then letting $\|f\|_{1,\rho} = 1$, Eq. (12) can be rewritten as

$$u(t)^{1+2/\nu} \leq -\frac{C}{2} \dot{u}(t), \quad (14)$$

which is equivalent to $\dot{v}(t) \geq 1$, where $v(t) = \frac{C\nu}{4} u(t)^{-2/\nu}$. Solving for v and recasting the result in terms of u yields

$$u(t) \leq \left(\frac{\nu C}{4t} \right)^{\nu/2}. \quad (15)$$

Taking the supremum over all functions f with $\|f\|_{1,\rho} = 1$ yields

$$\|S_t - \rho\|_{1 \rightarrow 2,\rho} \leq \left(\frac{\nu C}{4t} \right)^{\nu/4}. \quad (16)$$

□

Note that by duality of $\mathbb{L}_p$ norms and reversibility of the semigroup, one immediately also gets a bound on the $\mathbb{L}_{2 \rightarrow \infty}(\rho)$ norm for the semigroup, which was necessary to obtain the mixing time bound of Eq. (6). Furthermore, invoking the Riez-Thorin interpolation theorem (see, e.g., Ref. 1), the bound can be extended to give

$$\|S_t - \rho\|_{1 \rightarrow \infty,\rho} \leq \left(\frac{\nu C}{4t} \right)^{\nu/2}. \quad (17)$$

Some semigroups are not ultracontractive for all times but relax more rapidly for short times than a Poincaré inequality would suggest. In these cases, a different type of Nash inequality is more useful that zeros out the short time convergence of the semigroup.

Theorem 2 (Nash II). Let $\mathcal{L} : \mathcal{M}_d \rightarrow \mathcal{M}_d$ be a primitive reversible Liouvillian. If for all $f \in \mathcal{A}_d$, $\mathcal{L}$ satisfies a type II Nash inequality

$$\|f\|_{2,\rho}^{2+4/\nu} \leq C(\mathcal{E}(f)) + \frac{1}{T} \|f\|_{2,\rho}^2 \|f\|_{1,\rho}^{4/\nu} \quad (18)$$

for some positive constants $\nu, C, T \in \mathbb{R}^+$, then for all $t \leq T$,

$$\|S_t\|_{1 \rightarrow 2,\rho} \leq e^{t/T} \left(\frac{\nu C}{4t} \right)^{\nu/4}. \quad (19)$$

Proof. The proof is very similar to that of Theorem 1. Again we fix f satisfying $\|f\|_{1,\rho} = 1$, but now define

$$u(t) = e^{-2t/T} \|S_t(f)\|_{2,\rho}^2. \quad (20)$$

Then,

$$\dot{u}(t) = -2e^{-2t/T} (\mathcal{E}(S_t(f)) + \frac{1}{T} \|S_t(f)\|_{2,\rho}^2). \quad (21)$$

Eq. (18) now reads

$$u^{1+2/\nu}(t) \leq -\frac{C}{2} \dot{u}(t). \quad (22)$$

Using the same argument as in the proof of Theorem 1, we get

$$u(t) \leq \left(\frac{\nu C}{4t} \right)^{\nu/2}, \quad (23)$$

which implies

$$\|S_t\|_{1 \rightarrow 2,\rho} \leq e^{t/T} \left(\frac{\nu C}{4t} \right)^{\nu/4}. \quad (24)$$

□

The type II Nash inequality says something strong about the mixing at short times but does not provide any information about the asymptotic behavior. However, the ultracontractive bound in Eq. (19) is sufficient to obtain the improved mixing time bound of Eq. (6), provided that $T \geq \nu/(4\lambda)$, where λ is the spectral gap of $\mathcal{L}$.

Ultracontractivity (i.e., $\|S_t\|_{2 \rightarrow \infty,\rho} \leq 1$) is related to Nash inequalities in the sense that one is the infinitesimal formulation of the other, as we will see in the next theorem. This behavior is very reminiscent of the relationship between hypercontractivity and Log-Sobolev inequalities^{26,13} or the contraction of the $\|S_t\|_{2 \rightarrow 2,\rho}$ norm and the Poincaré inequality, Eq. (11). One can show (cf. Theorems 1 and 2) that ultracontractivity not only follows from a Nash inequality but also that the converse is true. This fact ensures that the mixing time bounds obtained from Nash inequalities are close to optimal for good constants (C, ν, T) , since the contractive bound on the $\mathbb{L}_{2 \rightarrow \infty}(\rho)$ norm eliminates the dependence on any generic pre-factor.

Theorem 3. Let $\mathcal{L} : \mathcal{M}_d \rightarrow \mathcal{M}_d$ be a primitive reversible Liouvillian, and suppose that the semigroup $S_t \equiv e^{t\mathcal{L}}$ satisfies

$$\|S_t - \rho\|_{1 \rightarrow 2,\rho} \leq \left(\frac{C}{t} \right)^{\nu/4}, \quad (25)$$

for all $t \geq 0$ and some positive constants $\nu, C \in \mathbb{R}^+$, then

$$\text{Var}_\rho(f)^{1+2/\nu} \leq C' \mathcal{E}(f) \|f\|_{1,\rho}^{4/\nu}, \quad (26)$$

for all $f \in \mathcal{A}_d$, where $C' = 2^{2+4/\nu} C$. Furthermore, if for all $t \leq T$,

$$\|S_t\|_{1 \rightarrow 2,\rho} \leq \left(\frac{C}{t} \right)^{\nu/4}, \quad (27)$$

for some positive constants $\nu, C, T \in \mathbb{R}^+$, then

$$\|f\|_{2,\rho}^{2+4/\nu} \leq C' \left(\mathcal{E}(f) + \frac{1}{T} \|f\|_{2,\rho}^2 \right) \|f\|_{1,\rho}^{4/\nu}, \quad (28)$$

for all $f \in \mathcal{A}_d$, where again $C' = 2^{2+4/\nu} C$.

Proof. We will only prove that Eq. (25) implies Eq. (26). The proof that Eq. (27) implies Eq. (28) is very similar, and a classical version of it can be found in Ref. 20.

Fix f with $\|f\|_{1,\rho} = 1$ and write, for $0 \leq t \leq T$,

$$\text{Var}_\rho(f) = \text{Var}_\rho(f_t) - \int_0^t ds \frac{d}{ds} \text{Var}_\rho(f_s) \quad (29)$$

$$= \text{Var}_\rho(f_t) + 2 \int_0^t ds \mathcal{E}(f_s) \quad (30)$$

$$\leq (C/t)^{\nu/2} + 2t\mathcal{E}(f). \quad (31)$$

The inequality follows from the ultracontractivity assumption Eq. (25) and from the observation that for all $t \geq 0$ and $f \in \mathcal{A}$, $\mathcal{E}(f) \geq \mathcal{E}(f_t)$. To see the latter, note that by primitivity, $\sqrt{-\mathcal{L}}$ is well defined and commutes with $\mathcal{L}$, and thus also with S_t . Then, by reversibility,

$$\mathcal{E}(f_t) = \langle f_t, -\mathcal{L}(f_t) \rangle_\rho \quad (32)$$

$$= \|\sqrt{-\mathcal{L}} e^{t\mathcal{L}}(f)\|_{2,\rho}^2 \quad (33)$$

$$= \|e^{t\mathcal{L}} \sqrt{-\mathcal{L}}(f)\|_{2,\rho}^2 \quad (34)$$

$$\leq \|\sqrt{-\mathcal{L}}(f)\|_{2,\rho}^2 = \mathcal{E}(f), \quad (35)$$

where the last inequality follows from contractivity of the $\mathbb{L}_{2 \rightarrow 2}(\rho)$ norm. Solving for the minimum of the right hand side of Eq. (31), we recover the Nash I inequality

$$\text{Var}_\rho(f)^{2+4/\nu} \leq B \mathcal{E}(f) \|f\|_{1,\rho}^{4/\nu}, \quad (36)$$

with

$$B = 2C(1 + 2/\nu)(1 + d/2)^{2/\nu} \leq 2^{2+4/\nu} C. \quad (37)$$

□

As will be evident in Sec. IV, the converse theorem is more useful for Nash II inequalities, as it is often much simpler to obtain a bound of the form Eq. (27) for some range of t up to a cutoff T , than for all real t .

A. Lower bound on the Log-Sobolev constant

Nash inequalities are functional inequalities involving $\mathbb{L}_p$ norms and Dirichlet forms. Many other functional inequalities (Poincaré, Sobolev, Log-Sobolev) have been extensively studied, both in the classical and in the quantum setting. Here, we relate the constants appearing in the Nash inequality with the Log-Sobolev constant, as defined in Refs. 13, 26, and 23.

Theorem 4. Let $\mathcal{L} : \mathcal{M}_d \rightarrow \mathcal{M}_d$ be the generator of a primitive reversible semigroup. If for all $f \in \mathcal{A}_d$, $\mathcal{L}$ satisfies

$$\text{Var}_\rho(f)^{1+2/\nu} \leq C \mathcal{E}(f) \|f\|_{1,\rho}^{4/\nu} \quad (38)$$

for some positive constants $\nu, C \in \mathbb{R}^+$, then the Log-Sobolev constant α_2 is bounded below by

$$\alpha_2 \geq \frac{2}{\nu C}. \quad (39)$$

If $\mathcal{L}$ satisfies

$$\|f\|_{2,\rho}^{2(1+2/\nu)} \leq C \left(\mathcal{E}(f) + \frac{1}{T} \|f\|_{2,\rho}^2 \right) \|f\|_{1,\rho}^{4/\nu}, \quad (40)$$

for some positive constants $\nu, C, T \in \mathbb{R}^+$, with $\nu C/4 \leq T$ and has spectral gap λ , then the Log-Sobolev constant α_2 is bounded below by

$$\alpha_2 \geq \frac{\lambda}{2(1 + \lambda t_0 + \nu/4 \log(\frac{\nu C}{4t_0}))} \quad (41)$$

for any $0 < t_0 \leq T$.

Proof. From the bound in Eq. (6), we get that for $t_0 = \nu C/4 \leq T$,

$$\|S_t\|_{2 \rightarrow \infty, \rho} \leq 1. \quad (42)$$

Then by Theorem 5 in Ref. 23, we simply get

$$\alpha_2 \geq \frac{\lambda}{2(1 + \lambda t_0 + \nu/4 \log(\frac{\nu C}{4t_0}))}. \quad (43)$$

The proof of Eq. (39) is essentially identical, but using a variant of Theorem 5 in Ref. 23 that can be found (in classical form) in Ref. 20 Theorem 2.2.13. $\square$

The Log-Sobolev constant α_2 lower bounds the spectral gap λ for primitive reversible channels (see Ref. 13), which completes the chain of inequalities between Nash, Log-Sobolev, and Poincaré constants.

Finally, we point out that the Nash inequalities are implied by non-commutative Sobolev inequalities. Indeed, consider the following non-commutative Sobolev inequalities:

$$\|f - \rho(f)\|_{2\nu/(\nu-1), \rho}^2 \leq C \mathcal{E}(f), \quad (44)$$

$$\|f\|_{2\nu/(\nu-2), \rho}^2 \leq C \left(\mathcal{E}(f) + \frac{1}{T} \|f\|_{2,\rho}^2 \right). \quad (45)$$

By Hölders inequality,

$$\|f\|_2^{2(1+2/\nu)} \leq \|f\|_{2\nu/(\nu-2)}^2 \|f\|_1^{4/\nu}, \quad (46)$$

and we immediately see that Eq. (44) implies a type I Nash inequality, and Eq. (45) implies a type II Nash inequality. Unlike in the classical setting, the converse is not known in the non-commutative setting.

B. Bounds on higher eigenvalues

One particular appeal of the Nash inequalities is that they imply bounds on the entire spectrum (i.e., higher eigenvalues), rather than just on the spectral gap. Let $\mathcal{L}$ be a reversible Liouvillian and let $0 \leq \lambda_0 \leq \lambda_1 \leq \dots \leq \lambda_{n-1}$ be the eigenvalues of $-\mathcal{L}$, and define the counting function

$$N(s) = \#\{i \in \{0, 1, \dots, n-1\} | \lambda_i \leq s\}, \quad s \geq 0. \quad (47)$$

N is a step function with $N(s) = 1$ for $0 \leq s < \lambda_1$ (for primitive S_t). It is easy to relate the counting function N to the eigenvalues of $-\mathcal{L}$. Define the one parameter function

$$\zeta(t) = \sum_i e^{-t\lambda_i}. \quad (48)$$

Since for $\lambda_i \leq 1/t$, we get $e^{-t\lambda_i} \geq e^{-1}$, and it follows that

$$N(1/t) \leq e\zeta(t). \quad (49)$$

Thus if we can find an upper bound on $\zeta(t)$, we can bound the counting function. Here, we show that $\zeta(t)$ can be bounded using Nash inequalities.

Theorem 5. Let $\mathcal{L} : \mathcal{M}_d \rightarrow \mathcal{M}_d$ be a primitive reversible Liouvillian. If for all $f \in \mathcal{A}_d$, $\mathcal{L}$ satisfies

$$\text{Var}_\rho(f)^{1+2/\nu} \leq C\mathcal{E}(f)\|f\|_{1,\rho}^{4/\nu} \quad (50)$$

for some positive constants $\nu, C \in \mathbb{R}^+$, then

$$N(s) \leq 1 + e(\nu Cs/2)^{\nu/2} \quad (51)$$

for all $s \geq 0$. If the generator satisfies a type II Nash inequality

$$\|f\|_{2,\rho}^{2(1+2/\nu)} \leq C \left(\mathcal{E}(f) + \frac{1}{T} \|f\|_{2,\rho}^2 \right) \|f\|_{1,\rho}^{4/\nu}, \quad (52)$$

then

$$N(s) \leq e^3(\nu Cs/2)^{\nu/2} \quad (53)$$

for all $s \geq 1/T$.

Proof. We will prove the bound for type II Nash inequalities. The proof of the bound for type I is very similar. We will use a mapping between quantum channels and matrices, whereby $S(f) = \sum_k E_k f E_k^\dagger$ gets mapped to $\hat{S} = \sum_k E_k \otimes \bar{E}_k$. Similarly, matrices X get mapped onto states as $|X\rangle = X \otimes \mathbb{1}|\Omega\rangle$, where $|\Omega\rangle = \sum_j |j, j\rangle$ is proportional to the maximally entangled state.

Now define the map $\Gamma_\rho(f) = \rho^{1/2} f \rho^{1/2}$ and the symmetrized semigroup as $\tilde{S}_t = \Gamma_\rho^{1/2} S_t \Gamma_\rho^{-1/2} = \Gamma_\rho^{-1/2} S_t^* \Gamma_\rho^{1/2}$. Since $\tilde{S}_t$ and S_t have the same spectrum, we get

$$\sum_j e^{-t\lambda_j} = \text{tr} [\hat{S}_t] \quad (54)$$

$$= \sum_j \langle K_j | \hat{S}_t | K_j \rangle \quad (55)$$

$$= \sum_j \text{tr} [K_j \tilde{S}_t(K_j)] \quad (56)$$

$$= \sum_j \langle L_j, S_t(L_j) \rangle_\rho \quad (57)$$

$$= \sum_j \|S_{t/2}(L_j)\|_{2,\rho}^2, \quad (58)$$

where $\{|K_j\rangle\}$ form some orthonormal basis, and $L_j = \Gamma_\rho^{-1/2}(K_j)$. Then, from a type II Nash inequality, we have

$$\|S_{t/2}\|_{1 \rightarrow 2,\rho}^2 \leq e^{2t/T} \left(\frac{\nu C}{2t} \right)^{\nu/2} \equiv \epsilon(t), \quad (59)$$

hence,

$$\sum_j e^{-t\lambda_j} \leq \epsilon(t) \sum_j \|L_j\|_{1,\rho}^2 \quad (60)$$

$$= \epsilon(t) \sum_j \text{tr} [\rho^{1/2} K_j]^2 \quad (61)$$

$$= \epsilon(t) \sum_j \langle \sqrt{\rho} | K_j \rangle \langle K_j | \sqrt{\rho} \rangle = \epsilon(t) \text{tr} [\rho]. \quad (62)$$

Thus, we get that whenever a type II Nash inequality holds, the eigenvalue counting function satisfies Eq. (53) for all $s \geq 1/T$. In order to show the bound in Eq. (51), one simply needs to use the identity $\|S_t(h) - \rho\|_{2,\rho}^2 = \|S_t(h)\|_{2,\rho}^2 - 1$. $\square$

Clearly, if $M(s)$ is a continuous and monotonously increasing function such that $N(s) \leq M(s)$, $s \geq 1/T$, then

$$\lambda_i \max\{s : N(s) \leq i\} \geq M^{-1}(i+1) \quad (63)$$

for all $i > M(1/T) - 1$. Hence, we obtain the following.

Corollary 6. Let $\mathcal{L} : \mathcal{M}_d \rightarrow \mathcal{M}_d$ be a primitive reversible Liouvillian, which satisfies a type I Nash inequality, then

$$\lambda_j \geq \frac{2j^{2/\nu}}{e^{2/\nu} \nu C}. \quad (64)$$

If $\mathcal{L}$ satisfies a type II Nash inequality, then

$$\lambda_j \geq \frac{2(j+1)^{2/\nu}}{e^{6/\nu} \nu C}, \quad (65)$$

for all $j > e^3 \left(\frac{\nu C}{2T}\right)^{\nu/2} - 1$.

IV. EXAMPLES

We provide a number of examples which illustrate the strengths and the weaknesses of the Nash inequalities. The general intuition about Nash inequalities is that they are good at identifying local uniform mixing for short times but are not very good at bounding global or long time mixing properties. For this, Poincaré and Log-Sobolev inequalities are typically better. As we will see in Example 3, the Nash inequalities provide the right mixing time estimate for a particle diffusing on a grid. This is consistent with classical results that have not proven very useful for many body systems but have provided the best mixing time results for models where information needs to spread across the grid before disappearing (see Ref. 7). In particular, the systems for which a Nash inequality is the right tool do not have a constant gap.

A. The depolarizing semigroup

As our first example, we prove a Nash I inequality for the depolarizing semigroup: $\mathcal{L}(f) = \gamma(\text{tr}[f\rho] \mathbb{1} - f)$. Its fixed point is ρ . The Nash inequality is saturated for projectors $f = |\psi\rangle\langle\psi|$.

Let $(C\gamma)^{\nu/4} \geq \|\rho^{-1}\|^{1/2}$, then

$$\text{Var}_\rho(f) \leq \|f\|_{2,\rho}^2 \leq (C\gamma)^{\nu/2} \|f\|_{1,\rho}^2. \quad (66)$$

Recalling that $\mathcal{E}(f) = \gamma \text{Var}_\rho(f)$ for the depolarizing semigroup, we recover the Nash I inequality

$$\text{Var}_\rho(f)^{1+2/\nu} \leq C \mathcal{E}(f) \|f\|_{1,\rho}^{4/\nu}. \quad (67)$$

Then, maximizing $C\gamma/4$ subject to $\|\rho^{-1}\|^{1/2} \leq (C\gamma)^{\nu/2}$ yields a mixing time of

$$\tau_{\text{mix}} = \frac{e \log(\|\rho^{-1}\|)}{2\gamma^2}. \quad (68)$$

Clearly, this bound provides no improvement on the one obtained from the spectral gap alone.

B. Unital qubit channels

Next, we consider reversible unital qubit semigroups and show that the Nash inequalities can be easily obtained as a function of the parameters of the King-Ruskai representation of qubit channels.¹⁶ Indeed, any qubit density operator can be represented as

$$\rho = \frac{1}{2}(\mathbb{1} + \vec{w} \cdot \vec{\sigma}), \quad (69)$$

where $\vec{w} \in \mathbb{R}^3$ and $|\vec{w}| \leq 1$ is a real vector and $\vec{\sigma} = (\sigma_X, \sigma_Y, \sigma_Z)$ are the Pauli matrices. Pure states are characterized by the sub-manifold satisfying $|\vec{w}| = 1$. A quantum channel $S : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ acts on the state ρ as

$$S(\rho) = \frac{1}{2}(\mathbb{1} + (\vec{s} + \Delta\vec{\omega}) \cdot \vec{\sigma}), \quad (70)$$

for some vector $\vec{s}$ and matrix Δ .

It can then be seen that the quantum channel S is unital if and only if $\vec{s} = 0$ and is reversible if and only if Δ is Hermitian. Let's now consider the case of a unital reversible qubit semigroup $S_t : \mathbb{C}^2 \rightarrow \mathbb{C}^2$. Then, the action of the semigroup on a state ρ is $S_t(\rho) = \frac{1}{2}(\mathbb{1} + (\Delta_t \vec{w}) \cdot \vec{\sigma})$, where $\Delta_t = e^{tL}$ is now also a semigroup. Denote the eigenvalues of $\Delta_t : (\lambda_1^t = e^{-tl_1}, \lambda_2^t = e^{-tl_2}, \lambda_3^t = e^{-tl_3})$. Then given that S_t is Hermitian and unital, $\lim_{t \rightarrow \infty} \Delta_t = 0$, which implies that (l_1, l_2, l_3) are all real and strictly positive.

We can now estimate the $\mathbb{L}_{1 \rightarrow 2}(\mathbb{1}/2)$ norm of S_t ,

$$\|S_t\|_{1 \rightarrow 2, \mathbb{1}/2}^2 = 2 \sup_{\text{tr}[\rho]=1} \text{tr}[\rho S_{2t}(\rho)] \quad (71)$$

$$= \sup_{|\vec{w}|=1} (1 + \vec{w} \cdot \Delta_{2t} \vec{w}) \quad (72)$$

$$= 1 + e^{-2tl_{\min}} \leq \left(\frac{1}{l_{\min} t} \right)^{1/2}, \quad (73)$$

for $t \leq 1/2l_{\min}$, where $l_{\min} = \min\{l_1, l_2, l_3\}$. Then, from Theorem 3, we get the Nash II inequality

$$\|f\|_{2,\rho}^6 \leq \frac{2^6}{l_{\min}} \left(\mathcal{E}(f) + \frac{1}{2l_{\min}} \|f\|_{2,\rho}^2 \right) \|f\|_{1,\rho}^4. \quad (74)$$

We might be tempted to extend the Nash II inequality to the case of a tensor power of reversible unital qubit channels. This can be done by invoking the multiplicativity of Shatten $1 \rightarrow 2$ norms for unital channels, proved in Refs. 14 and 15. Since for any unital qubit channel N ,

$$\|S^{\otimes N}\|_{1 \rightarrow 2} = \|S\|_{1 \rightarrow 2}^N, \quad (75)$$

we get for the unital reversible semigroup described above that

$$\|S_t^{\otimes N}\|_{1 \rightarrow 2, \mathbb{1}/2^N}^2 \leq \left(\frac{2}{l_{\min} t} \right)^{N/2}. \quad (76)$$

Eq. (76) then immediately leads to a Nash II inequality with constants $\nu = N$, $C' = 2^{2+4/N}/l_{\min}$, and $T = (2l_{\min})^{-1}$. Note that for $N > 2$, the constants will inevitably violate the assumption that $\nu(4\lambda)^{-1} \leq T$, which is a necessary condition to infer the mixing time bound from the constants in the Nash II inequality in Theorem 2. This is a simple example of the pathological behavior of Nash inequalities under multiplication. As a matter of fact, the ill behavior under the tensor product operation of Nash Inequalities and in turn Sobolev inequalities was a central motivation to introduce Log-Sobolev inequalities.⁹ It is known that this semi-group has in fact a system size independent Log-Sobolev constant α_2 .^{15,17,21}

C. Single particle diffusion and coherent hopping

As our last example, we consider a finite chain of length N with periodic boundary conditions. We assume that a single particle can hop dissipatively in both directions with equal rates. Moreover, we allow for additional coherent transport with a single particle Hamiltonian H . This model is defined on the Hilbert space $\mathbb{C}^{N+1}$ and can be viewed as the one-particle restriction of the quantum mechanical version of the symmetric exclusion process as considered in Ref. 8. We will see that the particular choice of the Hamiltonian H does not affect the convergence rate so that the equilibration bound holds for a large class of coherent hopping models.

The Lindblad operators that mediate the diffusive hopping are defined by $L_m = |m\rangle\langle m \oplus_N 1|$ for all $m = 0, \dots, N$. We denote by $\oplus_N$ the addition modulo N in $\mathbb{Z}_N$. Moreover, we allow for the coherent transport by any Hamiltonian H on $\mathbb{C}^{N+1}$, such as, for example, $H = \sum_m (L_m + L_m^\dagger)$. The semi-group can be written as

$$\mathcal{L}_C(f) = i[H, f] + \mathcal{L}(f), \quad \text{where} \quad \mathcal{L}(f) = \frac{1}{2} \sum_{m=0}^N [L_m, [f, L_m^\dagger]] + [L_m^\dagger, [f, L_m]]. \quad (77)$$

The stationary state of $\mathcal{L}_C$ is the maximally mixed state $\rho = (N+1)^{-1}\mathbb{1}$. For any hermitian $f \in \mathcal{M}_{N+1}(\mathbb{C})$ we have that $-i\text{tr}[f^\dagger[H, f]] = 0$, which implies the bound $\mathcal{E}_C(f) = -(N+1)^{-1}\text{tr}[f^\dagger\mathcal{L}_C(f)]$ by the form $\mathcal{E}(f) = -(N+1)^{-1}\text{tr}[f^\dagger\mathcal{L}(f)]$. That way, we have $\mathcal{E}_C(f) \leq \mathcal{E}(f)$ and the Nash II inequality Eq. (28) automatically holds for $\mathcal{L}_C$ with the same constants (C, ν, T) as for $\mathcal{L}$.

We now follow the same approach as in the previous example and bound the $\mathbb{L}_{1 \rightarrow 2}(\mathbb{1}/(N+1))$ norm of the semi-group $S_t = \exp(t\mathcal{L})$ that is generated by the Liouvillian $\mathcal{L}$ in Eq. (77). We note (see Ref. 25) that the supremum in the definition of $\|S_t\|_{1 \rightarrow 2} = \sup_f \|S_t\|_2 \|f\|_1^{-1}$ for Shatten norms $\|f\|_p^p = \text{tr}[|f|^p]$ is always attained for rank one projectors $f = |\psi\rangle\langle\psi|$. Hence, we obtain for the $\mathbb{L}_{1 \rightarrow 2}(\mathbb{1}/(N+1))$ norm

$$\|S_t\|_{1 \rightarrow 2, \mathbb{1}/(N+1)}^2 = (N+1) \sup_{\langle\psi|\psi\rangle=1} \langle\psi|S_{2t}(|\psi\rangle\langle\psi|)|\psi\rangle \quad (78)$$

$$= (N+1) \sup_{\langle\psi|\psi\rangle=1} \sum_{m,l=0}^N e^{-2t\lambda_{ml}} |\langle\psi|v_{ml}|\psi\rangle|^2, \quad (79)$$

where λ_{ml} are the eigenvalues of $-\mathcal{L}$ and v_{ml} are its (right) eigenvectors. From basic Fourier analysis, the eigenvectors and eigenvalues can all be evaluated analytically, yielding

$$\begin{aligned} 0 \leq m \leq N, \quad \lambda_{m0} &= -2 \left(1 - \cos\left(\frac{2\pi m}{N+1}\right) \right), & v_{m0} &= \frac{1}{\sqrt{N+1}} \sum_{k=0}^N e^{i\frac{2\pi mk}{N+1}} |k\rangle\langle k|, \\ l \neq 0, \quad \lambda_{ml} &= -2, & v_{ml} &= |m\rangle\langle m \oplus l|. \end{aligned} \quad (80)$$

Then, Eq. (79) can be evaluated explicitly as

$$\|S_t\|_{1 \rightarrow 2, \mathbb{1}/(N+1)}^2 = (N+1) \sum_{m=0}^N (e^{-2t\lambda_{m0}} |\langle\psi|v_{m0}|\psi\rangle|^2 + \sum_{l=1}^N e^{-t\lambda_{ml}} |\langle\psi|v_{ml}|\psi\rangle|^2) \quad (81)$$

$$\leq \sum_{m=0}^N (e^{-2t\lambda_{m0}} + (N+1) \sum_{l=1}^N e^{-2t\lambda_{ml}}), \quad (82)$$

by observing that $\sum_m |\langle\psi|v_{ml}|\psi\rangle|^2 \leq 1$ for all $l \neq 0$, which follows from simple linear algebra. We get the expression for $l = 0$ that $\sum_{k=0}^N \exp(i\frac{2\pi mk}{N+1}) |\psi_k|^2 \leq \langle\psi|\psi\rangle$, where $|\psi\rangle := \sum_i \psi_i |i\rangle$. We may now bound the $\mathbb{L}_{1 \rightarrow 2}(\mathbb{1}/(N+1))$ norm of the semigroup as

$$\|S_t\|_{1 \rightarrow 2, \mathbb{1}/(N+1)}^2 \leq 1 + \sum_{m=1}^N e^{-2t(1-\cos(\frac{2\pi m}{N+1}))} + (N+1)^2 e^{-4t}, \quad (83)$$

since we consider the supremum over normalized states $|\psi\rangle$. By using a trick from Ref. 7, we may bound the exponential of the cosine by a quadratic function and bound the sum in Eq. (83) by a Gaussian integral which then yields

$$\sum_{m=1}^N e^{-2t(1-\cos(\frac{2\pi m}{N+1}))} \leq 2 \sum_{m=1}^N e^{-4t \frac{m^2}{(N+1)^2}} \leq 2e^{-\frac{4t}{(N+1)^2}} \left(1 + \sqrt{\frac{(N+1)^2}{4t}} \right). \quad (84)$$

Finally, we observe that for $t \leq T = (N+1)^2/16$, this expression can be bounded by the polynomially decaying function

$$\|S_t\|_{1 \rightarrow 2, \mathbb{1}/(N+1)}^2 \leq \left(1 + 2e^{-\frac{4t}{(N+1)^2}} \left(1 + \sqrt{\frac{(N+1)^2}{4t}} \right) + (N+1)^2 e^{-4t} \right) \leq \sqrt{\frac{8(N+1)^2}{t}}. \quad (85)$$

The constants in the ultracontractive bound can be read off directly, $C = 8(N+1)^2$ and $\nu = 1$ for $T = (N+1)^2/16$. Due to the comparison argument with $\mathcal{E}(f)_C \geq \mathcal{E}(f)$ and by virtue of Theorem 3, we can state the Nash II inequality for the semigroup generated by $\mathcal{L}_C$,

$$\|f\|_{2, \mathbb{1}/(N+1)}^6 \leq 2^{10} (N+1)^2 \left(\mathcal{E}(f) + \frac{16}{(N+1)^2} \|f\|_{2, \mathbb{1}/(N+1)}^2 \right) \|f\|_{1, \mathbb{1}/(N+1)}^4. \quad (86)$$

We can therefore infer a mixing time bound that behaves as

$$\tau_{\text{mix}} = O((N+1)^2), \quad (87)$$

which gives the exact mixing time, since a matching lower bound can be obtained by inputting a rank one trial function. By comparison, the spectral gap bound estimate yields $|\lambda_1| \geq (N+1)^{-2}$ that would scale together with the pre-factor $\log(\|\rho^{-1}\|)$ as $\tau_{\text{mix}} = O((N+1)^2 \log(N+1))$ and therefore, the Nash inequality completely removes the pre-factor contribution.

ACKNOWLEDGMENTS

M.J.K. was supported by the Carlsbergfond and the Villum foundation. K.T. was supported by the Institute for Quantum Information and Matter, a NSF Physics Frontiers Center with support of the Gordon and Betty Moore Foundation (Grant Nos. PHY-0803371 and PHY-1125565).

- ¹ Beigi, S., “Sandwiched Rényi divergence satisfies data processing inequality,” *J. Math. Phys.* **54**(12), 122202 (2013).
- ² Carbone, R. and Sasso, E., “Hypercontractivity for a quantum Ornstein–Uhlenbeck semigroup,” *Probab. Theory Relat. Fields* **140**(3-4), 505–522 (2008).
- ³ Carlen, E. A., Kusuoka, S., and Stroock, D. W., “Upper bounds for symmetric Markov transition functions,” in *Annales de l’IHP Probabilités et Statistiques* (Gauthier-Villars, 1987), Vol. 23, pp. 245–287.
- ⁴ Cubitt, T. S., Lucia, A., Michalakis, S., and Perez-Garcia, D., “Stability of local quantum dissipative systems,” *Comm. Math. Phys.* **337**, 1275 (2015); preprint [arXiv:1303.4744](https://arxiv.org/abs/1303.4744) (2013).
- ⁵ Davies, E. B., *Heat Kernels and Spectral Theory* (Cambridge University Press, 1990), Vol. 92.
- ⁶ Davies, E. B. and Simon, B., “Ultracontractivity and the heat kernel for Schrödinger operators and Dirichlet Laplacians,” *J. Funct. Anal.* **59**(2), 335–395 (1984).
- ⁷ Diaconis, P. and Saloff-Coste, L., “Nash inequalities for finite Markov chains,” *J. Theor. Probab.* **9**(2), 459–510 (1996).
- ⁸ Eisler, V., “Crossover between ballistic and diffusive transport: The quantum exclusion process,” *J. Stat. Mech.: Theory Exp.* **2011**(06), P06007.
- ⁹ Gross, L., “Logarithmic Sobolev inequalities,” *Am. J. Math.* **97**, 1061–1083 (1975).
- ¹⁰ Kastoryano, M. J. and Brandao, F. G. S. L., “Quantum Gibbs samplers: The commuting case,” preprint [arXiv:1409.3435](https://arxiv.org/abs/1409.3435) (2014).
- ¹¹ Kastoryano, M. J. and Eisert, J., “Rapid mixing implies exponential decay of correlations,” *J. Math. Phys.* **54**(10), 102201 (2013).
- ¹² Kastoryano, M. J., Reeb, D., and Wolf, M. M., “A cutoff phenomenon for quantum Markov chains,” *J. Phys. A: Math. Theor.* **45**(7), 075307 (2012).
- ¹³ Kastoryano, M. J. and Temme, K., “Quantum logarithmic Sobolev inequalities and rapid mixing,” *J. Math. Phys.* **54**, 052202 (2013); e-print [arXiv:1207.3261](https://arxiv.org/abs/1207.3261).
- ¹⁴ King, C., “Additivity for unital qubit channels,” *J. Math. Phys.* **43**(10), 4641–4653 (2002).
- ¹⁵ King, C., “Hypercontractivity for semigroups of unital qubit channels,” *Commun. Math. Phys.* **328**(1), 285–301 (2014).
- ¹⁶ King, C. and Ruskai, M. B., “Minimal entropy of states emerging from noisy quantum channels,” *IEEE Trans. Inf. Theory* **47**(1), 192–209 (2001).
- ¹⁷ Montanaro, A. and Osborne, T. J., “Quantum boolean functions,” preprint [arXiv:0810.2435](https://arxiv.org/abs/0810.2435) (2008).
- ¹⁸ Muller-Hermes, A., Reeb, D., and Wolf, M. M., “Quantum subdivision capacities and continuous-time quantum coding,” *IEEE Trans. Inf. Theory* **61**, 565 (2014).
- ¹⁹ Pisier, G. and Xu, Q., “Non-commutative Lp-spaces,” in *Handbook of the Geometry of Banach Spaces* (Elsevier, 2003), Vol. 2, pp. 1459–1517.
- ²⁰ Saloff-Coste, L., “Lectures on finite Markov chains,” in *Lectures on Probability Theory and Statistics* (Springer, 1997), pp. 301–413.
- ²¹ Temme, K., “Thermalization time bounds for Pauli stabilizer Hamiltonians,” preprint [arXiv:1412.2858](https://arxiv.org/abs/1412.2858) (2014).
- ²² Temme, K., Kastoryano, M. J., Ruskai, M. B., Wolf, M. M., and Verstraete, F., “The χ^2 -divergence and mixing times of quantum Markov processes,” *J. Math. Phys.* **51**(12), 122201 (2010).
- ²³ Temme, K., Pastawski, F., and Kastoryano, M. J., “Hypercontractivity of quasi-free quantum semigroups,” *J. Phys. A: Math. Theor.* **47**(40), 405303 (2014).
- ²⁴ Varopoulos, N. T., “Hardy–Littlewood theory for semigroups,” *J. Funct. Anal.* **63**(2), 240–260 (1985).
- ²⁵ Watrous, J., “Notes on super-operator norms induced by Schatten norms,” preprint [arXiv:quant-ph/0411077](https://arxiv.org/abs/quant-ph/0411077) (2004).
- ²⁶ Zegarlinski, B., “Hypercontractivity in non-commutative Lp spaces,” in *Stochastic Processes, Physics, and Geometry, Max Planck Institute for Mathematics in the Sciences, Leipzig, January 18-22, 1999* (American Mathematical Society, 2000), Vol. 28, p. 323.
- ²⁷ S_t will act on observables (the Heisenberg picture), and S_t^* on states (the Schrödinger picture). The conjugation is with respect to the Hilbert–Schmidt inner product.