

SUPPLEMENTARY MATERIAL FOR:

Title: Synthesis of Metallointercalator-DNA Conjugates on a Solid Support
Authors: R. Erik Holmlin, Peter J. Dandliker, and Jacqueline K. Barton*
Affiliation: Division of Chemistry and Chemical Engineering
California Institute of Technology
Pasadena, CA 91125

FIGURE CAPTIONS

Figure S1. (A) Schematic illustration of the N-hydroxysuccinimidyl (NHS) ester of $[\text{Rh}(\text{phi})_2(\text{bpy}')]^{3+}$. (B) Matrix assisted laser desorption ionization (MALDI) time of flight (TOF) mass spectrum of $[\text{Rh}(\text{phi})_2(\text{bpy}'\text{-NHS})]^{3+}$.

Figure S2. HPLC chromatograms of an unmodified (upper trace) and an aminoalkylated (lower trace) 15-mer monitored at 260 nm. HPLC Conditions: C18 stationary phase; 50 mM ammonium acetate, pH 6.5/acetonitrile mobile phase; 0-15 % acetonitrile in 30 min. The aminoalkylated DNA shows a later retention time, consistent with increased hydrophobicity introduced by the linker.

Figure S3. HPLC chromatograms for the 3',5'-*bis*-heterometallated oligonucleotide pictured in Figure 9 monitored at 260 nm (top), 390 nm (middle), and 470 (nm) bottom. The 3',5'-*bis*-heterometallated oligonucleotide is the major product, and all eight diastereomers of this material coelute. HPLC Conditions: C18 stationary phase; 50 mM ammonium acetate, pH 6.5/acetonitrile mobile phase; 0-50 % acetonitrile in 75 min.

Supplementary Material




Figure S1

2

Supplementary Material




Figure S2




Figure S3