

Ternary H₂SO₄ - H₂O - NH₃ neutral and charged nucleation rates for a wide range of atmospheric conditions

Andreas Kürten, Federico Bianchi, Joao Almeida, Jonathan Duplissy, Eimear M. Dunne, Martin Breitenlechner, Arnaud P. Praplan, Ismael K. Ortega, Oona Kupiainen, Linda Rondo, Sebastian Ehrhart, Jasper Kirkby, Joachim Curtius, and CLOUD Collaboration

Citation: [AIP Conference Proceedings](#) **1527**, 310 (2013); doi: 10.1063/1.4803265

View online: <http://dx.doi.org/10.1063/1.4803265>

View Table of Contents: <http://scitation.aip.org/content/aip/proceeding/aipcp/1527?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Multi-species nucleation rates in CLOUD](#)

AIP Conf. Proc. **1527**, 326 (2013); 10.1063/1.4803269

[Variations of longwave downwelling irradiance \(LDI\) due to different atmospheric factors in Moscow](#)

AIP Conf. Proc. **1531**, 580 (2013); 10.1063/1.4804836

[Assessment of aerosol hygroscopic growth using an elastic LIDAR and BRAMS simulation in urban metropolitan areas](#)

AIP Conf. Proc. **1531**, 360 (2013); 10.1063/1.4804781

[General approach to the formulation and solution of the multi-parameter inverse problems of atmospheric remote sensing](#)

AIP Conf. Proc. **1531**, 240 (2013); 10.1063/1.4804751

[Hyperspectral retrieval of surface reflectances: A new scheme](#)

AIP Conf. Proc. **1531**, 51 (2013); 10.1063/1.4804705

Ternary H₂SO₄-H₂O-NH₃ Neutral and Charged Nucleation Rates for a Wide Range of Atmospheric Conditions

Andreas Kürten^a, Federico Bianchi^b, Joao Almeida^c, Jonathan Duplissy^d, Eimear M. Dunne^e, Martin Breitenlechner^f, Arnaud P. Praplan^d, Ismael K. Ortega^d, Oona Kupiainen^d, Linda Rondo^a, Sebastian Ehrhart^a, Jasper Kirkby^c, Joachim Curtius^a, and the CLOUD collaboration

^a*Institute for Atmospheric and Environmental Sciences, Goethe University of Frankfurt, 60438 Frankfurt am Main, Germany*

^b*Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen, Switzerland*

^c*CERN, 1211 Geneva, Switzerland*

^d*Department of Physics, University of Helsinki, FI-00014 Helsinki, Finland*

^e*Finnish Meteorological Institute, Kuopio Unit, 70100 Kuopio, Finland*

^f*Institute of Ion Physics and Applied Physics, University of Innsbruck, 6020 Innsbruck, Austria*

Abstract. The formation of new particles for the ternary system involving sulfuric acid, water vapor and ammonia has been studied in detail. The nucleation rates were obtained from experiments at the CERN CLOUD chamber which allows the measurement of new particle formation under very well defined conditions. Some of its key features are the suppression of contaminants at the technological limit and a very precise control of a wide range of temperatures, trace gas concentrations and nucleation rates. The effect of ionizing radiation on the ternary nucleation rates was investigated by using the CERN proton synchrotron beam (beam conditions), natural galactic cosmic rays (gcr conditions) as well as the high voltage clearing field inside the chamber to suppress the effect of charges (neutral conditions). The dependence of the nucleation rate on ion concentration, sulfuric acid and ammonia concentration as well as temperature was studied extensively. This way, an unprecedented set of data was collected giving insight into the role of neutral and charged ternary NH₃ nucleation and the relative importance of the different parameters.

Keywords: Chamber study, CLOUD experiment, New particle formation, Ammonia.

PACS: 92.60.Mt, 64.60.Q-

TERNARY H₂SO₄-H₂O-NH₃ NUCLEATION

New particle formation is an important atmospheric process affecting cloud formation, visibility and climate as well as air chemistry and human health. Generally, atmospheric boundary layer nucleation events cannot be explained by the binary nucleation of sulfuric acid and water vapor. Therefore, it had been suggested that ammonia could be an important compound which lowers the nucleation barrier when added as a ternary substance¹⁻⁴. Indeed, a recent CLOUD (Cosmics Leaving Outdoor Droplets) experiment has shown that ternary nucleation involving NH₃ is significantly enhancing the formation of new particles by a factor of 10 to 1000 over the nucleation

rates for the binary system⁵. However, the addition of ammonia can still not account for the high nucleation rates observed in the boundary layer at temperatures of 5°C or warmer. In these regions ternary organic vapors are most likely required to explain the high nucleation rates^{6,7}.

Nevertheless, at high latitudes or in the upper troposphere where temperatures are colder and other ternary vapors occur at too low concentrations, ammonia can still play an important role. Therefore, on a global scale the effect of ammonia for new particle formation can be significant and needs to be studied. Ammonia is also a compound of interest because its emissions are, similar to SO₂, strongly linked with anthropogenic activity. Ammonia emissions from livestock waste, volatilization of NH₃-based-fertilizers, biomass burning and crops comprise a big part of the total emissions. It is thought that these have led to a doubling of the total ammonia emissions since pre-industrial times and that a significant further increase will occur in the future^{8,9}.

With respect to the climate discussion it is of great interest to elucidate the impact of ionizing radiation on new particle and CCN formation. Recent results from CLOUD have shown that galactic cosmic rays can lead to a significant enhancement of the nucleation rates for the binary and the ternary NH₃ system under some conditions (FIGURE 1)⁵. It is, however, necessary to extend the available data set for a wider range of tropospheric conditions – especially with respect to colder temperatures. In a second step this data can then be used in models to investigate the sensitivity of the nucleation rates and the formation of CCN due to the different parameters on a global scale.




FIGURE 1. Nucleation rate as function of the ammonia mixing ratio (data from ref. 5). Temperature was 278 K and [H₂SO₄] was 6.3 × 10⁷ cm⁻³. Nucleation rates were measured at neutral (J_n), galactic cosmic ray (J_{gr}) and CERN pion beam conditions (J_{ch}).

EXPERIMENTAL METHODS

Nucleation rate measurements presented have been performed during the CLOUD3, CLOUD5 and CLOUD7 experiments at CERN between October 2009 and December 2012. The CLOUD chamber is designed to study the effect of ionizing radiation on particle formation, growth and activation to ice or cloud droplets under extremely well controlled conditions. It consists of a 3 m diameter electro-polished stainless steel cylinder with a volume of 26.1 m³. The chamber is surrounded by a thermal housing, and air circulating in the space between the chamber and the insulation was used to control the internal temperature between 208 and 300 K during the ternary NH₃ nucleation experiments.

Sulfuric acid has been measured by Chemical Ionization Mass Spectrometry (CIMS) at concentrations between $\sim 1 \times 10^6$ and $\sim 1 \times 10^9$ cm⁻³ during nucleation events. To achieve a homogenous mixing the sulfuric acid production was initiated in-situ through illumination of the chamber with a UV fibre-optic system. The ammonia originated from a gas bottle and was further diluted with nitrogen to achieve the desired concentration. The diluted NH₃ flow was then introduced into the CLOUD chamber where it was rapidly mixed into the chamber air by means of two magnetically driven stainless steel fans. The instrumentation used to measure the ammonia concentrations (between several pptv and ~ 1000 pptv) comprised of a Long Path Absorption Photometer (LOPAP)¹⁰, an Ion Chromatograph (IC)¹¹ and a Proton-Transfer Mass Spectrometer (PTR-MS)¹². The determination of the [NH₃] at concentrations lower than several pptv and at temperatures lower than the freezing point of water is challenging due to instrumental limitations. Under these conditions the ammonia concentrations were mainly derived from estimating the ammonia background (when no ammonia was intentionally flown into the chamber) as well as from calibration measurements which relate the amount of ammonia flown into the chamber and its measured concentration at 5°C.

TABLE 1 summarizes the parameter space which was covered at CLOUD for the ternary NH₃ system. Several hundred individual nucleation rate measurements were obtained from these experiments.

TABLE 1. Parameter space covered by the CLOUD experimental runs. ^aThe lowest ammonia concentrations under background conditions were not directly measured (see text for details). ^bDifferent relative humidities (RH) were tested but it was found that the nucleation rates were not very sensitive to changes in the humidity for the range studied.

Parameter	range
Temperature	208 to 300 K
[H ₂ SO ₄]	$\sim 1 \times 10^6$ to 1×10^9 cm ⁻³
[NH ₃]	^a $< 10^{-2}$ to $\sim 10^3$ pptv
^b RH	10 to 100%
Ion concentration	0 (neutral) to $\sim 5 \times 10^3$ cm ⁻³

THEORETICAL CALCULATIONS

Recently, a lot of progress has been made in describing the formation of new particles from first principles through quantum chemical calculations^{13,14}. Such calculations have been made also for the ternary system involving ammonia. The results from the ACDC model¹³ are compared to the experimentally determined nucleation rates. In addition, calculations have been made which yield the nucleation rate at the kinetic limit for a given concentration of sulfuric acid. These results yield the upper limit of the nucleation rate and are used as a cross-check for the experimental values. Furthermore, they are used to identify the conditions where nucleation is limited by the kinetics of the colliding clusters.

ACKNOWLEDGMENTS

We would like to thank CERN for supporting CLOUD with important technical and financial resources, and for providing a particle beam from the CERN Proton Synchrotron. This research has received funding from the EC Seventh Framework Programme (Marie Curie Initial Training Network "CLOUD-ITN" no. 215072, MC-ITN "CLOUD-TRAIN" no. 316662, and ERC-Advanced "ATMNUCLE" grant no. 227463), the German Federal Ministry of Education and Research (project nos. 01LK0902A and 01LK1222A), the Swiss National Science Foundation (project nos. 200020_135307 and 206620_130527), the Academy of Finland (Center of Excellence project no. 1118615), the Academy of Finland (135054, 133872, 251427, 139656, 139995, 137749, 141217, 141451), the Finnish Funding Agency for Technology and Innovation, the Nessling Foundation, the Austrian Science Fund (FWF; project no. P19546 and L593), the Portuguese Foundation for Science and Technology (project no. CERN/FP/116387/2010), the Swedish Research Council, Vetenskapsrådet (grant 2011-5120), the Presidium of the Russian Academy of Sciences and Russian Foundation for Basic Research (grants 08-02-91006-CERN and 12-02-91522-CERN), and the U.S. National Science Foundation (grants AGS1136479 and CHE1012293).

REFERENCES

1. D. J. Coffman and D. A. Hegg, *J. Geophys. Res.* **100**, D4, 7147–7160 (1995).
2. S. M. Ball, D. R. Hanson and F. L. Eisele, *J. Geophys. Res.* **104**, D19, 23,709–23,718 (1999).
3. P. Korhonen et al., *J. Geophys. Res.* **104**, D21, 26,349–26,353 (1999).
4. D. R. Hanson and F. L. Eisele, *J. Geophys. Res.* **107**, D12, 4158, 10.1029/2001JD001100 (2002).
5. J. Kirkby et al., *Nature* **476**, 429–433 (2011).
6. A. Metzger et al., *Proc. Natl. Acad. Sci.*, doi/10.1073/pnas.0911330107 (2010).
7. M. Chen et al., *Proc. Natl. Acad. Sci.* **109**, 46, 18713–18718 (2012).
8. S. V. Krupa, *Environ. Pollut.* **124**, 179–221 (2003).
9. L. Clarisse et al., *Nature Geoscience*, DOI: 10.1038/NGEO551 (2009).
10. F. Bianchi et al., *Atmos. Meas. Tech.* **5**, 1719–1725 (2012).
11. A. P. Praplan et al., *Atmos. Meas. Tech.* **5**, 2161–2167 (2012).
12. M. Norman et al., *Atmos. Chem. Phys.* **9**, 2635–2645 (2009).
13. M. J. McGrath et al., *Atmos. Chem. Phys.* **12**, 2345–2355 (2012).
14. I. K. Ortega et al., *Atmos. Chem. Phys.* **12**, 225–235 (2012).