

Mapping local Hamiltonians of fermions to local Hamiltonians of spins

F Verstraete¹ and J I Cirac²

¹ Institute for Quantum Information, Caltech, Pasadena, CA, USA

² Max-Planck-Institut für Quantenoptik, Hans-Kopfermann-Straße 1, Garching, D-85748, Germany

E-mail: verstrae@caltech.edu and Ignacio.Cirac@mpq.mpg.de

Received 25 August 2005

Accepted 17 September 2005

Published 29 September 2005

Online at stacks.iop.org/JSTAT/2005/P09012

[doi:10.1088/1742-5468/2005/09/P09012](https://doi.org/10.1088/1742-5468/2005/09/P09012)

Abstract. We show how to map local fermionic problems onto local spin problems on a lattice in any dimension. The main idea is to introduce auxiliary degrees of freedom, represented by Majorana fermions, which allow us to extend the Jordan–Wigner transformation to dimensions higher than one. We also discuss the implications of our results in the numerical investigation of fermionic systems.

Keywords: rigorous results in statistical mechanics, Hubbard model (theory), spin chains, ladders and planes (theory), density matrix renormalization group calculations

Many of the interesting strong correlation quantum phenomena that appear in Nature can be accounted for in terms of discretized models, i.e., in lattices. The corresponding effective theories often deal either with fermions, which can hop from one lattice site to another one as well as interact among them, or spins, which sit at different lattice sites. For local Hamiltonians in one spatial dimension, the Jordan–Wigner transformation (JWT) [1] allows us to map a *local* theory of fermions onto a *local* theory of spins, and thus those systems exhibit related physical phenomena. In higher dimensions, however, this transformation gives rise to nonlocal interactions between spins, and thus fermions and spins are expected to exhibit very different behaviour, at least when they are described in terms of local Hamiltonians.

In this paper we will show how one can always exactly map a local theory of fermions to the low energy sector of a local theory of spins in a lattice of arbitrary dimension. To achieve this, we introduce extra degrees of freedom in the form of Majorana fermions, which interact locally with the original ones, and which allow us to perform a JWT in higher dimensions than one. Similar ideas were previously used by Bravyi and Kitaev [2] in the context of simulating fermionic quantum-computational circuits with qubits; there they showed how local fermionic gates can be simulated with local gates on qubits if the quantum states are restricted to living in a specific sector of the Hilbert space that is determined by nonlocal gauge conditions. Also, Ball [3] has shown how these gauge conditions can be transformed into local terms of the Hamiltonian, except at the boundary where nonlocal terms remain. We present a mapping which has the advantage that the low energy sector is completely determined by local terms in the spin Hamiltonian. The spin Hamiltonians obtained are exactly of the type studied by Wen *et al* [4], where the opposite question was studied, i.e. how fermionic behaviour could originate out of the low energy sector of a bosonic Hamiltonian.

The question analysed in this paper is clearly of fundamental interest. However, it may also have important implications in the numerical investigations of fermionic lattice problems, e.g. the Fermi–Hubbard model in two dimensions. In fact, our original motivation was to develop a method to represent fermionic problems in terms of local spin problems, so that we could use the algorithms we have recently developed to deal with those systems in dimensions larger than one [5]. In that case, a direct numerical simulation in terms of the fermionic degrees of freedom was not a good choice, since in Fock space one always obtains nonlocal terms (which are directly related to the Jordan–Wigner transformation), and ground states of generic systems with long range interactions do not have to obey e.g. the area law [6] and other characteristic properties that allow for efficient simulations: ground states of local Hamiltonians are very special in the sense that they are completely determined by their local reduced density operators, and hence a variational method will work well if it allows one to approximate local properties well [7]. This is one of the reasons that it is important that the low energy sector of the spin Hamiltonian obtained is determined by local terms, and does not contain nonlocal gauge conditions. Apart from that, our results may also prove useful for other standard simulation methods, such as quantum Monte Carlo ones, since there one also effectively simulates spin Hamiltonians instead of fermionic ones directly.

From the mathematical point of view, the natural formalism for describing a state of fermions is second quantization. Within this language, the most general state of N

fermions can be parametrized as

$$|\psi\rangle = \sum_{i_1, i_2, \dots, i_N=0}^1 C_{i_1, i_2, \dots, i_N} \left(\hat{a}_1^\dagger\right)^{i_1} \left(\hat{a}_2^\dagger\right)^{i_2} \cdots \left(\hat{a}_N^\dagger\right)^{i_N} |\Omega\rangle$$

where \hat{a}_i^\dagger are creation operators obeying the usual anticommutation relations, and where $|\Omega\rangle$ denotes the vacuum. Note that, from the moment we consider second quantization, we have to enumerate the fermions and remember this ordering. After this choice, the associated Hilbert space (or Fock space) in which we can describe quantum states becomes a tensor product of spin 1/2s or qubits via the identification

$$|i_1, i_2, \dots, i_N\rangle \equiv \left(\hat{a}_1^\dagger\right)^{i_1} \left(\hat{a}_2^\dagger\right)^{i_2} \cdots \left(\hat{a}_N^\dagger\right)^{i_N} |\Omega\rangle.$$

So it is indeed simple to represent a collection of N fermions in a Hilbert space of N spins [8]. When trying to simulate fermionic systems, one has to work with the vector C_{i_1, \dots, i_N} which is defined on this Hilbert space of spins.

The obvious question is now of course what fermionic Hamiltonians and creation/annihilation operators look like in this spin representation. Let us e.g. consider the representation of the annihilation operator \hat{a}_k :

$$\langle\Omega| \left[\hat{a}_N^{j_N} \cdots \hat{a}_1^{j_1}\right] \hat{a}_k \left[\hat{a}_1^{\dagger i_1} \cdots \hat{a}_N^{\dagger i_N}\right] |\Omega\rangle = \delta_{j_k, 0} \delta_{i_k, 1} \prod_{n \neq k} \delta_{j_n, i_n} (-1)^{i_n}.$$

This can readily be derived by using the anticommutation relations. The effective Hamiltonian described in spin space can now easily be calculated. The transformation that we carried out is exactly equivalent to the JWT which maps commuting spin operators to anticommuting fermionic creation/annihilation operators. From this point of view, the Jordan–Wigner transformation is not just a very useful trick but a natural consequence of the formalism of second quantization. The Hamiltonian as defined on the Hilbert space of spins is thus obtained by carrying out a JWT on the fermionic Hamiltonian.

Let us illustrate this transformation for a specific example. Due to superselection rules, physical observables always contain an even number of creation/annihilation operators, and thus in the following we will consider fermionic observables of this sort. Here we recall how typical operators transform (we neglected constant terms):

$$\begin{aligned} \hat{a}_k^\dagger \hat{a}_k &\rightarrow \sigma_k^z \\ \hat{a}_k^\dagger \hat{a}_{k+n} + \text{c.c.} &\rightarrow \sigma_k^x S_{k,n} \sigma_{k+n}^x + \sigma_k^y S_{k,n} \sigma_{k+n}^y \end{aligned}$$

where $S_{k,n} = \otimes_{l=k+1}^{k+n-1} \sigma_l^z$ is the string operator, which is the most distinctive feature of this transformation and arises when considering terms for hopping between distant fermions.

The central question of this paper is now the following: let us assume we have a system of fermions on some lattice with a specific geometry; is it always possible to map a local Hamiltonian of fermions to a local Hamiltonian of spins in this way? The answer to this question could clearly depend on the geometry and the type of interaction. Let us first consider the one-dimensional case; one immediately sees that one can always choose the ordering of the fermions such that locality is preserved by the JWT. The situation seems to be very different in two dimensions. Let us, e.g., consider a translational invariant model with nearest neighbour interactions and hopping terms on a square lattice. Due to the string of σ^z operators, there exists no ordering of the fermions such that all terms for




Figure 1. Enumerating fermions on a square 2D lattice.

local hopping in both horizontal and vertical directions map to local operators in the spin representation. The same problem arises in all geometries different from a one-dimensional chain. In the following we will show how locality can nevertheless always be achieved by introducing extra degrees of freedom in the form of extra fermions.

For simplicity, let us consider a two-dimensional square lattice of spinless fermions with N columns and open boundary conditions. We enumerate the fermions as in figure 1 and consider interaction terms of the form

$$\hat{a}_i^\dagger \hat{a}_j \tag{1}$$

$$\hat{a}_i^\dagger \hat{a}_i \hat{a}_j^\dagger \hat{a}_j \tag{2}$$

where either $i = j$ or i and j label two geometrically neighbouring fermions. The JWT will map all operators of the form (2) to local spin operators, but the hopping terms of the form (1) will only result in local spin terms when $j = i \pm 1$ (i.e. all horizontal hopping terms and half of the vertical ones at the edges). So there is clearly only a problem with the vertical hopping terms.

The idea is now to introduce an extra fermion \hat{b}_i for each mode \hat{a}_i , and impose that these extra fermions are in the ground state $|\chi\rangle$ of the Hamiltonian

$$\mathcal{H}_{\text{aux}} = \sum_{\{i,j\}} \hat{P}_{ij} \equiv \sum_{\{i,j\}} -(\hat{b}_i + \hat{b}_i^\dagger)(\hat{b}_j - \hat{b}_j^\dagger),$$

where the set $\{i, j\}$ includes only pairs (i, j) that correspond to the directed edges in figure 2(a) from vertex i to j .³ It can readily be verified that all the \hat{P}_{ij} commute, $\hat{P}_{ij}^2 = \mathbb{I}$,

³ Note that if the number of columns N is odd, it is necessary to take an auxiliary lattice with $N - 1$ columns so as to have an even number of columns to close the loops in figure 2(a).




Figure 2. Two possible graphs specifying different Hamiltonians \mathcal{H}_{aux} for the auxiliary fermions.

and that the gap of the Hamiltonian \mathcal{H}_{aux} is 2. Its unique ground state has the property that

$$\forall (i, j) \in \{i, j\}, \quad \hat{P}_{ij}|\chi\rangle = |\chi\rangle.$$

Actually, we can define the operators $\hat{c}_i = (\hat{b}_i + \hat{b}_i^\dagger)$ and $\hat{d}_i = -i(\hat{b}_i - \hat{b}_i^\dagger)$ as two Majorana fermions making up the usual fermions [2].

Next we consider the system consisting of the original fermions \hat{a}_i and of the auxiliary system, together; geometrically, we assume that each site is occupied by two fermions, the original \hat{a}_i and the new one $\hat{b}_{i'}$. We leave the Hamiltonian on the auxiliary modes \mathcal{H}_{aux} unchanged, but alter the vertical hopping terms of the physical Hamiltonian in the following way:

$$\hat{a}_i^\dagger \hat{a}_j \rightarrow \hat{a}_i^\dagger \hat{a}_j \hat{P}_{i'j'} = \hat{a}_i^\dagger \hat{a}_j (i\hat{c}_{i'} \hat{d}_{j'}).$$

Clearly, all these vertical hopping terms commute with \mathcal{H}_{aux} , and hence the \hat{P}_{ij} are constants of motion (i.e. +1), and as a result the ground state of the full system will be the tensor product of the original ground state and $|\chi\rangle$.⁴ But the advantage of the new formulation is immediately clear when the fermions are reordered following $1, 1', 2, 2', \dots$ when doing second quantization. As a result, the hopping terms in the vertical direction will now correspond to quartic interaction terms which will remain local on performing a JW transformation: the string operator of k' will cancel that of k . Note that the configuration of the auxiliary fermions shown in figure 2(a) is only one possible choice and many other loop structures would lead to similar results (one possibility would e.g. be to have only loops with two vertical edges such as in figure 2(b)).

⁴ Strictly speaking, this is only true if the energy gap in the auxiliary system is greater than the energy that could potentially be gained by attributing negative weights to vertical hopping terms; this can always be guaranteed by multiplying the auxiliary Hamiltonian by a big enough factor.

But the problem is not quite solved yet, because the JW transformation will turn \mathcal{H}_{aux} into a nonlocal Hamiltonian. Note however that the JW does not map the vertical hopping at the alternating edges of the lattice hopping into nonlocal ones. Consider e.g. the pair $(N, N + 1)$ and the associated $\hat{P}_{N,N+1}$ at such an edge (here N is the number of columns; $N = 4$ in figure 1); let us now make the following substitution in \mathcal{H}_{aux} :

$$\begin{aligned}\hat{P}_{N-1,N+2} &\rightarrow \hat{P}_{N-1,N+2}\hat{P}_{N,N+1} \\ \hat{P}_{N-2,N+3} &\rightarrow \hat{P}_{N-2,N+3}\hat{P}_{N-1,N+2} \\ &\vdots \\ \hat{P}_{1,2N} &\rightarrow \hat{P}_{1,2N}\hat{P}_{2,2N-1}\end{aligned}$$

and similarly for all other rows. Due to the fact that all \hat{P} commute, this substitution does not affect the ground state and the gap of \mathcal{H}_{aux} . The advantage of using this new Hamiltonian is however obvious: the JW transformation will map it to a local Hamiltonian with local plaquette terms: the string of σ^z operators generated by $\hat{c}_{i'}$ will be cancelled by those of $\hat{c}_{(i+1)'}$. So we have indeed managed to transform a local Hamiltonian of fermions into a local Hamiltonian of spins! One could argue that this indicates that fermionic systems potentially represent the low energy sector of bosonic systems (see e.g. [4]).

Let us illustrate the kinds of terms that will appear in the spin Hamiltonian for the case of the square $N \times N$ lattice of spinless fermions. Without loss of generality, we assume N to be even. We arrange the auxiliary fermions as in figure 2(a), and imagine that we have two superimposed layers of $N \times N$ qubits, the first layer corresponding to the *physical* qubits and the second one to the auxiliary ones. We label the Pauli spin operators acting on the first layer as $X_{k,l}, Y_{k,l}, Z_{k,l}$ and those acting on the second layer as $\tilde{X}_{k,l}, \tilde{Y}_{k,l}, \tilde{Z}_{k,l}$ with $1 \leq k \leq N$ labelling the column and l the row. Let us systematically show what all terms in a fermionic form will look like in the spin picture.

First of all we observe that the following terms transform trivially:

$$\begin{aligned}\hat{a}_{k,l}^\dagger \hat{a}_{k,l} &\rightarrow Z_{k,l} - \mathbb{I} \\ \left(\hat{a}_{k,l}^\dagger \hat{a}_{k,l}\right) \left(\hat{a}_{k',l'}^\dagger \hat{a}_{k',l'}\right) &\rightarrow (Z_{k,l} - \mathbb{I})(Z_{k',l'} - \mathbb{I}).\end{aligned}$$

Horizontal hopping terms transform as

$$\hat{a}_{k,l}^\dagger \hat{a}_{k+1,l} + \hat{a}_{k+1,l}^\dagger \hat{a}_{k,l} \rightarrow (X_{k,l}X_{k+1,l} + Y_{k,l}Y_{k+1,l})\tilde{Z}_{k,l}.$$

Vertical hopping terms become

$$\begin{aligned}\hat{a}_{k,l}^\dagger \hat{a}_{k,l+1} + \hat{a}_{k,l+1}^\dagger \hat{a}_{k,l} &\rightarrow (X_{k,l}X_{k,l+1} + Y_{k,l}Y_{k,l+1})(-1)^{l+1} \left(\tilde{X}_{k,l}\tilde{Y}_{k,l+1}\right) \quad (k = \text{odd}) \\ &\rightarrow (X_{k,l}X_{k,l+1} + Y_{k,l}Y_{k,l+1})(-1)^{l+1} \left(\tilde{Y}_{k,l}\tilde{X}_{k,l+1}\right) \quad (k = \text{even})\end{aligned}$$

depending on whether k is even or odd.

The gauge conditions can be imposed by including the following six-body local terms in the spin Hamiltonian:

$$\begin{aligned}\sum_{l=\text{odd}, k=\text{odd}} (Z_{k+1,l}Z_{k,l+1}) \left(\tilde{Y}_{k,l}\tilde{Y}_{k+1,l}\tilde{Y}_{k,l+1}\tilde{Y}_{k+1,l+1}\right) \\ \sum_{l=\text{odd}, k=\text{even}} (Z_{k+1,l}Z_{k,l+1}) \left(\tilde{X}_{k,l}\tilde{X}_{k+1,l}\tilde{X}_{k,l+1}\tilde{X}_{k+1,l+1}\right)\end{aligned}$$

$$\sum_{l=\text{even}, k=\text{odd}} (Z_{k,l} Z_{k+1,l+1}) \left(\tilde{Y}_{k,l} \tilde{Y}_{k+1,l} \tilde{Y}_{k,l+1} \tilde{Y}_{k+1,l+1} \right) \\ \sum_{l=\text{even}, k=\text{even}} (Z_{k,l} Z_{k+1,l+1}) \left(\tilde{X}_{k,l} \tilde{X}_{k+1,l} \tilde{X}_{k,l+1} \tilde{X}_{k+1,l+1} \right).$$

It can easily be checked that these terms commute and also commute with the hopping terms. Finally, boundary conditions are required to fix the gauge uniquely (note that an ordering of the fermions and the corresponding JWT was crucial to make these terms local):

$$\sum_{k=\text{odd}} \tilde{X}_{k,1} Z_{k+1,1} \tilde{X}_{k+1,1} \\ \sum_{k=\text{odd}} \tilde{X}_{k,N} Z_{k,N} \tilde{X}_{k+1,N} \\ \sum_{k=\text{odd}} \tilde{X}_{N,k} Z_{N,k+1} \tilde{X}_{N,k+1} \\ \sum_{k=\text{even}} \tilde{Y}_{1,k} Z_{1,k+1} \tilde{Y}_{1,k+1}.$$

The sum of all these terms (of course with the appropriate prefactors of the original model) is the local spin Hamiltonian whose low energy sector is equivalent to the original fermionic model. Let us try to get some insight into all these terms. The hopping terms give rise to the well known $XX + YY$ interactions, but these are mediated by a \tilde{Z} term in the horizontal case and a $\tilde{X}\tilde{Y}$ term in the vertical case; it is intriguing that the noncommuting nature of these extra terms should be related to the topological features of fermionic systems. Let us now have a closer look at the gauge conditions. Let us first assume that all spins in the first layer are completely polarized in the Z direction; this would correspond to a completely trivial Hamiltonian for the original fermions. The qubits in the second layer would then interact via plaquette interactions of the form $XXXX$ or $YYYY$, as all ZZ expectation values are equal to 1. It can readily be checked that the effective Hamiltonian acting on the second layer is then exactly equal to the celebrated toric code Hamiltonian of Kitaev [9] if one rotates the lattice over $\pi/4$: the topological features of the toric code states seem to be exactly the ones needed to mediate the desired mapping from local interactions of fermions to spins. It would be very interesting to investigate this connection between the topological features of toric code states and emergent fermions further, and to arrive at a more intuitive explanation for the proposed mathematical construction.

This construction can readily be generalized to fermions with spin and any regular lattice geometry; in e.g. a 3D cubic lattice, one would need two extra fermions per site (i.e. four Majorana fermions). The Hamiltonian characterizing the auxiliary fermions can in this 3D case still be made local by multiplying four commuting $c_i d_j$ terms with each other. A triangular lattice can e.g. be obtained by embedding it into a square lattice; in this particular case some fermions would not be coupled to any others, and extra Majorana fermions guiding extra horizontal hopping terms have to be introduced. One can also include long range hopping terms by multiplying them with multiple factors \hat{P}_{ij} . A 2D square lattice of fermions with an additional spin $1/2$ degree of freedom could either

be treated by considering two layers of the spinless system, or by working with spin $3/2$ objects instead of qubits.

An interesting problem arises when the same analysis is carried out on systems with periodic boundary conditions. Let us e.g. consider the case of a 1D ring; here we can add two Majorana fermions to each original fermion, and use the link between the first unpaired Majorana fermion and the last one as a tool to make the term for the original interaction between the first and last fermions quartic. However, the JW transformation will make the interaction between these two Majorana fermions nonlocal, and there does not seem to be a way around this: the topology of the ring forces us to have one nonlocal term in the final spin Hamiltonian⁵. Similarly, in the case of a cylinder one nonlocal term will appear, and in the case of a torus two nonlocal terms. In other words, we can speculate that the ground state of the corresponding spin system has the property that it cannot be the unique ground state of a local spin Hamiltonian: the associated local Hamiltonian (i.e. without a nonlocal term) may possibly have a twofold-degenerate ground state, and a nonlocal operator which is topologically nontrivial will be needed to specify the sector. This phenomenon has indeed been identified in the case of the toric code states [9] of Kitaev, and is related to topological quantum order [10].

Instead of just considering regular lattices, one can ask the question of whether a similar mapping is possible in the case of a star-like geometry which e.g. occurs in multichannel Kondo-related impurity problems [11, 12]. Let us therefore consider an impurity coupled to two different kinds of fermions each having an extra spin degree of freedom. Enumerating the different fermions as in figure 3 (here fermions 6 and 7 correspond to the impurity), one finds that again a long range string operator will appear between fermions of the second type and the impurity (i.e. between $\{6, 17\}$ and $\{7, 18\}$). As in the previous case, we introduce two pairs of Majorana fermions and multiply the problematic hopping terms with the appropriate Majorana operators. This will indeed make the hopping terms local after the JWT. The problem in this case however is that after the JW there is no way to impose the identities $ic_k d_k = +1$ for both pairs in a local way. But here one can easily see that there is no need to impose that constraint: the effective Hamiltonian after the JW is block diagonal, one sector corresponding to $ic_k d_k = +1$ and the other one to $ic_k d_k = -1$; the second block is equivalent to the first one under local unitary operations. Indeed, the local transformation $a_i \rightarrow \exp(i\pi a_i^\dagger a_i) a_i$ on all fermions of one type will leave the terms for interaction between them invariant, and will only lead to a negative hopping term when coupling them to the impurity. So the ground states in the two sectors are related by local unitaries: no gauge fixing interaction term needs to be imposed, and when doing simulations on the spin Hamiltonian one can always check the sector one ends up in. So it is again enough to diagonalize a local spin Hamiltonian.

In conclusion, we have shown how it is possible to map a local problem of fermions onto a local problem of spins by using auxiliary Majorana fermions. This leads to a two-dimensional version of the Jordan–Wigner transformation. We believe that, apart from its fundamental interest, this map may become very useful in the context of numerical description of fermionic systems, since in that case local spin Hamiltonians seem to be

⁵ Note that this nonlocal term does not necessarily present a problem for the numerical simulation of these systems. In e.g. the XY chain, one just has to solve the problem in the sectors with even and odd numbers of particles independently.




Figure 3. Enumerating fermions in the case of a two-channel Kondo-like impurity problem; 1–5 represent the spin up fermions of the first channel, 6–7 the impurity, 8–12 the spin down fermions of the first channel, and 13–17 and 18–22 respectively the spin up and down fermions of the second channel.

better approximated. An open question is whether one can use this mapping to obtain certain interesting spin Hamiltonians exactly by transforming it back into free fermions, as is done in one-dimensional investigations and in the recent work of Kitaev [13].

Acknowledgments

We thank S Bravyi, J Preskill and G Vidal for discussions. After completion of this work, we became aware of the work by Ball [3], and we thank X Wen for bringing this work to our attention. This work was supported by the Gordon and Betty Moore Foundation, European projects CONQUEST, TOPQIP and QUPRODIS, and the DFG (SFB 631).

References

- [1] Jordan P and Wigner E, 1928 *Z. Phys.* **47** 631
- [2] Bravyi S and Kitaev A, 2002 *Ann. Phys., NY* **298** 210
- [3] Ball R C, 2004 *Preprint cond-mat/0409485*
- [4] Wen X-G, 2002 *Phys. Rev. Lett.* **88** 11602
Levin M and Wen X-G, 2003 *Phys. Rev. B* **67** 245316

- Wen X-G, 2003 *Phys. Rev. D* **68** 024501
Levin M and Wen X-G, 2005 *Phys. Rev. B* **71** 045110
Levin M A and Wen X-G, 2004 *Preprint* [cond-mat/0407140](#)
Wen X G, 2005 *Ann. Phys., NY* **316** 1
- [5] Verstraete F and Cirac J I, 2004 *Preprint* [cond-mat/0407066](#)
[6] Srednicki M, 1993 *Phys. Rev. Lett.* **71** 666
Holzney C, Larsen F and Wilczek F, 1994 *Nucl. Phys. B* **424** 443
Fiola M *et al*, 1994 *Phys. Rev. D* **50** 3987
Vidal G *et al*, 2003 *Phys. Rev. Lett.* **90** 227902
Plenio M B *et al*, 2005 *Phys. Rev. Lett.* **94** 060503
Hamma A, Ionicioiu R and Zanardi P, 2005 *Phys. Lett. A* **337** 22
- [7] Verstraete F and Cirac J I, 2005 *Preprint* [cond-mat/0505140](#)
[8] Zanardi P, 2002 *Phys. Rev. A* **65** 042101
Zanardi P, Lidar D and Lloyd S, 2004 *Phys. Rev. Lett.* **92** 060402
- [9] Kitaev A, 2003 *Ann. Phys., NY* **303** 2
- [10] Wen X G and Niu Q, 1990 *Phys. Rev. B* **41** 9377
- [11] Wilson K G, 1975 *Rev. Mod. Phys.* **47** 773
- [12] Verstraete F, Weichselbaum A, Schollwöck U, Cirac J I and von Delft J, 2005 *Preprint* [cond-mat/0504305](#)
[13] Kitaev A, 2005 *Preprint* [cond-mat/0506438](#)