THE MID-INFRARED EVOLUTION OF THE FU ORIONIS DISK

JOEL D. GREEN^{1,2}, OLIVIA C. JONES¹, LUKE D. KELLER³, CHARLES A. POTEET¹, YAO-LUN YANG²,
 WILLIAM J. FISCHER^{4,6}, NEAL J. EVANS II², BENJAMIN A. SARGENT¹, AND LUISA M. REBULL⁵

¹ Space Telescope Science Institute, Baltimore, MD 21218, USA

² Department of Astronomy, The University of Texas at Austin, Austin, TX 78712, USA

³ Department of Physics & Astronomy, Ithaca College, Ithaca, NY, USA

⁴ Goddard Space Flight Center, Greenbelt, MD 20771, USA

⁵ IPAC, Pasadena, CA 91125, USA

Received 2016 June 28; revised 2016 September 1; accepted 2016 September 5; published 2016 November 10

ABSTRACT

We present new SOFIA-FORCAST observations obtained in 2016 February of the archetypal outbursting low-mass young stellar object FU Orionis, and we compare the continuum, solid-state, and gas properties with mid-infrared data obtained at the same wavelengths in 2004 with *Spitzer*-IRS. In this study, we conduct the first mid-infrared spectroscopic comparison of an FUor over a long time period. Over a 12-year period, *UBVR* monitoring indicates that FU Orionis has continued its steady decrease in overall brightness by $\sim 14\%$. We find that this decrease in luminosity occurs only at wavelengths $\lesssim 20 \mu\text{m}$. In particular, the continuum shortward of the silicate emission complex at $10 \mu\text{m}$ exhibits a $\sim 12\%$ ($\sim 3\sigma$) drop in flux density but no apparent change in slope; both the *Spitzer* and SOFIA spectra are consistent with a 7200 K blackbody. Additionally, the detection of water absorption is consistent with the *Spitzer* spectrum. The silicate emission feature at $10 \mu\text{m}$ continues to be consistent with unprocessed grains, unchanged over 12 years. We conclude that either the accretion rate in FU Orionis has decreased by $\sim 12\text{--}14\%$ over this time baseline or the inner disk has cooled, but the accretion disk remains in a superheated state outside the innermost region.

Key words: accretion, accretion disks – infrared: stars – planets and satellites: formation – stars: formation – stars: protostars – stars: variables: T Tauri, Herbig Ae/Be

1. INTRODUCTION

The low-mass pre-main-sequence star FU Orionis (hereafter, FU Ori) brightened by $\Delta B = 6$ mag in 1936 (Herbig 1977) and subsequently has declined slowly ($\sim 0.013 \text{ mag yr}^{-1}$; Kenyon et al. 2000) to the present day (further examined in this work). This burst in brightness established a class of variable star (FU Orionis objects, or “FUors”; Hartmann & Kenyon 1996; Hartmann 1998) that ~ 10 other stars have since mimicked. Paczynski (1976) first proposed that FUors are the result of a sudden cataclysmic accretion of material from a reservoir that had built up in the circumstellar disk surrounding a young stellar object. In this model, the accretion rate rises from the typical rate for a T Tauri star ($\dot{M} \lesssim 10^{-7} M_{\odot} \text{ yr}^{-1}$) up to $\dot{M} \sim 10^{-4} M_{\odot} \text{ yr}^{-1}$, and then decays over an e-fold time of $\sim 10\text{--}100$ years (Hartmann & Kenyon 1985; Lin & Papaloizou 1985, pp. 981–1072; Bell & Lin 1994). Over the entire outburst, the star could accrete $\sim 0.01 M_{\odot}$ of material, roughly the entire mass of the minimum-mass solar nebula or a typical T Tauri disk (Andrews & Williams 2005). On average, FUor outbursts should occur (or recur) about five times per star formed in the local region of the Milky Way (Hartmann & Kenyon 1996); there have been ~ 10 FUors observed within a distance of 1.5 kpc in the last 80 years, compared with the star-formation rate in the same volume (about three per 50 years). These bursts also modify the protoplanetary disk chemistry and require a very different model than simple magnetospheric accretion (e.g., Green et al. 2006; Quanz et al. 2007; Zhu et al. 2007; Romanova et al. 2009; Cieza et al. 2016), usually invoked for accreting young stars. If FUors represent a short-duration stage that all young stars undergo, then

understanding their properties is vital to models of planet formation and the evolution of protoplanetary gas-rich disks.

The luminosity outburst is due to a sudden increase in the rate at which material accretes from the disk onto the star. Models explain the rise in accretion rate with the development of instabilities in the disk, but they differ as to where the instabilities originate. “Inside-out” models are those in which the instability develops at several astronomical units, where the lower temperatures, and therefore lower ionization fraction, render the magnetorotational instability as an inefficient accretion mechanism (e.g., Zhu et al. 2009b; Martin et al. 2012). “Outside-in” models are those in which gravitationally bound fragments develop at large radii (tens of astronomical units or more) due to infall from a protostellar envelope, then migrate inward through the disk, ultimately generating an outburst when they accrete onto the star (e.g., Vorobyov & Basu 2015). A third option that has been posited is an external perturber—the interaction of the disk with an exterior massive object that alters the balance of accretion and triggers the cascading flow (Bonnell & Bastien 1992), although recent studies suggest the refill rate is too slow to occur on few co-orbit timescales (Green et al. 2016). The wavelength dependence of any postoutburst variability can distinguish among these classes of models.

Dust grain growth, processing, and sedimentation are tracked by mid-infrared (mid-IR) dust features. Processing and grain growth in the disk can begin as early as the protostellar stage, when the disk is heavily extinguished. Most silicate features around low-mass young stars are consistent with both grain growth and high crystallinity fractions (e.g., Olofsson et al. 2009; Watson et al. 2009). Because grains from the ISM enter the disk as amorphous material, high crystalline fractions in young stars indicate that dust processing occurs early on.

⁶ NASA Postdoctoral Program Fellow.

Table 1
Observing Log

Obs Instrument	AORIDs	Filter(s)	On Source Int. Time	Num. Coadded Exp.	Total Exp. Time
FORCAST Imaging	0401461, 040146165	F056/F253	60	1	184
FORCAST Imaging	0401462, 041046166	F077/F315	60	1	184
FORCAST Imaging	0401463, 040146163	F111/F348	147 ^a	1	~400 ^a
FORCAST Grism	04014616	G063	200	2	550 ^b
FORCAST Grism	04014617	G111	3600	5	9225 ^c
FORCAST Grism	04014618	G227	1000	1	2577
IRS Stare	3569920	SL/SH/LH	18	1	...

Notes. Observations were performed over two different flights, 272 (2016 February 04) and 279 (2016 February 18).

^a The F111 band was observed in seven separate integrations; six were 13 s on source and one was 69 s on source. We report the combined exposure time and approximate overhead.

^b The exposure on Flight 272 exhibited ~50% worse S/N than the exposure on Flight 279 and was not used in the final spectrum presented in this work.

^c The two exposures on Flight 272 exhibited 10% higher continuum than the three exposures on Flight 279, further misaligned with photometric measurement, and were not used in the final spectrum presented in this work.

EX Lup, the archetype of a smaller magnitude burst from a T Tauri disk, exhibited crystalline grain production during its outburst (Ábrahám et al. 2009), due to enhanced heating. However, these crystalline grains disappeared during quiescence, likely due to X-ray amorphization of the grains, or dust transport either radially or vertically. In contrast, FUors show no evidence for processed grains during outburst, only grain growth (Quanz et al. 2007). No preoutburst mid-IR spectra of FUors exist to constrain their historical grain properties, and it may be that the silicate grains are modified during the outburst. For protoplanetary disks, the 10 and 20 μm silicate features probe optically thin dust at specific temperatures on the disk surface; as the disk heats and then cools, the radii probed by these features change accordingly and may evolve during the burst (e.g., Hubbard & Ebel 2014). As the disk cools, we might expect additional changes to occur, as grains are recirculated, modified, merged, or destroyed. During an FUor outburst, the temperature at 1 au may approach 1000 K or even higher; this becomes comparable to the temperature to form crystalline silicates or remove volatiles, which can affect habitable-zone planet formation (Hubbard & Ebel 2014). Farther out in the envelope, episodic increases in temperature can cause irreversible chemical processes, such as the observed conversion of mixed CO–CO₂ ice into pure CO₂ (Kim et al. 2012; Poteet et al. 2013).

Mid-IR spectroscopic variability has been only sparsely examined, although mid-IR evolution during bursts has been modeled (e.g., Bell & Lin 1994; Zhu et al. 2007; Johnstone et al. 2013). The biggest difference between the mid-IR SED of FU Ori and that of more typical young stellar objects is the contribution of the superheated inner disk. The innermost regions of the disk are expected to reach temperatures of 6000–8000 K, comparable to the stellar surface but over a vastly greater emitting area (the inner radius of the disk; Hartmann & Kenyon 1996; Hartmann 1998). Therefore the innermost disk radii are no longer coupled to any dust, which would be destroyed or ejected out to a significant fraction of an astronomical unit and would not be an opacity source. Thus, FU Ori’s SED is dominated at *all wavelengths* by disk emission, even in the optical/near-IR, and changes at optical/near-IR wavelengths would reflect changes in the superheated disk rather than the central protostar.

FU Ori ($D = 450\text{--}500$ pc; Hartmann & Kenyon 1996) is a wide ($0''.5$) binary system consisting of a outbursting young stellar object modeled as a $0.3 M_{\odot}$ with a $0.04 M_{\odot}$ circumstellar

disk (Zhu et al. 2007) at an A_V of ~ 1.8 (Zhu et al. 2010) and a more massive young companion ($M > 0.5 M_{\odot}$; Reipurth & Aspin 2004; Wang et al. 2004; Hales et al. 2015). The companion (FU Ori S) is at 227 au projected separation (Pueyo et al. 2012) at an A_V of 10 ± 2 (Pueyo et al. 2012).

In this work, we present new mid-IR photometry and spectroscopy of FU Ori obtained with the Stratospheric Observatory for Infrared Astronomy (SOFIA) in 2016, and we compare with previous observations with the *Spitzer Space Telescope* in 2004. We find that the continuum flux density at wavelengths shorter than 8 μm (arising from the hot inner disk) has decreased by 12%–14%, dominating the change in bolometric luminosity. The continuum at wavelengths greater than 10 μm has decreased only 7% within uncertainties. Additionally, we find that the H₂O vapor absorption resulting from the superheated disk photosphere remains present. Finally, we note no significant changes in the 10 μm silicate emission feature, which shows no evidence of crystallinity. Taken together, these observations suggest that the decrease in viscous dissipation energy over this period has occurred predominantly in the innermost section of the disk, which is changing rapidly.

2. OBSERVATIONS

2.1. SOFIA-FORCAST Spectroscopy

We observed FU Ori with the Faint Object infraRed CAmera for the SOFIA Telescope (FORCAST; Herter et al. 2012) on board the SOFIA (Young et al. 2012) on two separate flights (on 2016 February 4 and 18, flights 272 and 279, respectively); see Table 1 for the Observing Log. The observations were performed in C2N (two-position chop with nod) mode, using the $4''.7$ slit, and the NMC (nod-match-chop) pattern in which the chops and nod are in the same direction and have the same amplitude. In each case, an off-source calibrator was selected from an ensemble of observations from cycles 2–4, using the observation closest in zenith angle and altitude to the science target (A. Helton 2016, private communication). We did not use dithering. The FORCAST grism was used in three spectral bands, G063 ($\sim 5\text{--}8 \mu\text{m}$), G111 ($\sim 9\text{--}13 \mu\text{m}$), and G227 ($\sim 18\text{--}25 \mu\text{m}$), from 5 to 25 μm to achieve a spectral resolving power of $R \sim 200$. The achieved signal-to-noise ratios (S/Ns) for the individual grism segments were 30 (G063, measured from 5.7 to 6.3 μm), 46 (G111, measured from 11 to 12 μm), and 22 (G227, measured from 21 to

26 μm). The total observation time was 12,904 s (3.58 hr), including imaging, which totaled approximately 0.33 hr (see below).

The extracted SOFIA-FORCAST spectra are the best calibrated version of the Level 3 pipeline products, as identified in the SOFIA Data Cycle System. The G227 observation appears in only one exposure/AOR, but the G063 was observed on two separate occasions (one on each of two flights), and G111 was observed in five separate exposures to obtain a sufficient signal-to-noise ratio (three exposures on Flight 279 and two exposures on Flight 272). Notes on each individual grism observation follow.

G063. The Flight 279 G063 spectrum shows 50% higher noise than the Flight 272 spectrum, and we decided to exclude the low S/N data; this did not significantly affect the absolute flux density in the G063 band.

G111. However, the two Flight 272 G111 spectra are both 10% higher in flux density than the three Flight 279 spectra, and we excluded the *higher flux density* G111 spectra from the analysis. Additionally, the G111 spectrum includes several bands dominated by atmospheric ozone. Although the SOFIA pipeline attempts to correct for this using off-positions, the telluric correction cannot reliably remove the ozone feature because the ozone layer is inhomogeneous and the SOFIA beam tracks in and out of higher density regions on timescales shorter than the background-removal process. Thus we masked this section of spectrum ($\sim 9.3\text{--}10.05\ \mu\text{m}$) and did not consider it in this work.

G227. The G227 spectrum showed the highest overall uncertainty. The spectrum was rereduced using a custom pipeline (W. Vacca 2016, private communication). A preliminary version of a new asteroid response curve along with a different background region than that used for the original pipeline extraction was applied. The result shifted the G227 spectrum down $\sim 7\%$ from the default, in between the three longest-wavelength SOFIA photometric bands.

2.2. *Spitzer-IRS Spectroscopy*

FU Ori was observed by the InfraRed Spectrograph (Houck et al. 2004) on board the *Spitzer Space Telescope* (Werner et al. 2004) as part of the IRS_Disks guaranteed-time observing program (Houck et al. 2004) on 2004 March 4 (Table 1). The observations (AORID 3569920) were performed in Staring Mode, with two nods spaced by one-third of the slit width, and off-positions acquired only for the Short-Low (SL; $5.3\text{--}14\ \mu\text{m}$, $R \sim 60\text{--}120$) modules. The Short-High (SH; $10\text{--}20\ \mu\text{m}$, $R \sim 600$) and Long-High (LH; $20\text{--}38\ \mu\text{m}$, $R \sim 600$) modules did not include off-positions and were extracted as full aperture products (Green et al. 2006). To confirm that the IRS spectral reduction was consistent with the most recent calibration, we retrieved the data from the Cornell Atlas of *Spitzer* IRS Sources (CASSIS LRV7 and HRV1; Lebouteiller et al. 2011, 2015). The spectrum was then scaled to SL, adjusting SH fluxes upward by 2% and LH fluxes downward by 3%. Details on the original data reduction are published in Green et al. (2006) using the SMART package (Higdon et al. 2004).

The absolute calibration uncertainty of *Spitzer-IRS* is typically as large as 10%. However, the point-to-point (relative) uncertainty is $\sim 3\%$. After cross-calibrating SL with SH and LH as noted above, the absolute flux uncertainty is only the uncertainty in SL, which is 10%; however, the IRAC photometry, which covers the same bands, has an uncertainty

floor of only 3% (Carey et al. 2012). Thus we take the final uncertainty in absolute flux calibration for *Spitzer* to be 3%.

2.3. *Photometry*

We acquired contemporaneous FORCAST imaging in F056/F253, F077/F315, and F111/F348 in order to calibrate the spectral flux densities, characterize extended emission, and provide 2016 continuum levels from 25 to 35 μm . All imaging observations were acquired during Flight 272. A signal-to-noise ratio of >10 on source was achieved in exposures of 60 s or less in all six bands on each source. The SOFIA-FORCAST photometry was obtained from the Level 3 pipeline data products. The measured image quality of the SOFIA imaging ranges between $2''.7$ and $3''$ for F056, F077, and F111, and ranges between $3''.4$ and $3''.6$ for F253, F315, and F348; FORCAST only achieves the diffraction limit $<34\ \mu\text{m}$ due to telescope jitter (SOFIA Observer's Handbook, Figures 2–7). For consistency, the photometry were extracted uniformly using aperture photometry, in a $10''$ source annulus with a sky annulus $20''\text{--}40''$ away from the source. The F111 photometry is a weighted average of seven observations (one longer exposure was 69 s on source, and the other six short exposures were 13 s on source). The average flux density of the six short exposures is within 1% of the flux density measured from the long exposure. Images for each band are shown in Figure 1.

There also appears to be some structure in the background emission, making the extraction less certain. However, the source appears pointlike in all bands. The absolute flux uncertainty of the level 3 G063 and G227 grism products is 5%, and the G111 product is 10% (Herter et al. 2013). However, all of the photometric flux densities are within 3% of the level 3 spectroscopy. Thus we take the combined uncertainty of the absolute flux calibration between *Spitzer* and SOFIA (in quadrature) to be $\sim 4.2\%$. The background “structure” in some of the imaging appears significant at first glance; however, the *WISE* band 4 image (next section) shows no sign of structure. A more detailed extraction of F253 using a flat-field model (J. DeBuizer 2016, private communication) indicated that the background “structure” was due to flat-fielding.

2.4. *Photometry from Earlier Epochs*

In this work, we compare photometry close in time to the 2004 IRS observations with those close to the 2016 FORCAST observations (American Association of Variable Star Observers/AAVSO *BVR* from 2016, and FORCAST imaging from this work). Additional photometry is presented, including *UBVR* photometry previously presented in Green et al. (2006) (from 2004), 2MASS *JHK* (from 1998; Skrutskie et al. 2006), *WISE* (all four bands; 2010; Cutri et al. 2013), and *Spitzer-IRAC* bands 2 and 3 (2006). The *WISE* W2 band appears anomalously high (perhaps due to uncertainties related to emission from $\text{H}_2\text{ S}(9)$ at $4.6\ \mu\text{m}$). Due to these uncertainties, we find no convincing detection of background dust emission.

The *Spitzer-IRAC* photometry was observed but saturated beyond recovery in bands 1 and 4. Bands 2 ($4.5\ \mu\text{m}$) and 3 ($5.8\ \mu\text{m}$) were extracted as *Spitzer* Enhanced Imaging Products (SEIP⁷). We note that the FWHM is $1''.72$ ($4.5\ \mu\text{m}$) and $1''.88$ ($5.8\ \mu\text{m}$), as indicated by the IRAC Instrument Handbook

⁷ <http://irsa.ipac.caltech.edu/data/SPITZER/Enhanced/SEIP/overview.html>

Figure 1. The 5.6, 7.7, 11.1, 25.3, 31.5, and 34.8 μm broadband images of FU Ori from SOFIA-FORCAST, at arcmin scales to show the surrounding background, at the same absolute intensity scale. The apparent difference in centroid position is due to differences in the WCS pipeline processing.

(Table 2.1). The SEIP provides two extracted aperture sizes of diameters $3''.8$ and $5''.8$. The $5''.8$ diameter extraction window is a much better match with the IRS data, although in contrast to the similar-wavelength *WISE* band 2, the IRAC band 2 point is

low by 20% compared to models (possibly due to partial saturation). We note that if this difference between IRAC (2006) and *WISE* (2010) were due to intrinsic luminosity changes in the FU Ori disk, we would expect the *WISE* band to

Figure 2. The 0.3–38 μm SED of FU Ori, highlighting the decrease in luminosity between 2004 and 2016. The (2004) *Spitzer*-IRS spectroscopy (red line) and pre-2011 photometry (red squares; see the text for details) shortward of 8 μm are fit by a dereddened ($E(B - V) = 0.7$; Mathis 1990) 7200 K modified blackbody (Kenyon et al. 1988; Kenyon & Hartmann 1995; red dot-dashed line); the model is scaled assuming a distance of 450 pc to FU Ori. The (2016) SOFIA-FORCAST spectroscopy (black line) and photometry (black circles), as well as 2016 *BVR* AAVSO photometry, are fit by the same 7200 K modified blackbody curve but reduced in intensity by 12% (black dashed line).

be the lower of the two, but we find the reverse. For these reasons, we exclude both points from the analysis.

3. ANALYSIS

3.1. Continuum

First we consider changes in the continuum flux density. The (red) photometry presented in Figure 2 comes from various times between 1998 and 2010, but we used the AAVSO *UBVR* and decline rate in *B* and *V* ($0.015 \text{ mag yr}^{-1}$) measured for FU Ori in previous work (Kenyon et al. 2000; Green et al. 2006) to confirm that FU Ori has continued to decline in the *B* and *V* bands. Specifically, we averaged the *B* and *V* magnitudes measured in 2016 from the AAVSO, 11.178 ± 0.078 and 9.844 ± 0.156 , respectively. Photometry contemporaneous with the *Spitzer* spectroscopy (Green et al. 2006), taken in 2004 October, indicated *B* and *V* magnitudes of 11.01 ± 0.02 and 9.66 ± 0.02 , respectively. These data are shown in Table 2. Extrapolating from 2004 October to 2016 March indicates decline rates of ~ 0.015 and $0.016 \text{ mag yr}^{-1}$, respectively. This translates to a decrease of 14%–15% in the optical bands over the 12 years between the *Spitzer* and SOFIA observations.

Next we consider the mid-IR bands by comparing the common portions of the 5–28 μm data from *Spitzer* and SOFIA directly; the ratio is shown in Figure 3. We observe a $\sim 12\%$ decrease from 5 to 8 μm and a $\sim 7\%$ decrease from 10 to 28 μm ; these ratios are shown in Figure 3.

Finally we consider the entire 0.1–100 μm spectrum. We began by fitting the *Spitzer* 5–8 μm continuum, along with the *UBVR* and *WISE W1* band similar to Green et al. (2006), using the modified blackbody models from Kenyon & Hartmann (1995). We found that any of the 6590–7200 K models were equivalent fits to the observed 5–8 μm slope. There are larger differences between the models near 1 μm , but lacking 2016 data in the near-IR, the uncertainties are too great to distinguish between them. For the purpose of this comparison, we adopt the

7200 K model as in Green et al. (2006). We then fit the rebinned SOFIA continuum at the same anchor wavelengths and found that the same 7200 K model, but with 12% lower flux, matched the continuum to within uncertainties.

We found that the 6590–7200 K model slopes were not sufficiently distinguishable in either the optical/near-IR photometry or in the 5–8 μm spectroscopy. To get an idea of the uncertainties in the temperature, we used standard photometric colors from Table A.5 of Kenyon & Hartmann (1995). To illustrate this, Figure 4 shows three modified blackbody models reddened to an A_V of 1.8 (Mathis 1990), similar to FU Ori: 7200, 6890, and 7200 K but decreased by 12%; the difference between cooling from 7200 to 6890 K, rather than decreasing the accretion rate by 12%, is indistinguishable at wavelengths $> 1 \mu\text{m}$. Given that the *UBVR* photometry we include shows a decrease comparable to the IR, it is most plausible to explain the continuum flux change by depletion of the innermost region of the disk (or equivalently a decreased accretion rate); however, the mid-IR flux decrease can be explained by cooling (a decrease in the inner disk temperature). The cooling time in centrally irradiated T Tauri disks is very short, either a few years down to a few days depending upon the disk geometry (D’Alessio et al. 1999). Additionally, for typical FUor system parameters, Johnstone et al. (2013) found a cooling time in FUor disks on the order of a few weeks (albeit long compared with the dust-photon coupling timescale). Assuming the outer disk of FU Ori remains similar to a T Tauri disk, the cooling timescale is too rapid to explain the slow rate of decay.

Typically the draining of the gas within a disk would be noted in a decrease of the flux of the corotating molecular gas, as was seen in the postoutburst emission lines of EX Lup (Banzatti et al. 2015). Unlike T Tauri disks (Salyk et al. 2011), the CO lines are seen in absorption in FUors, even at 4.6 μm (Smith et al. 2009); they attribute the absorption to a

Table 2
UBVR 2004–2016 Photometry

JD	Calendar Date	Mag	δ Mag	Filter	Observer
2453278.46	2004 Sep 29	11.697	0.04	U	Ibrahimov
2453280.45	2004 Oct 1	11.706	0.04	U	Ibrahimov
2453281.472	2004 Oct 2	11.787	0.04	U	Ibrahimov
2453283.481	2004 Oct 4	11.788	0.04	U	Ibrahimov
2453284.483	2004 Oct 5	11.772	0.04	U	Ibrahimov
2453299.491	2004 Oct 20	11.809	0.04	U	Ibrahimov
2453278.46	2004 Sep 29	11.015	0.02	B	Ibrahimov
2453280.45	2004 Oct 1	10.971	0.02	B	Ibrahimov
2453281.472	2004 Oct 2	10.988	0.02	B	Ibrahimov
2453283.481	2004 Oct 4	11.031	0.02	B	Ibrahimov
2453284.483	2004 Oct 5	11.024	0.02	B	Ibrahimov
2453299.491	2004 Oct 20	11.022	0.02	B	Ibrahimov
2453309.476	2004 Oct 30	11.009	0.02	B	Ibrahimov
2453310.46	2004 Oct 31	10.999	0.02	B	Ibrahimov
2453311.499	2004 Nov 1	11.025	0.02	B	Ibrahimov
2457447.283	2016 Feb 28	11.122	0.087	B	DUBF
2457447.283	2016 Feb 28	11.2	0.08	B	DUBF
2457447.283	2016 Feb 28	11.095	0.086	B	DUBF
2457447.284	2016 Feb 28	11.134	0.081	B	DUBF
2457447.284	2016 Feb 28	11.078	0.094	B	DUBF
2457458.345	2016 Mar 10	11.191	0.067	B	DUBF
2457458.345	2016 Mar 10	11.209	0.066	B	DUBF
2457458.346	2016 Mar 10	11.143	0.069	B	DUBF
2457458.346	2016 Mar 10	11.148	0.067	B	DUBF
2457461.302	2016 Mar 13	11.187	0.05	B	DUBF
2457461.302	2016 Mar 13	11.272	0.045	B	DUBF
2457461.302	2016 Mar 13	11.186	0.051	B	DUBF
2457461.303	2016 Mar 13	11.162	0.053	B	DUBF
2457461.303	2016 Mar 13	11.179	0.048	B	DUBF
2457465.292	2016 Mar 17	11.249	0.114	B	DUBF
2457465.293	2016 Mar 17	11.188	0.119	B	DUBF
2457465.293	2016 Mar 17	11.173	0.119	B	DUBF
2457465.293	2016 Mar 17	11.32	0.101	B	DUBF
2457470.349	2016 Mar 22	11.092	0.069	B	DUBF
2457470.349	2016 Mar 22	11.28	0.056	B	DUBF
2457470.35	2016 Mar 22	11.148	0.063	B	DUBF
2457470.35	2016 Mar 22	11.169	0.063	B	DUBF
2457470.35	2016 Mar 22	11.35	0.054	B	DUBF
2457479.36	2016 Mar 31	11.177	0.072	B	DUBF
2457479.36	2016 Mar 31	11.308	0.074	B	DUBF
2457479.36	2016 Mar 31	11.24	0.069	B	DUBF
2457479.36	2016 Mar 31	11.197	0.07	B	DUBF
2457479.361	2016 Mar 31	11.026	0.085	B	DUBF
2457480.321	2016 Apr 01	11.171	0.078	B	DUBF
2457480.321	2016 Apr 01	11.005	0.096	B	DUBF
2457480.321	2016 Apr 01	11.206	0.077	B	DUBF
2457480.321	2016 Apr 01	11.092	0.087	B	DUBF
2457480.322	2016 Apr 01	11.191	0.074	B	DUBF
2453278.46	2004 Sep 29	8.485	0.015	R	Ibrahimov
2453280.45	2004 Oct 1	8.47	0.015	R	Ibrahimov
2453281.472	2004 Oct 2	8.466	0.015	R	Ibrahimov
2453283.481	2004 Oct 4	8.488	0.015	R	Ibrahimov
2453284.483	2004 Oct 5	8.504	0.015	R	Ibrahimov
2453299.491	2004 Oct 20	8.493	0.015	R	Ibrahimov
2453309.476	2004 Oct 30	8.483	0.015	R	Ibrahimov
2453310.46	2004 Oct 31	8.486	0.015	R	Ibrahimov
2453311.499	2004 Nov 1	8.477	0.015	R	Ibrahimov
2457448.274	2016 Feb 29	8.447	0.012	R	DUBF
2457448.274	2016 Feb 29	8.434	0.012	R	DUBF
2457448.274	2016 Feb 29	8.429	0.012	R	DUBF
2457448.274	2016 Feb 29	8.451	0.012	R	DUBF
2457448.274	2016 Feb 29	8.428	0.012	R	DUBF
2457451.324	2016 Mar 03	8.445	0.038	R	DUBF
2457451.324	2016 Mar 03	8.413	0.038	R	DUBF

Table 2
(Continued)

JD	Calendar Date	Mag	δ Mag	Filter	Observer
2457451.325	2016 Mar 03	8.469	0.029	R	DUBF
2457451.325	2016 Mar 03	8.498	0.023	R	DUBF
2457451.325	2016 Mar 03	8.473	0.024	R	DUBF
2457458.346	2016 Mar 10	8.542	0.012	R	DUBF
2457458.346	2016 Mar 10	8.546	0.012	R	DUBF
2457458.347	2016 Mar 10	8.558	0.012	R	DUBF
2457458.347	2016 Mar 10	8.548	0.012	R	DUBF
2457458.347	2016 Mar 10	8.557	0.012	R	DUBF
2457461.303	2016 Mar 13	8.525	0.01	R	DUBF
2457461.303	2016 Mar 13	8.54	0.01	R	DUBF
2457461.304	2016 Mar 13	8.516	0.011	R	DUBF
2457461.304	2016 Mar 13	8.543	0.01	R	DUBF
2457461.304	2016 Mar 13	8.53	0.01	R	DUBF
2457464.315	2016 Mar 16	8.588	0.017	R	DUBF
2457464.315	2016 Mar 16	8.601	0.018	R	DUBF
2457464.316	2016 Mar 16	8.557	0.018	R	DUBF
2457464.316	2016 Mar 16	8.587	0.017	R	DUBF
2457464.316	2016 Mar 16	8.593	0.017	R	DUBF
2457465.294	2016 Mar 17	8.528	0.015	R	DUBF
2457465.294	2016 Mar 17	8.579	0.016	R	DUBF
2457465.294	2016 Mar 17	8.599	0.016	R	DUBF
2457465.294	2016 Mar 17	8.578	0.015	R	DUBF
2457465.294	2016 Mar 17	8.563	0.015	R	DUBF
2457470.35	2016 Mar 22	8.554	0.012	R	DUBF
2457470.351	2016 Mar 22	8.534	0.013	R	DUBF
2457470.351	2016 Mar 22	8.543	0.012	R	DUBF
2457470.351	2016 Mar 22	8.556	0.013	R	DUBF
2457470.351	2016 Mar 22	8.543	0.012	R	DUBF
2457479.361	2016 Mar 31	8.514	0.011	R	DUBF
2457479.361	2016 Mar 31	8.501	0.01	R	DUBF
2457479.361	2016 Mar 31	8.512	0.011	R	DUBF
2457479.362	2016 Mar 31	8.477	0.011	R	DUBF
2457479.362	2016 Mar 31	8.46	0.011	R	DUBF
2457480.322	2016 Apr 01	8.487	0.01	R	DUBF
2457480.322	2016 Apr 01	8.502	0.009	R	DUBF
2457480.322	2016 Apr 01	8.493	0.009	R	DUBF
2457480.322	2016 Apr 01	8.505	0.01	R	DUBF
2457480.322	2016 Apr 01	8.483	0.01	R	DUBF
2457484.309	2016 Apr 05	8.448	0.012	R	DUBF
2457484.309	2016 Apr 05	8.456	0.011	R	DUBF
2457484.309	2016 Apr 05	8.43	0.012	R	DUBF
2457484.309	2016 Apr 05	8.482	0.012	R	DUBF
2457484.309	2016 Apr 05	8.466	0.011	R	DUBF
2453278.46	2004 Sep 29	9.654	0.02	V	Ibrahimov
2453280.45	2004 Oct 1	9.629	0.02	V	Ibrahimov
2453281.472	2004 Oct 2	9.639	0.02	V	Ibrahimov
2453283.481	2004 Oct 4	9.682	0.02	V	Ibrahimov
2453284.483	2004 Oct 5	9.677	0.02	V	Ibrahimov
2453299.491	2004 Oct 20	9.676	0.02	V	Ibrahimov
2453309.476	2004 Oct 30	9.659	0.02	V	Ibrahimov
2453310.46	2004 Oct 31	9.661	0.02	V	Ibrahimov
2453311.499	2004 Nov 1	9.652	0.02	V	Ibrahimov
2457447.282	2016 Feb 28	9.746	0.016	V	DUBF
2457447.282	2016 Feb 28	9.79	0.015	V	DUBF
2457447.283	2016 Feb 28	9.781	0.015	V	DUBF
2457447.283	2016 Feb 28	9.771	0.015	V	DUBF
2457447.283	2016 Feb 28	9.775	0.016	V	DUBF
2457448.272	2016 Feb 29	9.796	0.018	V	DUBF
2457448.273	2016 Feb 29	9.819	0.017	V	DUBF
2457448.273	2016 Feb 29	9.783	0.018	V	DUBF
2457448.273	2016 Feb 29	9.811	0.018	V	DUBF
2457448.273	2016 Feb 29	9.738	0.018	V	DUBF
2457451.322	2016 Mar 03	9.822	0.046	V	DUBF
2457451.322	2016 Mar 03	9.788	0.04	V	DUBF

Table 2
(Continued)

JD	Calendar Date	Mag	δ Mag	Filter	Observer
2457451.323	2016 Mar 03	9.803	0.035	V	DUBF
2457451.323	2016 Mar 03	9.855	0.031	V	DUBF
2457451.323	2016 Mar 03	9.838	0.03	V	DUBF
2457458.344	2016 Mar 10	9.856	0.019	V	DUBF
2457458.344	2016 Mar 10	9.838	0.02	V	DUBF
2457458.344	2016 Mar 10	9.867	0.018	V	DUBF
2457458.345	2016 Mar 10	9.889	0.018	V	DUBF
2457458.345	2016 Mar 10	9.846	0.019	V	DUBF
2457461.301	2016 Mar 13	9.88	0.014	V	DUBF
2457461.301	2016 Mar 13	9.887	0.014	V	DUBF
2457461.301	2016 Mar 13	9.89	0.013	V	DUBF
2457461.302	2016 Mar 13	9.888	0.014	V	DUBF
2457461.302	2016 Mar 13	9.903	0.014	V	DUBF
2457464.313	2016 Mar 16	9.899	0.032	V	DUBF
2457464.313	2016 Mar 16	9.912	0.031	V	DUBF
2457464.314	2016 Mar 16	9.875	0.031	V	DUBF
2457464.314	2016 Mar 16	9.913	0.031	V	DUBF
2457464.314	2016 Mar 16	9.944	0.03	V	DUBF
2457465.291	2016 Mar 17	9.902	0.03	V	DUBF
2457465.291	2016 Mar 17	9.926	0.027	V	DUBF
2457465.291	2016 Mar 17	9.926	0.027	V	DUBF
2457465.292	2016 Mar 17	9.976	0.026	V	DUBF
2457465.292	2016 Mar 17	9.917	0.028	V	DUBF
2457470.348	2016 Mar 22	9.912	0.018	V	DUBF
2457470.348	2016 Mar 22	9.889	0.018	V	DUBF
2457470.348	2016 Mar 22	9.861	0.018	V	DUBF
2457470.348	2016 Mar 22	9.878	0.018	V	DUBF
2457470.349	2016 Mar 22	9.905	0.017	V	DUBF
2457479.358	2016 Mar 31	9.806	0.028	V	DUBF
2457479.358	2016 Mar 31	9.858	0.026	V	DUBF
2457479.359	2016 Mar 31	9.778	0.029	V	DUBF
2457479.359	2016 Mar 31	9.802	0.03	V	DUBF
2457479.359	2016 Mar 31	9.825	0.027	V	DUBF
2457480.32	2016 Apr 01	9.796	0.024	V	DUBF
2457480.32	2016 Apr 01	9.832	0.022	V	DUBF
2457480.32	2016 Apr 01	9.794	0.024	V	DUBF
2457480.32	2016 Apr 01	9.822	0.024	V	DUBF
2457480.32	2016 Apr 01	9.795	0.025	V	DUBF
2457484.307	2016 Apr 05	9.855	0.038	V	DUBF
2457484.307	2016 Apr 05	9.74	0.042	V	DUBF
2457484.307	2016 Apr 05	9.843	0.036	V	DUBF
2457484.307	2016 Apr 05	9.761	0.037	V	DUBF
2457484.307	2016 Apr 05	9.807	0.036	V	DUBF

Note. *UBVR* photometry from 2004 to 2016 used in this work.

surrounding envelope of $0.1 M_{\odot}$, which FU Ori does not possess (e.g., Green et al. 2013).

3.2. Dust Features

The $10 \mu\text{m}$ silicate emission feature in the *Spitzer*-IRS spectrum showed no evidence of crystalline grains but some evidence for grain growth during the disk lifetime (Quanz et al. 2007). In order to compare the SOFIA feature, we used a spline fit to anchor the spectrum at $8 \mu\text{m}$, between the SiO and water vapor features, at $13 \mu\text{m}$ (the long-wavelength edge of the G111 spectrum), and beyond $20 \mu\text{m}$ (centered within the G253 spectrum).

Two different spline fits to both the *Spitzer* and SOFIA spectra are shown in Figure 5 and described below. We first tried independent spline fitting on each spectrum, using 5.6,

7.5 , 13.6 , 14.9 , 19.3 , 21.3 , and $24.4 \mu\text{m}$ anchor points for the *Spitzer* data; and 6.1 , 7.4 , 13.5 , 19.3 , 21.3 , and $24.4 \mu\text{m}$ for the SOFIA data. We identified the anchor continuum points separately, and unlike the IRS spectrum, the FORCAST spectrum does not have continuous wavelength coverage from 5 to $38 \mu\text{m}$, nor over the 9 – $13 \mu\text{m}$ complete silicate complex. Thus the fits are slightly different, and the residuals appear to have shifted (Figure 5, top panel). Specifically, the $10 \mu\text{m}$ silicate feature *appears* broader (to longer wavelengths) and taller. To check whether this difference was due to the differences in spline fitting between the two spectra, we performed a second spline fitting and subtraction using exactly the same anchor points for both the SOFIA-FORCAST and the *Spitzer*-IRS spectrum. Because we were limited to wavelength points common to both spectra, we used only six anchor points rather than the original seven in the IRS spectrum. The results are shown in the bottom of Figure 5. The features are virtually unchanged given the S/N of the SOFIA spectrum; we see no evidence of further dust processing or grain growth in the last 12 years, and we attribute any apparent differences in the feature to incomplete spectral coverage in the FORCAST bands.

Unfortunately, we lack sufficient spectral coverage to compare the $20 \mu\text{m}$ silicate features, but we find them consistent at wavelengths greater than $20 \mu\text{m}$, and we attribute any differences at shorter wavelengths to edge effects in the G253 spectrum.

3.3. Gaseous Absorption Features: H_2O

The blended rovibrational absorption lines of H_2O , originally noted in Green et al. (2006), are still present in the SOFIA-FORCAST spectrum (Figure 6). The short-wavelength shoulder marks the boundary between the blended R-branch lines ($\lambda < 6.3 \mu\text{m}$) and the blended P-branch lines ($\lambda > 6.3 \mu\text{m}$). The blended P-branch absorption lines appear unchanged over the last 12 years. However, the relatively shallower blended R-branch absorption detected in the *Spitzer*-IRS spectrum is not clearly detected. In order to compare the features directly, we apply a -0.5 Jy offset to the IRS spectrum, shown in Figure 7. Differences between the R- and P-branch lines were noted, for example, in Orion-BN/KL (González-Alfonso et al. 1998), who found the R branch in absorption and the P branch in emission. However, further observations with higher S/N are needed to confirm any changes.

Additional absorption attributed to SiO ($8 \mu\text{m}$) by Green et al. (2006) may be present in the FORCAST spectrum, but without full spectral coverage or improved sensitivity, we cannot determine whether this feature has changed in detail.

4. DISCUSSION

FUors are generally understood as large accretion bursts, although models differ on where the infalling material begins its descent. Binarity and outburst behavior has been examined in several studies (e.g., Bonnell & Bastien 1992; Reipurth et al. 2002; Millan-Gabet et al. 2006; Green et al. 2016). As noted earlier, FU Ori is itself a $0''.5$ binary, and disk–binary interactions could promote an episodic or burst mode of accretion (Reipurth & Aspin 2004). Assuming the central star mass of FU Ori is $M \sim 0.3 M_{\odot}$ (Zhu et al. 2009a), the integrated accretion luminosity (using model parameters from Green et al. 2006) corresponds to a total accreted mass of

Figure 3. Ratio of the flux density of the SOFIA-FORCAST spectrum of FU Ori (2016) divided by the *Spitzer*-IRS spectrum (2004). For comparison, the blue line indicates a decrease of 12%, and the red line indicates a decrease of 7%.

Figure 4. Three models using standard photometric colors of young stars (reddened to an A_V of 1.8) from Kenyon & Hartmann (1995): 7200 K (black solid line), 6890 K (red dot-dashed line), and 7200 K decreased by 12% (0.12 mag) (blue dashed line). The magnitude scale has an arbitrary offset. Cooling from 7200 to 6890 K is virtually indistinguishable from a 12% decrease in flux at wavelengths $>1 \mu\text{m}$, but cooling would not result in a significant decrease in $UBVR$ flux density.

$M \sim 0.018 M_{\odot}$. Furthermore, FU Ori was the target of a recent ALMA study that separated the two components (Hales et al. 2015) and demonstrated that the remnant envelope is tenuous at best, indicating that there is no external reservoir for resupplying the disk material in repeated bursts. They hypothesized that the more extinguished southern component could act as the mass reservoir for resupply. Flattened envelope models can explain the long-wavelength SED and provide long-term resupply potential, although whether sufficient mass exists for resupply is unclear, judging by the far-IR excess

reported by Green et al. (2013), who estimated a combined disk/envelope mass of $0.05 M_{\odot}$ from the *Herschel*-SPIRE $>350 \mu\text{m}$ excess. Higher-resolution observations from the Submillimeter Array further contained the size and mass of such an envelope plus disk to $0.05 M_{\odot}$ within 450 au or less (M. Dunham 2016, private communication, Green et al. 2016).

Regardless of the origin of the infalling material, our study suggests that at the mature stages of outburst, the infalling material accretes from the inside out. The decrease in only the short-wavelength continuum, without change to spectral shape,

Figure 5. Top left: independent spline fits to the SOFIA and *Spitzer* spectra of FU Ori. The IRS (red) spectrum is fit by the green solid spline, and the FORCAST (black) spectrum is fit by the blue spline, including an interpolation between 14 and 18 μm (blue dashed line). The spline anchor points are marked with blue and green diamonds. The spline anchor points were determined independently for the two data sets (see text). Top right: the resulting spline-fit subtracted IRS (red) and FORCAST (black) spectra. Bottom: spline-fit subtracted IRS and FORCAST spectra with five *identical* spline anchor points, highlighting the consistency of the silicate feature (see text).

Figure 6. Comparison of *Spitzer*-SL (red) and SOFIA G063 and G111 (black) spectra covering 5–14 μm , along with 7200 K modified blackbody models as in Figure 2. The SOFIA spectrum has been smoothed to low resolution for easier comparison with the SL data. The water absorption features can be clearly seen in both spectra, marked by the blue shaded regions (González-Alfonso et al. 1998).

indicates a decrease in density of the hot material, rather than a decrease in temperature. Combined with the lack of change in the silicate dust features after removing continuum effects, we suggest that the luminosity decrease in FU Ori is due primarily to this draining effect, leading to a decrease in the accretion rate.

Because the peak temperature in the disk still appears to be about 7200 K, the infall radius (centrifugal radius) is quite close to the central star. Zhu et al. (2007) modeled FU Ori as a superheated disk with a maximum effective temperature of 6420 K at $r_{\text{in}} = 5 R_{\odot}$, decreasing to 4660 K at $3 r_{\text{in}}$ and eventually to 977 K at $30 r_{\text{in}}$ (0.7 au). They added a

Figure 7. Zoom-in of *Spitzer*-SL (red) and SOFIA G063 (black) spectra covering 5–8 μm . The blue shaded regions indicate the wavelengths of the R- and P-branch H_2O lines (González-Alfonso et al. 1998). In this figure, the *Spitzer* spectrum has been offset -0.5 Jy to align with the SOFIA data. The P-branch water absorption complex can be clearly seen in both spectra, but the R-branch lines appear weak in the SOFIA spectrum.

second component from a more typical accretion disk >1 au. Supposing over the past 12 years that FU Ori accreted only material at a temperature >977 K, the accreted material would all be within ~ 0.7 au, where the free-fall time is much shorter than 12 years. It may be refilled from surrounding regions, but the maximum refill rate is capped by the upper limit to the change in IR excess beyond $20 \mu\text{m}$, which arises from annuli beyond the boundaries of the superheated disk at ~ 1 au (Zhu et al. 2009a). It appears that the outer disk does not react as quickly to this decreased luminosity on a 12 year timescale; within the uncertainties in scatter, the continuum level $<20 \mu\text{m}$ has changed by $\sim 7\%$.

However, we note that we cannot rule out cooling in the inner disk from this change in the SED. If this is due to cooling of the inner disk rather than a decrease in the accretion rate, and the $5\text{--}8 \mu\text{m}$ is dominated by the hot disk as models suggest, then the temperature decrease from a 12% decrease in luminosity would be only 4% (i.e., from 7200 to 6900 K). This is a small effect on the slope and is consistent with our data in the mid-IR (Figure 4). Potentially, a cooling pattern and decreased central accretion could be distinguished with high-fidelity IR continuum measurements, increased coverage in the optical to near-IR bands, or by changes in the width or shape of CO lines from differing disk annuli. Bell et al. (1995) noted that the decrease in the SED of FUor V1057 Cyg was least significant between 1 and $10 \mu\text{m}$; between 2004 and 2016, at least, FU Ori does not seem to follow this pattern, maintaining a decay rate similar to the optical bands out to at least $8 \mu\text{m}$. This difference may reflect the relatively small amount of material surrounding FU Ori or the overall slower decay rate compared with that of V1057 Cyg. An alternate explanation is that the FU Ori disk is geometrically thicker due to the extreme nature and duration of the accretion event (e.g., Hartmann et al. 2011 and references therein) and is beginning to settle as the midplane begins to cool.

5. CONCLUSIONS

We have performed the first comparison of multiepoch mid-IR spectra of an FUor to examine the changes occurring in the system during an outburst. We compared the 2004 SED and IR spectrum of FU Orionis with newly acquired (2016) SOFIA-FORCAST spectrum and photometry to identify changes over the intervening 12-year period. After establishing cross-calibration between the various measurements, we find a 12% ($\sim 3\sigma$) decrease in the $5\text{--}14 \mu\text{m}$ continuum without a significant change in shape. We find lesser changes ($\lesssim 7\%$) to the SED beyond $10 \mu\text{m}$ and no significant change to the silicate dust or the rovibrational water absorption from the disk photosphere. From this, we posit a decrease in gas at the highest temperatures and therefore the innermost regions of the disk, either due to cooling or depletion via accretion processes.

We thank L. Andrew Helton, Sachin Shenoy, James DeBuizer, William Vacca, Joseph Adams, Paule Sonnen-trucker, Andrea Banzatti, Miguel Requena-Torres, Klaus Pontoppidan, and Alexander Hubbard for helpful discussions of our results. We thank the SOFIA Science Center (USRA) for data reduction and analysis support. This work is based in part on observations made with the NASA/DLR Stratospheric Observatory for Infrared Astronomy (SOFIA). SOFIA is jointly operated by the Universities Space Research Association, Inc. (USRA), under NASA contract NAS2-97001, and the Deutsches SOFIA Institut (DSI) under DLR contract 50 OK 0901 to the University of Stuttgart. Partial support for this work was provided by NASA through award #NAS2-97001, SOF 04-0146 issued by USRA. We acknowledge the variable star observations from the AAVSO International Database contributed by observers worldwide and used in this work, and in particular observer “DUBF” Franky Dubois. The work of WJF was supported by an appointment to the NASA Postdoctoral Program at Goddard Space Flight Center, administered by the Universities Space Research Association under contract with

NASA. OCJ acknowledges NASA grant NNX14AN06G. This research has made use of NASA's Astrophysics Data System Bibliographic Services. This research has made use of the VizieR catalog access tool, CDS, Strasbourg, France.

REFERENCES

- Ábrahám, P., Juhász, A., Dullemond, C. P., et al. 2009, *Natur*, **459**, 224
- Andrews, S. M., & Williams, J. P. 2005, *ApJ*, **631**, 1134
- Banzatti, A., Pontoppidan, K. M., Bruderer, S., Muzerolle, J., & Meyer, M. R. 2015, *ApJL*, **798**, L16
- Bell, K. R., & Lin, D. N. C. 1994, *ApJ*, **427**, 987
- Bell, K. R., Lin, D. N. C., Hartmann, L. W., & Kenyon, S. J. 1995, *ApJ*, **444**, 376
- Bonnell, I., & Bastien, P. 1992, *ApJL*, **401**, L31
- Carey, S., Ingalls, J., Hora, J., et al. 2012, *Proc. SPIE*, **8442**, 84421Z
- Cieza, L. A., Casassus, S., Tobin, J., et al. 2016, arXiv:1607.03757
- Cutri, R. M., et al. 2013, yCat, 2328
- D'Alessio, P., Calvet, N., Hartmann, L., Lizano, S., & Cantó, J. 1999, *ApJ*, **527**, 893
- González-Alfonso, E., Cernicharo, J., van Dishoeck, E. F., Wright, C. M., & Heras, A. 1998, *ApJL*, **502**, L169
- Green, J. D., Evans, N. J., II, Kóspál, Á., et al. 2013, *ApJ*, **772**, 117
- Green, J. D., Hartmann, L., Calvet, N., et al. 2006, *ApJ*, **648**, 1099
- Green, J. D., Kraus, A. L., Rizzuto, A. C., et al. 2016, arXiv:1605.03270
- Hales, A. S., Corder, S. A., Dent, W. R. D., et al. 2015, *ApJ*, **812**, 134
- Hartmann, L. (ed.) 1998, *Accretion Processes in Star Formation* (Cambridge: Cambridge Univ. Press), 32
- Hartmann, L., & Kenyon, S. J. 1985, *ApJ*, **299**, 462
- Hartmann, L., & Kenyon, S. J. 1996, *ARA&A*, **34**, 207
- Hartmann, L., Zhu, Z., & Calvet, N. 2011, arXiv:1106.3343
- Herbig, G. H. 1977, *ApJ*, **217**, 693
- Herter, T. L., Adams, J. D., De Buizer, J. M., et al. 2012, *ApJL*, **749**, L18
- Herter, T. L., Vacca, W. D., Adams, J. D., et al. 2013, *PASP*, **125**, 1393
- Higdon, S. J. U., Devost, D., Higdon, J. L., et al. 2004, *PASP*, **116**, 975
- Houck, J. R., Roellig, T. L., van Cleve, J., et al. 2004, *ApJS*, **154**, 18
- Hubbard, A., & Ebel, D. S. 2014, *Icar*, **237**, 84
- Johnstone, D., Hendricks, B., Herczeg, G. J., & Bruderer, S. 2013, *ApJ*, **765**, 133
- Kenyon, S. J., & Hartmann, L. 1995, *ApJS*, **101**, 117
- Kenyon, S. J., Hartmann, L., & Hewett, R. 1988, *ApJ*, **325**, 231
- Kenyon, S. J., Kolotilov, E. A., Ibragimov, M. A., & Mattei, J. A. 2000, *ApJ*, **531**, 1028
- Kim, H. J., Evans, N. J., II, Dunham, M. M., Lee, J.-E., & Pontoppidan, K. M. 2012, *ApJ*, **758**, 38
- Lebouteiller, V., Barry, D. J., Goes, C., et al. 2015, *ApJS*, **218**, 21
- Lebouteiller, V., Barry, D. J., Spoon, H. W. W., et al. 2011, *ApJS*, **196**, 8
- Lin, D. N. C., & Papaloizou, J. 1985, in *Protostars and Planets II*, ed. D. C. Black & M. S. Matthews
- Martin, R. G., Lubow, S. H., Livio, M., & Pringle, J. E. 2012, *MNRAS*, **423**, 2718
- Mathis, J. S. 1990, *ARA&A*, **28**, 37
- Millan-Gabet, R., Monnier, J. D., Akeson, R. L., et al. 2006, *ApJ*, **641**, 547
- Olofsson, J., Augereau, J.-C., van Dishoeck, E. F., et al. 2009, *A&A*, **507**, 327
- Paczynski, B. 1976, in *IAU Symp. 73, Structure and Evolution of Close Binary Systems*, ed. P. Eggleton, S. Mitton, & J. Whelan, 75
- Poteet, C. A., Pontoppidan, K. M., Megeath, S. T., et al. 2013, *ApJ*, **766**, 117
- Pueyo, L., Hillenbrand, L., Vasisht, G., et al. 2012, *ApJ*, **757**, 57
- Quanz, S. P., Henning, T., Bouwman, J., et al. 2007, *ApJ*, **668**, 359
- Reipurth, B., & Aspin, C. 2004, *ApJL*, **608**, L65
- Reipurth, B., Hartmann, L., Kenyon, S. J., Smette, A., & Bouchet, P. 2002, *AJ*, **124**, 2194
- Romanova, M. M., Ustyugova, G. V., Koldoba, A. V., & Lovelace, R. V. E. 2009, *ASSP*, **13**, 153
- Salyk, C., Blake, G. A., Boogert, A. C. A., & Brown, J. M. 2011, *ApJ*, **743**, 112
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, **131**, 1163
- Smith, R. L., Pontoppidan, K. M., Young, E. D., Morris, M. R., & van Dishoeck, E. F. 2009, *ApJ*, **701**, 163
- Vorobyov, E. I., & Basu, S. 2015, *ApJ*, **805**, 115
- Wang, H., Apai, D., Henning, T., & Pascucci, I. 2004, *ApJL*, **601**, L83
- Watson, D. M., Leisenring, J. M., Furlan, E., et al. 2009, *ApJS*, **180**, 84
- Werner, M. W., Roellig, T. L., Low, F. J., et al. 2004, *ApJS*, **154**, 1
- Young, E. T., Becklin, E. E., Marcum, P. M., et al. 2012, *ApJL*, **749**, L17
- Zhu, Z., Hartmann, L., Calvet, N., et al. 2007, *ApJ*, **669**, 483
- Zhu, Z., Hartmann, L., & Gammie, C. 2009a, *ApJ*, **694**, 1045
- Zhu, Z., Hartmann, L., Gammie, C., & McKinney, J. C. 2009b, *ApJ*, **701**, 620
- Zhu, Z., Hartmann, L., Gammie, C. F., et al. 2010, *ApJ*, **713**, 1134