

**DISCOVERY OF TETRAGONAL ALMANDINE, $(\text{Fe,Mg,Ca,Na})_3(\text{Al,Si,Mg})_2\text{Si}_3\text{O}_{12}$,
A NEW HIGH-PRESSURE MINERAL IN SHERGOTTY**

Chi Ma^{1,*}, Oliver Tschauner². ¹Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125, USA; ²Department of Geoscience and High Pressure Science and Engineering Center, University of Nevada, Las Vegas, NV 89154, USA; *Email: chi@gps.caltech.edu.

Introduction: During a nanomineralogy investigation of the Shergotty meteorite, we have identified a new shock-induced high-pressure silicate, majoritic almandine with a tetragonal $I4_1/a$ structure, in an impact melt pocket. Field-emission scanning electron microscope, electron back-scatter diffraction, electron microprobe and synchrotron diffraction were used to characterize its chemical composition, structure, and associated phases. Tetragonal majorite ($\text{Mg}_3(\text{SiMg})\text{Si}_3\text{O}_{12}$) with the $I4_1/a$ structure was synthesized [e.g., 1] and found later in the Tenham chondrite [2,3]. Tetragonal almandine-pyrope phase (TAPP) with the $I-42d$ structure was reported to occur as inclusions in lower-mantle diamonds [4]. We present here the first occurrence of tetragonal almandine with the $I4_1/a$ structure in a shocked meteorite from Mars as a new high-pressure phase.

Occurrence, Chemistry, and Crystallography: Tetragonal almandine in Shergotty occurs as aggregates of subhedral crystals, 0.8 – 2.5 μm in diameter, along with stishovite in the central region of a shock melt pocket (Fig. 1). The shock melt pocket is $\sim 450 \times 1000 \mu\text{m}^2$ in size, surrounded by pyroxene and maskelynite. The rock consists of mainly clinopyroxene and maskelynite, plus fayalite, silica phases, Ti-bearing magnetite, ilmenite, baddeleyite, chlorapatite, merrillite, and pyrrhotite.

The mean chemical composition of tetragonal almandine by electron microprobe analysis is (wt%) SiO_2 48.17, Al_2O_3 13.09, FeO 18.40, CaO 9.25, MgO 7.16, Na_2O 2.85, MnO 0.55, TiO_2 0.39, Cr_2O_3 0.08, K_2O 0.07, total 100.01, giving rise to an empirical formula of $(\text{Fe}_{1.16}\text{Ca}_{0.75}\text{Mg}_{0.61}\text{Na}_{0.42}\text{Mn}_{0.03}\text{K}_{0.01})(\text{Al}_{1.16}\text{Si}_{0.63}\text{Mg}_{0.19}\text{Ti}_{0.02})\text{Si}_3\text{O}_{12}$. The general formula is $(\text{Fe,Mg,Ca,Na})_3(\text{Al,Si,Mg})_2\text{Si}_3\text{O}_{12}$. The end-member formula is $\text{Fe}_3\text{Al}_2\text{Si}_3\text{O}_{12}$. Electron back-scatter diffraction indicated this phase has a garnet-related structure. Synchrotron X-ray diffraction revealed that this garnet has actually a tetragonal structure ($I4_1/a$) with unit cell dimensions: $a = 11.585(9) \text{ \AA}$, $c = 11.63(4) \text{ \AA}$, $V = 1561(7) \text{ \AA}^3$, and $Z = 8$.

Origin and Significance: Tetragonal almandine is the polymorph of cubic almandine, a new high-pressure garnet mineral, formed by shock metamorphism via the Shergotty impact event on Mars. It apparently crystallized from Fe-rich shock-induced melt under high-pressure.

References: [1] Angel R.J. et al. 1989. *American Mineralogist* 74:509–512. [2] Xie Z. & Sharp T.G. 2007. *Earth and Planetary Science Letters* 254:433–445. [3] Tomioka N. et al. 2016. *Science Advances* 2:e1501725. [4] Harris J. et al. 1997. *Nature* 387:486–488.




Fig. 1. Back-scatter electron image showing tetragonal almandine with stishovite in a Shergotty melt pocket.