

Catalytic enantioselective stereoablative reactions: an unexploited approach to enantioselective catalysis†

Justin T. Mohr, David C. Ebner and Brian M. Stoltz*

Received 20th July 2007, Accepted 30th August 2007

First published as an Advance Article on the web 1st October 2007

DOI: 10.1039/b711159m

Approaches to the preparation of enantioenriched materials *via* catalytic methods that *destroy* stereogenic elements of a molecule are discussed. Although these processes often decrease overall molecular complexity, there are several notable advantages including material recycling, enantiodivergence and convergence, and increased substrate scope. Examples are accompanied by discussion of the critical design elements required for the success of these methods.

Since the inception of enantioselective catalytic methodology, the prevailing strategic approach has relied on inducing chirality into a prochiral atom by the generation of new asymmetric centers or axes (Fig. 1a). While this tactic has proven extremely effective, the number of viable prochiral functional groups is relatively limited.

The Arnold and Mabel Beckman Laboratories of Chemical Synthesis, Division of Chemistry and Chemical Engineering, California Institute of Technology, 1200 E. California Boulevard, MC 164-30, Pasadena, CA, 91125, USA. E-mail: stoltz@caltech.edu; Fax: (+1) 626-564-9297

† The HTML version of this article has been enhanced with colour images.




Fig. 1 Strategies for enantioselective catalysis.

Justin T. Mohr received his A.B. degree in chemistry from Dartmouth College in Hanover, NH, USA in 2003. In the same year, he joined the labs of Professor Brian Stoltz at Caltech, where he has worked toward his Ph.D. as a Lilly Fellow. His research interests include new methods of C–C bond formation, enantioselective catalysis, and total synthesis of natural products.

David C. Ebner received his B.S. in chemistry and B.A. in mathematics from the University of St. Thomas in St. Paul, MN, USA in 2002. He is currently pursuing his Ph.D. at Caltech in the labs of Professor Brian Stoltz. He has been awarded the National Defense Science and Engineering Fellowship and National Science Foundation Graduate Fellowship. His research interests involve the development of catalytic asymmetric reactions and their application to the total synthesis of natural products.

Brian M. Stoltz obtained his B.S. in chemistry and B.A. in German from Indiana University of Pennsylvania in 1993. He then earned his Ph.D. in 1997 in the labs of Professor John L. Wood at Yale University. Following an NIH postdoctoral fellowship in the laboratories of Professor E. J. Corey at Harvard University, he joined the faculty at Caltech in 2000, where he is currently a Professor of Chemistry. His research focuses on the design and implementation of new strategies for the synthesis of complex molecules possessing important biological properties, in addition to the development of new synthetic methods including asymmetric catalysis and cascade processes.



Justin T. Mohr



David C. Ebner



Brian M. Stoltz


An alternative approach to the production of enantioenriched materials is to begin with a racemic mixture and subsequently eliminate the intrinsic stereochemistry from a portion or all of this mixture. The scope of this approach to enantioselective catalysis is as wide as the number of chiral molecules in existence. While inherently a complexity minimizing process, this approach has proven to be valuable in the synthesis of chiral building blocks and more complex synthetic targets.

In 2005, we defined the term “stereoablation” in the context of an enantioconvergent reaction.¹ Our initial definition was “the conversion of a chiral molecule to an achiral molecule,” based on the Oxford English Dictionary definition for ablation: “the action or process of carrying away or removing; removal.”² Upon further consideration of the importance of such methods in enantioselective chemical transformations, we have seen fit to expand the scope of this definition to include reactions where an existing stereocenter in a molecule is destroyed, but the intermediate molecule need not be wholly achiral.³ This revised definition thereby includes many other important advances. To date, few stereoablative strategies have been exploited for enantioselective catalysis, although notable exceptions include metal π -allyl alkylations⁴ and many dynamic kinetic resolutions.⁵ In this Emerging Area highlight, recent examples of novel approaches to asymmetric catalytic methods for stereoablation will be discussed. We hope to demonstrate that this is an important, though underutilized, method of asymmetric synthesis.⁶

When considering catalytic enantioselective stereoablative reactions, two possible regimes arise: one in which the stereoablative step is the enantioselective step (Fig. 1b), and one in which stereoablation precedes the enantioselective step (Fig. 1c). In the first case, a catalyst must selectively react with one enantiomer or enantiotopic group of the substrate to provide enantioenrichment. In the second case, a nonselective stereoablation is required before the enantioselective step.


Kinetic resolution is of the former type, selectively transforming one enantiomer of a racemic mixture to product. In addition to making enantiomer isolation a trivial process, stereoablative approaches often have the added capability of converting the achiral product back into a racemic starting material mixture by a relatively straightforward procedure. Recycling this material minimizes the waste common to many kinetic resolutions due to discard of the undesired enantiomer.

Recently, the Stoltz laboratory has reported an oxidative kinetic resolution (OKR) of secondary alcohols (Scheme 1).^{7–10} Utilizing molecular oxygen as the terminal oxidant, [Pd(nbd)Cl₂] (nbd = norbornadiene) and (–)-sparteine catalyze the oxidation of alcohol (+)-**1** to achiral ketone **2**, leaving unreacted alcohol (–)-**1** of high ee. Selective stereoablation by β -hydride elimination of a Pd-alkoxide to form product ketone has been shown to be enantiodetermining by extensive mechanistic^{7d,8} and computational^{7f} studies. To date, a wide variety of secondary alcohols have been successfully resolved with this catalyst system.




Scheme 1 Stoltz's oxidative kinetic resolution of secondary alcohols.

Additionally, ketone **4**, obtained in the resolution of alcohol (±)-**3**, can be recycled by reduction to racemic (±)-**3** in quantitative yield, allowing greater than 50% overall yield of the enantioenriched alcohol after a second resolution (Scheme 2).




Scheme 2 Stoltz's stereoablative kinetic resolution with recycle.

When other stereocenters are present in the alcohol, enantioenriched ketones can be obtained. In the Stoltz synthesis of (+)-amurensinine (**7**), racemic alcohol (±)-**5** was resolved successfully using the [(sparteine)PdCl₂] catalyst system (Scheme 3).¹¹ In addition to highly enantioenriched alcohol (–)-**5**, diketone (–)-**6a** was obtained in 79% ee. This diketone presumably arises from overoxidation of the initial ketone product ((+)-**6b**, R = H₂) in the presence of O₂. In fact, the monoketone (+)-**6b** could be isolated with 77% ee at shorter reaction times, albeit with lower ee of alcohol (–)-**5**. Importantly, the products (–)-**6a** and (+)-**6b** have the opposite configuration at C(5), potentially providing access to (–)-amurensinine. In general, OKR of alcohols with multiple stereocenters can provide enantioenriched product ketones as well as alcohols, opening the door to enantiodivergent synthetic strategies.



Scheme 3 OKR in the Stoltz synthesis of (+)-amurensinine.


Oxidative resolution of sulfoxides has also been demonstrated. Unlike alcohol oxidation, in which C–H bond *cleavage* is stereoablative, in sulfoxide oxidation S–O bond *formation* leads to stereoablation. Of particular note is an example by Jackson and coworkers. It was found that a racemic mixture of sulfoxides was effectively resolved with a vanadium catalyst and diiodide ligand (*R*)-**9** (Scheme 4).¹² The high selectivity in sulfoxide oxidative kinetic resolution led them to investigate a tandem enantioselective sulfide oxidation followed by sulfoxide resolution. Treatment of sulfide **11** with their oxidative conditions provided sulfoxide (*R*)-**8** in 70% yield and >99.5% ee, along with achiral sulfone **10**. The combined effect of the two processes allows the synthesis of highly enantioenriched sulfoxides with higher yields than a typical kinetic resolution. Additionally, coupling two enantioselective reactions has the potential to bring poorly selective methods into



Scheme 4 Jackson's oxidation of sulfoxides and sulfides.


the synthetically useful range because of the enhanced yield and product enantiopurity relative to the individual steps.

An unusual example of stereoablative kinetic resolution has been reported by Noyori *et al.*¹³ Hydrogenation of allylic alcohols with chiral catalyst $[(S)\text{-BINAP}]\text{Ru}(\text{OAc})_2$ results in kinetic resolution by symmetrizing one enantiomer of substrate (Scheme 5). This reductive kinetic resolution (RKR) is capable of resolving racemic alcohols such as $(\pm)\text{-12}$ with exceptionally high selectivity factors, providing achiral alcohol **13** as the byproduct. Although $(R)\text{-12}$ is obtained in high ee, there is currently no simple, direct method for recycling **13** back to $(\pm)\text{-12}$. Nonetheless, this RKR process provides a complementary method to the previously described OKR, using a reductive gas instead of an oxidative gas.




Scheme 5 Noyori's reductive kinetic resolution.

In addition to byproduct recycling, greater than 50% yield in a stereoablative process can be achieved by performing a desymmetrization. Such reactions utilize substrates that contain two enantiotopic functional groups, one of which selectively reacts with a chiral catalyst. Stoltz and Ferreira have reported a desymmetrization of meso diol **14** using their Pd-catalyzed oxidation conditions to obtain ketoalcohol $(+)\text{-15}$ in 72% yield with 95% ee (Scheme 6).^{7a}

Scheme 6 Stoltz's desymmetrization of meso diol **14**.

Catalytic enantioselective processes have also been employed in the desymmetrization of epoxides. Andersson and Södergren have reported the use of chiral diamine **17** in the rearrangement of epoxides to allylic alcohols.¹⁴ Treatment of cyclohexene oxide (**16**) with 5 mol% **17** in the presence of LDA as the stoichiometric base provided $(R)\text{-2-cyclohexenol}$ (**12**) in 96% ee (Scheme 7). Selective removal of one of the enantiotopic protons in the starting material accompanies destruction of one of the stereocenters of the epoxide in the elimination step. While there have been several other




Scheme 7 Andersson's epoxide desymmetrization.

examples of catalytic asymmetric epoxide desymmetrization, this system has the largest reported substrate scope, with five allylic alcohols formed with good to excellent ee.


A second type of stereoablative enantioselective catalysis consists of stereoablation followed by enantioselective bond formation. In these enantioconvergent processes, both enantiomers of a racemic mixture are converted to an achiral intermediate, which is converted subsequently to an enantioenriched product in a separate process (Fig. 1c). It is critical to avoid kinetic resolution in the stereoablative step in order to ensure good yield in a reasonable time.

A prominent type of enantioconvergent catalysis is dynamic kinetic resolution (DKR) of racemic alcohols. A particularly elegant system was developed by Bäckvall *et al.*, wherein an achiral metal catalyst (**18**) capable of rapid stereomutation *via* the corresponding ketone was coupled with selective acylating enzyme CALB (Scheme 8).¹⁵ The rates of these two simultaneous reactions are critical to the success of the process. The rate of stereomutation must be considerably greater than the rate of acylation in order to maintain an optimal 1 : 1 mixture of alcohol enantiomers for the enzymatic resolution. While kinetic resolution by acylation is a common approach to obtaining enantioenriched alcohols, the pairing of the stereoablative Ru catalyst and the acylation enzyme increases the overall efficiency of the reaction, as it allows yields greater than 50%. However, systems such as this are rare because the two concurrent catalytic reactions must tolerate one another.



Scheme 8 Bäckvall's dynamic kinetic resolution of alcohols.


To avoid catalyst incompatibility, it is desirable to identify a single catalyst system capable of both the stereoablative step and enantioselective bond-forming step. In the realm of alcohol oxidation, Williams and Adair recently reported a deracemization of secondary alcohols utilizing a bifunctional Ru catalyst (Scheme 9).¹⁶ This system uses a single catalyst to perform a nonselective stereoablative oxidation followed by an enantioselective reduction. Exposure of racemic alcohol mixture $(\pm)\text{-20}$ to a catalyst formed *in situ* from $[\text{RuCl}_2(\text{benzene})_2]$, phosphine **21**, and $(R,R)\text{-DPEN}$ (**22**) with cyclohexanone as a hydrogen acceptor



Scheme 9 Williams' deracemization of benzylic alcohols.


produces achiral ketone **23**. Pressurization of the reaction with H_2 promotes enantioselective hydrogenation to the enantioenriched alcohol (*S*)-**20**. While the demonstrated substrate scope of this reaction is still limited, the system overcomes the shortcoming of low yields of kinetic resolution processes, providing benzylic alcohols in 82–97% yield. The unique ability of the Ru catalyst to operate *via* two distinct mechanisms is critical to the success of this method. According to the principle of microscopic reversibility, the nonselective transfer dehydrogenation must also be nonselective in the reverse reaction, and therefore cannot complete the deracemization. However, introduction of an atmosphere of H_2 opens a different, highly selective mechanistic pathway leading to alcohols of high ee.

Recently, Stoltz *et al.* established that racemic mixtures of allyl β -ketoesters are efficiently converted to enantioenriched α -quaternary cycloalkanones in an enantioconvergent process mediated by Pd and phosphinooxazoline (PHOX) ligands (Scheme 10).¹ The mechanism is presumed to proceed through a Pd-enolate (**26**) formed by deallylation and stereoablative loss of CO_2 from (\pm)-**24**. No significant kinetic resolution of the racemic starting materials was observed, and, coupled with the high chemical yield and enantioselectivity, these reactions represent an efficient method for the generation of enantioenriched building blocks for synthesis.¹⁷



Scheme 10 Stoltz's stereoablative enantioconvergent allylation.


An interesting extension of this enantioconvergent method is the combination of a reactive allyl enol carbonate moiety with a latent allyl β -ketoester (**28**, Scheme 11). In the course of this reaction, a new stereocenter is first generated *via* decomposition of the allyl enol carbonate to reveal a Pd-enolate which undergoes enantioselective allylation. It is important that the catalyst be able to effectively overcome the inherent stereochemical preference of the substrate since the starting material is a racemic mixture. If the catalyst is unable to overcome the substrate preference, then a poor product d.r. will result. Notably, in this reaction, a 7 : 3 d.r. was obtained with Ph_3P as ligand, while an enhanced d.r. of 4 : 1 was observed with (*S*)-*t*-BuPHOX (**25**) as ligand. In the second step of this double-allylation reaction, the newly revealed ketone in **29** activates the allyl ester toward decarboxylation and formation



Scheme 11 Stoltz's cascade asymmetric allylation generating two quaternary stereocenters.

of Pd-enolate **30**. Catalyst control over the configuration of the second stereocenter leads to a Horeau type enhancement¹⁸ of the overall ee of the product. In this case, product (*-*)-**31** forms in 92% ee.

A second stereoablative reaction has been reported with the Pd-PHOX catalyst system. In this case, the putative Pd-enolate (**26**) is trapped with an alternate electrophile: a proton (Scheme 12).¹⁹ Again, the enantiopure catalyst is involved in both the bond-breaking and bond-forming steps, although the exact mechanistic course of the reaction remains unclear.²⁰ The divergent reactivity of the enolate intermediate toward different electrophiles highlights the effectiveness and convenience of these stereoablative reactions. While the stereoablative step in both reactions is likely identical, two different structural motifs (α -quaternary and α -tertiary ketones) are both available from a common starting material.²¹




Scheme 12 Stoltz's stereoablative enantioconvergent protonation.


Catalyst design in catalytic enantioconvergent processes is especially important in cases such as the enantioselective decarboxylative allylation and protonation reactions described above. Since the catalyst is intimately involved in both the stereoablative (C–C bond-breaking) and enantioselective (C–C or C–H bond-forming) steps, it is critical that the first step be insensitive to substrate stereochemistry.

Analogous enolate methods are known in which stoichiometric reagents are used in the stereoablative step.²² Importantly, kinetic resolution of the starting material is avoided by employing an achiral reagent (*e.g.*, *sec*-BuLi) for this process. Among these is the asymmetric Li-enolate protonation method of Vedejs and Kruger, wherein a catalytic amount of a chiral amine (**34**) coupled with slow addition of stoichiometric phenylacetic acid derivative **35** leads to amide (*R*)-**33** in high ee (Scheme 13).²³

A unique, metal-free approach to stereoablation was developed by Hénin and Muzart *et al.*²⁴ In this work, a light initiated Norrish Type II fragmentation is employed to eliminate the stereocenter present in tetralone **36** and access intermediate enol **37** (Scheme 14). Subsequently, amino alcohol **38** mediates




Scheme 13 Vedejs' enantioselective enolate protonation.



Scheme 14 Muzart's photolytic stereoablative process.


tautomerization to the enantioenriched product (*R*)-**32**. Other amino alcohols provide higher levels of conversion and yield at the cost of enantioselectivity.

A recent report of a stereoablative enantioconvergent process for cross-coupling was detailed by Fu *et al.* in 2005. In the reaction, a racemic α -bromo amide or benzylic bromide is treated with catalytic Ni, enantiopure (*i*-Pr)-Pybox ligand, and an alkylzinc reagent to create an enantioenriched tertiary stereocenter (Scheme 15).²⁵ Although the mechanistic details have not been fully elucidated, it has been hypothesized that the racemic bromide (**39**) initially decomposes to a radical intermediate (**40**), negating the stereochemistry of the starting material. Subsequent combination of the carbon-centered radical with the Ni catalyst and Negishi-type coupling provides (*-*)-**42** and completes the catalytic cycle.



Scheme 15 Fu's enantioconvergent Negishi coupling.

As a final example, Trost and Ariza have reported an intermolecular system where both the electrophilic and nucleophilic partners are racemic (**43** and **44**, Scheme 16).²⁶ It is proposed that (\pm)-**43** is converted to an achiral η^3 -allyl ligand bound to Pd (**45**), which is subsequently attacked by deprotonated azlactone **46**, forming product **47** with excellent enantio- and diastereocontrol. The remarkable stereochemical control in this work is made possible by two separate stereoablative steps.



Scheme 16 Trost's doubly-stereoconvergent allylic alkylation.

Conclusions

Although to date the primary focus during the development of enantioselective catalysis has been the creation of new stereocenters on prochiral substrates, asymmetric catalysis is not limited to the selective construction of new stereocenters. The selective destruction of stereogenic elements is also a viable, and increasingly important, technique that is beginning to show its utility in synthetic applications. This approach has several advantages including easily recycled byproducts, easily accessible racemic or meso starting materials, entries into enantioconvergent catalytic processes, and opportunities for enantiodivergent synthesis. As these new methods become more prominent and are further developed by the synthetic community they will surely play a pivotal role in the construction of enantiopure materials.

Acknowledgements

We thank NIH-NIGMS (R01 GM 080269-01), Eli Lilly (pre-doctoral fellowship to JTM), NDSEG (pre-doctoral fellowship to DCE), and NSF (pre-doctoral fellowship to DCE) for funding.

References

- J. T. Mohr, D. C. Behenna, A. M. Harned and B. M. Stoltz, *Angew. Chem., Int. Ed.*, 2005, **44**, 6924–6927.
- The Oxford English Dictionary*, ed. J. Simpson and E. Weiner, Oxford University Press, Oxford, 2nd edn, 1989.
- We suggest the following as a concise, inclusive definition for stereoablation: a chemical process where an existing stereogenic element in a molecule is eliminated.
- For a recent review, see: B. M. Trost, *J. Org. Chem.*, 2004, **69**, 5813–5837.
- For a recent review, see: H. Pellissier, *Tetrahedron*, 2003, **59**, 8291–8327.
- Recently, the related topic of racemization in asymmetric synthesis has been reviewed, see: F. F. Huerta, A. B. E. Minidis and J.-E. Bäckvall, *Chem. Soc. Rev.*, 2001, **30**, 321–331.
- (a) E. M. Ferreira and B. M. Stoltz, *J. Am. Chem. Soc.*, 2001, **123**, 7725–7726; (b) J. T. Bagdanoff, E. M. Ferreira and B. M. Stoltz, *Org. Lett.*, 2003, **5**, 835–837; (c) J. T. Bagdanoff and B. M. Stoltz, *Angew. Chem., Int. Ed.*, 2004, **43**, 353–357; (d) R. M. Trend and B. M. Stoltz, *J. Am. Chem. Soc.*, 2004, **126**, 4482–4483; (e) D. D. Caspi, D. C. Ebner, J. T. Bagdanoff and B. M. Stoltz, *Adv. Synth. Catal.*, 2004, **346**, 185–189; (f) R. J. Nielson, J. M. Keith, B. M. Stoltz and W. A. Goddard, III, *J. Am. Chem. Soc.*, 2004, **126**, 7967–7974; (g) D. C. Ebner, Z. Novák and B. M. Stoltz, *Synlett*, 2006, 3533–3539.
- A similar system was simultaneously reported, see: M. S. Sigman and D. R. Jensen, *Acc. Chem. Res.*, 2006, **39**, 221–229, and references cited therein.
- The selectivity factor s was determined using the equation: $s = k_{\text{fast}}/k_{\text{slow}} = \ln[(1 - C)(1 - ee)]/\ln[(1 - C)(1 + ee)]$, where C = conversion, see: H. B. Kagan, J. C. Fiaud, in *Topics in Stereochemistry*, ed. E. L. Eliel, Wiley & Sons, New York, 1988, vol. 18, pp. 249–330.
- For a recent review of alcohol oxidation, including enantioselective examples, see: J. B. Arterburn, *Tetrahedron*, 2001, **57**, 9765–9788.
- U. K. Tambar, D. C. Ebner and B. M. Stoltz, *J. Am. Chem. Soc.*, 2006, **128**, 11752–11753.
- C. Drago, L. Caggiano and R. F. W. Jackson, *Angew. Chem., Int. Ed.*, 2005, **44**, 7221–7223.
- M. Kitamura, I. Kasahara, K. Manabe, R. Noyori and H. Takaya, *J. Org. Chem.*, 1988, **53**, 708–710.
- M. J. Södergren and P. G. Andersson, *J. Am. Chem. Soc.*, 1998, **120**, 10760–10761.
- (a) A. L. E. Larsson, B. A. Persson and J.-E. Bäckvall, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1211–1212; (b) B. A. Persson, A. L. E. Larsson, M. L. Ray and J.-E. Bäckvall, *J. Am. Chem. Soc.*, 1999, **121**, 1645–1650; (c) B. Martín-Matute, M. Edin, K. Bogár and J.-E. Bäckvall, *Angew. Chem., Int. Ed.*, 2004, **43**, 6535–6539.
- G. R. A. Adair and J. M. J. Williams, *Chem. Commun.*, 2007, 2608–2609.

- 17 For related stereoablative, enantioconvergent allylation reactions, see: (a) M. Nakamura, A. Hajra, K. Endo and E. Nakamura, *Angew. Chem., Int. Ed.*, 2005, **44**, 7248–7251; (b) B. M. Trost, R. N. Bream and J. Xu, *Angew. Chem., Int. Ed.*, 2006, **45**, 3109–3112; (c) E. C. Burger, B. R. Baron and J. A. Tunge, *Synlett*, 2006, 2824–2826; (d) S. R. Schulz and S. Blechert, *Angew. Chem., Int. Ed.*, 2007, **46**, 3966–3970.
- 18 (a) J. P. Vigneron, M. Dhaenes and A. Horeau, *Tetrahedron*, 1973, **29**, 1055–1059; (b) V. Rautenstrauch, *Bull. Soc. Chim. Fr.*, 1994, **131**, 515–524; (c) S. E. Baba, K. Sartor, J. Poulin and H. Kagan, *Bull. Soc. Chim. Fr.*, 1994, **131**, 525–533.
- 19 J. T. Mohr, T. Nishimata, D. C. Behenna and B. M. Stoltz, *J. Am. Chem. Soc.*, 2006, **128**, 11348–11349.
- 20 For related Pd-catalyzed systems for enantioselective protonation, see: (a) F. Hénin and J. Muzart, *Tetrahedron: Asymmetry*, 1992, **3**, 1161–1164; (b) S. J. Abouldhoda, F. Hénin, J. Muzart, C. Thorey, W. Behnen, J. Martens and T. Mehler, *Tetrahedron: Asymmetry*, 1994, **5**, 1321–1326; (c) S. J. Abouldhoda, S. Létinois, J. Wilken, I. Reiners, F. Hénin, J. Martens and J. Muzart, *Tetrahedron: Asymmetry*, 1995, **6**, 1865–1868; (d) J. Muzart, F. Hénin and S. J. Abouldhoda, *Tetrahedron: Asymmetry*, 1997, **8**, 381–389; (e) M. A. Baur, A. Riahi, F. Hénin and J. Muzart, *Tetrahedron: Asymmetry*, 2003, **14**, 2755–2761; (f) M. Sugiura and T. Nakai, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 2366–2368; (g) Y. Hamashima, H. Somei, Y. Shimura, T. Tamura and M. Sodeoka, *Org. Lett.*, 2004, **6**, 1861–1864.
- 21 Despite the proposed common intermediate, experimental evidence suggests that the mechanism of C–H bond-formation in the enantioselective protonation is substantially different from the mechanism of C–C bond-formation in the enantioselective allylation. See: (a) ref. 19; (b) J. A. Keith, D. C. Behenna, J. T. Mohr, S. Ma, S. C. Marinescu, J. Oxaard, B. M. Stoltz and W. A. Goddard, III, *J. Am. Chem. Soc.*, 2007, **129**, 11876–11877.
- 22 Although many catalytic systems for enolate functionalization are known in the literature, the majority access the intermediate enolate from the corresponding achiral silyl enol ether in order to exclude Lewis basic amines. For representative examples, see: (a) A. Yanagisawa, T. Kikuchi, T. Watanabe, T. Kuribayashi and H. Yamamoto, *Synlett*, 1995, 372–374; (b) ref. 20f; (c) Y. Yamashita, K. Odashima and K. Koga, *Tetrahedron Lett.*, 1999, **40**, 2803–2806.
- 23 E. Vedejs and A. W. Kruger, *J. Org. Chem.*, 1998, **63**, 2792–2793.
- 24 (a) F. Hénin, J. Muzart, J.-P. Pete, A. M'boungou-M'passi and H. Rau, *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 416–418; (b) F. Hénin, A. M'boungou-M'passi, J. Muzart and J.-P. Pete, *Tetrahedron*, 1994, **50**, 2849–2864.
- 25 (a) C. Fischer and G. C. Fu, *J. Am. Chem. Soc.*, 2005, **127**, 4594–4595; (b) F. O. Arp and G. C. Fu, *J. Am. Chem. Soc.*, 2005, **127**, 10482–10483.
- 26 B. M. Trost and X. Ariza, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 2635–2637.