

Measurement and calculation of the two-dimensional backscattering Mueller matrix of a turbid medium: errata

Brent D. Cameron

Biomedical Engineering Program, Texas A&M University, College Station, Texas 77843-3120

M. J. Raković

*Center for Theoretical Physics, Department of Physics, Texas A&M University, College Station, Texas 77843-4242, and
Institute of Physics, University of Belgrade, P.O. Box 57, 11001 Belgrade, Yugoslavia*

Mehrübe Mehrübeoğlu

Department of Electrical Engineering, Texas A&M University, College Station, Texas 77843-3128

George W. Kattawar

Center for Theoretical Physics, Department of Physics, Texas A&M University, College Station, Texas 77843-4242

Sohi Rastegar, Lihong V. Wang, and Gerard L. Coté

Biomedical Engineering Program, Texas A&M University, College Station, Texas 77843-3120

In our recent Letter,¹ several typographical errors were present. On p. 487, in Fig. 2, the equations for the following Mueller matrix elements should read as $S_{14} = (RO - LO)$, $S_{22} = (HH + VV) - (HV + VH)$, $S_{23} = (PH + MV) - (PV + MH)$, $S_{24} = (RH + LV) - (RV + LH)$, $S_{32} = (HP + VM) - (HM + VP)$, $S_{33} = (PP + MM) - (PM + MP)$, $S_{34} = (RP + LM) - (RM + LP)$, $S_{41} = (OR - OL)$, $S_{42} = (HR + VL) - (HL + VR)$, $S_{43} = (PR + ML) - (PL + MR)$, and $S_{44} = (RR + LL) - (RL + LR)$. Also on p. 487, in the left-hand column, line 10 from the top should read as follows: $mf p' = 1/[\mu_a + \mu_s(1 - g)]$, was 0.957 cm.

Reference

1. B. D. Cameron, M. J. Raković, M. Mehrübeoğlu, G. W. Kattawar, S. Rastegar, L. V. Wang, and G. L. Coté, *Opt. Lett.* **23**, 485 (1998).