

# Coronal properties of the luminous radio-quiet quasar QSO B2202–209

E. S. Kammoun,<sup>1</sup>★ G. Risaliti,<sup>2,3</sup> D. Stern,<sup>4</sup> H. D. Jun,<sup>4</sup> M. Graham,<sup>5</sup> A. Celotti,<sup>1,6,7</sup>  
E. Behar,<sup>8</sup> M. Elvis,<sup>9</sup> F. A. Harrison,<sup>5</sup> G. Matt<sup>10</sup> and D. J. Walton<sup>4,5</sup>

<sup>1</sup>SISSA, via Bonomea 265, I-34135 Trieste, Italy

<sup>2</sup>Dipartimento di Fisica e Astronomia, Università di Firenze, via G. Sansone 1, I-50019 Sesto Fiorentino (Firenze), Italy

<sup>3</sup>INAF – Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125 Firenze, Italy

<sup>4</sup>Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA

<sup>5</sup>California Institute of Technology, 1200 East California Boulevard, Pasadena, CA 91125, USA

<sup>6</sup>INAF – Osservatorio Astronomico di Brera, via Bianchi 46, I-23807 Merate, Italy

<sup>7</sup>INFN – Sezione di Trieste, via Valerio 2, I-34127 Trieste, Italy

<sup>8</sup>Department of Astronomy, University of Maryland, College Park, MD 20742, USA

<sup>9</sup>Harvard Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

<sup>10</sup>Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, via della Vasca Navale 84, I-00146 Roma, Italy

Accepted 2016 November 7. Received 2016 November 5; in original form 2016 July 23

## ABSTRACT

We present an analysis of the joint *XMM–Newton* and *NuSTAR* observations of the radio-quiet quasar QSO B2202–209. Using an optical observation from the Hale Telescope at the Palomar Observatory, we revise the redshift of the source from the previously reported  $z = 1.77$  to  $z = 0.532$ , and we estimate the mass of the central black hole,  $\log(M_{\text{BH}}/M_{\odot}) = 9.08 \pm 0.18$ . The X-ray spectrum of this source can be well described by a power law of photon index  $\Gamma = 1.82 \pm 0.05$  with  $E_{\text{cut}} = 152_{-54}^{+103}$  keV, in the rest frame of the source. Assuming a Comptonization model, we estimate the coronal temperature to be  $kT_e = 42 \pm 3$  keV and  $kT_e = 56 \pm 3$  keV for a spherical and a slab geometry, respectively. The coronal properties are comparable to the ones derived for local active galactic nuclei, despite a difference of around one order of magnitude in black hole mass and X-ray luminosity ( $L_{2-10} = 1.93 \times 10^{45}$  erg s<sup>−1</sup>). The quasar is X-ray loud, with an unusually flat observed optical-to-X-ray spectral slope  $\alpha_{\text{OX}} = 1.00 \pm 0.02$ , and has an exceptionally strong optical [O III] line. Assuming that both the X-ray emission and the [O III] line are isotropic, these two extreme properties can be explained by a nearly edge-on disc, leading to a reduction in the observed ultraviolet continuum light.

**Key words:** galaxies: active – galaxies: nuclei – quasars: individual: QSO B2202–209 – X-rays: galaxies.

## 1 INTRODUCTION

It is generally thought that the primary hard X-ray continuum emission arising around accreting black holes [active galactic nuclei (AGN) and X-ray binaries], is due to Compton up-scattering of ultraviolet (UV)/soft X-ray disc photons off a hot ( $\sim 10^9$  K), trans-relativistic medium, usually referred to as the X-ray corona (e.g. Shapiro, Lightman & Eardley 1976; Haardt & Maraschi 1993; Petrucci et al. 2001a,b). The resulting spectrum, known as the ‘primary’ emission, can be described by a power law with a high-energy exponential cutoff that depends on the Compton  $y$ -parameter of the medium, which is a function of the electron temperature ( $kT_e$ ) and the optical depth ( $\tau$ ) of the corona. These electrons are thought to be heated and confined by magnetic fields emerging from the

ionized accretion disc (e.g. Galeev, Rosner & Vaiana 1979; Haardt & Maraschi 1991; Merloni & Fabian 2001).

In addition to being directly detected by an observer at infinity, the primary emission will irradiate and be ‘reflected’ by the accretion disc (George & Fabian 1991). X-rays incident on the disc will be subjected to Compton scattering and photoelectric absorption followed either by Auger de-excitation or by fluorescent line emission (see e.g. Lightman & White 1988; George & Fabian 1991). The resulting reflection spectrum is characterized by the iron K  $\alpha$  emission line at  $\sim 6.4$ – $7.0$  keV (depending on the ionization state of the disc) and a broad component peaked at around 20–30 keV, known as ‘Compton hump’. It should be noted that the accretion disc is not the only reflector able to reprocess the primary emission in AGN. Many sources reveal the presence of narrow Fe K  $\alpha$  emission lines (e.g. Bianchi et al. 2009) that could be explained by reflection from distant, dense material such as the narrow-line region or the putative molecular

\* E-mail: ekammoun@sissa.it

torus invoked in AGN unification models (Ghisellini, Haardt & Matt 1994).

Several lines of evidence suggest that the corona is compact, and located close to the black hole. Spectral-timing and reverberation studies, for example, are suggestive of a physically small corona that lies  $3\text{--}10 r_g$  [where  $r_g = GM/c^2$  is the gravitational radius of a black hole (BH) with mass  $M$ ] above the central BH (e.g. Fabian et al. 2009; Emmanoulopoulos et al. 2014; Parker et al. 2014; Gallo et al. 2015). It has been inferred from X-ray microlensing analyses of some bright lensed quasars that the hard X-rays are emitted from compact regions with half-light radii less than  $6 r_g$  (Morgan et al. 2008; Chartas et al. 2009; Reis & Miller 2013). Moreover, eclipses of the X-ray source have also placed constraints on the size of the hard X-ray emitting regions,  $r \lesssim 10 r_g$  (e.g. Risaliti et al. 2007; Maiolino et al. 2010; Sanfrutos et al. 2013). Analyses of the emissivity profile of the accretion disc (Wilkins & Fabian 2011, 2012) suggest an extended corona a few gravitational radii from the accretion disc. More recently, Wilkins et al. (2015) and Wilkins & Gallo (2015) explained the long and short time-scale variabilities seen in Mrk 335 in the context of an ‘aborted jet launch’ as proposed earlier by Ghisellini, Haardt & Matt (2004). However, the coronal temperature and optical depth have remained poorly constrained due to the lack of high-quality X-ray measurements extending above 10 keV. Complex spectral components, including reflection from the accretion disc as well as from distant matter, contribute to the whole X-ray spectrum, making a precise determination of the cutoff energy challenging.

The unprecedented sensitivity of *NuSTAR* (Harrison et al. 2013) covering the 3–79 keV band, has provided for the first time the capability to measure spectral parameters in a precise and robust way including both the Compton reflection component and the high-energy cutoff ( $E_{\text{cut}} \sim 100\text{--}180$  keV) in some of the brightest local AGN, with luminosities of the order of  $10^{42\text{--}44}$  erg s $^{-1}$  in the *NuSTAR* energy band (e.g. Ballantyne et al. 2014; Brenneman et al. 2014; Marinucci et al. 2014; Baloković et al. 2015; Matt et al. 2015). Combining the high sensitivity of *XMM-Newton* (Jansen et al. 2001) at soft X-ray energies with that of *NuSTAR* at hard energies, we are able to investigate the coronal properties of more distant and more luminous quasars. This allows us to understand better the coronal physics and the dependence of the corona on luminosity and redshift.

In this work, we study the X-ray spectrum provided by a coordinated *XMM-Newton* and *NuSTAR* observation of the radio-quiet quasar (RQQ) QSO B2202–209 (hereafter B2202, also known as PB 5062). The redshift of the source was estimated by Reboul et al. (1987) to be  $z = 1.77$ , implying a luminosity of  $L_{2\text{--}10} \simeq 3 \times 10^{46}$  erg s $^{-1}$ , which would make this source erroneously, as we will show, the most luminous RQQ within its redshift (Eitan & Behar 2013). However, by analysing its X-ray spectrum, we found the need of an additional absorber at an intermediate redshift of 0.53. Reboul et al. (1987) determined the redshift of the source by identifying a broad emission line observed at  $\sim 4290$  Å with a C IV  $\lambda 1549$  line. However, instead identifying the line with Mg II  $\lambda 2800$  implies a redshift of 0.532, compatible with the redshift of the X-ray absorber. This discrepancy motivated us to obtain a higher quality optical spectrum using the Hale Telescope at Palomar Observatory, presented in the next section. The Palomar data show that the source is actually situated at a redshift of 0.532. The main aim of this project is to measure the coronal properties of the source. The measurements of  $E_{\text{cut}}$  for local radio-quiet AGN are based on the deviation (of the order of 10–30 per cent) from a power law at the highest energy part of the *NuSTAR* spectrum. Considering the

redshift of B2202, particularly for the previously assumed  $z = 1.77$ ,  $E_{\text{cut}}$  is shifted to lower observed energies, helping compensate for the cosmological dimming.

This paper is structured as follows: Section 2 describes the observations and the data reduction; Section 3 presents the spectral analysis. Finally, in Section 4, we provide a brief summary of our main results and discuss their implications. The following cosmological parameters are assumed:  $\Omega_M = 0.27$ ,  $\Omega_\Lambda = 0.73$  and  $H_0 = 70$  km s $^{-1}$  Mpc $^{-1}$ .

## 2 OBSERVATIONS AND DATA REDUCTION

### 2.1 Palomar observations

We obtained an optical spectrum of B2202 using the dual-beam Double Spectrograph on the 200-inch Hale Telescope at Palomar Observatory. The 900 s spectrum, obtained on UT 2016 May 28 in photometric conditions, used the 1.5 arcsec wide slit, the 5500 Å dichroic to split the light, the 600  $\ell$  mm $^{-1}$  grating on the blue arm ( $\lambda_{\text{blaze}} = 4000$  Å; spectral resolving power  $R \equiv \lambda/\Delta\lambda \sim 1200$ ), and the 316  $\ell$  mm $^{-1}$  grating on the red arm ( $\lambda_{\text{blaze}} = 7500$  Å;  $R \sim 1800$ ). We processed the data using standard techniques within IRAF, and calibrated the spectrum using an archival sensitivity function obtained in 2016 February using the same instrument configuration and observing conditions.

The processed spectrum, shown in the upper panel of Fig. 1, shows the source to be an AGN at  $z = 0.532$  with many standard emission features identified, including broad Mg II  $\lambda 2800$ , broad hydrogen Balmer emission lines, and narrow [O III]  $\lambda\lambda 4959, 5007$ ; the redshift has been determined from the latter features. In particular, the new data clearly demonstrate that the line at  $\sim 4290$  Å previously identified as C IV by Reboul et al. (1987) is indeed Mg II.

### 2.2 X-ray observations


B2202 was observed simultaneously by *XMM-Newton* and *NuSTAR*, on 2015 November 06–07 (Obs. IDs 0764370201 and 60101030002, respectively). The basic observation details are presented in Table 1. Here we summarize our data-reduction procedures.

#### 2.2.1 XMM-Newton

The *XMM-Newton* data were reduced using SAS v.15.0.0 and the latest calibration files. We followed the standard procedure for reducing the data of the EPIC-pn (Strüder et al. 2001) and the two EPIC-MOS (Turner et al. 2001) CCD cameras, all operating in

**Table 1.** Net exposure times, and count rates estimated from the background-subtracted data in the 0.5–10 keV range for *XMM-Newton*, and the 4–30 keV range for *NuSTAR*.

Instrument	Net exposure (ks)	Count Rate (count s $^{-1}$ )
EPIC-PN	60.5	$0.501 \pm 0.003$
MOS 1	69.7	$0.152 \pm 0.001$
MOS 2	69.4	$0.156 \pm 0.001$
FPMA	106.8	$0.0233 \pm 0.0005$
FPMB	106.5	$0.0227 \pm 0.0005$




**Figure 1.** Upper panel: optical spectrum obtained using the Palomar DBSP. Lower panels: X-ray background-subtracted source spectra (continuous lines) and background spectra (dots). Left-hand panel: the spectra extracted from the EPIC-pn (black), EPIC MOS1 (blue) and EPIC MOS2 (red) instruments. Right-hand panel: the spectra extracted from the FPMA (black) and FPMB (red) modules.

full frame mode with a thin filter for the EPIC-pn and a medium filter for the EPIC-MOS cameras. The EPIC-pn and EPIC-MOS data were processed using `EPPROC` and `EMPROC`, respectively. Source spectra and light curves were extracted from a circular region of radius  $\sim 40$  arcmin. The corresponding background spectra and light curves were extracted from an off-source circular region located on the same CCD chip, with a radius approximately twice that of the source to ensure a high signal-to-noise ratio. We filtered out periods with strong background flares estimated to be around 7.5 ks. We corrected the light curves for the background count rate using `EPICLCCORR`. The extracted light curves did not show any significant spectral variability. Response matrices were produced using the `FTOOLS` `RMFGEN` and `ARFGEN`. We rebinned the observed spectra, shown in the bottom left panel of Fig. 1, using the `SAS` task `SPECGROUP` to have a minimum S/N of 5 in each energy bin. The MOS1 and MOS2 spectra are consistent, so we combined them using the `SAS` command `COMBINE`.

### 2.2.2 *NuSTAR*

We reduced the *NuSTAR* data following the standard pipeline in the *NuSTAR* Data Analysis Software (`NUSTARDAS` v1.4.1), and instrumental responses from *NuSTAR* `CALDB` v20151008. We cleaned the unfiltered event files with the standard depth correction, which significantly reduces the internal background at high energies, and South Atlantic Anomaly passages were excluded from our analysis. We extracted the time-averaged source and background spectra from circular regions of radii 40 and 100 arcsec, respectively, for both focal plane modules (FPMA and FPMB) using the `HEASOFT` task `NUPRODUCT`, and requiring a minimum of 50 counts per bin. The extracted spectra are presented in the bottom right panel of Fig. 1. The spectra extracted from both modules are consistent with each other. The background starts to dominate the source above  $\sim 30$  keV. For that reason, we decided to analyse the *NuSTAR* data in the observed 4–30 keV energy range, which corresponds to the




**Figure 2.** Upper panel: observed spectra from EPIC-pn (blue) and *NuSTAR* (green) plotted together with the best-fitting model (black line) composed of an absorbed cutoff power law. Lower panel: data/model ratio for the EPIC-pn (blue), EPIC-MOS (red) and *NuSTAR* (green) data.

6–46 keV energy range in the rest frame of the source. The data from FPMA and FPMB are analysed jointly in this work, but they are not combined together.

### 3 SPECTRAL ANALYSIS

Throughout this work, spectral fitting was done using *XSPEC* v12.9 (Arnaud 1996). Unless stated otherwise, uncertainties are listed at the 90 per cent confidence level ( $\Delta\chi^2 = 2.71$ ). We included a variable constant, for each instrument, in order to account for the residual uncertainties in the flux calibration between the various detectors, fixing the constant for the EPIC-pn data to unity. We considered the EPIC-pn and the merged EPIC-MOS spectra in the 0.5–10 keV range, and the FPMA/B spectra in the 4–30 keV band, as mentioned above.

First, we fitted the spectra with a simple, absorbed power law with an exponential high-energy cutoff (`cutoffpl` model in *XSPEC*). We fixed the Galactic absorption to the equivalent hydrogen column density in the line of sight towards B2202,  $N_{\text{H}} = 5.52 \times 10^{20} \text{ cm}^{-2}$  (Kalberla et al. 2005). Furthermore, we considered an intrinsic neutral absorber at the redshift of the source (`zwaabs`). The model fit is unacceptable ( $\chi^2/\nu = 1.32$ ), mostly due to the need of an extra ionized absorption component. For that reason, we add a partial covering absorption by partially ionized material (`zxipcf`) at the redshift of the source. The fit improved ( $\chi^2/\nu = 1.12$ ). However, a significant deficit could be detected in the residuals at energy  $\sim 1.8$  keV, most probably due to inaccuracies in modelling the Si absorption in the CCD detectors. We modelled this deficit using a Gaussian absorption profile with a free centroid energy, width and normalization (in the observed frame). The fit improves significantly by adding this line ( $\Delta\chi^2/\Delta\nu = 25/3$ , *F*-test null probability =  $4.14 \times 10^{-5}$ ), resulting in a  $\chi^2/\nu = 1.05$ . The final model and data/model ratio are presented in Fig. 2. We performed a Monte Carlo Markov Chain (MCMC) analysis to estimate the errors on the parameters, using the Goodman–Weare algorithm (Goodman & Weare 2010) with a chain of 50 000 elements, and discarding the first 5000 elements as part of the ‘burn-in’ period. We found a high-energy cutoff  $E_{\text{cut}} = 99_{-35}^{+67}$  keV in the observed frame that corresponds to an energy of  $152_{-54}^{+103}$  keV in the rest frame of the source,



**Figure 3.** Upper panel:  $\Gamma - E_{\text{cut}}$  contour plot (in the rest frame) for the cutoff power-law model. Lower panel:  $kT_e - \tau$  contour plot (in the rest frame) for the Comptonization model assuming a spherical geometry and a slab geometry. We plot the 68 per cent (dotted red lines), 95 per cent (dashed green lines) and 99.7 per cent (solid blue lines) confidence levels.

with a photon index  $\Gamma = 1.82 \pm 0.05$ . In the upper panel of Fig. 3, we present the contour plots derived from the MCMC analysis showing the constraints on the photon index  $\Gamma$  and the high-energy cutoff  $E_{\text{cut}}$  in the rest frame of the source. Moreover, we identified a partially ionized absorber ( $\log [\xi (\text{erg s}^{-1} \text{ cm}^{-1})] = 0.39_{-0.75}^{+0.80}$ ) situated at the redshift of the source covering  $\sim 30$  per cent of the source and having a column density  $N_{\text{H}} = 2.32_{-0.88}^{+0.83} \times 10^{23} \text{ cm}^{-2}$ . In addition, we identified a neutral absorber at the rest frame of the source with a column density  $N_{\text{H}} = 1.4_{-0.20}^{+0.22} \times 10^{21} \text{ cm}^{-2}$ . The best-fitting parameters are reported in Table 2.

The measurement of  $E_{\text{cut}}$  could be affected by the presence of a reflection component in the spectrum. In order to test the stability of our measurement, we replaced the power-law model by the neutral reflection model `PEXMON` (Nandra et al. 2007), considering an intermediate inclination of  $60^\circ$  and solar abundances. The fit improves by  $\Delta\chi^2 = -6$  for one additional free parameter. We found a low reflection fraction ( $R_{\text{frac}} = 0.31_{-0.21}^{+0.24}$ ) and a low high-energy cutoff of  $E_{\text{cut}} = 69_{-25}^{+47}$  keV in the rest frame of the source, but consistent within the error bars with the value found assuming a simple cutoff power-law model. In addition, we tested the possibility of having relativistic reflection using the `relxill` model (Dauser et al. 2013, 2016; García et al. 2014). The fit was

**Table 2.** Best-fitting parameters obtained assuming a cutoff power-law model, and a Comptonization model for both spherical and slab geometries.

Parameter	Cutoff	Compton (spherical)	Compton (slab)
	zwabs		
$N_{\text{H}} (10^{21} \text{ cm}^{-2})$	$1.40^{+0.22}_{-0.20}$	$1.40^{+0.13}_{-0.12}$	$1.38^{+0.15}_{-0.13}$
	zxcipcf		
$N_{\text{H}} (10^{23} \text{ cm}^{-2})$	$2.32^{+0.83}_{-0.88}$	$2.32^{+0.68}_{-0.55}$	$2.43^{+0.77}_{-0.45}$
$\log [\xi (\text{erg s}^{-1} \text{ cm}^{-1})]$	$0.39^{+0.80}_{-0.75}$	$0.40^{+0.79}_{-0.47}$	$0.36^{+0.76}_{-0.35}$
CF (per cent)	$30^{+6}_{-5}$	$30^{+4}_{-3}$	$29^{+4}_{-3}$
	cutoffpl		
$\Gamma$	$1.82 \pm 0.05$	–	–
$E_{\text{cut}}^a$ (keV)	$99^{+67}_{-35}$	–	–
Norm ( $\times 10^{-4}$ )	$5.26^{+0.54}_{-0.41}$	–	–
	compTT		
$kT_e^b$ (keV)	–	$42 \pm 3$	$56 \pm 3$
$\tau$	–	$2.40^{+0.18}_{-0.16}$	$0.63^{+0.06}_{-0.05}$
Norm ( $\times 10^{-4}$ )	–	$4.23^{+0.45}_{-0.41}$	$3.07^{+0.34}_{-0.28}$
	Gaussian (absorption)		
$E$ (keV)	$1.8 \pm 0.04$	$1.8 \pm 0.04$	$1.8 \pm 0.03$
$\sigma$ (eV)	$72^{+55}_{-40}$	$72^{+47}_{-34}$	$73^{+54}_{-38}$
Norm ( $\times 10^{-6}$ )	$4.75^{+1.77}_{-2.34}$	$4.73^{+1.47}_{-1.87}$	$4.79^{+1.58}_{-2.17}$
$\chi^2/\text{d.o.f.}$	363/344	363/344	363/344
	1.05	1.05	1.05

Notes. <sup>a</sup>The cutoff energy is reported in the observed frame.

<sup>b</sup>The electron temperature is reported in the rest frame of the source.

statistically accepted ( $\chi^2/\nu = 1.04$ ), and  $E_{\text{cut}}$  was consistent with the values determined above. However, neither the reflection fraction nor the ionization parameter of the disc could be well constrained.

Finally, assuming that the power-law spectrum is obtained by Comptonization of soft photons arising from the accretion disc by hot electrons in a corona, we substituted the cutoff power law with the Comptonization model `compTT` (Titarchuk 1994). This model allows us to determine the coronal temperature and optical depth assuming either a spherical or a slab geometry. We fixed the temperature of the seed photons, assumed to follow a Wien distribution law, to  $kT = 20$  eV. We obtained a fit statistically similar to the one obtained for the cutoff power-law model. We performed an MCMC analysis similar to the one mentioned previously, and found the coronal temperature and optical depth to be  $kT_e = 42 \pm 3$  keV,  $\tau = 2.40^{+0.18}_{-0.16}$  (spherical geometry) and  $kT_e = 56 \pm 3$  keV,  $\tau = 0.63^{+0.06}_{-0.05}$  (slab geometry). The electron temperatures are in agreement with the estimation of  $E_{\text{cut}}$ , within the error bars, obtained assuming a high-energy cutoff power-law model ( $E_{\text{cut}} \simeq 2\text{--}3kT_e$ , e.g. Petrucci et al. 2001b). The optical depth is larger for the spherical geometry. This is expected because the optical depth estimated assuming a spherical geometry is the radial (effective) depth, while the slab optical depth is the vertical one that should be lower than the effective one. The  $kT_e - \tau$  contour plots for both geometries are presented in the lower panel of Fig. 3. The best-fitting parameters are shown in Table 2. The observed 2–10 keV flux is  $F_{2-10} = (1.43 \pm 0.07) \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ , while the unabsorbed 2–10 keV luminosity, in the rest frame, is  $L_{2-10} = (1.93 \pm 0.07) \times 10^{45} \text{ erg s}^{-1}$ .

B2202 was also observed by *XMM-Newton* in 2001 May (Obs. ID 12440301, hereafter obs. 1), for  $\sim 30$  ks. We extracted the EPIC-

**Table 3.** Full widths at half-maximum and equivalent widths of the broad and narrow H  $\beta$  lines and the [O III] $\lambda\lambda 4959, 5007$  doublet seen in the Palomar spectra.

Line	FWHM ( $\text{km s}^{-1}$ )	EW ( $\text{\AA}$ )
H $\beta_{\text{broad}}$	$6940 \pm 1033$	$128.4 \pm 5.9$
H $\beta_{\text{narrow}}$	$961 \pm 49$	$18.4 \pm 8.5$
[O III] $\lambda 4959$	$1144 \pm 77$	$49 \pm 3$
[O III] $\lambda 5007$	$995 \pm 50$	$145.7 \pm 3.4$

pn/MOS data in a similar way to that described in Section 2.2.1. We fit the EPIC spectra of this observation by an absorbed power-law model, similar to the best-fitting model presented above, but without considering either the high-energy cutoff or the absorption feature at  $\sim 1.8$  keV. The fit is good ( $\chi^2/\nu = 0.97$ ) and the parameters of both neutral and ionized absorbers are consistent, within the error bars, with the ones found in 2015. However, we found a steeper power law with a photon index  $\Gamma = 1.97^{+0.05}_{-0.06}$ . During obs. 1, the source was  $\sim 1.5$  times brighter than in 2015 (unabsorbed luminosity,  $L_{2-10} = 2.92 \times 10^{45} \text{ erg s}^{-1}$ ) showing a behaviour often observed in AGNs, where the X-ray spectrum gets steeper when the source is brighter.

## 4 DISCUSSION AND CONCLUSIONS

We present joint *XMM-Newton* and *NuSTAR* observations of the luminous quasar B2202. The 0.5–30 keV spectrum of this source could be fitted with an absorbed (a neutral absorber and a partially ionized absorber at the redshift of the source) power law ( $\Gamma \sim 1.82$ ) with an exponential high-energy cutoff ( $E_{\text{cut}} \sim 153$  keV, at the rest frame of the source). No strong reflection features have been identified in the spectrum.

In order to derive the black hole mass ( $M_{\text{BH}}$ ) of B2202, we fit the H  $\beta$ -line region with the `IDL` package `MPFIT` (Markwardt 2009). Approximating the uncertainties to the flux using the standard deviation of the spectrum outside the emission lines, we fit the rest frame 4200–5600  $\text{\AA}$  including a power-law continuum, broad and narrow Gaussian components for the H  $\beta$  line, narrow [O III] $\lambda\lambda 4959, 5007$  doublet, narrow He II  $\lambda 4686$  and the blended H  $\gamma$  and [O III] $\lambda 4363$ . The contribution from the Fe II complex is negligible for this target and thus was ignored. The full widths at half-maximum (FWHM) and the equivalent widths (EW) of the H  $\beta$  lines and the [O III] $\lambda\lambda 4959, 5007$  doublet are presented in Table 3. Using the width of the broad H  $\beta$  line and the monochromatic 5100  $\text{\AA}$  luminosity ( $L_{5100}$ ), we get  $\log(M_{\text{BH}}/M_{\odot}) = 9.08 \pm 0.18$  from the Jun et al. (2015) estimator. This implies an Eddington luminosity of  $L_{\text{Edd}} = (1.56 \pm 0.28) \times 10^{47} \text{ erg s}^{-1}$ . Note that  $L_{5100}$  is derived directly from the spectrum, that was checked to yield consistent  $i$ -band magnitude to that from the SDSS<sup>1</sup> (Ahn et al. 2014) imaging, within a 10 per cent difference. We applied then the bolometric correction to  $L_{5100}$  suggested by Marconi et al. (2004),  $L_{\text{bol}}/\nu_{\text{B}}L_{\text{B}} = 7.9 \pm 2.9$ , and we estimate the bolometric luminosity of this source to be  $L_{\text{bol}} = (5.72 \pm 2.15) \times 10^{45} \text{ erg s}^{-1}$ . This means that the source is accreting at  $\sim 3.66$  per cent of its Eddington limit. However, by applying the bolometric correction to  $L_{2-10}$  (equation 21 in Marconi et al. 2004), we obtain a higher bolometric luminosity,  $L_{\text{bol}} = 1.79 \times 10^{47} \text{ erg s}^{-1} \simeq 1.15 L_{\text{Edd}}$ . Moreover, we estimated the monochromatic luminosity of the source at 2500  $\text{\AA}$  to be  $L_{\nu}(2500 \text{ \AA}) = (1.06 \pm 0.02) \times 10^{45} \text{ erg s}^{-1}$ , which implies an optical-to-X-ray spectral slope


<sup>1</sup> <http://skyserver.sdss.org/dr10/en/home.aspx>

$\alpha_{\text{OX}} = -0.384 \times \log [L_{2\text{keV}}/L_{2500}] = 1.01 \pm 0.02$ . This is a strong outlier in the  $\alpha_{\text{OX}}$ –luminosity relation ( $\alpha_{\text{OX}} \sim 1.6$  for typical quasars of the same luminosity, Lusso & Risaliti 2016). However, the large EW of  $[\text{O III}]\lambda 5007$  can be indicative of a high-inclination angle  $\theta$  between the disc axis and the line of sight. In fact, we observe the projected EW along the line of sight,  $\text{EW}_O = \text{EW}^* / \cos \theta$ , where  $\text{EW}_O$  is the observed EW, and  $\text{EW}^*$  is the EW as measured in a face-on disc. By considering an average value of  $\text{EW}^*$ , for all quasars, to be  $\sim 11 \text{ \AA}$  (Risaliti, Salvati & Marconi 2011; Bisogni, Marconi & Risaliti 2017), we obtain  $\theta \simeq 85^\circ$ . This means that the disc is observed nearly edge-on and can explain the high X-ray loudness of the source. In other terms, having such a high inclination of the disc with respect to the line of sight, the intrinsic UV luminosity should be higher than the observed one. Instead, the inclination will not affect the X-ray primary that is thought to be emitted isotropically. If we assume an intrinsic  $L_{5100}$  to be larger than the observed value by a factor of  $1/\cos \theta$ , then this leads to an intrinsic  $\alpha_{\text{OX, int}} \simeq 1.43$  comparable to other sources. In addition, using the  $[\text{O III}]\lambda 5007$  luminosity  $[L_{[\text{O III}]}] = (2.07 \pm 0.32) \times 10^{43} \text{ erg s}^{-1}$ , we estimated the expected X-ray luminosity, following the correlation found by Panessa et al. (2006), to be  $L_X = (3.2 \pm 0.6) \times 10^{45} \text{ erg s}^{-1}$ . The estimated value of  $L_X$  is in agreement within a factor of 1.65 with the value derived from the X-ray analysis in Section 3. Given the isotropic nature of the X-ray primary emission, the high inclination of the disc and the agreement between the X-ray and the  $[\text{O III}]\lambda 5007$  measurements, this is suggestive that, for this source, the measurement of  $L_{\text{bol}}$  using the X-ray luminosity is more reliable compared to the bolometric correction estimated using  $L_{5100}$ .

Interestingly, we found that the coronal properties are in agreement with the ones determined for local less luminous and less massive Seyfert galaxies (e.g. Fabian et al. 2015; Lubiński et al. 2016, and references therein). Lubiński et al. (2016) analysed the hard

X-ray spectra of 28 Seyfert galaxies based on observations with *INTEGRAL* (Winkler et al. 2003). The values of the electron temperature, photon index and reflection fraction that we found are in good agreement with the median values of their full sample ( $\langle kT_e \rangle = 48^{+57}_{-14} \text{ keV}$ ,  $\langle \Gamma \rangle = 1.81^{+0.18}_{-0.05}$  and  $\langle R_{\text{frac}} \rangle = 0.32^{+0.33}_{-0.28}$ , respectively). Lubiński et al. (2016) suggest that the small value of  $R_{\text{frac}}$  is indicative of a small solid angle under which the corona is seen from the disc. In other terms, this means a reduction in the flux of seed photons emitted by the disc that are cooling the corona. Moreover, they suggest the need of an additional process (such as synchrotron self-Compton) that is able to explain the efficient cooling of a compact corona, close to the BH and separated geometrically from the accretion disc. Note that the low amount of reflection is in agreement with the X-ray Baldwin effect (e.g. Iwasawa & Taniguchi 1993) whereby the equivalent width of the Fe K  $\alpha$  line diminishes with X-ray luminosity. In fact, we modelled the Fe K  $\alpha$  line by adding a Gaussian in emission, to the high-energy cutoff power-law model, and we estimated its EW to be  $\leq 40 \text{ eV}$ . However, in our case, the high inclination of the disc can explain the small values of  $R_{\text{frac}}$  and  $\text{EW}(\text{Fe K } \alpha)$ , since most of the reflected light coming from the accretion disc will not be emitted in the direction of the observer. Instead, the weak reflection can be associated with sparse clouds, most likely to be in the broad-line region, covering a small fraction of the solid angle. The similarities in both spectral shape and cutoff energy between B2202 and local AGNs are therefore indicative of a universal process determining the spectrum of the X-ray emission, with the strength of the coupling between disc and corona affecting only the overall normalization of such spectra.

We plot in Fig. 4 the observed multi-epoch spectral energy distribution (SED) of the source by adding, to the optical and X-ray spectra analysed in this paper, the *XMM–Newton* spectrum obtained



**Figure 4.** The observed SED of the source, obtained using photometric data from *WISE* in 2010 (magenta triangles) and *2MASS* in 2000 (cyan squares), in addition to the Palomar Observatory spectrum obtained in 2016 (red dotted line) and corrected for the Galactic extinction and for the Oxygen A-band absorption, the *XMM–Newton* spectrum obtained in 2001 (red dots) and the joint *XMM–Newton* and *NuSTAR* observations in 2015 (blue points). The typical radio-quiet quasar SEDs (Elvis et al. 1994) is also plotted (black solid line) for comparison.

in 2001 and the archival photometric data obtained from the Two Micron All Sky Survey<sup>2</sup> (2MASS, observed in 2000; Skrutskie et al. 2006), and the *Wide-field Infrared Survey Explorer*<sup>3</sup> (WISE, observed in 2010; Wright et al. 2010). We compare the observed SED to a typical SED of an RQQ (Elvis et al. 1994) trying to match the X-ray spectra obtained in 2015. Fitting the SED properly, taking into account the complications and the long time-scale variability of the source, is beyond the scope of this paper. However, it is obvious that the optical spectrum is fainter and redder compared to a standard SED, but this is expected given the high inclination of the accretion disc, which leads the emission from the host galaxy, usually negligible, to show up. Moreover, interestingly, the IR emission appears to be low compared to the standard quasar SED. Adding to this the fact that the X-ray spectra are barely affected by absorption, and the small amount of reflection, this implies a particular system in which we observe the accretion disc nearly edge-on with no evidence of any Compton thick parsec scale reprocessor, indicating a small covering fraction by the torus, if any.

This source was chosen on the basis of the previously reported redshift that mistakenly let it be considered one of the most luminous RQQs. However, our results on a less extreme quasar demonstrate that joint *NuSTAR* and *XMM–Newton* observations with a moderate exposure time are capable of making good measurements of coronal properties for quasars at cosmological redshifts. This significantly extends the previous work done primarily on relatively local and lower luminosity Seyferts (e.g. Fabian et al. 2015; Lubinowski et al. 2016).

## ACKNOWLEDGEMENTS

This research made use of data from the *NuSTAR* mission, a project led by the California Institute of Technology, managed by the Jet Propulsion Laboratory, and funded by NASA, *XMM–Newton*, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA, and the Hale Telescope at Palomar Observatory. This research has made use of the *NuSTAR* Data Analysis Software (NUSTARDAS) jointly developed by the ASI Science Data Center (ASDC, Italy) and the California Institute of Technology (USA). We like to acknowledge S.G. Djorgovski for providing the Palomar observations. We thank the anonymous referee for comments and suggestions, which significantly contributed to improving the quality of the manuscript. EB received funding from the European Union Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 655324.

Facilities: *NuSTAR*, Palomar: DBSP, *XMM–Newton*.

## REFERENCES

Ahn C. P. et al., 2014, *ApJS*, 211, 17  
 Arnaud K. A., 1996, in Jacoby G. H., Barnes J., eds, *ASP Conf. Ser. Vol. 101, Astronomical Data Analysis Software and Systems V*. Astron. Soc. Pac., San Francisco, p. 17  
 Ballantyne D. R. et al., 2014, *ApJ*, 794, 62  
 Baloković M. et al., 2015, *ApJ*, 800, 62  
 Bianchi S., Guainazzi M., Matt G., Fonseca Bonilla N., Ponti G., 2009, *A&A*, 495, 421  
 Bisogni S., Marconi A., Risaliti G., 2017, *MNRAS*, 464, 385  
 Brenneman L. W. et al., 2014, *ApJ*, 788, 61  
 Chartas G., Kochanek C. S., Dai X., Poindexter S., Garmire G., 2009, *ApJ*, 693, 174

Dauser T., Garcia J., Wilms J., Böck M., Brenneman L. W., Falanga M., Fukumura K., Reynolds C. S., 2013, *MNRAS*, 430, 1694  
 Dauser T., Garcia J., Walton D. J., Eikmann W., Kallman T., McClintock J., Wilms J., 2016, *A&A*, 590, A76  
 Eitan A., Behar E., 2013, *ApJ*, 774, 29  
 Elvis M. et al., 1994, *ApJS*, 95, 1  
 Emmanoulopoulos D., Papadakis I. E., Dovčiak M., McHardy I. M., 2014, *MNRAS*, 439, 3931  
 Fabian A. C. et al., 2009, *Nature*, 459, 540  
 Fabian A. C., Lohfink A., Kara E., Parker M. L., Vasudevan R., Reynolds C. S., 2015, *MNRAS*, 451, 4375  
 Galeev A. A., Rosner R., Vaiana G. S., 1979, *ApJ*, 229, 318  
 Gallo L. C. et al., 2015, *MNRAS*, 446, 633  
 García J. et al., 2014, *ApJ*, 782, 76  
 George I. M., Fabian A. C., 1991, *MNRAS*, 249, 352  
 Ghisellini G., Haardt F., Matt G., 1994, *MNRAS*, 267, 743  
 Ghisellini G., Haardt F., Matt G., 2004, *A&A*, 413, 535  
 Goodman J., Weare J., 2010, *Commun. Appl. Math. Comput. Sci.*, 5, 65  
 Haardt F., Maraschi L., 1991, *ApJ*, 380, L51  
 Haardt F., Maraschi L., 1993, *ApJ*, 413, 507  
 Harrison F. A. et al., 2013, *ApJ*, 770, 103  
 Iwasawa K., Taniguchi Y., 1993, *ApJ*, 413, L15  
 Jansen F. et al., 2001, *A&A*, 365, L1  
 Jun H. D. et al., 2015, *ApJ*, 806, 109  
 Kalberla P. M. W., Burton W. B., Hartmann D., Arnal E. M., Bajaja E., Morras R., Pöppel W. G. L., 2005, *A&A*, 440, 775  
 Lightman A. P., White T. R., 1988, *ApJ*, 335, 57  
 Lubinowski P. et al., 2016, *MNRAS*, 458, 2454  
 Lusso E., Risaliti G., 2016, *ApJ*, 819, 154  
 Maiolino R. et al., 2010, *A&A*, 517, A47  
 Marconi A., Risaliti G., Gilli R., Hunt L. K., Maiolino R., Salvati M., 2004, *MNRAS*, 351, 169  
 Marinucci A. et al., 2014, *MNRAS*, 440, 2347  
 Markwardt C. B., 2009, in Bohlender D. A., Durand D., Dowler P., eds, *ASP Conf. Ser. Vol. 411, Astronomical Data Analysis Software and Systems XVIII*. Astron. Soc. Pac., San Francisco, p. 251  
 Matt G. et al., 2015, *MNRAS*, 447, 3029  
 Merloni A., Fabian A. C., 2001, *MNRAS*, 328, 958  
 Morgan C. W., Kochanek C. S., Dai X., Morgan N. D., Falco E. E., 2008, *ApJ*, 689, 755  
 Nandra K., O’Neill P. M., George I. M., Reeves J. N., 2007, *MNRAS*, 382, 194  
 Panessa F., Bassani L., Cappi M., Dadina M., Barcons X., Carrera F. J., Ho L. C., Iwasawa K., 2006, *A&A*, 455, 173  
 Parker M. L. et al., 2014, *MNRAS*, 443, 1723  
 Petrucci P. O., Merloni A., Fabian A., Haardt F., Gallo E., 2001a, *MNRAS*, 328, 501  
 Petrucci P. O. et al., 2001b, *ApJ*, 556, 716  
 Reboul H., Vanderriest C., Fringant A. M., Cayrel R., 1987, *A&A*, 177, 337  
 Reis R. C., Miller J. M., 2013, *ApJ*, 769, L7  
 Risaliti G., Elvis M., Fabbiano G., Baldi A., Zezas A., Salvati M., 2007, *ApJ*, 659, L111  
 Risaliti G., Salvati M., Marconi A., 2011, *MNRAS*, 411, 2223  
 Sanfrutos M., Miniutti G., Agís-González B., Fabian A. C., Miller J. M., Panessa F., Zoghbi A., 2013, *MNRAS*, 436, 1588  
 Shapiro S. L., Lightman A. P., Eardley D. M., 1976, *ApJ*, 204, 187  
 Skrutskie M. F. et al., 2006, *AJ*, 131, 1163  
 Strüder L. et al., 2001, *A&A*, 365, L18  
 Titarchuk L., 1994, *ApJ*, 434, 570  
 Turner M. J. L. et al., 2001, *A&A*, 365, L27  
 Wilkins D. R., Fabian A. C., 2011, *MNRAS*, 414, 1269  
 Wilkins D. R., Fabian A. C., 2012, *MNRAS*, 424, 1284  
 Wilkins D. R., Gallo L. C., 2015, *MNRAS*, 449, 129  
 Wilkins D. R., Gallo L. C., Grupe D., Bonson K., Komossa S., Fabian A. C., 2015, *MNRAS*, 454, 4440  
 Winkler C. et al., 2003, *A&A*, 411, L1  
 Wright E. L. et al., 2010, *AJ*, 140, 1868

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.

<sup>2</sup> <http://irsa.ipac.caltech.edu/Missions/2mass.html>

<sup>3</sup> <http://irsa.ipac.caltech.edu/Missions/wise.html>