

## 2MASS STUDIES OF DIFFERENTIAL REDDENING ACROSS THREE MASSIVE GLOBULAR CLUSTERS

DAVID R. LAW,<sup>1</sup> STEVEN R. MAJEWSKI,<sup>1</sup> MICHAEL F. SKRUTSKIE,<sup>1</sup> JOHN M. CARPENTER,<sup>2</sup> AND HINA F. AYUB<sup>1</sup>

Received 2003 May 19; accepted 2003 June 19

### ABSTRACT

*J*-, *H*-, and *K<sub>s</sub>*-band data from the Two Micron All Sky Survey (2MASS) are used to study the effects of differential reddening across the three massive Galactic globular clusters  $\omega$  Centauri, NGC 6388, and NGC 6441. Evidence is found that variable extinction may produce false detections of tidal tails around  $\omega$  Centauri. We also investigate what appears to be relatively strong differential reddening toward NGC 6388 and NGC 6441 and find that differential extinction may be exaggerating the need for a metallicity spread to explain the width of the red giant branches for these two clusters. Finally, we consider the implications of these results for the connection between unusual, multipopulation globular clusters and the cores of dwarf spheroidal galaxies (dSph).

*Key words:* Galaxy: structure — globular clusters: general —  
globular clusters: individual ( $\omega$  Centauri, NGC 6388, NGC 6441)

### 1. INTRODUCTION

The giant Galactic globular cluster  $\omega$  Centauri (NGC 5139, hereafter  $\omega$  Cen) is not only the most massive globular cluster in the Milky Way, but its stars have also been shown to have a large spread in both age and metallicity (see, for instance, many contributions in van Leeuwen, Hughes, & Piotto 2002). In light of these discoveries, it has been proposed (Lee et al. 1999; Majewski et al. 2000)—and now dynamically modeled (Tsujiyama & Shigeo 2003)—that  $\omega$  Cen may be the nucleus of a disrupting dSph, similar to the Galactic globular cluster M54. The second largest globular cluster, M54 is currently believed to be the core of the Sgr dSph (Da Costa & Armandroff 1995; Bassino & Muzzio 1995; Sarajedini & Layden 1995; Layden & Sarajedini 2000) and lies at a nucleation point for the several different metallicity populations of Sgr (Sarajedini & Layden 1995; Majewski et al. 2003). The recent observation of apparent tidal tails around  $\omega$  Cen by Leon, Meylan, & Combes (2000, hereafter LMC00) bolsters the notion of this cluster being the remnant of a larger, now disrupted parent object, and with  $\omega$  Cen and M54 as models, Ree et al. (2002), Lee et al. (1999, 2002), and Yoon et al. (2000) have proposed that other clusters with evidence of multiple populations, such as clusters with bimodal horizontal branches, may also be the nuclei of disrupting dSph satellite galaxies. Two clusters proposed to be M54 analogs (Ree et al. 2002) are NGC 6388 and NGC 6441, the third and fifth largest globular clusters in the Milky Way, respectively, since both have observed characteristics suggesting internal metallicity spreads. Moreover, high mean metallicity clusters like these (see Table 1) should have only a red stub of a horizontal branch (HB), but NGC 6388 and NGC 6441 also have extended blue HB tails (Piotto et al. 1997, hereafter P97). A *bimodal* (or extended) metallicity distribution may explain both this curious HB morphology (P97) as well as the unusually wide red giant branches (RGBs) observed in these clusters.

Because it is difficult to understand how multiple generations of stars might be created in a stellar system of initial mass on the order of  $10^5$ – $10^6 M_{\odot}$  (which is not massive enough to retain significant amounts of gas after an initial starburst), it is possible that the multipopulation clusters are the remnants of *formerly* larger stellar systems, for example, dSph galaxies, where multiepoch star formation histories are commonly found. The verification of the existence of additional multiple population clusters would increase the need to formulate a general process—e.g., the disruption of initially larger systems—that leads to a more substantial *class* of these objects.

These scientific questions are complicated by the presence of nonnegligible differential reddening in the fields of some the most interesting examples of potential multimodal clusters. In this paper we concentrate on two effects of differential reddening relevant to the globular cluster evolution picture outlined above: the potential creation of false signatures of both tidal tails around, and of multiple metallicities within, globular clusters. These issues are explored in case studies of the three massive globular clusters  $\omega$  Cen, NGC 6388, and NGC 6441.

Tidal tail searches around Galactic satellites can be particularly vulnerable to differential reddening effects because search techniques typically rely on the detection of star-count overdensities. Lines of sight with a lower dust column density will naturally reveal more stars to a given magnitude limit. This may create an apparent overdensity of stars in regions with lower extinction that could be mistaken for, or overemphasize, density enhancements from tidal tails. In the case of the reported tidal tails around  $\omega$  Cen by LMC00, the contours of the detected tails curiously and closely correspond to contours of the prominent *IRAS* 100  $\mu$ m dust emission (and therefore, presumably, optical extinction by dust) in this field, as these authors have observed. Because LMC00 did not correct for differential extinction in their optical star counts (S. Leon 2002, private communication), at minimum there must be some dust-shaping of the morphology of mapped  $\omega$  Cen tidal tails. We reevaluate the possible impact of differential reddening on large field studies of  $\omega$  Cen by comparing a customized tidal tail search on raw 2MASS data with one performed on 2MASS data

<sup>1</sup> Department of Astronomy, University of Virginia, P.O. Box 3818, Charlottesville, VA 22903; drlaw@alumni.virginia.edu, srm4n@virginia.edu, skrutskie@virginia.edu, hfa9b@virginia.edu.

<sup>2</sup> Department of Astronomy, California Institute of Technology, MS 105-24, Pasadena, CA 91125; jmc@astro.caltech.edu.

TABLE 1  
CLUSTER PARAMETERS FROM HARRIS (1996)

Cluster	[Fe/H]	Core Radius (arcmin)	Tidal Radius (arcmin)
47 Tucanae.....	-0.76	0.44	47.25
$\omega$ Centauri.....	-1.62	2.58	44.85
NGC 6388.....	-0.60	0.12	6.21
NGC 6441.....	-0.53	0.11	8.00

differentially dereddened according to the dust maps of Schlegel, Finkbeiner, & Davis (1998).

The cause of the observed width of the RGB in NGC 6388 and NGC 6441 has been variously explained as differential reddening and/or the presence of multiple populations of different metallicity, with no clear consensus emerging. P97 originally concluded that differential reddening on scales greater than  $8''$  was unlikely for NGC 6388 but possible for NGC 6441, although only to a few hundredths of a magnitude. Later evidence indicating the presence of strong differential reddening on small scales across both clusters was found by Raimondo et al. (2002) using *Hubble Space Telescope* (*HST*) WFPC2 small area surveys similar to those used by P97. Pritzl et al. (2001, 2002) also found differential reddening in NGC 6441 and NGC 6388 using Cerro Tololo Inter-American Observatory *B*- and *V*-band photometry, although they note that in NGC 6388 the differential reddening probably covers a smaller magnitude range. If differential reddening were significant, the reddening vector could reproduce the sloping red HBs and spread the RGB stars perpendicular to the unreddened RGBs, inflating their apparent width. Numerical experiments performed by Raimondo et al. (2002) indicate that differential reddening on the order of  $\Delta E(B-V) \approx 0.1$  would be substantial enough to cause an unreddened HB of similar metallicity (e.g., that of 47 Tucanae) to resemble a sloping red stub similar to that of NGC 6388 or NGC 6441. Nevertheless, Raimondo et al. (2002) concluded that reddening is not the reason for the strange HBs since their sloping nature is present even in small area subsamples exhibiting a thin RGB, which suggests that a second effect is at work.

Pritzl et al. (2001, 2002) and Ree et al. (2002) argue that variable metallicity may adequately explain the observed features of the NGC 6388 and NGC 6441 color-magnitude diagrams (CMDs), and recent stellar evolutionary models (e.g., Sweigart & Catelan 1998) confirm that a metallicity spread can naturally create a sloping HB. Specifically, Ree et al. (2002) find that two distinct populations separated by

1.2 Gyr and 0.15 dex in [Fe/H] can reproduce both the observed wide RGBs and tilted HBs in the clusters. However, Pritzl et al. (2001, 2002) argue that the simple two-population model fails to reproduce the observed HB slope and that a more general metallicity *spread* is necessary.

Because selective extinction at the 2MASS wavelengths *J* ( $1.25 \mu\text{m}$ ), *H* ( $1.65 \mu\text{m}$ ), and *K<sub>s</sub>* ( $2.17 \mu\text{m}$ , hereafter simply referred to as *K*) is roughly 1/10 that in visual bands, the magnitude of differential reddening at these wavelengths is commensurately reduced. Thus, we apply the 2MASS point-source catalog to test whether (1) differential reddening affects interpretations of an *infrared* search for tidal tails around  $\omega$  Cen and (2) the NGC 6388 and NGC 6441 RGBs are broadened at near-infrared wavelengths, where differential reddening will have less impact on the width of the cluster RGBs.

## 2. DATA

*J*-, *H*-, and *K<sub>s</sub>*-band data were drawn from the 2MASS all-sky point-source working survey database as of 2002 July. Initial values of the mean field extinction  $E(B-V)$  and distance modulus ( $m - M$ ) for each cluster were obtained from the Harris compilation as posted on the World Wide Web on 1999 June 22 (see Harris 1996).<sup>3</sup> These were converted to the required infrared extinction parameters using the standard extinction models of Cardelli, Clayton, & Mathis (1989). Table 2 gives a complete list of the adopted extinction parameters.

## 3. $\omega$ CENTAURI: A 2MASS TALE

### 3.1. Tidal Tail Search Method

The LMC00 study of optical star counts in the  $\omega$  Cen field makes no corrections for differential reddening because magnitude calibrations were not available for the majority of clusters in their study (S. Leon 2002, private communication). As Figures 1*a* and 1*b* illustrate, the proposed tidal tails of  $\omega$  Cen are from enhanced star counts whose pattern appears to follow regions of lower dust absorption on the *IRAS* 100  $\mu\text{m}$  map. This effect is particularly noticeable on the western edge of the cluster. LMC00 note and discuss the possibility that the star counts are affected by reddening but determine that the tidal tails are of a sufficiently large signal-to-noise ratio (S/N) that differential reddening should not significantly affect their tidal tail search.

<sup>3</sup> See <http://physun.physics.mcmaster.ca/Globular.html>.

TABLE 2  
EXTINCTION PARAMETERS

Cluster	$A_V$	$(m - M)_V$	$(m - M)_K$	$E(B-V)$	$E(J-K)$	$E(J-H)$	$E(H-K)$
47 Tucanae.....	0.124	13.37	13.26	0.04	0.021	0.011	0.009
$\omega$ Centauri.....	0.299	13.92	13.63	0.12	0.060	0.033	0.027
NGC 6388.....	1.240	16.54	15.44	0.40	0.209	0.109	0.090
NGC 6441.....	1.395	16.62	15.38	0.45	0.234	0.123	0.102

NOTE.—Values for  $A_V$ ,  $(m - M)_V$ , and  $E(B-V)$  are from Harris 1996, all other parameters were calculated according to Cardelli et al. 1989. Note that differential dereddening in the field of  $\omega$  Cen is performed according to the Schlegel et al. 1998 reddening maps, which give an average color-excess of  $E(B-V) = 0.13$  in a  $4^\circ \times 4^\circ$  field around the cluster.




FIG. 1.—(a) Smoothed-function surface density plot of  $\omega$  Cen showing tidal tails (using logarithmic contours) found by LMC00. The dotted line describes the region within which  $n_T$  is defined (§ 3.1), the arrows indicate the cluster proper motion (*dotted arrow*), the direction of the Galactic center (*dashed arrow*), the direction perpendicular to the Galactic plane (*solid arrow*), and a length of 100 pc at the distance of  $\omega$  Cen (*horizontal double arrow*). (b) IRAS 100  $\mu$ m chart overlaid with LMC00 tidal tail contours, darker shading indicates a greater amount of dust along the line of sight. The total range of color excess is on the order of  $\Delta E(B-V) = 0.18$  mag. (c) Linear contoured surface density plot of  $\omega$  Cen showing apparent tidal tails in the nondereddened 2MASS data. Dashed lines denote all contours beneath the mean background level. (d) Surface density plot of  $\omega$  Cen after differential dereddening; all stages of the analysis are identical to those used to create panel (c). Panels (a) and (b) are reproduced from LMC00 courtesy of S. Leon and the European Southern Observatory.

By comparing star-count patterns obtained using raw 2MASS data with those from 2MASS data differentially dereddened according to the dust maps of Schlegel et al. (1998), we intend to reevaluate whether uncorrected differential extinction can produce a false tidal tail signal. The 2MASS data have a 99% completeness limit at  $J = 16$  mag in the direction of  $\omega$  Cen, yielding almost 500,000 stars within a  $6^\circ \times 6^\circ$  box centered on the cluster. Because a photometric search for diffuse tidal tails presents considerable challenges, not the least of which is substantial field star contamination in this low-latitude field, a two-stage method is employed to minimize field star contamination and probe for star-count excesses as deeply into the surrounding field as possible: First, a coarse filtering mask in color-color-magnitude (CCM) space is used to extract only those stars whose CCM location is similar to that of stars observed at radii less than the cluster core radius. Although this coarse mask does not take into account foreground/background contamination of the cluster core, it filters out a significant number of stars whose CCM locations are significantly different from those in the cluster. Second, we then apply a signal-to-noise filter similar to that used by

Odenkirchen et al. (2001) to separate cluster stars from field stars and emphasize star count overdensities in the potential tidal tails.

In the first stage of filtering, a filtering mask is applied in CCM space to preselect those CCM locations containing stars within  $100''$  of the center of the cluster (approximately  $\frac{2}{3}$  the core radius  $r_c$ ; see Table 1). We divide this irregularly shaped region in  $(J, J-H), (J-K)$  space into a  $50 \times 50 \times 50$  grid in CCM space bounded by the limits of the CCM region; this grid size is selected to achieve a balance between coarse binning and low number statistics. With the individual cells described by indices  $(i, j, k)$  respectively, we define  $n_C(i, j, k)$  as the number of stars within  $100''$  of the cluster center that fall within each grid cell in CCM space. To obtain an estimate of the number of field stars included in  $n_C(i, j, k)$ , we represent the field star distribution  $n_F(i, j, k)$  by the stars within an annulus  $100.0 \leq r \leq 150.0$  (roughly 3 times the tidal radius of the cluster). We assume that the distribution of field stars is relatively even across the cluster and scale the field mask by the ratio  $q$  of the areas over which the field and core regions were summed. The selection mask  $P$  is defined as the ratio of the core mask to the sum of

the core and scaled field masks,

$$P(i, j, k) = \frac{n_C(i, j, k)}{n_C(i, j, k) + q^{-1}n_F(i, j, k)}. \quad (1)$$

Thus,  $P(i, j, k)$  will have a value near 1 for regions of CCM space in which the core sample  $n_C(i, j, k)$  is not substantially contaminated with field stars and will decrease as contamination becomes more severe. The complete data set is filtered using this CCM selection mask: a star is rejected from the data set if the corresponding  $P(i, j, k)$  is less than some minimum value  $P_{\min}$ . This algorithm is a slightly more sophisticated means of eliminating stars unrelated to  $\omega$  Cen than simple preliminary cuts in the color-magnitude (CM) plane.

We find that a value of  $P_{\min} = 0.9$  allows us to optimally select cluster stars over field stars and appears to best preserve the CMD of the cluster center. This preliminary filtering rejects roughly 90% of the stars in the original data set.

In the second stage of filtering, we apply a modified version of Grillmair et al.'s (1995) method in the CM plane as described by Odenkirchen et al. (2001). This second stage eliminates field stars more effectively than the first stage, but we find that the method does not perform well unless a coarser preselection of stars is performed first to reduce the data set to a size that the more computationally complex second-stage routine can handle. With some recycling of notation, we define a grid in  $(J, J-K)$  space, with indices  $(j, k)$ , respectively, and find the S/N of probable cluster stars to field stars in each CM grid cell. We count the number of stars  $n_C(i, j, k)$  that fall within each grid cell and lie within  $r = 20'$  of the cluster center (chosen to correspond to the region in which LMC00 observe circular cluster surface density contours; see Fig. 1a) and the number of stars  $n_F(j, k)$  that fall within the CM grid cell and lie between  $r = 100'$  and  $r = 150'$  from the cluster center. The field star population is assumed to be uniform, and the expected number of field stars of a given  $(j, k)$  that fall within  $r = 20'$  of the cluster core is subtracted from  $n_C(j, k)$ . With  $q$  representing the ratio of the area of sky on which  $n_F$  and  $n_C$ , respectively, have been defined, the local S/N of cluster to field stars for each grid cell  $(j, k)$  is, by standard Poissonian statistics,

$$s(j, k) = \frac{n_C(j, k) - q^{-1}n_F(j, k)}{\sqrt{n_C(j, k) + q^{-2}n_F(j, k)}}. \quad (2)$$

It is possible to use  $s(j, k)$  as a mask for selection in the CM plane by eliminating all  $(j, k)$  grid cells with  $s(j, k)$  less than some limiting value  $s_{\lim}$ , which is chosen such that the number of stars in the tidal tail region (as reported by LMC00) is a maximum relative to the field star population. The new matrix  $n_T(j, k)$  represents the total number of stars in each  $(j, k)$  grid cell for the region<sup>4</sup> of the cluster in which tidal tails have been reported by LMC00 (see Fig. 1a, *dashed lines*). Starting from  $s_{\lim} = s_{\max}$  [where  $s_{\max}$  is the maximum value of all  $s(j, k)$ ] and iterating downward to  $s_{\lim} = 0$ , we calculate  $N_{\text{tail}} = \sum_{j,k} n_T(j, k)$  and  $N_{\text{field}} = \sum_{j,k} n_F(j, k)$  for all  $(j, k)$  such that  $s(j, k) \geq s_{\lim}$ . Hence,  $N_{\text{field}}$  and  $N_{\text{tail}}$ , respectively are the total number of field stars and cluster stars in LMC00's tidal tail regions whose locations in

$(J, J-K)$  space satisfy  $s(5 > j, k) \geq s_{\lim}$ . The optimal value of  $s_{\lim}$  is that which maximizes  $N_{\text{tail}}/N_{\text{field}}$ ; hence this final selection mask filters the data set for all stars that fall within cells in CM space such that  $s(J, J-K) \geq s_{\lim}$ .

The positions of stars from this filtered set are then converted to a smooth density distribution by application of a two-dimensional Gaussian kernel with an adaptive half-maximum width (e.g., Silverman 1986) for each star set to the distance of the 30th nearest neighbor in the plane of the sky. Each of these Gaussians is evaluated at finely spaced grid points in the plane of the sky, and these values are summed to create an image of relative stellar densities at each location on the grid (Figs. 1c and 1d).<sup>5</sup>

We find that this S/N optimization method worked efficiently to reduce the 2MASS data set by a further 80%–90% and brought out star-count overdensities around the cluster that have the appearance of tidal tails. Focusing the definition of  $n_T$  on the regions found by LMC00 to possess tidal tails is considerably more effective at bringing these apparent tails out of the background than defining  $n_T$  more generally; however, this particular approach obviously is not useful for a general survey of previously unknown tidal tail searches because it requires a priori knowledge of where tails may be found. We verified that the overdensities found by this method are not merely artifacts of the definition of  $n_T$  in the region of the LMC00-expected tails by redefining  $n_T$  in locations rotated around the cluster core by  $90^\circ$  (that is, protruding east-west as opposed to north-south). Using this definition of  $n_T$ , no star-count overdensities were detected around the cluster, which supports the idea that the apparent tails found with the first  $n_T$  definition had not been an artificial product of the processing method and that the method of sampling  $n_T$  in the tail region is a crucial step in bringing those tails out of the background noise.

### 3.2. Results

In the analysis of the raw 2MASS data described above, star-count excesses to the north and south of the cluster are found. As Figure 1c shows, to within the limits imposed by the smoothing kernel, the locations of the 2MASS tails correspond to that of the tails found by LMC00.

The same analysis has been repeated with 2MASS data dereddened according to the extinction maps of Schlegel et al. (1998).<sup>6</sup> The Schlegel et al. maps have a resolution of approximately 6'1, which corresponds to about 1500 cells of reddening data over a  $4^\circ \times 4^\circ$  field of view, and show differential reddening on the order of  $\Delta E(B-V) = 0.18$  mag over that field of view. In contrast to what is found for the raw 2MASS data, the results of the analysis on the dereddened 2MASS data do not reveal any elongation of the circular cluster contour lines or any overdensities resembling tidal tails (Fig. 1d). Figure 2 plots radial surface density profiles for both original and dereddened 2MASS data, and shows that the surface density of the dereddened cluster falls off smoothly with radius, like a King (1966) profile. In contrast, the surface density of stars in the nondereddened cluster data appears to drop at  $\log(r) = 1.8$  ( $r \approx 60'$ )—corresponding to the low star count regions at this radius to the east

<sup>4</sup> We describe this region as the two sections of an annulus (of inner radius  $50'$  and outer radius  $100'$ ) located directly north and south of the circle described by the inner radius.

<sup>5</sup> The reduced field of view of these figures is chosen to correspond to that of LMC00.

<sup>6</sup> We have used the programs available on the World Wide Web at <http://www.astro.princeton.edu/~schlegel/dust/index.html>.




FIG. 2.—(a) Radial surface density profile of  $\omega$  Centauri, error bars are derived from Poisson statistics. Data are shifted vertically to match the LMC00 profile. (b) Radial profile reproduced from LMC00, figure courtesy of S. Leon and the European Southern Observatory. A power law is fitted to the data in the external parts, while the inner surface density profile comes from the data by Trager, King, & Djorgovski (1995). The vertical arrow indicates the cluster tidal radius.

and west of the cluster—and rise again at  $\log(r) = 2.0$  ( $r = 100'$ )—which corresponds to high star count regions to the north and south of the cluster (Fig. 1c).

Given the fidelity with which the apparent infrared and optical tails follow the regions of lower reddening on the *IRAS* 100  $\mu\text{m}$  map (Fig. 1b, particularly on the western and southern edges of the cluster), and the disappearance of the infrared tails in the dereddened cluster data, we speculate that by analogy the  $\omega$  Cen tidal tails previously detected with the optical star counts, which are more affected by dust extinction, could have been artifacts of differential reddening. We must also point out, however, that our test of the

2MASS database is not completely analogous to the LMC00 analysis: because of the  $J = 16$  mag completeness limit of 2MASS we are unable to probe to the same effective depth as the LMC00 study. The differences in relative depth are demonstrated by the fact that while LMC00 observe reasonably circular contours out to a radius of about  $45'$  (roughly the tidal radius, see Table 1), we find circular contours only out to about  $40'$  before we begin to observe distinct elongation of the contours into potential tidal tails (Fig. 1). Thus, the LMC00 counts reach fainter surface brightness limits than does 2MASS and are sensitive to more diffuse, lower density tails than are we. Nevertheless, we regard the 2MASS test presented here as a cautionary example demonstrating that a proper accounting of differential reddening is an important step in mapping any tidal tails in the  $\omega$  Cen (or any other heavily obscured) field.

#### 4. DIFFERENTIAL REDDENING IN THE FIELDS OF NGC 6388 AND NGC 6441

##### 4.1. Processing

To investigate the effects of differential reddening on the width of the RGB in the clusters NGC 6388 and NGC 6441, we use the uniform photometry of 2MASS to look for spatial patterns in the distribution of red giant star colors within each cluster. 47 Tucanae (NGC 104, hereafter 47 Tuc) is adopted as a control case for a cluster of similar mean metallicity but with little differential reddening [average  $E(B-V) \approx 0.02$ ]. We also compare our results with the  $\omega$  Cen data—although this cluster has an overall  $[\text{Fe}/\text{H}]$  much lower than the other clusters, it nonetheless provides a useful example of the effects of *variable metallicity* within a cluster over spatial scales with relatively little differential reddening on the scale of arcminutes [ $\Delta E(B-V) = 0.01$  mag, note that the previous section explored extinction variations in  $\omega$  Cen on the scale of *degrees*].

##### 4.1.1. $J-K$ Color Distributions

Analysis of RGB color *spreads* at different magnitudes in one cluster as well as differences between clusters is simplified if we remove the mean RGB color-magnitude relation for each cluster (i.e., make the center of the RGB locus for each cluster appear vertical in the color-magnitude diagram). Since the  $[\text{Fe}/\text{H}]$  of the unreddened cluster 47 Tuc is similar to that of NGC 6388 and NGC 6441 to within about 0.25 dex (see Table 1), the shape, height, and position of the RGBs for the three clusters in a reddening-corrected, absolute color-magnitude space should be similar. Hence, we use the well-defined 2MASS CMD of 47 Tuc to “straighten” the RGB of the clusters in our sample (Fig. 3a). We convert apparent to absolute magnitudes using the distance moduli in Table 2 and consider only that part of the RGB with absolute magnitude  $M_K \leq -2$  to minimize contamination from cluster HB as well as field stars. A Legendre polynomial is fitted to the 47 Tuc RGB, with a rejection threshold of  $2\sigma$  iterated until convergence. The final polynomial is converted to power series coefficients up to third order, which are used to straighten the RGBs of all four clusters. We define an adjusted  $J-K$  color given by

$$(J-K)_A = J-K - E(J-K) - (a + bM_K + cM_K^2), \quad (3)$$

where  $E(J-K)$  values are given in Table 2, and the




FIG. 3.—RGB of 47 Tuc: (a) before and (b) after removing the mean color-magnitude relation

coefficients  $a$ ,  $b$ , and  $c$  are power series coefficients given in Table 3. Similar equations are derived for the  $(J-H, K)$  and  $(H-K, K)$  CMDs.

Using the adjusted values, histograms of colors readily show that the RGBs of NGC 6388 and NGC 6441 (Fig. 4) are wider than those of 47 Tuc and even that of  $\omega$  Cen, which is well established to have a metal spread from  $[\text{Fe}/\text{H}] \approx -2.2$  to  $[\text{Fe}/\text{H}] \approx -0.7$  (Vanture, Wallerstein, & Suntzeff 2002). The latter result is somewhat surprising, given the fact that the metal-rich RGB of 47 Tuc provides a poor approximation of the metal-poor RGB of  $\omega$  Cen and produces a clear tilt in the “adjusted”  $\omega$  Cen RGB (Fig. 4). The Figure 4 histograms are divided into five  $(J-K)_A$  color bins in order to study the spatial distribution of RGB stars by color. The five color bins from the lowest value of  $(J-K)_A$  (most blue) to the highest (most red) are referred to as bins A–E, respectively. Because globular clusters are dynamically mixed on timescales that are short compared to chemical enrichment, there should be no significant difference in the spatial distribution of the stars in these five bins. For example, we show below (§ 4.2.1) that neither 47 Tuc nor  $\omega$  Cen exhibit any such spatial variation. On the other hand, if differential extinction were significant, the distributions of stars by color would be uneven across the face of the cluster as a reflection of reddening variations. This is a test we now apply to NGC 6388 and NGC 6441.

TABLE 3  
RGB STRAIGHTENING COEFFICIENTS

Color	$a$	$b$	$c$
$J-K$ .....	0.5436	-0.0361	0.0105
$J-H$ .....	0.5388	0.0066	0.0120
$H-K$ .....	0.0513	-0.0153	0.0021

NOTE.—Here  $a$ ,  $b$ , and  $c$  are the coefficients of the zeroth-, first-, and second-order terms, respectively.

Various methods of dividing the sample population into color bins were explored. The results are not very sensitive to which method is chosen. The following technique ensures similar treatment of each cluster: bin ranges are assigned to each cluster using a  $(J-K)_A$  histogram of a subsample of the purest cluster stars, defined by an angular radius at which the density of stars decreased to a limiting density given by

$$\text{limiting density} = (1/e^3)(\text{central density} - \text{outer density}), \quad (4)$$

(where the outer density is determined as the density of stars at the edge of the field of view). In all cases, this resulted in a sample of stars within a radius of between roughly  $1'$  and  $3'$ , which is outside the core radius but well within the tidal radius of each cluster (see Table 1). Using a color histogram with binning width 0.02 mag, the central three bins are defined within the  $(J-K)_A$  range from the central peak to the colors at which counts decrease to  $1/e$  of the central peak. Bin C is centered at the peak of the histogram and has  $\frac{1}{3}$  of the width of  $B + C + D$ . The range of bin A is determined as the color range blueward of bin B up to where the counts first drop to 1 (to eliminate extreme colored stars from the sample). Likewise, bin E is defined as the range redward of bin D up to where the counts first drop to 1 (Fig. 5). Field star contamination (or contamination by non-RGB stars in the cluster), not differential reddening, is assumed to be the reason for stars that lie very far from the fiducial RGB, and the described culling of the data set promises to retain only those stars that make up the bulk of the RGB and that could potentially have been scattered to their current locations in the CMD by differential reddening. These color bin definitions are applied to the full  $300''$  data set. Similar binnings were created for  $(J-H)_A$  and  $(H-K)_A$  histograms, although only the  $(J-K)_A$  and  $(J-H)_A$  distributions were found to be useful in our analysis (see below).




FIG. 4.—Histograms of the RGB for all four clusters. The field of view is  $300''$ . Note the width of NGC 6388 and NGC 6441. The  $\omega$  Cen RGB is canted to the left because the color-magnitude relation used to straighten the RGBs was derived from the much more metal-rich 47 Tuc. Histograms in  $(J-H)_A$  are similar in character to the  $(J-K)_A$  distributions shown here.

#### 4.1.2. $J-H$ and $H-K$ Colors

Before proceeding further, we apply one additional filter designed to preserve reddening effects and reduce noise from astrophysical and photometric scatter. This filter takes advantage of the fact that the degree of scatter due to red-

dening in one color should be correlated to that in another color. A Legendre polynomial with an iterative  $2\sigma$  rejection threshold (similar to that used earlier in § 4.1.1) was fitted to the cluster  $(J-K)_A$  versus  $(J-H)_A$  color-color diagram (Fig. 6), and color-color cuts were performed to eliminate stars at greater than  $2\sigma$  from the Legendre fit. This




FIG. 5.—Histogram of the RGB of NGC 6388. The selection criteria for the color bins illustrated are discussed in § 4.1.1.

independently derived fit for each cluster is very close to the fiducial reddening vector for standard extinction models with  $R_V = 3.1$ , lending further evidence to support the presence of differential reddening across each cluster. Figures 6 and 7 show color-color plots of  $(J-K)_A$  versus  $(J-H)_A$  and  $(H-K)_A$  color, respectively, with the adopted outlier cuts shown. (See Table 4 for color-color elimination parameters.)

Note that the dynamic range in color of  $(H-K)_A$  is small compared to the reddening effect sought, and it is difficult to determine which stars appear redder than average as a result of differential reddening instead of photometric errors or




FIG. 6.—Color-color diagram of the RGB of NGC 6388. Shown are the boundaries between  $(J-K)_A$  color bins (short-dashed lines) and the location of  $(J-K)$ ,  $(J-H)$  color-color cuts (long-dashed lines), as described in § 4.1.1. The fiducial reddening vector is given for comparison.




FIG. 7.—Same as Fig. 6, but for NGC 6388

astrophysical reasons. We therefore reject  $(H-K)_A$  color from our analysis, since it does not have the discriminating power required to pick out the required distribution from the noise of points in Figure 7.

4.2. Results and Discussion

4.2.1. Five Bin Results

Spatial plots of the stars in each of the  $(J-K)_A$  bins for 47 Tuc,  $\omega$  Cen, NGC 6388, and NGC 6441 are shown in Figures 8–11, respectively. As expected for a monometallic population with little reddening, 47 Tuc shows no spatial segregation of the stars in the five bins. Although giant stars in  $\omega$  Cen have been found to exhibit a spatial variation in metallicity (Jurcsik 1998), this variation is on scales of roughly  $20'$ ; in contrast, the mixed metallicity population in our smaller field of view shows no spatial segregation by RGB color. Both 47 Tuc and  $\omega$  Cen have few stars in bins A and E relative to their total population. In contrast, NGC 6388 exhibits a pronounced abundance of redder stars (bins D and E) in the northern and northwestern regions of the cluster compared to southern regions (Fig. 10), and the extreme bins A and E each contain significantly more stars relative to the total population of the cluster than these bins contained for 47 Tuc and  $\omega$  Cen. Comparison with the 47 Tuc and  $\omega$  Cen templates for monometallic and multimetallic clusters (respectively) having comparatively little differential reddening over small scales leads one to the

TABLE 4  
COLOR-COLOR ELIMINATION PARAMETERS

Cluster	$a$	$b$
47 Tucanae.....	0.012	0.779
$\omega$ Centauri.....	-0.019	0.632
NGC 6388.....	0.044	0.766
NGC 6441.....	0.047	0.755

NOTE.—The variables  $a$  and  $b$  are the coefficients of the zeroth- and first-order polynomial terms, respectively.  $(J-K)_A$  is the abscissa, and  $(J-H)_A$  is the ordinate.




Fig. 8.—Spatial distribution of stars within each bin for the cluster 47 Tucanae. Bin A:  $-0.201 \leq (J-K)_A < -0.082$ ; bin B:  $-0.082 \leq (J-K)_A < -0.031$ ; bin C:  $-0.031 \leq (J-K)_A < 0.011$ ; bin D:  $0.011 \leq (J-K)_A < 0.042$ ; bin E:  $0.042 \leq (J-K)_A < 0.098$ ; bin F: all previous bins. The  $(J-H)_A$  sky distribution is similar. The dashed circle denotes our field of view.

conclusion that it is unlikely that the NGC 6388 color spread is due primarily to metallicity effects. Rather, Figure 10 suggests that differential reddening on small scales is causing (1) a greater number of stars to appear in bins A and E and (2) large spatial variations by observed stellar color.

Likewise, NGC 6441 (Fig. 11) shows an enhancement of redder stars (bins D and E) in the southeastern quadrant of the field of view, supporting the view that it too is subject to

substantial differential reddening. Using isochrone fitting in subfields, Heitsch & Richtler (1999) similarly find that the southeast region of NGC 6441 has higher  $E(B-V)$  than the northwestern region by about 0.15 mag. Layden et al. (1999) concur with Heitsch & Richtler (1999) to within a few hundredths of a magnitude using RR Lyrae stars and CMDs in sectors of an annulus around the cluster center. However, in contrast to the distinct dichotomy in NGC 6388 (redder stars in the northwest, bluer stars in the




FIG. 9.—Spatial distribution of stars within each bin for the cluster  $\omega$  Centauri. Bin A:  $-0.22 \leq (J-K)_A < -0.183$ ; bin B:  $-0.183 \leq (J-K)_A < -0.129$ ; bin C:  $-0.129 \leq (J-K)_A < -0.075$ ; bin D:  $-0.075 \leq (J-K)_A < -0.021$ ; bin E:  $-0.021 \leq (J-K)_A < 0.138$ ; bin F: all previous bins.

southeast), the spatial distribution of stars in bins A–C in NGC 6441 appears relatively constant (with the possible exception of decreased counts in the southeastern quadrant of bin A), and the cluster presents the appearance of an inhomogeneously distributed redder population superposed on a centrally concentrated bluer population background. This effect may be an artifact of the severe contamination of the cluster by field stars since NGC 6441 is located in an extremely crowded field near the Galactic center (as vividly demonstrated by the washed out star-count map shown in

Fig. 12). Indeed, Heitsch & Richtler (1999) note the strong contamination of their own CMD of NGC 6441 by field stars that cover the lower part of the cluster RGB. In this case, strong contamination by blue field stars may contribute to the apparent width of the cluster RGB.

#### 4.2.2. Reddening Maps

Figure 12 shows 2MASS star-count density (Fig. 12a) and mean stellar color (Fig. 12b) maps for all four of our




Fig. 10.—Spatial distribution of stars within each bin for the cluster NGC 6388. Bin A:  $-0.341 \leq (J-K)_A < -0.101$ ; bin B:  $-0.101 \leq (J-K)_A < -0.03$ ; bin C:  $-0.03 \leq (J-K)_A < 0.03$ ; bin D:  $0.03 \leq (J-K)_A < 0.08$ ; bin E:  $0.08 \leq (J-K)_A < 0.179$ ; bin F: all previous bins.

clusters. In this figure, higher star counts and redder mean colors appear whiter (Table 5 provides the numerical values for each gray scale). The field of view of these maps is more than 100 times larger than that used for the analyses in the rest of this section since this method only provides coarse resolution maps which, however, are useful for understanding qualitatively the global character and degree of patchiness of the dust obscuration around each cluster. For example, assuming that the average intrinsic star color in a

given direction is relatively uniform over degree scales, the mean color maps can be interpreted as showing the general character of the reddening in the direction of each cluster. However, these maps have too coarse a gray scale to show color variations on the order of  $E(B-V) \leq 0.1$ .

Figure 12 shows that the mean stellar color in the direction of NGC 6388 is not only patchier than that in the direction of 47 Tuc or  $\omega$  Cen but that the number of redder stars appears to be greater to the north of the cluster than to the




FIG. 11.—Spatial distribution of stars within each bin for the cluster NGC 6441. Bin A:  $-0.422 \leq (J-K)_A < -0.16$ ; bin B:  $-0.16 \leq (J-K)_A < -0.047$ ; bin C:  $-0.047 \leq (J-K)_A < 0.047$ ; bin D:  $0.047 \leq (J-K)_A < 0.12$ ; bin E:  $0.12 \leq (J-K)_A < 0.198$ ; bin F: all previous bins.  $(J-H)_A$  distribution is similar. Stars are absent from the 2MASS catalog in a small region at the west of the field of view due to contamination by the bright star HD 161892.

south,<sup>7</sup> in agreement with the trend found above. Distinct regions of higher reddening and lower reddening are demarcated by a boundary that passes close to the center of the cluster. Differential reddening across this apparent dust boundary may produce the observed overabundance of

<sup>7</sup>Note that this remark applies to the central few pixels immediately around the cluster.

redder RGB stars in the north-northwestern regions of the cluster compared to the south-southeastern regions.

The NGC 6441 field mean color map also shows evidence of patchy extinction, although it is not obvious whether any of these patches could produce the observed reddening patterns near the cluster that are likely the case for NGC 6388. However, the presence of large-scale patchiness is sufficient to suggest the possibility of patchiness on the angular scales explored in § 4.2.1.




FIG. 12.—The 2MASS maps of all stars (i.e., both cluster and field stars) within a  $1^\circ$  radius of each cluster. In each case, panel (a) gives the star count density and panel (b) the mean stellar color. Higher star counts and redder colors are whiter (see Table 5 for scales). Vertical stripes in panels (a) are due to nightly variations in the sensitivity of the 2MASS survey. Note that the fields of view in this figure are on the order of square degrees, while in Figs. 8–11 the fields of view are on the order of square arcminutes. The larger pixel size for 47 Tuc was required to obtain a reasonable S/N because of low star counts in this region of high Galactic latitude.

TABLE 5  
STAR COUNT AND REDDENING PARAMETERS

Cluster	Minimum Counts	Maximum Counts	Minimum $J-K$	Maximum $J-K$
47 Tucanae.....	...	...	...	...
$\omega$ Centauri.....	0	5	0.4	1.0
NGC 6388.....	0	15	0.4	1.0
NGC 6441.....	0	30	0.4	1.2

NOTE.—Parameters define the scales used in Fig. 12.

4.2.3. Kernel-smoothed Color Distributions

Histograms of star colors within regions of greater reddening should be offset from those of subregions with lesser reddening. To study the color distributions in subsections of the cluster fields, which will have larger statistical noise, we apply a uniform-width-normalized Gaussian kernel (e.g., Silverman 1986) to the  $(J-K)_A$  color distribution of stars in NGC 6388 and NGC 6441. Figure 13 shows these smoothed distributions for the northwest and southeast quadrants of each cluster. The half-maximum width of the kernel is the minimum value that produces a clearly defined central peak in the distribution for a given cluster. NGC 6388 required this smoothing width to be only 0.02 mag, while NGC 6441 required a wider Gaussian of width 0.06 mag to adequately smooth out secondary peaks.

The central peak of the distribution of the northwestern quadrant of NGC 6388 is offset from that of the southeastern quadrant by roughly  $\Delta(J-K)_A = 0.061$  mag, and the peaks of the northwestern and southeastern quadrants of NGC 6441 are offset by roughly  $\Delta(J-K) = 0.072$  mag. Because the width of the kernel used for NGC 6441 is on the order of the offset between the peaks, the precision of the derived  $\Delta(J-K)_A$  is less for NGC 6441 than for NGC 6388. Nevertheless, the  $\Delta(J-K)_A$  results for both clusters are in agreement with the qualitative arguments for differential reddening presented earlier in §§ 4.2.1 and 4.2.2, and our value for the differential reddening across NGC 6441 is consistent with the more detailed optical extinction maps produced by Layden et al. (1999) and Heitsch & Richtler (1999) to within 0.02 mag in  $E(J-K)$ . If such an offset exists between the color distribution of stars in different regions of each cluster, then the artificial width produced by the superposition of these offset distributions may contribute to the thicker RGB of the entire cluster. Figure 14 simulates the effect of superposing two offset color distributions by artificially reddening the northern half of the monometallic cluster 47 Tuc by 0.061 mag in  $(J-K)_A$ ; it is observed that while

TABLE 6  
K-S STATISTICS FOR 47 TUCANAE

Quadrant	NE	NW	SE	SW
NE.....	...	0.599	0.228	0.256
NW.....	0.599	...	0.539	0.116
SE.....	0.228	0.539	...	0.629
SW.....	0.256	0.116	0.629	...

NOTE.—The K-S statistics are shown for correlation significance between the two given quadrants (probability that the two samples are drawn from the same parent distribution).

TABLE 7  
K-S STATISTICS FOR  $\omega$  CENTAURI

Quadrant	NE	NW	SE	SW
NE.....	...	0.583	0.034	0.590
NW.....	0.583	...	0.234	0.997
SE.....	0.034	0.234	...	0.260
SW.....	0.590	0.997	0.260	...

NOTE.—See Table 6 for description.

such a superposition can artificially inflate the width of the color distribution (Fig. 13, bottom), it is only by about half the amount required to recreate the width of the distributions observed in NGC 6388 and NGC 6441. The inability of this simulation to entirely explain the RGB widths of these clusters may be due to the presence of a large reddening spread (as opposed to the simplistic two-value model used here), or it may be indicative of a contributing effect of differential metallicity.

4.2.4. Kolmogorov-Smirnov Statistics

To test the significance of the differences in  $(J-K)_A$  color distribution in different quadrants of each cluster, we applied the Kolmogorov-Smirnov (K-S) test to compare spatial color distributions numerically. As a calibration of the method, we first compared the color distribution across the clusters 47 Tuc and  $\omega$  Cen. As Tables 6 and 7 show, the  $(J-K)_A$  color distribution of stars in, for example, the northwestern quadrant of the clusters correlates with the distribution of stars in the southeastern quadrant with significance levels of about 0.5 and 0.25 for 47 Tuc and  $\omega$  Cen, respectively. These significance levels are consistent with a probability that stars within the two regions could have been randomly drawn from one parent distribution. Tables 6 and 7 demonstrate that this relation significance is roughly similar for comparisons between all quadrants for both 47 Tuc and  $\omega$  Cen.

Correlations between the southeastern and southwestern, and northeastern and northwestern quadrants of NGC 6388 have significance levels of roughly 0.9 and 0.5, respectively. In contrast, the K-S test yields a probability of only 0.002 that the northwestern and southeastern quadrants share a similar color distribution (Table 8). Compare this to the variable metallicity cluster  $\omega$  Cen, which maintains a relatively high correlation significance of color distribution between all quadrants in spite of its metallicity spread. It appears unlikely that a metallicity spread is the dominant cause of the RGB width in NGC 6388 (we expect mixing in clusters in any case). Instead, the stars in these regions almost certainly come from statistically distinct color

TABLE 8  
K-S STATISTICS FOR NGC 6388

Quadrant	NE	NW	SE	SW
NE.....	...	0.533	$3 \times 10^{-6}$	0.005
NW.....	0.533	...	0.0001	0.006
SE.....	$3 \times 10^{-6}$	0.0001	...	0.471
SW.....	0.005	0.006	0.471	...

NOTE.—See Table 6 for description.




FIG. 13.—The  $(J-K)_A$  distributions for different regions of NGC 6388 and 6441, smoothed by normalized Gaussian kernels of half-maximum widths 0.02 and 0.06, respectively. The vertical scale for each panel is defined such that the integrated area under the curve is equal to the number of stars in the given quadrant. Note the offset between histograms in the northwest and southeast quadrants for both clusters.

populations, pointing again to the presence of strong differential reddening.

Application of the K-S test to the  $(J-K)_A$  color distribution in the southeastern and northwestern quadrants of NGC 6441 yields a near zero likelihood of correlation (Table 9). Qualitatively, this is not surprising because there are very few stars from bins D and E in the northwestern quadrant, while there is a considerably higher number of redder stars that lie in the southeastern quadrant. However, while these two quadrants are by orders of magnitude the least-correlated regions of this globular cluster, correlations between all other combinations of quadrants are only on

TABLE 9  
K-S STATISTICS FOR NGC 6441

Quadrant	NE	NW	SE	SW
NE.....	...	$4 \times 10^{-6}$	0.143	0.097
NW.....	$4 \times 10^{-6}$	...	$9 \times 10^{-7}$	0.001
SE.....	0.143	$9 \times 10^{-7}$	...	0.001
SW.....	0.097	0.001	0.001	...

NOTE.—See Table 6 for description.




FIG. 14.—The  $(J-K)_A$  distributions for 47 Tuc (solid line) and for 47 Tuc with  $\Delta(J-K)_A = 0.061$  mag of artificial differential reddening (dashed line). The Gaussian smoothing kernel for each curve has a half-maximum width of 0.02.

the order of 0.01–0.1, which suggests (as we have concluded in § 4.2.2) that reddening in NGC 6441 is patchy on smaller angular scales than in NGC 6388.

If differential reddening is producing the observed width in these clusters, the RGB histograms in subregions of each cluster should also be narrower than that of each cluster as a whole. Using *HST* WFPC2 visual-band data for NGC 6388 and NGC 6441, Raimondo et al. (2002) find this to be the case for both clusters. In summary, it appears very probable that differential reddening is artificially inflating the apparent thickness of the RGB in these two clusters, although the thickness may not be fully explained by this effect alone.

## 5. DISCUSSION AND CONCLUSIONS

Differential extinction complicates photometric studies of Galactic stellar populations, and even nearby globular clusters with  $E(B-V)$  as low as about 0.1 mag may be visibly affected. In our search for tidal tails around the giant globular cluster  $\omega$  Cen we have found that differentially dereddening 2MASS photometry largely eliminates apparent tidal tail candidates present in data that have not been corrected for extinction. Given the similarity of the false 2MASS tidal tail signature to the optical tails reported by LMC00, it would be beneficial to verify the diffuse optical star count distribution in the  $\omega$  Cen field with deep visual photometry that accounts for the  $\Delta E(B-V)$  variations.

We have also investigated the effects of differential reddening on the CMDs of the clusters NGC 6388 and NGC 6441, which exhibit abnormally large spreads in their RGBs. The analyses presented make use of spatial maps by stars in color bins, maps of mean stellar color, kernel-smoothed histograms of stellar color in subregions, and K-S statistical analyses of the distribution of  $(J-K)$  and  $(J-H)$  colors across the clusters. These tests indicate that the northwestern quadrant of NGC 6388 is subject to about  $\Delta E(B-V) \approx 0.12$  mag more reddening than the southeastern quadrant, and that the southeastern quadrant of NGC 6441 is more heavily reddened by about  $\Delta E(B-V) \approx 0.14$  mag than the northwestern quadrant. These findings are consistent with previous results from Heitsch & Richtler (1999) and Layden et al. (1999). As numerical experiments performed here and by Raimondo et al. (2002) have shown, this amount of differential reddening could produce anomalously wider red giant branches for these clusters (in addition to the tilt observed in their horizontal branches, as demonstrated by Raimondo et al. (2002)). Raimondo et al. (2002) have shown that the RGB in subregions of these clusters is thin (as expected if differential reddening were the cause of the width of the RGB across the entire cluster). Although we find that possible width inflation due to the superposition of thin RGBs from two *quadrants* subject to different amounts of differential extinction may not be sufficient to completely explain the observed width of the cluster RGBs, it may be possible that the RGB width could be explained more fully by a more realistic superposition of RGBs from smaller subregions. Therefore, if not accounting for the entire RGB-widening effect, at minimum differential reddening may be exaggerating metallicity-induced RGB spreads and the actual spread in age or metallicity within the clusters is likely not as large as has previously been suggested (e.g., Pritzl et al. 2001, 2002).

The number of known globular clusters showing clearly anomalous metallicity spreads remains small. With few bona fide members, this “class” of globular cluster can still comfortably be accommodated by origin pictures that are exceptional, e.g., along the lines of the M54-Sgr paradigm. However, even in the obvious case of the multipopulation  $\omega$  Cen, the M54 analogy may be less compelling than previously thought, and the picture that some globular clusters are the “residue” of dSph disruption more ambiguous.

The authors would like to thank S. Leon and G. Meylan for helpful discussion regarding the effects of reddening on  $\omega$  Centauri. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center, funded by the National Aeronautics and Space Administration and the National Science Foundation. This work has been undertaken with the support of NASA/JPL contract 1228235 as well as a David and Lucille Packard Foundation Fellowship.

## REFERENCES

- Bassino, L. P., & Muzzio, J. C. 1995, *Observatory*, 115, 256  
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245  
 Da Costa, G. S., & Armandroff, T. E. 1995, *AJ*, 109, 2533  
 Grillmair, C. J., Freeman, K. C., Irwin, M., & Quinn, P. J. 1995, *AJ*, 109, 2553  
 Harris, W. E. 1996, *AJ*, 112, 1487  
 Heitsch, F., & Richtler, T. 1999, *A&A*, 347, 455  
 Jurcsik, J. 1998, *ApJ*, 506L, 113  
 King, I. R. 1966, *AJ*, 71, 64  
 Layden, A. C., & Sarajedini, A. 2000, *AJ*, 119, 1760

- Layden, A. C., Ritter, L. A., Welch, D. L., & Webb, T. M. A. 1999, *AJ*, 117, 1313
- Lee, Y. W., Rey, S. C., Ree, C. H., Joo, J. M., Sohn, Y. J., & Yoon, S. J. 2002, in *ASP Conf. Ser. 265, Omega Centauri, a Unique Window into Astrophysics*, ed. F. van Leeuwen, J. D. Hughes, & G. Piotto (San Francisco: ASP), 305
- Lee, Y. W., Joo, J. M., Sohn, Y. J., Rey, S. C., Lee, H. C., & Walker, A. R. 1999, *Nature*, 402, 55L
- Leon, S., Meylan, G., & Combes, F. 2000, *A&A*, 359, 907
- Majewski, S. R., Patterson, R. J., Dinescu, D. I., Johnson, W. Y., Ostheimer, J. C., Kunkel, W. E., & Palma, C. 2000, *Proc. 35th Liege Intl. Astrophys. Colloq., The Galactic Halo, from Globular Clusters to Field Stars*, ed. A. Noels, P. Magain, D. Caro, E. Jehin, G. Parmentier, & A. A. Thoul (Leige: Institut d'Astrophys. Geophys.), 619
- Odenkirchen, M., et al. 2001, *ApJ*, 548, 165
- Piotto, G., et al. 1997, in *Advances in Stellar Evolution*, ed. R. T. Rood & A. Renzini (Cambridge: Cambridge Univ. Press), 84
- Pritzl, B. J., Smith, H. A., Catelan, M., & Sweigart, A. V. 2001, *AJ*, 122, 2600
- Pritzl, B. J., Smith, H. A., Catelan, M., & Sweigart, A. V. 2002, *AJ*, 124, 949
- Raimondo, G., Castellani, V., Cassisi, S., Brocato, E., & Piotto, G. 2002, *ApJ*, 569, 975
- Ree, C. H., Yoon, S. J., Rey, S. C., & Lee, Y. W. 2002, in *ASP Conf. Ser. 265, Omega Centauri, a Unique Window into Astrophysics*, ed. F. van Leeuwen, J. D. Hughes, & G. Piotto (San Francisco: ASP), 101
- Sarajedini, A., & Layden, A. C. 1995, *AJ*, 109, 1086
- Schlegel, D., Finkbeiner, D., & Davis, M. 1998, *ApJ*, 500, 525
- Silverman, B. W. 1986, *Density Estimation for Statistics and Data Analysis* (London: Chapman & Hall)
- Sweigart, A. V., & Catelan, M. 1998, *ApJ*, 501, L63
- Trager, S. C., King, I. R., & Djorgovski, S. 1995, *AJ*, 109, 218
- Tsujimoto, T., & Shigeyama, T. 2003, *ApJ*, 590, 803
- van Leeuwen, F., Hughes, J. D., & Piotto, G., eds. 2002, in *ASP Conf. Ser. 265, Omega Centauri, a Unique Window into Astrophysics* (San Francisco: ASP)
- Vanture, A. D., Wallerstein, G., & Suntzeff, N. B. 2002, *ApJ*, 569, 984
- Yoon, S. J., Lee, C. H., & Lee, Y. W. 2000, *BAAS*, 32, 41.02