Extreme Radio-wave Scattering Associated with Hot Stars

Mark A. Walker¹, Artem V. Tuntsov¹, Hayley Bignall², Cormac Reynolds², Keith W. Bannister³, Simon Johnston³,
 Jamie Stevens⁴, and Vikram Ravi⁵

¹ Manly Astrophysics, 15/41-42 East Esplanade, Manly, NSW 2095, Australia; Mark.Walker@manlyastrophysics.org

² CSIRO Astronomy and Space Science, 26 Dick Perry Avenue, Kensington, WA 6151, Australia; hbignall@gmail.com

³ CSIRO Astronomy and Space Science, P.O. Box 76, Epping, NSW 1710, Australia; Keith.Bannister@csiro.au

⁴ CSIRO Paul Wild Observatory, 1828 Yarrie Lake Road, Narrabri, NSW 2390, Australia

⁵ Cahill Center for Astronomy and Astrophysics, MC 249-17, California Institute of Technology, Pasadena, CA 91125, USA

Received 2017 March 22; revised 2017 April 26; accepted 2017 April 26; published 2017 June 27

Abstract

We use data on extreme radio scintillation to demonstrate that this phenomenon is associated with hot stars in the solar neighborhood. The ionized gas responsible for the scattering is found at distances up to 1.75 pc from the host star, and on average must comprise $\sim 10^5$ distinct structures per star. We detect azimuthal velocities of the plasma, relative to the host star, up to 9.7 km s^{-1} , consistent with warm gas expanding at the sound speed. The circumstellar plasma structures that we infer are similar in several respects to the cometary knots seen in the Helix and in other planetary nebulae. There the ionized gas appears as a skin around tiny molecular clumps. Our analysis suggests that molecular clumps are ubiquitous circumstellar features, unrelated to the evolutionary state of the star. The total mass in such clumps is comparable to the stellar mass.

Key words: circumstellar matter – ISM: structure – local interstellar matter – scattering – stars: massive

1. Introduction

The refractive index of ionized gas is very large at low frequencies, leading to profound effects on the propagation of radio waves through interstellar space—see the review by Rickett (1990). A useful reference model for many of the observed effects is distributed Kolmogorov turbulence in the diffuse, ionized interstellar medium (ISM; e.g., Armstrong et al. 1995). However, there are several phenomena, which collectively can be called “extreme scattering,” that are known to involve flux modulations and scattering angles much larger than can be accommodated in the Kolmogorov picture. Manifestations of extreme scattering include: multiple imaging of pulsars (Cordes & Wolszczan 1986; Rickett et al. 1997); extreme scattering events (ESEs; Fiedler et al. 1987, 1994; Bannister et al. 2016); intra-day variability of flat-spectrum radio quasars (IDVs; Kedziora-Chudczer et al. 1997; Dennett-Thorpe & de Bruyn 2000; Bignall et al. 2003); and parabolic arcs in pulsar “secondary” spectra (Stinebring et al. 2001; Cordes et al. 2006). These phenomena share at least one key characteristic: they all appear to require plasma pressures that are much greater than is typical of the diffuse ISM, and it therefore seems likely that they are somehow related. At present, there is no consensus on the cause of any of these phenomena, with suggestions ranging from current sheets seen edge-on (Goldreich & Sridhar 1995; Pen & Levin 2014), to the ionized skins of tiny, molecular clouds (Walker & Wardle 1998; Walker 2007).

Part of the difficulty in studying extreme scattering is that the phenomenon is transient, so one must first identify an active source, and then bring telescopes to bear on it. With a new observing program, dubbed ATESE, on the Australia Telescope Compact Array (ATCA), we have been able to identify sources undergoing extreme scattering, in a way that makes efficient use of telescope resources (Bannister et al. 2016). Although the ATESE project aims to find ESEs in progress, and then study them intensively, we have also identified new IDV sources (e.g., Tuntsov et al. 2017).

Most recently, the ATESE project discovered strong variability in the source PKS 1322–110 (Bannister et al. 2017). Subsequent follow-up with ATCA showed large-amplitude oscillations on a timescale of a few hours, so we now classify it as a (rapid, large-amplitude) IDV source. While considering optical follow-up of PKS 1322–110, we realized that it lies only $8'.5$ from Spica (α Vir)—the 16th brightest star in the sky. Only a handful of extreme scintillators are known, so it seemed improbable that this close positional coincidence would occur by chance. We were also aware of the suggestion of Dennett-Thorpe & de Bruyn (2002) that the IDV of J1819+3845 might be related to the foreground, bright star Vega. We therefore examined whether ionized gas associated with foreground stars provides a good explanation for the IDV phenomenon.

Although we continue to monitor PKS 1322–110, it will be some time before that data set rivals the coverage of published data on the two best-studied IDV sources: J1819+3845 (Dennett-Thorpe & de Bruyn 2000, 2002, 2003) and PKS 1257–326 (Bignall et al. 2003, 2006). Both of these sources displayed IDV over an interval of many years, revealing annual cycles in the scintillation timescale (which follow from the changing velocity of the Earth’s orbit around the Sun). The extensive, high-quality data for both sources provide tight constraints on the plasma responsible for the scattering, permitting stringent tests of any model that might be proposed, and they form the basis for our analysis.

This paper is organized as follows. In the next section, we present a rationale for selecting candidate stellar counterparts to extreme scintillators and apply it to the two well-studied IDV sources J1819+3845 and PKS 1257–326, yielding Vega and Alhakim, respectively, as likely stellar counterparts. The positions and motions of these stars suffice to predict the annual cycles of IDV, and we find that those predictions are a good match to the observed annual cycles. In Section 3, we show that it is highly improbable that this agreement is fortuitous. Section 4 considers the physical context in which

the plasma structures might arise and identifies the cometary knots in the Helix Nebula as the appropriate paradigm. Discussion and conclusions follow in Sections 5 and 6.

2. Hot Stars Foreground to IDV Sources

The discovery of IDV in PKS 1322–110 (Bannister et al. 2017), with Spica less than $9'$ away in the foreground, immediately suggests an association between the star and the ionized gas responsible for the scattering. The fact that Spica is a (binary) B-star further suggests that hot stars may be necessary to create the conditions for extreme scattering because B-type (and earlier) stars constitute a tiny fraction of all stars. Reinforcing that point is the idea that UV photons might be the agent that produces the plasma. We therefore adopt the hypothesis that material associated with hot, foreground stars is responsible for extreme scattering. The single example of Spica does not tell us which spectral types may give rise to extreme scattering. Initially, we include types O, B, and A in our definition of “hot” stars; we return to this issue later in the paper (Section 5).

To be more specific, our hypothesis is that IDV is caused by radially elongated plasma structures around hot stars. The rationale for specifying a radial orientation will become clear later. That the structures responsible for IDV must be elongated (i.e., statistically anisotropic) was pointed out by B. Rickett (1997, personal communication to M.A.W.) soon after IDV was discovered in PKS 0405–385 (Kedziora-Chudczer et al. 1997; Rickett et al. 2002). It was subsequently confirmed for⁶ J1819+3845 and PKS 1257–326. Indeed, the annual cycles of both of these sources are well approximated by a model in which the anisotropy is infinite—i.e., a one-dimensional model (Walker et al. 2009). We employ the one-dimensional approximation in this paper.

In order to identify which foreground stars could be responsible for the scintillation of a background radio source, we need to define the domain of influence of each star. Various physically motivated definitions are possible, in principle—e.g., based on the ionizing luminosity of the star. Our choice is the Hill surface, which defines the region within which material can remain bound to the star, in the presence of the Galactic tidal field. For the Sun, the Hill surface has principal semi-axes of approximately (1.42, 0.92, 0.74) pc, in cylindrical, Galactic coordinates (R, ϕ, Z) (Antonov & Latyshev 1972). The Galactic tidal field for local stars is similar to that experienced by the Sun, but the volume of the Hill surface scales with the stellar mass. In this paper, we therefore adopt a fiducial impact parameter of 2 pc as the domain of influence of local, massive stars.

The very rapid scintillations of J1819+3845 and PKS 1257–326 leave no doubt that the scattering material must be located at line-of-sight distances $\ll 1$ kpc from us (Dennett-Thorpe & de Bruyn 2000, 2003). Thus, a deep stellar catalog is not necessary for cross-matching; we employ the *Hipparcos* catalog (Perryman et al. 1997). Figure 1 shows the local *Hipparcos* stars along the line of sight to our two IDV sources, in cylindrical coordinates (ρ, z) relative to the axis defined by the direction to the radio source, and with the observer at the origin. We see that the line of sight to J1819+3845 pierces the

Figure 1. Locations of *Hipparcos* stars, in cylindrical coordinates, (ρ, z), around the line of sight to J1819+3845 (top) and PKS 1257–326 (bottom). The size of each point represents luminosity, and color indicates spectral type: blue $\leftarrow$ (O, B, A); green $\leftarrow$ (F); red $\leftarrow$ (G, K, M).

Hill surface of only one, local, hot star—Vega (α Lyr, A0V, $z = 7.7$ pc)—whereas for PKS 1257–326 there are two: Alhakim (ι Cen, A2V, $z = 18$ pc) and HD 112934 (A9V, $z = 55$ pc). That these lines of sight pass through the Hill surfaces of three hot stars with $z < 60$ pc is, in itself, slightly surprising. The cylindrical regions shown in Figure 1 encompass a volume $9\times$ greater than the volume of interest, and together they contain eight hot stars (blue points in the figure). We would therefore expect to see 0.9 hot stars within $\rho < 2$ pc from these lines of sight, whereas we find three.

Although three hot stars meet our selection criterion, for now we set aside HD 112934 and concentrate on Alhakim and Vega. The specific reasons for doing so will become clear in Section 5, where we argue that HD 112934 is probably the chance interloper we expected to find.

The coincidences between our radio sources and their foreground stars become much more remarkable when we consider all the constraints on the scattering plasma coming from the annual cycles of the IDV. These constraints localize the scattering material in three spatial dimensions, one angle and one velocity component, and are listed in Table 1. Corresponding quantities for the proposed counterpart stars are also given, as per the *Hipparcos* catalog. The quantities listed are as follows. The angle β is the orientation of the long axis of the (highly anisotropic) flux pattern, whereas β^* is the orientation of the line joining the star and the radio source. The quantity $v_{\perp}$ is the (negative of) the velocity component of the scattering plasma, perpendicular to the long axis of the flux pattern, whereas $v_{\perp}^*$ is the (negative of) the velocity component of the star, perpendicular to the same axis.⁷ The quantity $a_{\perp}$ is the length-scale of the flux pattern perpendicular to the long axis, determined from the product of the instantaneous perpendicular speed and the timescale of the scintillations.

⁶ It is also true of pulsar parabolic arcs—see Walker et al. (2004), Cordes et al. (2006), Briske et al. (2010)—which appear to be the counterpart, in pulsars, of the IDV phenomenon in quasars.

⁷ The velocity, $v_{\perp}$, as defined by Walker et al. (2009), is the steady component of the observer’s velocity through the flux pattern and in the context of the present discussion that corresponds to the negative of the relevant component of the plasma velocity, relative to the solar system barycenter.

Table 1
Properties of Scattering Plasma and of Nearby Stars

Radio Source	β (N $\rightarrow$ E) $^\circ$	$v_\perp$ (km s $^{-1}$)	$a_\perp$ (Mm)	z_s (pc)	β^* (N $\rightarrow$ E) $^\circ$	$v_\perp^*$ (km s $^{-1}$)	z^* (pc)	ρ^* (pc)	Star
J1819+3845	−97.35 (0.4)	19.67 (0.24)	29.5 (0.6)	10.9	−91.93 (0.00)	10.02 (0.07)	7.74 (0.03)	0.461 (0.002)	Vega
PKS 1257–326	126.60 (0.6)	20.50 (0.30)	44.9 (1.0)	13.5	134.50 (0.00)	22.4 (0.4)	17.9 (0.2)	1.75 (0.02)	Alhakim

Note. Columns 2, 3, and 4 are from Table 1 of Walker et al. (2009); they characterize the scattering plasma. Column 5 gives $z_s = a_\perp/\theta_\perp$, with $\theta_\perp$ as specified in Section 2. Columns 6, 7, 8, and 9 are the azimuth, perpendicular velocity component, and distances along and perpendicular to the line of sight for the host stars. Measurement uncertainties are given in parentheses; the uncertainty associated with z_s is predominantly systematic and is likely ~ 0.25 dex.

For a point-like radio source, the spatial flux pattern created by the scattering material would have structure on a wide range of scales, with large fluctuations on short timescales, for example. For a quasar, the source size has a smoothing effect on the observed flux pattern: all fluctuations on angular scales that are small compared to the source size are heavily damped, and $a_\perp = z_s \theta_\perp$, where $\theta_\perp$ is the angular size of the autocorrelation of the source intensity profile and z_s is the distance of the scattering material from the observer. To evaluate $\theta_\perp$, we model the radio emission as a synchrotron source near the self-Compton limit, mildly boosted toward the observer: a circularly symmetric Gaussian intensity profile, $\exp(-\theta^2/2\sigma^2)$, with a peak brightness temperature of 10^{13} K. Both radio sources have mean fluxes at 5 GHz, where the scintillation timescales were measured, of approximately 200 mJy, from which we determine $\sigma \simeq 13 \mu\text{as}$. The uncertainty in this estimate is likely ~ 0.25 dex, associated with ~ 0.5 dex uncertainty in the peak brightness temperature.

We require $\theta_\perp$ appropriate for the reported scintillation timescales, which follow different conventions depending on the author (e.g., the HWHM of the temporal autocorrelation of the flux measurements). The models reported by Walker et al. (2009) were fit to timescales corresponding to $\theta_\perp \simeq 1.35\sigma$ for J1819+3845 (Dennett-Thorpe & de Bruyn 2003, Table 1, column 8) and $\theta_\perp \simeq 1.67\sigma$ for PKS 1257–326 (Bignall et al. 2003). This model yields the estimate $z_s = a_\perp/\theta_\perp$, as given in column 5 of Table 1. The corresponding line-of-sight distance of the stars is given in column 8.

Using Table 1, we can compare the values of β , $v_\perp$, and z_s , describing the scattering plasma, with those of β^* , $v_\perp^*$, and z_s^* describing the positions and motions of the proposed counterpart stars. Three things are immediately evident from that comparison. First, the differences in orientations and velocity components are large compared to the uncertainties in the measurements. In Section 4, we discuss the interpretation of those differences. Although values for $a_\perp$ are determined quite precisely from the scintillation data, the same cannot be said of z_s because of the systematic uncertainty associated with our simplistic model of the radio source structure. Second, although our source model is unsophisticated, it suffices to demonstrate that the scattering material must be very local, as our counterpart stars indeed are. Third, the orientations of β are similar to the azimuths of the stars, β^* . It is less obvious whether the few km s $^{-1}$ differences between $v_\perp$ and $v_\perp^*$ are surprising. In the next section, we address that issue, and we assess the probability that Vega and Alhakim are not responsible for the observed scintillations, which is very small.

3. Probability of Chance Associations

Consider the possibility that the hot stars we have identified, close to the line of sight to IDV sources, have no causal

relationship to the scintillations. In that case, we are drawing randomly from among all such local stars, and we can use the *Hipparcos* catalog to estimate the chances of two random coincidences that are at least as close as the associations we have identified. We define “local,” here, to mean stars within 60 pc.

We assume that the position and velocity distributions are approximately independent, locally. We characterize the former by the local density of hot stars, $n = 4.0 \times 10^{-4}$ pc $^{-3}$. For the latter, we require the probability of matching specific velocity components, perpendicular to the line of sight. As the local, hot stars are found all over the sky, we take the relevant transverse velocity component to be in the same orientation, with respect to Galactic coordinates, as for the scintillation patterns.⁸ From Table 1, the velocity intervals of interest are (in km s $^{-1}$) $v_\perp = 19.7 \pm 9.7$ for J1819+3845 and $v_\perp = 20.5 \pm 1.9$ for PKS 1257–326, within which there are 85 and 21 local, hot stars in the *Hipparcos* catalog, respectively, out of a total of 365. Thus, we find the probability of a chance match in velocity, at least as close as we obtain for J1819+3845-Vega, to be $p_\perp = 0.233$, and for PKS 1257–326-Alhakim it is $p_\perp = 0.058$.

The number of stars that we expect to match the scattering plasma, in position and velocity, at least as well as our chosen stars, is just (with $\Delta\beta$ in radians)

$$N = n \Delta\beta \rho^2 \Delta z p_\perp. \quad (1)$$

The relevant intervals $\rho \leq \rho^*$, $\Delta\beta = 2|\beta^* - \beta|$, and $\Delta z = 2|z^* - z_s|$ can be determined from Table 1. We thus find that $N = 2.4 \times 10^{-5}$ for the J1819+3845-Vega pairing and $N = 1.7 \times 10^{-4}$ for PKS 1257–326-Alhakim. It is, therefore, unlikely that either of our proposed associations would have arisen as a random coincidence.

Taken together, the two proposed pairings are extremely unlikely to have arisen by chance. To evaluate the overall probability, we must consider how our original hypothesis—i.e., that the scattering material is radially oriented filaments, associated with hot stars—was arrived at. It was the link between Spica and PKS 1322–110 that triggered our investigation of a connection with hot stars; that piece of information is independent of the probabilities we have computed here, and thus has no impact upon them. However, the notion that the anisotropic structure of the scattering material is radially organized did not come from the PKS 1322–110-Spica connection. Rather, it was suggested to us by the fact that β , for J1819+3845, is similar to β^* for Vega. Thus,

⁸ In principle, a better estimate could be arrived at by using the three-dimensional vector component, parallel to that measured for the scintillation pattern. However, reliable radial velocity information is often not available for hot stars.

the overall probability of obtaining our result by chance is increased by the ratio $\pi/\Delta\beta$ for the J1819+3845-Vega association—a factor of 16.6. In conclusion, the probability of obtaining both results by chance is 6.8×10^{-8} .

Having established that there is a physical association between the foreground stars and the IDV, we now consider the origin of the plasma structures.

4. Nature of the Plasma Filaments

Perhaps the first physical picture that comes to mind is that the filamentary structures might arise in a wind from the central star. There are several problems with that idea. First, A-star winds are feeble. An upper limit on the mass loss from Vega, for example, is $\dot{M} \lesssim 4 \times 10^{-10} M_{\odot} \text{ yr}^{-1}$ (Lanz & Catala 1992), which corresponds to mean electron densities $n_e \lesssim 10^{-6} \text{ cm}^{-3}$ at 2 pc, if the wind velocity is 300 km s^{-1} . By contrast, the IDV phenomenon requires values $\sim 10 \text{ cm}^{-3}$ (Tuntsov et al. 2013); the stellar wind is not a natural source of such dense plasma. We would thus require the wind to be concentrated into radial streamers, covering a miniscule fraction of the sphere and with a density contrast $\gtrsim 10^7$. If the wind were a purely hydrodynamic phenomenon, such density contrasts should dissipate rapidly in a turbulent cascade. It is tempting to invoke magnetic fields, in a hydromagnetic wind, strong enough to confine the dense plasma. If so, the field lines and the plasma filaments ought to be azimuthally oriented at large radii (Parker 1958), whereas we require radial filaments. In short, it seems unlikely that the hot star is the source of the scattering plasma.

Having excluded the stellar wind, we are led to the idea that the source of the plasma is itself remote from the star, but close to the line of sight to the background radio source. The central stars are, however, a source of UV photons in abundance, so one possibility is that the plasma could be an ionized flow driven from the surface of dense, neutral clumps bound to the star. Such clumps have no counterpart in the theory of stellar structure and evolution. There is, however, at least one observed example of a strikingly similar situation: the cometary knots in the Helix Nebula (e.g., Meaburn et al. 1992; O’Dell & Handron 1996; Huggins et al. 2002; Meixner et al. 2005; Matsuura et al. 2009). Such structures appear to exist in other planetary nebulae (O’Dell et al. 2002), but are most clearly seen in the Helix, presumably because it is one of the closest planetaries.

Several points of similarity argue that the cometary knots in the Helix provide the correct paradigm for the environments of hot, main-sequence stars. First, the cometary knots are seen at distances of 0.2–0.4 pc from the central star of the nebula—see, particularly, Matsuura et al. (2009)—this is not much smaller than is required for the structures around Vega (0.45 pc). And it is possible that the Helix might contain yet more knots at greater distances from the central star. Being shielded from the stellar radiation by gas at smaller radii, they would be harder to see and could fall below the detection limit of observations to date. Second, extreme scattering is a phenomenon that is typically transient (Kedziora-Chudczer 2006; Lovell et al. 2008; de Bruyn & Macquart 2015), and is thus likely to be caused by discrete entities; the cometary knots in the Helix, although large in number, appear to be genuinely discrete. Third, the transverse scale of the cometary knots in the Helix is $\sim 10^2 \text{ au}$; this is very tiny by astronomical standards, but similar to the upper end of the range of dimensions inferred for

extreme scattering structures. In particular, the plasma concentration responsible for IDV in J1819+3845 is $\sim 10^2 \text{ au}$ in size (de Bruyn & Macquart 2015). Fourth, the tails of the cometary knots in the Helix are everywhere approximately radial, matching the orientation required for the IDV plasma anisotropies. Although the observed morphology of individual cometary knots corresponds to spatial scales that are $\gtrsim 10^5 \times$ greater than the plasma inhomogeneities responsible for IDV (i.e., $a_{\perp}$ in Table 1), the anisotropy on small scales may well be determined by the large-scale morphology and/or flow patterns of the plasma. Finally, the incidence of rapid, large-amplitude IDV requires $\sim 10^5$ plasma concentrations around each hot star—consistent with the number of cometary knots identified in the Helix (Matsuura et al. 2009). To see this, we calculate the probability of a randomly chosen line of sight intersecting one of the plasma concentrations responsible for IDV. The size of the latter has been estimated at $\sim 100 \text{ au}$ for J1819+3845 (de Bruyn & Macquart 2015), so we adopt a cross-section $\sim 10^4 \text{ au}^2$. With a hot-star number density of $4 \times 10^{-4} \text{ pc}^{-3}$, and a path length (scale height of hot stars in the Galactic disk) of $\sim 100 \text{ pc}$, we would have a sky-covering fraction of $\sim 10^{-8}$ if each hot star carried a single plasma concentration. The actual sky-covering fraction can be estimated from the results of Lovell et al. (2003), who found that J1819+3845 was the only rapid, large-amplitude variable in a sample of 710 compact radio quasars; hence, $\sim 10^5$ plasma concentrations are required around every hot star.

Returning now to Table 1, the paradigm provided by the Helix facilitates straightforward interpretations of the differences $|\beta - \beta^*|$ and $|v_{\perp} - v_{\perp}^*|$, both of which are several times larger than their measurement uncertainties. The observed differences in orientation can arise in two ways: the symmetry axes of the cometary tails may deviate from precisely radial orientations and the individual tails may have a non-zero opening angle. We note that foreshortening tends to increase the apparent deviation—e.g., by a factor of 2 at the median inclination of 60° —thus, the intrinsic deviation from radial is probably $\lesssim 4^\circ$ in both the symmetry axes and the tail opening angles. The observed differences in perpendicular velocity cannot be due to the azimuthal velocities of the neutral clumps, relative to the stars, as they could not then be bound to their hosts. Instead, we should ascribe the detected azimuthal velocity components to the motion of the plasma relative to the clumps. The relatively large value of 9.7 km s^{-1} , inferred for J1819+3845-Vega, is suggestive of free expansion of the plasma in the azimuthal direction, perpendicular to the cometary tails.

5. Discussion

The essential similarities between plasma structures in the Helix nebula and those around nearby stars indicate that neutral gas clumps, which are known to be molecular in the case of the Helix (Huggins et al. 1992), are a common circumstellar feature. Although we drew that connection in order to illuminate the origin of IDV, it also has implications for the Helix. If swarms of molecular clumps are present around main-sequence stars, they did not get there as a result of late-stage stellar evolution, as is often assumed for the Helix (e.g., García-Segura et al. 2006). We will argue that main-sequence stars of all types may carry populations of molecular clumps. The progenitor of the Helix could have been such a star, so it is no

longer appropriate to assume that the observed clumps are a product of stellar mass loss.

Where, then, did all of these clumps come from? Viable formation scenarios must precede the main sequence. One possibility is that clumps form contemporaneously with stars. Another possibility is that parsec-sized clusters of molecular clumps constitute the starting point and a star subsequently forms at the center of each cluster as a result of disruptive collisions between clumps. We cannot say which, if either, of these speculations is correct. There are two key considerations for any proposed scenario. First, the clumps must have long lifetimes—comparable to or greater than the main-sequence lifetime of an A-star—which strongly suggests that they are, individually, self-gravitating clouds. Second, the formation mechanism must be very efficient because the total mass in molecular clumps is large—as we now demonstrate.

In the case of the cometary knots in the Helix, estimates of the masses of individual, neutral clumps have yielded values of $\sim 2 \times 10^{-5} M_{\odot}$ (e.g., Meaburn et al. 1998), implying that their total mass in the Helix is $\gtrsim 1 M_{\odot}$. This estimate is very rough, as both clump masses and numbers are highly uncertain. Notwithstanding those uncertainties, it appears that the total mass is comparable to the stellar mass, and in the present work, we have argued that the situation is similar for local, hot stars. As this paper concerns itself with the interpretation of IDV, which involves a strong selection bias in favor of local, ionized material, we anticipate that stars of all spectral types, throughout the Galaxy, may also be surrounded by swarms of tiny molecular clumps. If so, then the total molecular clump mass constitutes a substantial fraction of the mass of the Galaxy.

That conclusion was also reached by Walker & Wardle (1998), via a different line of reasoning. They demonstrated that if ESEs are caused by the ionized skins of self-gravitating, neutral clouds, then the observed event rate requires that those clouds must form a substantial component of the Galaxy’s dark matter. Detailed models of such clouds confirm that they can have very low luminosities (M. Walker & M. Wardle 2017, in preparation).

A third line of evidence supports the view that the total mass of molecular clumps is large. Using recent results for the Oort constants, based on *Gaia* data (Bovy 2017) and the local mass density determined from *Hipparcos* data (Cr et al. 1998), we have re-evaluated the size and shape of the Hill surface, following Antonov & Latyshev (1972) and Heisler & Tremaine (1986). We find principal semi-axes of (1.37, 0.91, 0.73) pc for the Sun. Taking a mass of $2.5 M_{\odot}$ for Alhakim (HD 115892; Quanz et al. 2011), we find that the Hill surface extends only 70% of the angular distance to PKS 1257–326. We therefore require an additional mass $\gtrsim 5 M_{\odot}$, bound to Alhakim, in order to extend the Hill surface as far out as the PKS 1257–326 line of sight. That conclusion is specific to Alhakim, but, in combination with our other two estimates of molecular mass, it alludes to a likely conflict with contemporary cosmological models, in which most of the matter is non-baryonic.

How do our results sit in relation to ideas about the local ISM? It has been suggested that the IDV of J1819+3845 and PKS 1257–326 is connected with the boundaries of certain local clouds, whose properties have been inferred from absorption-line spectroscopy (Linsky et al. 2008). The absorption-line clouds have much larger sky-covering fractions than the plasma concentrations responsible for extreme scattering and are therefore

not directly identifiable with the cometary knots themselves. However, as the plasma continues to flow outward, away from the star, it presumably expands laterally until it becomes space filling. The resulting gas would have a low density, but the column density might nevertheless be high enough to be detectable in absorption-line studies. It might therefore be interesting to re-examine the absorption-line data to see whether they are consistent with, e.g., centro-symmetric velocity fields around local stars.

It is interesting to consider how the scintillations of background radio sources might change when the molecular cloud clusters are located around late-type stars, rather than the early types we have been concerned with so far. Lower photospheric temperatures combined with lower bolometric luminosities (for main-sequence stars) mean that the UV radiation fields are much smaller for the late types. Likewise, the late types are expected to have lower mechanical (wind) luminosities, at least while stars are on the main sequence. Consequently, the plasma skins are expected to be much less prominent in the case of molecular clumps around cooler stars. Presumably, this means that the phenomenon of extreme scattering is then restricted to clumps that lie well inside the star’s Hill surface, so that very small impact parameters are required in order for extreme scattering to be observed. At present, we lack a reliable theoretical basis for predicting plasma densities, and hence scattering characteristics, from the properties of the central star. In this paper, our focus has been on O-, B-, and A-type stars, implying that spectral types F, and later, are too cool to promote extreme scattering. That is a reasonable assumption, given the rapid change in ionizing luminosity near that boundary, but at present it is just an hypothesis.

Continuing to even cooler stars, we can presumably reach a regime where there are no molecular clumps that have plasma dense enough to cause ESEs and IDVs. However, molecular gas has a refractive index of its own, and self-gravitating molecular clouds can constitute strong lenses even without a plasma skin (Draine 1998). At frequencies well below the strong, far-UV absorptions of H_2 and He, this type of lensing is almost achromatic. Thus, the molecular clump populations around cool stars might reveal themselves by radio-wave scintillations that are frequency independent. In connection with that possibility, we note that weakly chromatic scintillations have been reported in the blazar PKS 1413+135 (Vedantham et al. 2017), for which the line of sight passes close to Arcturus (α Boo, K0III, $z = 11.3$ pc, $\rho = 1.1$ pc).

It is unclear what role, if any, the star HD 112934 plays in the observed scintillations of PKS 1257–326. We account for the IDV of PKS 1257–326 by the presence of Alhakim in the foreground: HD 112934 is not needed. Nor is it a good match to the scintillation characteristics of PKS 1257–326: $\beta^* = 10^\circ.7$, approximately 64° from β ; $v_{\perp}^* = 30.9 \pm 1.5$ km s $^{-1}$, yielding $p_{\perp} = 0.32$ and $z^* = 55 \pm 2$ pc, which is much larger than the 13.5 pc expected for the scattering material. These attributes all suggest that HD 112934 is a random interloper, unrelated to the behavior of PKS 1257–326.

What is surprising, though, is that HD 112934 is so close to the line of sight: its impact parameter is $\rho^* = 0.160 \pm 0.006$ pc. That is so small that the probability of this star being there by chance, computed using Equation (1), is only about twice that calculated for Alhakim, even though $\Delta\beta$, Δz , and $p_{\perp}$ are all much larger. However HD 112934 fits into the picture, it does

nothing to weaken our conclusion that hot, foreground stars are central to the IDV phenomenon.

Having established the association between foreground stars and two particular IDV sources, it is sensible to examine the possible role of foreground stars more broadly in extreme scattering phenomena. A thorough treatment is beyond the scope of this paper, but some brief commentary is appropriate. The two IDV sources we focused on were chosen because they offer precise localizations of the scattering plasma. No other known IDVs approach the level of precision of these two. Consequently, although it might be possible to identify plausible stellar hosts for the foreground plasma, we do not expect that associations could be made so confidently for any of the other known IDVs.

A good example is PKS 0405–385, whose IDV has been intensively studied (Kedziora-Chudczer et al. 1997; Kedziora-Chudczer 2006). Like J1819+3845 and PKS 1257–326, its variations are very rapid, so we may reasonably expect the responsible plasma to be very local. Indeed, as PKS 0405–385 is almost an order of magnitude brighter than either J1819+3845 or PKS 1257–326, its angular size should be expected to be $\sim 3\times$ larger, and if $v_\perp$ is not too different from the other two IDV then with a similar timescale we expect similar $a_\perp$, and hence z_s should be smaller by a factor ~ 3 . That is so close that Sirius (α CMa, A1V, $z = 2.1$ pc, $\rho = 1.8$ pc) is the only plausible hot stellar counterpart. To determine whether Sirius is indeed the host, we need measurements of β , $v_\perp$, and $a_\perp$ for the plasma causing the scintillations. Unfortunately, although an evolution of the scintillation timescale of PKS 0405–385 was noted by Kedziora-Chudczer (2006), it was based on two short periods of variability, and no annual cycle has been demonstrated for this source. We therefore cannot confirm Sirius as the host star.

The same is true for the other extreme scattering phenomena, and for the same reason, poor localization of the plasma. As an example, one of the best-studied manifestations of parabolic arcs is in the pulsar B0834+06 (Stinebring et al. 2001; Hill et al. 2005; Briskin et al. 2010). In that case, the orientation, on the sky, of the plasma anisotropy is determined to similar precision as for our two IDV sources (β in our Table 1). But the plasma velocity and line-of-sight distance both have large uncertainties—the latter being several hundred parsecs (Briskin et al. 2010). A subsequent parallax measurement for this pulsar (Liu et al. 2016) has substantially reduced these uncertainties.

We expect similar difficulties in the case of IDV that is less extreme than that exhibited by PKS 1257–326 and J1819+3845. Slower variations, with lower amplitude, are expected to arise when the scattering plasma is more distant, as a result of smoothing associated with the source size, and such variability should be relatively common. That is qualitatively consistent with the many cases of IDV reported by Lovell et al. (2008). Identifying the stellar host of the scattering plasma will be more difficult in those cases because the larger range in line-of-sight distance admits more potential counterparts. However, if firm identifications prove elusive, a statistical model may be worthwhile—using the known locations and velocities of hot stars in the solar neighborhood to predict the chances of observing IDV in each radio source.

In the introduction to this paper, we noted that IDVs and ESEs are likely to be related, but we did not say how. Now, armed with a picture of the plasma concentrations responsible for IDV, we can suggest a possible relationship. Consider a

cometary knot, such as in the Helix. As the width of the knot is comparable to the extent of the plasma responsible for IDV in J1819+3845 (de Bruyn & Macquart 2015), we suppose that IDV arises when the line of sight pierces a typical point, suggesting that microstructure is present throughout the plasma skin. The ESE phenomenon, however, does not seem to be caused by microstructure; instead, it requires a column-density curvature that is both strong and consistent in its sign over a broad spatial extent (Tuntsov et al. 2016). For a thin skin of plasma, the smooth component of the column density—i.e., its macrostructure—has a curvature that is strongest for lines of sight that are just inside the boundary of the structure, so that is where we might plausibly expect ESEs to arise.

Various types of studies would be useful in testing our ideas. Atomic emission lines, such as H α , are generically expected to arise from extreme scattering plasma, so their discriminating power is modest. Imaging in molecular emission lines, on the other hand, can directly address some of the key issues raised in this paper. Taking the Helix as our paradigm motivates observations in the H $_2$ rovibrational lines, for example, and the pure rotational lines of CO. We note, however, the factor of ~ 10 difference in photospheric temperatures between the upper main sequence and the central star of the Helix. Therefore, if it is ionizing photons that are responsible for creating the cometary knots, we expect that their optical/near-IR emission will appear very faint in comparison with any main-sequence host.

6. Conclusions

Position and velocity constraints on the plasma responsible for IDV in J1819+3845 and PKS 1257–326 place the material close to Vega and Alhakim, respectively—two nearby A-stars. The probability that this coincidence occurred by chance is very low, and we conclude that the relationship is causal: the scattering material must be associated with these stars. The size, location, orientation, and numbers of plasma concentrations all point to circumstellar environments that are fundamentally similar to the Helix Nebula, where it is known that the plasma is associated with dense molecular clumps. Therefore, we conclude that similar populations of molecular clumps are present around hot stars in the solar neighborhood. Bearing in mind the selection bias associated with IDV studies—i.e., rapid scintillation of radio quasars requires nearby, high-density plasma—we anticipate that circumstellar molecular clumps are likely to be ubiquitous across all stellar spectral types and throughout the Galaxy. If so, three independent estimates indicate that those clumps make a major contribution to the mass of the Galaxy.

Hats off to Ger de Bruyn and Jane Dennett-Thorpe for spotting, very early on, the possibility of a connection between Vega and J1819+3845. The referee’s thoughtful comments helped to sharpen our presentation.

References

- Antonov, V. A., & Latyshev, I. N. 1972, in IAU Symp. 45, The Motion, Evolution of Orbits, and Origin of Comets, ed. G. A. Chebotarev, E. I. Kazimirschak-Polonskaia, & B. G. Marsden (Dordrecht: Reidel), 341
- Armstrong, J. W., Rickett, B. J., & Spangler, S. R. 1995, *ApJ*, **443**, 209
- Bannister, K. W., Bignall, H. E., Johnston, S., et al. 2017, *ATel*, **10024**
- Bannister, K. W., Stevens, J., Tuntsov, A. V., et al. 2016, *Sci*, **351**, 354
- Bignall, H. E., Jauncey, D. L., Lovell, J. E. J., et al. 2003, *ApJ*, **585**, 653
- Bignall, H. E., Macquart, J. P., Jauncey, D. L., et al. 2006, *ApJ*, **652**, 1050

- Bovy, J. 2017, *MNRAS*, **468**, L63
- Briskin, W. F., Macquart, J.-P., Gao, J. J., et al. 2010, *ApJ*, **708**, 232
- Cordes, J. M., Rickett, B. J., Stinebring, D. R., & Coles, W. A. 2006, *ApJ*, **637**, 346
- Cordes, J. M., & Wolszczan, A. 1986, *ApJL*, **307**, L27
- Cr, M., Chereul, E., Bienaym, O., & Pichon, C. 1998, *A&A*, **329**, 920
- de Bruyn, A. G., & Macquart, J.-P. 2015, *A&A*, **574**, 125
- Dennett-Thorpe, J., & de Bruyn, A. G. 2000, *ApJL*, **529**, L65
- Dennett-Thorpe, J., & de Bruyn, A. G. 2002, *Natur*, **415**, 57
- Dennett-Thorpe, J., & de Bruyn, A. G. 2003, *A&A*, **404**, 113
- Draine, B. T. 1998, *ApJL*, **509**, L41
- Fiedler, R., Dennison, B., Johnston, K. J., Waltman, E. B., & Simon, R. S. 1994, *ApJ*, **430**, 581
- Fiedler, R. L., Dennison, B., Johnston, K. J., & Hewish, A. 1987, *Natur*, **326**, 675
- Garca-Segura, G., Lpez, J. A., Steffen, W., Meaburn, J., & Manchado, A. 2006, *ApJL*, **646**, L61
- Goldreich, P., & Sridhar, S. 1995, *ApJ*, **438**, 763
- Heisler, J., & Tremaine, S. 1986, *Icar*, **65**, 13
- Hill, A., Stinebring, D. R., Asplund, C. T., et al. 2005, *ApJL*, **619**, L171
- Huggins, P. J., Bachiller, R., Cox, P., & Forveille, T. 1992, *ApJL*, **401**, L43
- Huggins, P. J., Forveille, T., Bachiller, R., et al. 2002, *ApJL*, **573**, L55
- Kedziora-Chudczer, L. 2006, *MNRAS*, **369**, 449
- Kedziora-Chudczer, L., Jauncey, D. L., Wieringa, M. H., et al. 1997, *ApJL*, **490**, L9
- Lanz, T., & Catala, C. 1992, *A&A*, **257**, 663
- Linsky, J. L., Rickett, B. J., & Redfield, S. 2008, *ApJ*, **675**, 413
- Liu, S., Pen, U.-L., Macquart, J.-P., Briskin, W., & Deller, A. 2016, *MNRAS*, **458**, 1289
- Lovell, J. E. J., Jauncey, D. L., Bignall, H. E., et al. 2003, *AJ*, **126**, 1699
- Lovell, J. E. J., Rickett, B. J., Macquart, J.-P., et al. 2008, *ApJ*, **689**, 108
- Matsuura, M., Speck, A. K., McHunu, B. M., et al. 2009, *ApJ*, **700**, 1067
- Meaburn, J., Clayton, C. A., Bryce, M., et al. 1998, *MNRAS*, **294**, 201
- Meaburn, J., Walsh, J. R., Clegg, R. E. S., et al. 1992, *MNRAS*, **255**, 177
- Meixner, M., McCulloch, P., Hartman, J., Son, M., & Speck, A. 2005, *AJ*, **130**, 1784
- O'Dell, C. R., Balick, B., Hajian, A. R., Henney, W. J., & Burkert, A. R. 2002, *AJ*, **123**, 3329
- O'Dell, C. R., & Handron, K. D. 1996, *AJ*, **111**, 1630
- Parker, E. N. 1958, *ApJ*, **128**, 664
- Pen, U.-L., & Levin, Y. 2014, *MNRAS*, **442**, 3338
- Perryman, M. A. C., Lindegren, L., Kovalevsky, J., et al. 1997, *A&A*, **323**, L49
- Quanz, S. P., Kenworthy, M. A., Meyer, M. R., Girard, J. H. V., & Kasper, M. 2011, *ApJL*, **736**, L32
- Rickett, B. J. 1990, *ARAA*, **28**, 561
- Rickett, B. J., Kedziora-Chudczer, L., & Jauncey, D. L. 2002, *ApJ*, **581**, 103
- Rickett, B. J., Lyne, A. G., & Gupta, Y. 1997, *MNRAS*, **287**, 739
- Stinebring, D. R., McLaughlin, M. A., Cordes, J. M., et al. 2001, *ApJL*, **549**, L97
- Tuntsov, A. V., Bignall, H. E., & Walker, M. A. 2013, *MNRAS*, **429**, 2562
- Tuntsov, A. V., Stevens, J., Bannister, K. W., et al. 2017, *MNRAS*, in press (arXiv:1705.06051)
- Tuntsov, A. V., Walker, M. A., Koopmans, L. V. E., et al. 2016, *ApJ*, **817**, 176
- Vedantham, H. K., Readhead, A. C. S., Hovatta, T., et al. 2017, *ApJ*, submitted (arXiv:1702.05519)
- Walker, M. A. 2007, in ASP Conf. Ser. 365, SINS—Small Ionized and Neutral Structures in the Diffuse ISM, ed. M. Haverkorn & W. M. Goss (San Francisco, CA: ASP), 299
- Walker, M. A., de Bruyn, A. G., & Bignall, H. E. 2009, *MNRAS*, **397**, 447
- Walker, M. A., Melrose, D. B., Stinebring, D. R., & Zhang, C. M. 2004, *MNRAS*, **354**, 43
- Walker, M. A., & Wardle, M. J. 1998, *ApJL*, **498**, L125