

DIVISION OF THE HUMANITIES AND SOCIAL SCIENCES

# CALIFORNIA INSTITUTE OF TECHNOLOGY


PASADENA, CALIFORNIA 91125

## OBJECTIVE SUBJECTIVE PROBABILITIES

Kim C. Border  
Caltech

Paolo Ghirardato  
University of Iowa and Università di Torino

Uzi Segal  
Boston College



**SOCIAL SCIENCE WORKING PAPER 1228**

June 2005

# Objective Subjective Probabilities

Kim C. Border

Paolo Ghirardato

Uzi Segal

## Abstract

This note shows that if the space of events is sufficiently rich and the subjective probability measure of each individual is non-atomic, then there is a  $\sigma$ -algebra of events on which everyone will have the same probability, and moreover, the range of these probabilities is the entire segment  $[0, 1]$ .

JEL classification numbers: D800

Key words: Subjective probability, objective probability, Dubins–Spanier theorem, fair division

# Objective Subjective Probabilities

Kim C. Border\*

Paolo Ghirardato<sup>†</sup>

Uzi Segal<sup>‡</sup>

## 1 Introduction

An important assumption in the formal analysis of social choice theory is the existence of social lotteries, that is, lotteries whose outcomes are social policies.<sup>1</sup> Such lotteries can increase the fairness of the social allocation mechanism or solve disputes in a cheap, efficient manner. For a social lottery to work, it must be considered fair by all individuals in society. Formally, if society finds it optimal to randomize over the  $k$  pure social policies  $s_1, \dots, s_k$  by using the probability vector  $p = (p_1, \dots, p_k)$ , then everyone in society must agree that the mechanism is indeed using these probabilities.<sup>2</sup>

But do such mechanisms exist? Even in the model of Anscombe and Aumann [2], where each decision maker is assumed to face subjective horse lotteries and objective roulettes, it does not follow that all decision makers agree on what is objective. An Italian-speaking person, facing a die whose sides are marked *Uno, Tre, Cinque, Sette, Otto, Dieci* will consider the event “the die will show an odd number” to be objective, while a non Italian-speaking person will consider it subjective (or even ambiguous). Nothing in the assumptions and structure of the Anscombe–Aumann model implies agreement on what is a roulette lottery. The issue is even more critical in Savage’s [12] framework, where all events are assumed to be subjective.

Recently, Ghirardato et. al. [6] showed that even if probabilities don’t exist (that is, beliefs are ambiguous), it is still possible, under some assumptions, to obtain mixture-like operators over random variables. But these procedures are subjective, and cannot be jointly used. Machina [11], on the other hand, assumes that preferences are smooth

---

\*Division of the Humanities and Social Sciences, California Institute of Technology, Pasadena CA 91125. kcborder@caltech.edu

<sup>†</sup>Dipartimento di Statistica e Matematica Applicata, Università di Torino, Piazza Arbarello 8, 10122 Torino Italy, ghirardato@econ.unito.it and Department of Economics, University of Iowa, Iowa City IA 52242, paolo-ghirardato@uiowa.edu

<sup>‡</sup>Department of Economics, Boston College, Chestnut Hill MA 02467. segalu@bc.edu

<sup>1</sup>See Harsanyi [8], and more recently, Epstein and Segal [5], Broome [3], Kamm [9], or Karni and Safra [10].

<sup>2</sup>Unlike the cake division problem, where participants want to ensure receiving *at least* their fair share, in social choice individuals often wish also not to receive *more* than their fair share.

and proves that for each  $r \in [0, 1]$  there is a sequence of events  $E_n$  such that for each  $i$ ,  $\mu_i(E_n) \rightarrow r$ . Unfortunately, the limits of these sequences of events don't exist.<sup>3</sup> Moreover, from the social point of view it may be important for everyone to agree that an event has probability exactly  $\frac{1}{n}$ , not approximately  $\frac{1}{n}$ .

In this note we show that if the space of events is sufficiently rich and the subjective probability function of each individual is non-atomic, then there is a  $\sigma$ -algebra of events over which everyone will have the same probability function, and moreover, the range of these probabilities is the whole  $[0, 1]$  segment. In other words, even in a fully subjective world (for example, Savage's), there is a rich set of events that can be used for joint randomizations. We prove existence, but we don't know how to construct specific such  $\sigma$ -algebras. This does not annul the contribution of this note. Randomization in social choice theory plays an important theoretical role, but it doesn't necessarily follow that policy makers do actually randomize. Our aim is to close a theoretical gap that exists in the literature—if commonly accepted devices do not exist, then models using randomization to enhance fairness would become void. Theorem 1 shows that there are enough events over which decision makers agree.

## 2 A Theorem

**Theorem 1** *Let  $\mu_1, \dots, \mu_n$  be nonatomic, countably additive probability measures on a measurable space  $(S, \Sigma)$ . Then there is a sub- $\sigma$ -algebra  $\hat{\Sigma}$  of  $\Sigma$  on which all the measures agree, which is rich in the sense that for every real number  $r \in [0, 1]$ , it contains a set of (unanimous) measure  $r$ .*

*Proof:* We start by using a well known result of Dubins and Spanier, which is restated in the Appendix. According to their theorem, it is possible to partition  $S$  into two sets  $E_0$  and  $E_1 = E_0^c$  belonging to  $\Sigma$  such that  $\mu_i(E_0) = \mu_i(E_1) = 1/2$ , for all  $i = 1, \dots, n$ . Let  $A_1$  denote the  $\sigma$ -algebra generated by this partition, namely  $A_1 = \{\emptyset, E_0, E_1, S\} \subset \Sigma$ .

Repeating this operation, we can partition  $E_0$  into two disjoint sets  $E_{00}$  and  $E_{01}$ , and also partition  $E_1$  into two disjoint sets  $E_{10}$  and  $E_{11}$ , so that  $\mu_i(E_{b_1 b_2}) = 1/4$  for all  $i = 1, \dots, n$  and  $b_1 = 0, 1$  and  $b_2 = 0, 1$ . Let  $A_2$  denote the  $\sigma$ -algebra generated by  $\{E_{00}, E_{01}, E_{10}, E_{11}\}$ . Note that  $A_1 \subset A_2 \subset \Sigma$ .

Proceeding in this fashion, for each  $m$ , partition  $S$  into  $2^m$  pairwise disjoint sets  $E_{b_1 \dots b_m}$ , where each  $b_j \in \{0, 1\}$ , satisfying

$$E_{b_1 \dots b_{m-1} b_m} \subset E_{b_1 \dots b_{m-1}},$$

and

$$\mu_i(E_{b_1 \dots b_m}) = 1/2^m$$

---

<sup>3</sup>For example, let the event  $A_n$  be “the  $n$ -th digit to the right of the decimal point of the temperature tomorrow will be odd.” Then as  $n \rightarrow \infty$ , all individual beliefs regarding these events will converge to  $\frac{1}{2}$ . But we know of no sense of limit for which  $\lim A_n$  exists as an event of probability  $\frac{1}{2}$ .

for all  $i = 1, \dots, n$ . Letting  $A_m$  denote the  $\sigma$ -algebra generated by this partition, we have  $A_{m-1} \subset A_m \subset \Sigma$ .

Set  $A = \bigcup_{m=1}^{\infty} A_m \subset \Sigma$ . Then it is easy to verify that  $A$  is an algebra, but not a  $\sigma$ -algebra, and that all the measures  $\mu_1, \dots, \mu_n$  agree on  $A$ . Let  $\mu$  denote the common restriction of each  $\mu_i$  to  $A$ . Then, for any dyadic rational  $q = k/2^m$  in the unit interval there is a set  $E$  in  $A_m \subset A$  with  $\mu(E) = q$ .

Let  $\hat{\Sigma} = \sigma(A) \subset \Sigma$ , the  $\sigma$ -algebra generated by  $A$ . By the Carathéodory Extension Theorem (see Appendix),  $\mu$  has a unique extension to  $\hat{\Sigma}$ , which we again denote by  $\mu$ . Since this extension is unique, each  $\mu_i$  agrees with  $\mu$  on  $\hat{\Sigma}$ .

Moreover the range of  $\mu$  is all of  $[0, 1]$ . To see this, let  $r$  belong to the unit interval. Then  $r$  has a binary expansion  $r = \sum_{m=1}^{\infty} b_m/2^m$ , where each  $b_m$  is a binary digit (bit), 0 or 1. For each  $m$ , choose the set  $F_m \in A_m$  by

$$F_m = \begin{cases} \emptyset & \text{if } b_m = 0 \\ E_{\underbrace{0\dots 0}_{m-1}}1 & \text{if } b_m = 1. \end{cases}$$

Note that  $\mu(F_m) = b_m/2^m$ . By construction, the sets  $F_m$  are pairwise disjoint. (To see this, suppose  $F_k$  and  $F_m$  are nonempty with  $k < m$ . Then  $F_k = E_{\underbrace{0\dots 0}_{k-1}}1$  and  $F_m$  is a subset of  $E_{\underbrace{0\dots 0}_{k-1}}0$ , which is disjoint from  $F_k$ .) Thus the set  $F = \bigcup_{m=1}^{\infty} F_m$  belongs to  $\hat{\Sigma}$  and satisfies  $\mu(F) = r$ . ■

### 3 An Example

The following example shows that we cannot extend our result to more than a finite set of individuals. For consider the countably infinite set  $\{0, 1, 2, \dots\}$ . Let  $S = [0, 1]$  equipped with its Borel  $\sigma$ -algebra, and let  $\{q_i : i = 0, 1, 2, \dots\}$  be an enumeration of the rationals in  $[0, 1)$  where  $q_0 = 0$ . Assume that Person  $i$ 's subjective probability  $P_i$  has the density  $f_i$  given by

$$f_i(t) = \begin{cases} \frac{1}{2} & t < q_i \\ \frac{1 - q_i}{1 - q_i} & t \geq q_i \end{cases}$$

Note that  $P_0$  is just Lebesgue measure.

We now want to show that there is no event  $A$  such that  $P_i(A) = \frac{1}{2}$  for all  $i = 0, 1, \dots$ . Suppose  $A$  is such an event. Computing the probabilities according to person 0 and person  $i > 0$ , we obtain

1.  $\lambda(A \cap [0, q_i)) + \lambda(A \cap [q_i, 1]) = \frac{1}{2}$ .
2.  $\frac{1}{2}\lambda(A \cap [0, q_i)) + \frac{2-q_i}{2-2q_i}\lambda(A \cap [q_i, 1]) = \frac{1}{2}$ ,

where  $\lambda$  is ordinary Lebesgue measure. The solution of this system is  $\lambda(A \cap [0, q_i)) = q_i/2$  and  $\lambda(A \cap [q_i, 1]) = (1 - q_i)/2$ . For all rationals  $a$  and  $b$  it now follows, by taking  $q_i > b$ , that  $\lambda(A \cap [a, b]) = (b - a)/2$ .

It is well known (see for instance, Halmos [7, Theorem A in §16, p. 68]) that there is no Lebesgue measurable set (and so no Borel set) satisfying this property.

## Appendix

The following well known result is due to Dubins and Spanier [4]. It may also be found in Aliprantis and Border [1, Theorem 12.34, p. 445].

**Dubins–Spanier Theorem** *Let  $\mu_1, \dots, \mu_n$  be nonatomic probability measures on a measurable space  $(S, \Sigma)$ . Given  $\alpha_1, \dots, \alpha_m \geq 0$  with  $\sum_{j=1}^m \alpha_j = 1$ , there is a partition  $\{E_1, \dots, E_m\}$  of  $S$  satisfying  $\mu_i(E_j) = \alpha_j$  for all  $i = 1, \dots, n$  and  $j = 1, \dots, m$ .*

Dubins and Spanier also include a lesser known result, which is a slight sharpening of part of the Lyapunov Convexity Theorem. It shows that the family of events on which the measures agree is rich, but it does not show that it includes a rich  $\sigma$ -algebra.

**Dubins–Spanier [4, Lemma 5.3]** *Let  $\mu_1, \dots, \mu_n$  be nonatomic (countably additive) probability measures on a measurable space  $(S, \Sigma)$ . Then there is a subfamily  $\{E_\alpha : \alpha \in [0, 1]\}$  of  $\Sigma$  satisfying*

$$\mu_i(E_\alpha) = \alpha \quad \text{for all } i = 1, \dots, n,$$

and

$$\alpha < \beta \implies E_\alpha \subset E_\beta.$$

A complete statement of the Carathéodory Extension Theorem may be found in Aliprantis and Border [1, Theorem 9.22, p. 343]. For our purposes, we need only the following special case.

**Carathéodory Extension Theorem** *Let  $\mathcal{A}$  be an algebra of subsets of  $X$  and let  $\mu$  be a probability measure on  $\mathcal{A}$ . Then  $\mu$  has a unique extension to  $\sigma(\mathcal{A})$ , the  $\sigma$ -algebra generated by  $\mathcal{A}$ .*

## References

- [1] Aliprantis, C. D. and K. C. Border. 1999. *Infinite dimensional analysis: A hitchhiker's guide*, 2d. ed. Berlin: Springer–Verlag.
- [2] Anscombe, F. J. and R. J. Aumann. 1963. A definition of subjective probability. *Annals of Mathematical Statistics* 34(1):199–205.
- [3] Broome, J. 1984. Selecting people randomly. *Ethics* 95:38–55.
- [4] Dubins, L. E. and E. H. Spanier. 1961. How to cut a cake fairly. *American Mathematical Monthly* 68(1):1–17.
- [5] Epstein, L. and U. Segal. 1992. Quadratic social welfare functions. *Journal of Political Economy* 100:691–712.
- [6] Ghirardato, P., F. Maccheroni, M. Marinacci, and M. Siniscalchi. 2001. A subjective spin on roulette wheels. *Econometrica* 71:1897–1908.
- [7] Halmos, P. R. 1974. *Measure theory*. Number 18 in Graduate Texts in Mathematics. New York: Springer–Verlag. Reprint of the edition published by Van Nostrand, 1950.
- [8] Harsanyi, J. C. 1955. Cardinal welfare, individualistic ethics, and interpersonal comparisons of utility. *Journal of Political Economy* 63:309–321.
- [9] Kamm, F. M. 1993. *Morality, mortality: Death and whom to save from it (volume 1)*. New York: Oxford University Press.
- [10] Karni, E. and Z. Safra. 2002. Individual sense of justice: A utility representation. *Econometrica* 70:263–284.
- [11] Machina, M. J. 2004. Almost-objective uncertainty. *Economic Theory* 24:1–54.
- [12] Savage, L. J. 1954. *Foundations of statistics*. New York: John Wiley and Sons. Reprint. New York: Dover, 1972.