

Schrödinger operators in the twentieth century*

Barry Simon^{a)}

*Division of Mathematics 253-37, California Institute of Technology,
Pasadena, California 91125-3700*

(Received 4 February 2000; accepted for publication 24 February 2000)

This paper reviews the past fifty years of work on spectral theory and related issues in nonrelativistic quantum mechanics. © 2000 American Institute of Physics.
[S0022-2488(00)01906-X]

CONTENTS

I.	INTRODUCTION.....	3523
II.	MATHEMATICAL TOOLS AND ISSUES.....	3524
III.	SELF-ADJOINTNESS.....	3525
IV.	PROPERTIES OF EIGENFUNCTIONS, GREEN'S FUNCTIONS, SEMIGROUPS, AND ALL THAT.....	3529
V.	ONE-DIMENSIONAL DECAYING POTENTIALS.....	3531
VI.	INVERSE SPECTRAL THEORY.....	3534
VII.	ERGODIC POTENTIALS.....	3536
VIII.	TWO-BODY HAMILTONIANS.....	3538
IX.	<i>N</i> -BODY HAMILTONIANS.....	3540
X.	CONSTANT ELECTRIC AND MAGNETIC FIELDS.....	3543
XI.	COULOMB ENERGIES.....	3544
XII.	EIGENVALUE PERTURBATION THEORY.....	3545
XIII.	RESONANCES.....	3546
XIV.	THE QUASICLASSICAL LIMIT.....	3547
	REFERENCES.....	3548

I. INTRODUCTION

The twentieth century is the century of science. In a century that has seen special and general relativity, quantum electrodynamics and chromodynamics, a total revamping of our understanding of molecules and of the cosmos, plate tectonics, and the rise of microbiology, one can make the case that the most spectacular scientific development was the discovery of nonrelativistic quantum mechanics in the first quarter of the century. Its aftermath not only changed the physicist's view of matter, but it set the stage for the revolutions in chemistry, our understanding of stars, biology, and practical electronics.

In what is one of the more striking cases of serendipity, just as Heisenberg and Schrödinger were discovering the "new" quantum theory, von Neumann was developing the theory of unbounded self-adjoint operators and Weyl the representations of compact Lie groups—two subjects of great relevance to the mathematics underlying nonrelativistic quantum mechanics. In short order they produced books (von Neumann²⁷¹ and Weyl²⁷⁵) that used this mathematics to give a mathematical foundation to the framework of quantum mechanics. With later additions, notably by Bargmann, Wigner, and Mackey, the basic foundations are mathematically firm.

This is analogous to having formulated classical mechanics as Hamiltonian flows on symplec-

*Dedicated to Tosio Kato (1917–1999), father of the modern theory of Schrödinger operators.

^{a)}Electronic mail: bsimon@caltech.edu

tic manifolds. What remains is what might be called the second-level foundations—existence of solutions of the time-dependent Schrödinger equation (which is equivalent to self-adjointness of these operators) and general qualitative issues in dynamics. It is this subject, essentially born 50 years ago, that I will review here. The subject matter is vast with hundreds of contributors and thousands of papers. Each section of this paper is a proxy for what deserves a book or at least a very long review article. In attempting to overview such a vast area in a few pages, I have had to focus on the high points. No proofs are given and I have settled for usually quoting the initial or especially significant papers. I have no doubt that I have left out some important papers, and if so, I ask the forgiveness of the reader (and their authors!).

To keep this paper a reasonable size, I have focused almost entirely on the general basics of Schrödinger operators and some simple applications to atomic and molecular Hamiltonians. That means, among other areas, I have not considered general second-order operators on \mathbb{R}^n and on general manifolds (but see Davies and Safarov,⁵⁷ Davies,⁵⁵ and Kenig¹⁵⁴) nor have I considered some of the detailed papers on perturbations of Hamiltonians with periodic potential (see, e.g., Deift and Hempel⁵⁸ and Gesztesy and Simon⁹¹) nor the extensive literature on Dirac operators nor the considerable work on Schrödinger operators in a bounded region with some boundary conditions including subtle results on what happens at irregular boundary points (see Davies⁵⁵) nor the results on phenomena like the quantum Hall effect that apply and extend the general theory to results in condensed matter physics. While there are a few results about $-\Delta + V$ for cases where $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$, again there is a large literature we will not attempt to review. While Sec. X has a brief discussion of constant magnetic field, we have not attempted to discuss the recent extensive literature on nonconstant magnetic fields.

There is a companion piece to this one on open problems.²⁶⁰

II. MATHEMATICAL TOOLS AND ISSUES

The mathematics most relevant to the modern theory of Schrödinger operators is functional, real, harmonic, and complex analysis. In this section, we will briefly set the stage to fix notation. For more details, see Reed and Simon.^{214,211}

Quantum Hamiltonians are unbounded operators, defined on a dense subspace rather than the whole Hilbert space. Physics books tend to emphasize the symmetry (“Hermiticity”) of the Hamiltonian; that is, that $\langle H\varphi, \psi \rangle = \langle \varphi, H\psi \rangle$ for all φ, ψ in $D(H)$. But more important is a property called self-adjointness. The adjoint H^* of an operator H is defined to be the maximal operator so that $\langle H^*\varphi, \psi \rangle = \langle \varphi, H\psi \rangle$ for all $\psi \in D(H)$, $\varphi \in D(H^*)$. Hermiticity says only that H^* is an extension of H .

We say H is self-adjoint if $H = H^*$, H is called essentially self-adjoint if and only if H is symmetric and has a unique self-adjoint extension. This holds if and only if H^* is self-adjoint. Self-adjointness is important in the first place because if H is self-adjoint, one can form the unitary group e^{-itH} and so solve $i\dot{\varphi}_t = H\varphi_t$ (as $\varphi_t = e^{-itH}\varphi$) for initial conditions $\varphi \in D(H)$. Indeed, Stone’s theorem says that any one-parameter continuous unitary group is associated with a self-adjoint operator. Second, self-adjointness implies the spectral theorem. There is for each Borel set $A \subset \mathbb{R}$, a projection, $E_A(H)$, so that $H = \int \lambda dE_\lambda$ and $e^{-itH} = \int e^{-it\lambda} dE_\lambda$. One defines spectral measures $d\mu_\varphi^H$ by

$$\mu_\varphi^H(A) = (\varphi, E_A(H)\varphi) \quad (\text{II.1})$$

so that

$$\int e^{-it\lambda} d\mu_\varphi^H(\lambda) = (\varphi, e^{-itH}\varphi) \quad (\text{II.2})$$

and

$$\int \frac{d\mu_\varphi^H(\lambda)}{\lambda - z} = (\varphi, (H - z)^{-1} \varphi). \tag{II.3}$$

$\sigma(H)$, the spectrum of H , is precisely $\cup_\varphi \text{supp}(d\mu_\varphi^H)$.

Much of what we discuss in this paper involves two distinct decompositions of the spectrum of H . The first is

$$\sigma_{\text{disc}}(H) = \{\lambda \mid \lambda \text{ is an eigenvalue of finite multiplicity and an isolated point of } \sigma(H)\}$$

$$\sigma_{\text{ess}}(H) = \sigma(H) \setminus \sigma_{\text{disc}}(H).$$

Equivalently, $\lambda \in \sigma_{\text{disc}}(H)$ if and only if for some $\varepsilon > 0$, $\dim E_{(\lambda - \varepsilon, \lambda + \varepsilon)}(H)$ is finite and for all $\varepsilon > 0$, $E_{(\lambda - \varepsilon, \lambda + \varepsilon)}(H) \neq 0$. $\sigma_{\text{disc}}(H)$ captures the notion of bound states.

The second breakup involves the fact that any measure $d\mu$ on \mathbb{R} has a decomposition

$$d\mu = d\mu_{\text{pp}} + d\mu_{\text{ac}} + d\mu_{\text{sc}},$$

where $d\mu_{\text{pp}}$ is a pure point measure (sum of delta functions), $d\mu_{\text{ac}}$ is $F(\lambda)d\lambda$, with F a non-negative locally integrable density, and $d\mu_{\text{sc}}$ is a singular continuous measure (like the Cantor measure). I will define $\sigma_{\text{pp}}(H)$ to be the set of eigenvalues of H ; it is not the union of the supports of μ_{pp} because it may not be closed

$$\sigma_{\text{ac}}(H) = \bigcup_\varphi \text{supp}(d\mu_\varphi^H)_{\text{ac}},$$

$$\sigma_{\text{sc}}(H) = \bigcup_\varphi \text{supp}(d\mu_\varphi^H)_{\text{sc}}.$$

One often defines a refined set Σ_{ac} with $\bar{\Sigma}_{\text{ac}} = \sigma_{\text{ac}}(H)$, the essential support of the ac measure. Basically, the essential support of the a.c. measure $F(\lambda)d\lambda$ is $\{\lambda \mid F(\lambda) \neq 0\}$. It is defined modulo sets of Lebesgue measure zero. Σ_{ac} is the union of the essential support of $(d\mu_\varphi^H)_{\text{ac}}$ over a countable dense set of φ s.

III. SELF-ADJOINTNESS

The theory of Schrödinger operators was born with Kato's famous self-adjointness theorem for atomic Hamiltonians. His theorem abstracted states the following:

Theorem III.1: (Kato¹⁴⁴) Let $\mathcal{H} = L^2(\mathbb{R}^{3N})$ where $x \in \mathbb{R}^{3N}$ is written (x_1, \dots, x_N) with $x_i \in \mathbb{R}^3$. Let Δ_i be the Laplacian in x_i and let V_i, V_{ij} be functions on \mathbb{R}^3 in $L^2(\mathbb{R}^3) + L^\infty(\mathbb{R}^3)$. Let

$$H_0 = - \sum_{i=1}^N (2\mu_i)^{-1} \Delta_i, \tag{III.1}$$

$$V = \sum_{i=1}^N V_i(x_i) + \sum_{i < j} V_{ij}(x_i - x_j) \tag{III.2}$$

and let $H = H_0 + V$. Then H defined on $D(H_0)$ is self-adjoint and is essentially self-adjoint on $C_0^\infty(\mathbb{R}^{3N})$.

Remarks:

(1) See Reed and Simon²¹¹ for a proof.

(2) The basic idea of the proof is a perturbation theoretic one. There is a general theorem (the Kato–Rellich theorem) that if A is a self-adjoint operator and B is a symmetric operator with $D(B) \supset D(A)$ and for some $\alpha < 1$ and $\beta > 0$ and all $\varphi \in D(A)$, that

$$\|B\varphi\| \leq \alpha\|A\varphi\| + \beta\|\varphi\|, \tag{III.3}$$

then $A + B$ is self-adjoint on $D(A)$ and essentially self-adjoint on any domain of essential self-adjointness for A . If (III.3) holds, we will say B is A bounded. The infimum over all α is called the relative bound of B with respect to A .

(3) If one looks at a general bound of type (III.3) with $\alpha < 1$ where $A = -\Delta$ on $L^2(\mathbb{R}^k)$ and B is multiplication by V , then in terms of requirements that $V \in L^p_{loc}(\mathbb{R}^k)$, one needs

$$p \geq 2 \quad k = 1, 2, 3 \tag{III.4a}$$

$$p > 2 \quad k = 4 \tag{III.4b}$$

$$p \geq \frac{k}{2} \quad k \geq 5 \tag{III.4c}$$

by using Sobolev estimates (see, e.g., Cycon *et al.*⁵³).

(4) If $k = 3N$ and we use *only* the L^p requirements of Remark 3, Coulomb potentials stop working already at $N = 2$. Thus, for Kato's theorem, it is critical to use Sobolev estimates in subsets of variables as Kato did.

An industry developed in understanding when $-\Delta + V$ is essentially self-adjoint on $C^\infty_0(\mathbb{R}^n)$. An illustrative example is

Example: Let $H = -\Delta - c|x|^{-2}$ on $C^\infty_0(\mathbb{R}^k)$ with $n \geq 5$ (needed for $H\varphi \in L^2$ for all $\varphi \in C^\infty_0(\mathbb{R}^k)$). Then it can be seen (Reed and Simon,²¹¹ Example 4 in Sec. X.2) that if $c > c_0 = (n - 4)n/4$, then H is not self-adjoint on C^∞_0 . This is a quantum analog of the classical fact that if $V = -c|x|^{-2}$ for any $c > 0$, a set of initial conditions of positive measure falls into $x = 0$ in finite time ($c_0 > 0$ is a reflection of an uncertainty principle repulsion).

This example shows that for pure L^p requirements, one cannot do better than (III.4) since $|x|^{-2} \in L^p + L^\infty$ if $p < k/2$. But it turns out this is only so if V is allowed to have any sign. For $V \geq 0$, one can do much better. The best result of this genre is

Theorem III.2: (Leinfelder and Simader¹⁷³) Let $V \geq 0$, $V \in L^2_{loc}(\mathbb{R}^k)$, $\{a_j\}_{j=1}^k \in L^4_{loc}(\mathbb{R}^k)$ with $\nabla \cdot a \in L^2_{loc}(\mathbb{R}^k)$ (distributional derivatives). Then

$$H = \sum_{j=1}^k (i\partial_j - a_j)^2 + V \tag{III.5}$$

is essentially self-adjoint on $C^\infty_0(\mathbb{R}^k)$.

Remarks:

(1) For a proof, see Cycon *et al.*⁵³

(2) This is essentially a best possible result. If $a = 0$, H is defined on C^∞_0 if and only if $V \in L^2_{loc}$; so the result says for positive V , we have essential self-adjointness if and only if H is defined. Similarly, unless there are cancellations, $a_j \in L^4_{loc}$ and $\nabla \cdot a \in L^2_{loc}$ is required for H to be defined on C^∞_0 .

(3) It was Simon²³⁹ who first realized that for $V \geq 0$, there only needed to be local L^2 conditions. However, he required a global condition $\int |V(x)|^2 e^{-bx^2} dx < \infty$ for some $b > 0$. It was Kato¹⁵² who proved the general $a = 0$ result (and also allowed for smooth a 's). Kato's paper used the distributional inequality, now called Kato's inequality

$$\Delta|u| \geq \text{Re}(\text{sgn } u \Delta u) \tag{III.6}$$

that is also critical to the Leinfelder–Simader proof.

(4) (III.6) is essentially equivalent to the fact that $e^{t\Delta}$ is positivity preserving. The version of (III.6) with magnetic fields is equivalent to diamagnetic inequalities:

$$|(e^{-tH}\varphi)(x)| \leq (e^{t\Delta}|\varphi|)(x) \tag{III.7}$$

for the H of (III.5) (with $V \geq 0$). These ideas were discovered by Nelson,¹⁹⁸ Simon,^{241,247} and Hess *et al.*¹¹⁹

While there are best possible self-adjointness results for magnetic fields and positive potentials, the results for V 's which can be negative are not in such a definitive form. All the basic principles are understood but I am not aware of a single result that puts them all together (one of the best results is in Kato's paper¹⁵¹ although, as we will see, it is not quite optimal with regard to local singularities). So I will present the general principles that are understood in this case.

(a) $-|x|^2$ *borderline for behavior at infinity*. Negative potentials V of compact support for which $H = -\Delta + V$ is essentially self-adjoint on C_0^∞ normally obey a global estimate of the form (III.3) (with $A = -\Delta$, $B = V$) and, in particular, H is bounded from below. However, if V is not of compact support, it can go to minus infinity at infinity without destroying self-adjointness. More or less, the borderline for keeping self-adjointness is $-|x|^2$. For example, it can be proven (see, e.g., Reed and Simon,²¹¹ Theorem X.9) that $-(d^2/dx^2) - |x|^\alpha$ on $L^2(-\infty, \infty)$ is essentially self-adjoint on $C_0^\infty(-\infty, \infty)$ if and only if $\alpha \leq 2$. This is attractive since a classical particle with the same potential reaches infinity in finite time if and only if $\alpha > 2$. Nelson has examples (see Reed and Simon,²¹¹ p. 156) of $V(x)$ with $V(x) \leq -cx^4$ so $-(d^2/dx^2) + V(x)$ is still essentially self-adjoint and thus, the borderline will not be if and only if, but the general version of this is that if $V(x) \geq -cx^2$ in some averaged sense, then $-\Delta + V(x)$ will be essentially self-adjoint on C_0^∞ . The earliest version of this is Ikebe and Kato.¹³⁰ My favorite theorem of this genre is due to Faris and Lavine⁸⁰ (see Reed and Simon,²¹¹ Theorem X.38). In particular, Stark Hamiltonians where $V = \mathbf{c} \cdot \mathbf{x} + V_0$ are essentially self-adjoint for suitable V_0 . In any event, I will focus henceforth on cases where $-\Delta + V$ is not unbounded from below.

(b) *Stability of relative boundedness under adding $V \geq 0$ or a magnetic field*. Suppose $A \geq 0$. Then (III.3) holds for some $\alpha < 1$ if and only if

$$\lim_{\gamma \rightarrow \infty} \|B(A + \gamma)^{-1}\| < 1.$$

On the other hand, (III.7) implies that for $V \geq 0$, any \mathbf{a} and any multiplication operator W :

$$\|W(H + \gamma)^{-1}\| \leq \|W(-\Delta + \gamma)^{-1}\|$$

and so the second principle is that in studying the negative part of V , one can assume V is negative and then add back an arbitrary positive L^2_{loc} positive V . While this is true, it ignores situations where there are cancellations between the positive and negative parts which can occur (see, e.g., Combes and Ginibre⁴⁸).

(c) *Relative bounds need only hold uniformly locally*. The following proposition holds:

Proposition III.3: *Suppose V is a function on \mathbb{R}^d so that for some α, β and every y ,*

$$\|V\chi(\cdot - y)\varphi\| \leq \alpha\|-\Delta\varphi\| + \beta\|\varphi\|, \tag{III.8}$$

where χ is the characteristic function of the unit cube. Then for any $\tilde{\alpha} > \alpha$, there is some $\tilde{\beta}$ so that

$$\|V\varphi\| \leq \tilde{\alpha}\|-\Delta\varphi\| + \tilde{\beta}\|\varphi\|. \tag{III.9}$$

This result is proven by a variant of an idea of Sigal.²³¹ Find a ‘‘partition of unity’’ $\{j_\mu\}_\mu$ so that $\sum j_\mu^2 = 1$, each j_μ is supported in some unit cube (so $j_\mu\chi(\cdot - y_\mu) = j_\mu$ for some j_μ), and the j_μ 's are locally finite, $\sum(\nabla j_\mu)^2$ is uniformly bounded (the j_μ 's can be translates of a single j_μ) and $\sum|\Delta j_\mu|$ is uniformly bounded. If $H_0 = -\Delta$, we have (where C is related to $\|\sum(\nabla j_\mu)^2\|_\infty$ and $\|\sum(\Delta j_\mu)\|_\infty$)

$$\sum_\mu [j_\mu, [j_\mu, H_0^2]] \leq C(H_0 + 1)$$

and from this that

$$\sum \|H_0 j_\mu \varphi\|^2 \leq (1 + \varepsilon) \|H_0 \varphi\|^2 + C_\varepsilon \|\varphi\|^2. \tag{III.10}$$

Thus

$$\begin{aligned} \|V\varphi\|^2 &= \sum_\mu \|V\chi(\cdot - y_\mu) j_\mu \varphi\|^2 \\ &\leq (1 + \varepsilon) \alpha^2 \sum_\mu \|H_0 j_\alpha \varphi\|^2 + (1 + \varepsilon^{-1}) \beta^2 \|\varphi\|^2 \quad (\text{by III.8}) \\ &\leq (1 + \varepsilon)^2 \alpha^2 \|H_0 \varphi\|^2 + ((1 + \varepsilon^{-1}) \beta^2 + C_\varepsilon) \|\varphi\|^2 \quad (\text{by III.10}) \end{aligned}$$

which yields (III.9).

Proposition III.3 states that the proper condition on V to yield a $-\Delta$ bound is a uniform local condition.

(d) *Convolution results are the proper local condition.* As discussed earlier, L^p conditions on V do not properly control functions on subspaces. Explicitly, let $\pi: \mathbb{R}^k \rightarrow \mathbb{R}^l$ be a projection and $V(x) = W(\pi(x))$. Then for V to be $-\Delta$ bounded (assuming $k \geq l \geq 5$), we need $W \in L^p_{\text{loc}}(\mathbb{R}^l)$ for $p \geq l/2$ and so $V \in L^p_{\text{loc}}(\mathbb{R}^k)$ with $p \geq l/2$. But if V is not a function of a subset of variables, in general we need $p \geq k/2$. It is a discovery of Stummel²⁶² that by stating conditions in terms of convolution estimates, one can find conditions that respect subsets of variables. In particular, the following is a space S_ν introduced in Stummel:²⁶² Let V be a function on \mathbb{R}^ν ; we say $V \in S_\nu$ if and only if

$$\begin{aligned} \lim_{\alpha \downarrow 0} \left[\sup_x \int_{|x-y| \leq \alpha} |x-y|^{4-\nu} |V(y)|^2 d^\nu y \right] &= 0 \quad \text{if } \nu \geq 5, \\ \lim_{\alpha \downarrow 0} \left[\sup_x \int_{|x-y| \leq \alpha} \ln(|x-y|^{-1}) |V(y)|^2 d^\nu y \right] &= 0 \quad \text{if } \nu = 4 \\ \sup_x \int_{|x-y| \leq 1} |V(y)|^2 d^\nu y &< \infty \quad \text{if } \nu \leq 3. \end{aligned}$$

This class respects functions of subvariables in the sense that if $\pi: \mathbb{R}^k \rightarrow \mathbb{R}^l$ is a projection, $V(x) = W(\pi(x))$ and $W \in S_l$, then $V \in S_k$. Moreover, it is not hard to show (see, e.g., Cycon *et al.*⁵³) that if $V \in S_\nu$, then V is $-\Delta$ bounded with relative bound zero. Moreover (see Cycon *et al.*,⁵³ Theorem 1.9), if for some $a, b > 0$ and δ with $0 < \delta < 1$ and all $0 < \varepsilon < 1$ and $\varphi \in D(H_0)$

$$\|V\varphi\|^2 \leq \varepsilon \|\Delta\varphi\|^2 + a \exp(b\varepsilon^{-\delta}) \|\varphi\|^2, \tag{III.11}$$

then V is in S_ν . See Schechter²²⁶ for more on Stummel conditions.

(3) *The Kato class and going beyond relative boundedness.* In his inequality paper,¹⁵² Kato introduced a form analog K_ν of S_ν : Let V be a function on \mathbb{R}^ν ; we say $V \in K_\nu$ if and only if

$$\lim_{\alpha \downarrow 0} \left[\sup_x \int_{|x-y| \leq \alpha} |x-y|^{2-\nu} |V(y)| d^\nu y \right] = 0 \quad \text{if } \nu \geq 3, \tag{III.12a}$$

$$\lim_{\alpha \downarrow 0} \left[\sup_x \int_{|x-y| < \alpha} \ln(|x-y|^{-1}) |V(y)| d^\nu y \right] = 0 \quad \text{if } \nu = 2, \tag{III.12b}$$

$$\sup_x \int_{|x-y| \leq 1} |V(y)| d^\nu y < \infty \quad \text{if } \nu = 1. \tag{III.12c}$$

Then Kato¹⁵¹ proved if $\max(-V, 0) \in K_\nu$ and $V \in L^2_{\text{loc}}(\mathbb{R}^\nu)$, then $-\Delta + V$ is essentially self-adjoint on $C^\infty_0(\mathbb{R}^\nu)$. While it is not Kato's proof, this is intimately connected with the semigroup result discussed in Sec. IV. Defining the form sum H , one knows $\exp(-tH): L^2 \rightarrow L^\infty$ so $L^\infty \cap L^2 \cap D(H)$ is a domain of essential self-adjointness. It is not hard to then show $L^\infty \cap D(H)$, the L^∞ functions of compact support are a domain of essential self-adjointness. Then convolution allows one to get C^∞_0 approximations.

(f) *Logarithmic improvements.* Neither S_ν nor K_ν is quite the ideal space for essential self-adjointness. For example, if $\nu \geq 5$ and $V(x) = |x|^{-2}(1 + |\log|x||)^{-\alpha}$, V is in K_ν only if $\alpha > 1$, in S_ν only if $\alpha > \frac{1}{2}$, but $-\Delta$ bounded with relative bound zero if $\alpha > 0$.

Analogous to the issue of self-adjointness is a question of whether maximal and minimal forms agree. This is discussed in Kato¹⁵² and Simon²⁴⁸ (see Theorem 1.13 in Cycon *et al.*⁵³).

IV. PROPERTIES OF EIGENFUNCTIONS, GREEN'S FUNCTIONS, SEMIGROUPS, AND ALL THAT

I wrote a long review of these subjects 20 years ago (Simon²⁵⁰) and the situation has hardly changed since then, although there has been extensive interesting work on what happens for general elliptic operators and for bounded regions (see, e.g., Davies⁵⁵). So it will suffice to hit a few major themes. The basic theorem is

Theorem IV.1: *Let $V_+ \in L^1_{\text{loc}}(\mathbb{R}^\nu)$ and $V_- \in K_\nu$, the space of (III.12). Let $H = -\Delta + V$ as a form sum. Then for any $p \leq q, e^{-tH}$ maps L^p to L^q and for $t \leq 1$,*

$$\|e^{-tH}\|_{p,q} \leq Ct^{-\alpha}, \tag{IV.1}$$

where

$$\alpha = \frac{\nu}{2} \left(\frac{1}{p} - \frac{1}{q} \right). \tag{IV.2}$$

Remarks:

(1) Semigroup L^p bounds were first found by Davies,⁵⁴ Herbst and Sloan,¹¹⁸ and Kovalenko and Semenov¹⁶¹ with further developments by Carmona,⁴¹ Simon,²⁴⁶ and Aizenman and Simon.¹²

(2) In particular, it was Aizenman and Simon¹² who found that K_ν is the natural class for L^p bounds. Indeed, they not only proved Theorem IV.1 in this form but also showed that if $V \leq 0$ and $\exp(-tH)$ maps L^∞ to itself with $\lim_{t \downarrow 0} \|e^{-tH}\|_{\infty, \infty} = 1$, then $V \in K_\nu$.

(3) The result holds when magnetic fields are added (by a diamagnetic inequality).

(4) Most of these authors use a combination of path integral estimates and L^p interpolation theory. In particular, the Feynman–Kac and Feynman–Kac–Itô formulas (see Simon²⁴⁶ for extensive discussion) are useful tools in studying Schrödinger operators. See Simon²⁵⁹ for an extension to cases when $V(x) \geq -cx^2$.

(5) In fact, e^{-tH} takes L^p not only into L^∞ but into the continuous functions (see Simon,²⁵⁰ Theorem B.3.1).

(6) (IV.1)/(IV.2) are precisely the best results for $H = -\Delta$.

(7) This theorem says that H can be defined as the generator of a semigroup on each L^p space. The spectrum has been shown to be L^p independent in Hempel and Voight.¹¹³ For a general discussion of L^p Schrödinger operators, see Davies.⁵⁶

Once one has these estimates, they can be used to derive:

(a) *Sobolev estimates*: As in the free case if V obeys the conditions of Theorem IV.1, then $(H-z)^{-\nu}$ takes L^p to L^q if

$$p^{-1} - q^{-1} < \left(\frac{2\alpha}{\nu} \right). \tag{IV.3}$$

The result (see Simon,²⁵⁰ Theorem B.2.1) is obtained by integrating the semigroup bound. (IV.3) comes from (IV.2) and the requirement of integrability at $t=0$.

(b) *Integral kernels*: Bounded operators from L^1 to L^∞ have bounded integral kernels and so Theorem IV.1 can be used (see Simon,²⁵⁰ Theorem B.7.1) to prove e^{-tH} , $(H-z)^{-\alpha}$ ($\alpha > \nu/2$) are integral operators with continuous integral kernels. One can also show (Simon,²⁵⁰ Theorem B.7.2) that for $0 < \alpha < \nu/2$, $(H-z)^{-\alpha}$ is an integral operator with an integral kernel that is continuous away from $x=y$ with a precise singularity at $x=y$.

(c) *Eigenfunctions*: Since global eigenfunctions (i.e., $\varphi \in L^2$ that obey $H\varphi = E\varphi$) are in $\text{Ran}(e^{-tH})$, Theorem IV.1 implies such eigenfunctions are in L^∞ . In fact, all this can be done locally. Any eigenfunction (distributional solution of $H\varphi = E\varphi$) is automatically continuous and one can prove Harnack inequalities and subsolution estimates. This is discussed in detail in Aizenman and Simon¹² and Simon.²⁵⁰

We end this section with a discussion of some issues involving eigenfunctions. There is much literature on when Schrödinger operators have positive solutions. This was begun by Allegretto¹³ and Piepenbrink²⁰⁶ with later results by Agmon⁵ and Pinchover.²⁰⁷

Here is a typical theorem (Simon [Ref. 250, Theorem C.8.1]):

Theorem IV.2: *Let $V_- \in K_\nu$ and $K_+ \in K_\nu^{\text{loc}}$. Then $Hu = Eu$ has a nonzero distributional solution which is everywhere positive if and only if $\inf \text{spec}(H) \geq E$.*

There is also much literature on the issue of exponential decay of eigenfunctions. One result (see Simon²⁵⁰, Theorem C.3.1) says that any L^2 eigenfunction actually goes to zero pointwise—of interest only for eigenfunctions of embedded eigenvalues. For discrete spectrum, the decay is at least exponential under minimal regularity hypothesis on V . The original key papers on this theme are by O'Connor²⁰⁰ and Combes and Thomas.⁴⁷ From their ideas, one obtains (see Sec. C.3 of Simon);²⁵⁰

Theorem IV.3: *Let $V_- \in K_\nu$, $V_+ \in K_\nu^{\text{loc}}$ and let $H = -\Delta + V$ and let $Hu = Eu$ with $u \in L^2$. Then*

$$|u(x)| \leq C e^{-A|x|}, \tag{IV.4}$$

where:

- (i) For general E in the discrete spectrum, (IV.4) holds for some $A > 0$ and $C > 0$.
- (ii) If H has compact resolvent, then (IV.4) holds in the sense for any $A > 0$, there is a suitable $C > 0$.
- (iii) If $\Sigma_{\text{ess}} = \inf \sigma_{\text{ess}}(H)$ and $E < \Sigma_{\text{ess}}$, then (IV.4) holds in the sense that for any $A \leq \sqrt{E - \Sigma_{\text{ess}}}$, there is a suitable $C > 0$.

One can go beyond this to get fairly detailed behavior on decay in cases when H has compact resolvent or for N -body potentials. In one dimension, one can justify under some regularity conditions the WKB formula that states when $V(x) \rightarrow \infty$, eigenfunctions decay like

$$V(x)^{-1/4} \exp\left(-\int_a^x \sqrt{V(y) - E} dy\right). \tag{IV.5}$$

It was Agmon⁴ who realized the proper higher-dimensional analog for this involves what is now called the Agmon metric: $\rho(x)$ is the geodesic distance of x to 0 in the Riemannian metric

$\rho_{ij}(x) = \delta_{ij}(V(x) - E)_+ d^2x$. There is a related but more subtle definition for N -body systems. See Agmon⁴ and Deift *et al.*⁵⁹ for further discussions. See Simon²⁵³ and Helffer and Sjöstrand¹¹¹ for an application to tunneling probabilities.

Eigenfunctions play a critical role in explicit spectral representations of Schrödinger operators. The basic ideas go back to work of Browder,³⁹ Garding,⁸⁸ Gel'fand,⁸⁹ Kac,¹³⁷ and especially Berezanskii.^{29,30} See Sec. C.5 of Simon²⁵⁰ and Last and Simon¹⁷⁰ for some additional one-dimensional results.

Finally, we mention issues of cusps and nodes of eigenfunctions. Kato¹⁴⁸ has a famous paper on cusps at Coulomb singularities for atomic eigenfunctions. See Hoffmann-Ostenhof *et al.*^{108,120,121} for recent developments in this area.

V. ONE-DIMENSIONAL DECAYING POTENTIALS

One-dimensional Schrödinger operators

$$-\frac{d^2}{dx^2} + V(x) \tag{V.1}$$

on $L^2(-\infty, \infty)$ and $L^2(0, \infty)$ and their discrete analogs

$$hu(n) = u(n+1) + u(n-1) + V(u)u(n) \tag{V.2}$$

on $l^2(-\infty, \infty)$ and $l^2[0, \infty)$ have been heavily studied for two reasons. First, ordinary differential equation (ODE)/difference equation methods allow one to study them in much greater detail than one can the higher-dimensional analogs. Second, if $V(x) = V(|x|)$ is a spherically symmetric function on \mathbb{R}^p , then $-\Delta + V$ is unitarily equivalent to a direct sum of operators on $L^2(0, \infty)$ or the form (V.1) where the effective V 's have the form $V_l(x) = \kappa_l|x|^{-2} + V(x)$ for suitable κ_l 's. The details can be found, for example, in Reed and Simon,²¹¹ Example 4 to the Appendix for X.1.

The one-dimensional theory has been in and out of vogue. It was extensively studied from 1930 to 1950 with important contributions by Titchmarsh, Kodaira, Gel'fand, Hartman–Wintner, Levinson, Coddington–Levinson, and Jost. Significant developments during the next 25 years were mainly in the area of inverse spectral theory (a major exception was Weidmann's work,²⁷³ to be discussed shortly) which will be discussed in Sec. VI. From about 1975 starting with the work of Goldsheid *et al.*⁹⁸ and Pearson,²⁰⁴ this has been an active area with extensive study of the one-dimensional case, especially with long-range and with ergodic potentials.

One special feature of one dimension is that one can limit spectral multiplicities under very general conditions on V :

Theorem V.1:

- (a) Let $H = -(d^2/dx^2) + V(x)$ on $L^2(0, \infty)$ with fixed $hu(0) + u'(0) = 0$ boundary conditions and suppose H is essentially self-adjoint on $C_0^\infty[0, \infty)$. Then H has simple spectrum (multiplicity 1).
- (b) Let $H = -(d^2/dx^2) + V(x)$ on $L^2(-\infty, \infty)$ and suppose H is essentially self-adjoint on $C_0^\infty(-\infty, \infty)$. Then
 - (i) The absolutely continuous spectrum of H is of multiplicity at most 2.
 - (ii) The singular spectrum of H is of multiplicity 1.

Remarks:

- (1) All one needs for local regularity of V is $V \in L^1[0, R]$ for all $R > 0$ or $L^1_{loc}(-\infty, \infty)$.
- (2) The result holds even if H is not essentially self-adjoint (V limit circle at $\pm\infty$) so long as a boundary condition is imposed at ∞ or at $-\infty$.
- (3) The only subtle part of the result is that the singular continuous spectrum is simple on the real line. This is a theorem of Kac;^{138,139} see also Berezanskii.^{29,30} My preferred proof is due to Gilbert.^{95,96}

In this section, we will discuss the case where $V(x) \rightarrow 0$ at infinity. In Sec. VI, we will discuss inverse spectral theory, and in Sec. VII, we will discuss ergodic potentials. (These two subjects are mainly one dimensional.) The issue of the asymptotic eigenvalue distribution when $V \rightarrow \infty$ as $\pm\infty$ is discussed in Sec. XIV on the quasiclassical limit.

This section will discuss (V.1)/(V.2) in situations where $V(x)$ (or $V(n)$) goes to zero (at least in an average sense) as $x \rightarrow \infty$ (or $n \rightarrow \infty$). The interesting thing is that there are three natural breaks in behavior. Expressed in terms of $|x|^{-\alpha}$ behavior, they are

- (i) At $\alpha=2$, we shift between a finite number of bound states ($\alpha>2$) or an infinite number ($\alpha<2$) at least if $V(x)<0$.
- (ii) At $\alpha=1$ ($V \in L^1$), we shift between a pure scattering situation for positive energies ($\alpha > 1$) and the possibility of positive energy bound states ($\alpha < 1$).
- (iii) At $\alpha = \frac{1}{2}$, ($V \in L^2$), we shift from there being a.c. spectrum for almost everywhere positive energy ($\alpha > \frac{1}{2}$) to at least the possibility of very different spectrum.

(i) and (ii) have been known since the earliest days of quantum mechanics. The $\alpha = \frac{1}{2}$ borderline first occurred in Simon²⁵¹ who found that random decay potentials had point spectrum when $\alpha < \frac{1}{2}$. Delyon *et al.*⁶⁴ then showed if $\alpha = \frac{1}{2}$, there may be some nonpoint spectrum. As we will see, subsequent results confirmed this borderline.

The negative spectrum for decaying potentials is easy: So long as $\int_x^{x+1} |V(y)| dy \rightarrow 0$, H is bounded below and has $[0, \infty)$ as essential spectrum by Weyl's criterion (see, e.g., Reed and Simon,²¹³ Sec. XIII.4), which means that $(-\infty, 0)$ has only discrete eigenvalues of finite multiplicity, which can only accumulate at energy 0. Indeed, by Theorem V.1, the point spectrum is of multiplicity 1. Once these basics are established for the discrete spectrum, a number of detailed questions about it arise:

(a) *Is σ_{disc} finite or infinite?* The borderline, as mentioned above, is r^{-2} decay. Explicitly, one has Bargmann's bound²⁴ that the number of eigenvalues on a half line with $u(0)=0$ boundary conditions is bounded by $\int x|V(x)| dx$ and on a whole line by $1 + \int_{-\infty}^{\infty} |x||V(x)| dx$ (see Simon²⁴⁰ for a review of bounds on the number of bound states). On the other hand, if $\lim_{x \rightarrow \infty} |x|^2 V(x) \leq -\frac{1}{4}$, one can prove that H has an infinity of bound states (see, e.g., Reed and Simon,²¹³ Theorem XIII.6).

(b) *If σ_{disc} is infinite, how does $\lim_{\lambda \uparrow 0} \dim E_{(-\infty, \lambda)}(H)$ diverge?* This is a quasiclassical limit and discussed in Sec. XIV.

(c) *Bounds on moments of eigenvalues.* Lieb and Thirring,¹⁸⁶ motivated in part by their work on the stability of matter,¹⁸⁵ initiated extensive study on the best constant $L_{\gamma,1}$ in

$$\sum_j |e_j|^\gamma \leq L_{\gamma,1} \int |V(x)|^{\gamma+1/2} dx,$$

which holds if $\gamma \geq \frac{1}{2}$. Here $\{e_j\}$ are the negative eigenvalues of H . For $\gamma \geq \frac{3}{2}$, the constant $L_{\gamma,1}$ is known to be quasiclassical (Aizenman and Lieb).⁹ For $\gamma \in [\frac{1}{2}, \frac{3}{2})$, it is known that $L_{\gamma,1}$ is strictly larger than the quasiclassical result.¹⁸⁶ It is conjectured to be the optimal value for a single bound state, as explained in Lieb and Thirring,¹⁸⁶ but this is still open (except at $\gamma = \frac{1}{2}$ (Hundertmark *et al.*¹²⁵).

(d) *Is there a bound state for weak coupling?* In one (and two) dimensions, H has bound states even for very weak coupling. The result (Simon²⁴²) is that if $\int |x||V(x)| dx < \infty$ and $\int V(x) dx \leq 0$ and $V \neq 0$, then H always has a bound state and the binding energy of $-\Delta + \mu V$ is $\sim c \mu^2$ as $\mu \downarrow 0$ (if $\int V(x) dx < 0$; it is $\sim c \mu^4$ if $\int V(x) = 0$).

As for positive energies, the situation is simple if $V \in L^1$:

Theorem V.2: *Let $V \in L^1(-\infty, \infty)$ or $L^1(0, \infty)$. Then $HE_{(0, \infty)}(H)$ is unitarily equivalent to $-d^2/dx^2$ (on $L^2(-\infty, \infty)$ or $L^2(0, \infty)$ with $u(0)=0$ boundary conditions).*

Remarks:

- (1) This result is essentially due to Titchmarsh.²⁶⁷
- (2) In terms of $r^{-\alpha}$ falloff, $V \in L^1$ means $\alpha > 1$.

(3) Using scattering theoretic ideas, one can prove wave operators exist and are complete (see Sec. VIII).

(4) This says there is no point of singular continuous spectrum at positive energies and that the a.c. spectrum has essential support $(0, \infty)$ with multiplicity 2 or 1.

(5) We have stated the result for $u(0)=0$ boundary condition for simplicity; it holds for all boundary conditions at 0.

As for slower decay than L^1 , if one has control of derivatives, one can still conclude the positive spectrum is purely absolutely continuous. The simplest result of this genre is

Theorem V.3: (Weidmann²⁷³) *Let $V = V_1 + V_2$ where V_1 is in L^1 , $V_2(x) \rightarrow 0$ as $x \rightarrow \pm\infty$, and V_2 is of bounded variation. Then, $HE_{(0,\infty)}(H)$ is unitarily equivalent to $-d^2/dx^2$ (on $L^2(-\infty, \infty)$ or on $L^2(0, \infty)$ with $u(0)=0$ boundary conditions).*

Remarks:

- (1) V_2 of bounded variation with $V_2 \rightarrow 0$ at infinity essentially says that $-dV_2/dx \in L^1$; in fact, any V_2 of bounded variation can be written $V_3 + V_4$ with $V_3 \in L^1$ and V_4 a C^1 function with $dV_4/dx \in L^1$.
- (2) Pure power potentials $r^{-\alpha}$ for any $\alpha > 0$ are included in this theorem; indeed, any monotone function $V(x)$ with $V(x) \rightarrow 0$ as $x \rightarrow \infty$ is of bounded variation.

For a short proof of Theorems V.2/V.3, see Simon.²⁵⁶ Both theorems can be understood as coming from the fact that all solutions of $-u'' + Vu = \lambda u$ with $\lambda > 0$ are bounded. That such a conclusion implies the spectrum is purely absolutely continuous was first indicated by Carmona⁴² (who required some kind of uniformity in λ). Important later developments that capture this idea are due to Gilbert and Pearson,⁹⁷ Last and Simon,¹⁷⁰ and Jitomirskaya and Last.¹³⁵ The tools in those papers are also important for the proofs of the results of Sec. VII.

Once one allows decay slower than $r^{-1-\epsilon}$ for both V and V' , the conclusion of Theorems V.2/V.3 can fail because of embedded point spectrum. The original examples of this were found by von Neumann and Wigner.²⁷² Basically, if $V(x) = \gamma|x|^{-1} \sin(x)$ for x large and $\gamma > 1$, then $-u'' + Vu = \frac{1}{4}u$ has a solution which is L^2 at infinity (see, e.g., Theorem XI.67 in Reed and Simon²¹³). By adjusting V at finite x , one can arrange for any boundary condition one wants at $x=0$. In fact, if one allows slightly slower decay than $|x|^{-1}$, one can arrange dense point spectrum. Naboko¹⁹⁷ and Simon²⁵⁷ have shown that for any sequence $\{\lambda_n\}_{n=1}^\infty$ of energies in $(0, \infty)$ (Naboko has a mild restriction on the λ 's) and any $g(r)$ obeying $\lim_{r \rightarrow \infty} rg(r) = \infty$, there is a $V(x)$ obeying:

- (i) $|V(x)| \leq g(|x|)$ for x large;
- (ii) $-u'' + Vu = \lambda_n u$ has a solution L^2 at infinity and obeying a prescribed boundary condition at $x=0$.

Remark: It is an interesting open question about whether there exist potentials decaying faster than $|x|^{-1/2-\epsilon}$ with dense singular continuous spectrum (rather than dense point spectrum).

The interesting fact is that even though potentials of Naboko–Simon type have dense point spectrum, they may also have lots of a.c. spectrum. The best result is:

Theorem V.4: (Deift and Killip⁶⁰) *Let $V \in L^2$. Then the essential support of the a.c. spectrum of $H = -(d^2/dx^2) + V$ is $[0, \infty)$.*

Remarks:

- (1) In terms of $r^{-\alpha}$ decay, this result requires $\alpha > \frac{1}{2}$.
- (2) This result is optimal in that it is known for any Orlicz space strictly larger than L^2 in terms of behavior at infinity, there are V 's whose associated H has no a.c. spectrum.
- (3) The first result of this genre was found by Kiselev¹⁵⁶ who proved the conclusion of this theorem for $|V(x)| \leq Cx^{-3/4-\epsilon}$. There were subsequent improvements of this by Kiselev,¹⁵⁷ Christ and Kiselev,⁴⁶ and Remling.²¹⁸
- (4) Killip¹⁵⁵ has a partially alternate proof of Theorem V.4.

Once the decay is allowed to be slower than $r^{-1/2}$, one can have much different spectrum in $[0, \infty)$:

(i) If W is a suitable family of random homogeneous potentials and $V(x) = |x|^{-\alpha}W(x)$ with $\alpha < \frac{1}{2}$, then H has only dense point spectrum in $(0, \infty)$. This was first proven in the discrete case by Simon²⁵¹ and later in the continuum case by Kotani and Ushiroya.¹⁶⁰

(ii) Generic potentials decaying like $|x|^{-\alpha}$ ($\frac{1}{2} > \alpha > 0$) produce singular continuous spectrum as discovered by Simon.²⁵⁵ For example, in $\{V \in C(\mathbb{R}) \mid \sup_x |x|^\alpha |V(x)| \equiv \|V\|_\alpha\}$ viewed as a complete metric space in $\|\cdot\|_\alpha$, a dense G_δ of V 's are such that $-d^2/dx^2 + V(x)$ has purely singular continuous spectrum on $[0, \infty)$.

(iii) Much more is known in the borderline $\alpha = \frac{1}{2}$ case, at least for the discrete Schrödinger operator (V.2). For example, if a_n are independent, identically distributed random variables uniformly distributed in $[-1, 1]$ and $V(n) = \mu n^{-1/2}a_n$, then for suitable coupling constants μ and energies E in $[-2, 2]$, the spectral measures have fractional Hausdorff dimension with an exactly computable local dimension. This is discussed in Kiselev *et al.*¹⁵⁸ There are earlier results on this model by Delyon *et al.*⁶⁴ and Delyon.⁶²

(iv) A very different class of decaying potentials was studied by Pearson.²⁰⁴ His potentials are of the form

$$V(x) = \sum_{n=1}^{\infty} a_n W(x - x_n), \tag{V.3}$$

where $W \geq 0$, $a_n \rightarrow 0$, and $x_n \rightarrow \infty$ very rapidly so the bumps are sparse. He showed that for suitable a_n, x_n , the corresponding H has purely singular spectrum—providing the first explicit examples of such spectrum. Strong versions of his results were found by Remling²¹⁷ and Kiselev *et al.*¹⁵⁸ In particular, the latter authors proved if $(x_{n+1}/x_n) \rightarrow \infty$ (e.g., $x_n = n!$), then potentials of the form (V.3) lead to H 's with purely singular spectrum if $\sum a_n^2 = \infty$ and to ones with purely a.c. spectrum if $\sum a_n^2 < \infty$.

VI. INVERSE SPECTRAL THEORY

One area related to Schrödinger operators, especially in one dimension, is the question of inverse theory: How does one go from spectral or scattering information to the potential. There is much literature, including three books I would like to refer the reader to: Chadan and Sabatier,⁴⁵ Levitan,¹⁷⁶ and Marchenko.¹⁹⁰ I will only touch some noteworthy ideas here.

In one dimension, a key role is played by the Weyl m function and the associated spectral measure $d\rho$. Given a potential V so that H is self-adjoint with $u(0) = 0$ boundary conditions, for each z with $\text{Im } z > 0$, there is a solution $u(x; z)$ of $-u'' + Vu = zu$ which is L^2 at infinity. The m function is defined by

$$m(z) = \frac{u'(0; z)}{u(0, z)}. \tag{VI.1}$$

$\text{Im } m(z) > 0$ in $\text{Im } z > 0$ so by the Herglotz representation theorem

$$m(z) = B + \int d\rho(\lambda) \left[\frac{1}{\lambda - z} - \frac{\lambda}{1 + \lambda^2} \right] \tag{VI.2}$$

for a suitable constant B . $d\rho$ is called the spectral measure for H . One can recover $d\rho$ from m by

$$\frac{1}{\pi} \text{Im } m(\lambda + i\varepsilon) d\lambda \rightarrow d\rho(\lambda) \tag{VI.3}$$

weakly as $\varepsilon \downarrow 0$ and (VI.2) allows the recovery of m from $d\rho$ given the known asymptotics (Atkinson,¹⁵ Gesztesy and Simon⁹³)

$$m(-\kappa^2) = -\kappa + o(1) \tag{VI.4}$$

as $|\kappa| \rightarrow \infty$ with $\delta < \text{Arg} \kappa < (\pi/2) - \delta$. $d\rho$ really is a spectral measure for let $\tilde{\varphi}(x, \lambda)$ solve $-\tilde{\varphi}'' + V\tilde{\varphi} = \lambda \tilde{\varphi}$ with boundary conditions $\tilde{\varphi}(0, \lambda) = 0$, $\tilde{\varphi}'(0, \lambda) = 1$, and define for $f \in C_0^\infty(0, \infty)$

$$(Uf)(\lambda) = \int \tilde{\varphi}(x, \lambda) f(x) dx. \tag{VI.5}$$

Then U is a unitary map of $L^2(0, \infty, dx)$ to $L^2(\mathbb{R}, d\rho(\lambda))$; in particular,

$$\int |(Uf)(\lambda)|^2 d\rho(\lambda) = \int |f(x)|^2 dx \tag{VI.6}$$

or formally

$$\int \varphi(x, \lambda) \varphi(y, \lambda) d\rho(\lambda) = \delta(x - y). \tag{VI.7}$$

Moreover, $(UHf)(\lambda) = \lambda(Uf)(\lambda)$. $d\rho$ and its equivalent function m is therefore close to spectral information. One way of seeing this explicitly is if $V(x) \rightarrow \infty$. In that case, m is meromorphic, the poles of m are precisely the eigenvalues of H with $u(0) = 0$ boundary conditions and by definition of m , the zeros are precisely the eigenvalues with $u'(0) = 0$ boundary conditions. m is uniquely determined by these two sets of eigenvalues.

In many ways, the fundamental result in inverse theory is the following one:

Theorem VI.1: (Borg³⁷–Marchenko¹⁸⁸) *m determines q , that is, if q_1 and q_2 have equal m 's, then $q_1 = q_2$.*

Recently, the following local version of the Borg–Marchenko theorem was proven

Theorem VI.2: *Let q_1 and q_2 be potentials and m_1 and m_2 their m functions. Then $q_1 = q_2$ on $[0, a]$ if and only if*

$$|m_1(-\kappa^2) - m_2(-\kappa^2)| = O(e^{-2a\kappa})$$

as $\kappa \rightarrow \infty$ for κ obeying $\delta \leq \arg \kappa \leq \pi/2 - \delta$.

Remarks:

(1) This result was first proven by Simon²⁵⁸ when q_1 and q_2 are bounded from below.

(2) The general result which even allows q_i to be limit circle at infinity was first obtained by Gesztesy and Simon.⁹³

(3) A simple proof of Theorem VI.2 was subsequently obtained by Gesztesy and Simon.⁹⁴

Given the uniqueness result, it is natural to ask about concrete methods of determining q given m . There are two approaches for the general case. The first is due to Gel'fand and Levitan⁹⁰ and depends on the orthogonality relation (VI.7), while the other, due to Simon,²⁵⁸ is a kind of continuum analog of the continued fraction approach to solving the moment problem.

The Gel'fand–Levitan approach depends on a representation of the solutions φ due to Povzner²⁰⁸ and Levitan:¹⁷⁵

$$\varphi(x, \lambda) = \frac{\sin(kx)}{k} + \int_0^x K(x, y) \frac{\sin(ky)}{k} dy, \tag{VI.8}$$

where $\lambda = k^2$. In essence, (VI.7) leads to a linear Volterra integral equation for K whose kernel is determined by ρ . Once one has K , one can determine V from (VI.8) and $-\varphi'' + V\varphi = \lambda \varphi$ or from more direct relations of K to V .

The approach of Simon depends on a representation of m as a Laplace transform

$$m(-\kappa^2) = -\kappa - \int_0^a A(\lambda)e^{-2\kappa\lambda}d\lambda + O(e^{-2a\kappa}), \tag{VI.9}$$

which determines A given m (there is also a direct relation of A to ρ given in Gesztesy and Simon⁹³). One can introduce a second variable and function $A(x, \alpha)$ so $A(x=0, \alpha) \equiv A(\alpha)$. A obeys

$$\frac{\partial A}{\partial x} = \frac{\partial A}{\partial \alpha} + \int_0^\beta A(x, \beta)A(x, \alpha - \beta)d\beta \tag{VI.10}$$

and

$$\lim_{\alpha \downarrow 0} A(x, \alpha) \equiv V(x). \tag{VI.11}$$

In this approach, m determines $A(x=0, \cdot)$ by (VI.9); the differential equation (VI.10) determines $A(x, \alpha)$, and then (VI.11) determines V .

Inverse spectral theory is connected to inverse scattering for short-range potentials since $d\rho$ on $[0, \infty)$ is determined by scattering data. Scattering data also determine the positions of the negative eigenvalues. One needs to supplement that with the weight of the pure points at these negative eigenvalues known as norming constants. Marchenko^{190,189} has an approach to inverse scattering related to the Gel'fand–Levitán approach by using a different representation than (VI.8). When $\int_0^\infty x|V(x)|dx < \infty$, Levin¹⁷⁴ has proven that in $\text{Im } k > 0$, there is a solution $\psi(x, k)$ of $-\psi'' + V\psi = k^2\psi$ given by

$$\psi(x, k) = e^{ikx} + \int_x^\infty \tilde{K}(x, y)e^{iky}dy.$$

Krein^{162–164} also developed an approach to inverse problems. A different approach to inverse scattering is due to Deift and Trubowitz.⁶¹ For another approach to inverse problems, see Melin.¹⁹⁵ Inverse theory for periodic potentials also has an extensive literature starting with Dubrovin *et al.*,⁷⁰ Its and Matveev,¹³² McKean and van Moerbeke,¹⁹³ McKean and Trubowitz,¹⁹² and Trubowitz.²⁶⁸

As for higher-dimensional inverse scattering, these scattering data overdetermine the potential. For example, for short-range V 's, the scattering amplitude at fixed momentum transfer approaches the Fourier transform of V at large energy, so the large energy asymptotics of scattering determine V . There is considerable literature on recovering V from partial scattering data, which we will not try to summarize here.

One reason for the interest in inverse theory is the connection it sets up between spectral theory of Schrödinger operators and the analysis of certain nonlinear partial differential equations like KdV (see Dodd *et al.*,⁶⁸ Novikov *et al.*,¹⁹⁹ and Belokolos *et al.*²⁶).

VII. ERGODIC POTENTIALS

Let Ω be a compact metric space with probability measure $d\gamma$ and T_t with $t \in \mathbb{R}^v$ or T_n with $n \in \mathbb{Z}^v$ be an ergodic family of measure-preserving transformations. Let $f: \Omega \rightarrow \mathbb{R}$ be continuous. For $\omega \in \Omega$, define

$$V_\omega(x) = f(T_x\omega) \tag{VII.1}$$

and

$$H_\omega = -\Delta + V_\omega. \tag{VII.2}$$

Note: To allow unbounded V 's as seen, for example, in Gaussian random potentials, one wants to extend this picture to either allow f to be discontinuous and/or take values in $\mathbb{R} \cup \{\infty\}$, and/or allow Ω to be noncompact; for simplicity, we will discuss this model for motivation.

H_ω is a family of Schrödinger operators, not a single one, but by the ergodicity and an obvious translation covariance $V_{T_y\omega}(x) = V_\omega(x+y)$, many spectral properties occur with the probability one. So one can speak of typical properties. In particular, it is known that the full spectrum Σ , the essential support of the absolutely continuous spectrum Σ_{ac} , the closure of the point spectrum $\overline{\Sigma}_{pp}$, and the singular continuous spectrum Σ_{sc} are a.e. constant in ψ (see, e.g., Theorems 9.2 and 9.4 in Cycon *et al.*⁵³ for proofs; the result for Σ and $\overline{\Sigma}_{pp}$ is due to Pastur²⁰² and the other results to Kunz and Souillard¹⁶⁵). Note only $\overline{\Sigma}_{pp}$ is a.e. constant; Σ_{pp} , the actual set of eigenvalues is not.

Examples:

(1) Let $\Omega = [a, b]^{Z^v}$ and let $d\gamma$ be the infinite product of normalized Lebesgue measure on $[a, b]$. Let $(T_m\omega)_n = \omega_{n+m}$. The corresponding discrete Schrödinger operator is called the Anderson model and is typical of random potential models.

(2) If Ω is a compact Abelian group with Z^v or \mathbb{R}^v as a dense subgroup, $d\gamma$ is the Haar measure and T_x is the group translate, then V is a periodic or almost periodic function. A frequently discussed example is

$$V(n) = \lambda \cos(\pi \alpha n + \theta), \tag{VII.3}$$

where α is irrational, θ runs in $[0, 2\pi]$ (which is Ω), and λ is a parameter. The corresponding discrete Schrödinger operator is called the almost Mathieu model.

The simplest example of this framework—which is atypical in many ways—is the periodic potential. The basic facts in this case go back to physics literature at the start of quantum mechanics (Bloch, Brillouin, Kramer, and Wigner) and, in one dimension, to work on Hill's equation (Floquet, Lyapunov, Hamel, and Haupt). A critical early mathematical paper on the multidimensional case is Gel'fand.⁸⁹ The key result is that for periodic V 's with a mild local regularity condition, $H = -\Delta + V$ has purely absolutely continuous spectrum. This result is discussed in detail in Reed and Simon,²¹² Sec. XIII.16). The only subtle part of the argument is to eliminate the possibility of what are called flatbands, a result of Thomas.²⁶⁴

In the mathematical physics literature, the period from 1975 onwards has seen enormous interest in the study of almost periodic and random models and special cases thereof. Three books that discuss this are part of Carmona and Lacroix,⁴⁴ Cycon *et al.*,⁵³ and Pastur and Figotin.²⁰³ We will only touch some of the general principles, leaving the details—especially of detailed models—to the books and the vast literature. We will make references to the Lyapunov exponent without defining it; see Cycon *et al.*,⁵³ Sec. 9.3.

For random potentials, the most interesting results concern localization. While the spectrum is typically an interval (e.g., for the Anderson model in ν dimensions, it is $[a - 2\nu, b + 2\nu]$), the spectrum is pure point with eigenvalues dense in the interval and exponentially decaying eigenfunctions.

In one dimension, localization was first rigorously proven by Goldsheid *et al.*⁹⁸ with a later alternative by Kunz and Souillard.¹⁶⁵ Following an idea of Kotani,¹⁵⁹ Simon and Wolff,²⁶¹ and Delyon *et al.*⁶³ found another proof. Typical is

Theorem VII.1: *For the one-dimensional Anderson model, the spectrum is $[a - 2, b + 2]$ and is pure point with probability one with eigenfunctions decaying at the Lyapunov rate.*

Carmona *et al.*⁴³ and Shubin *et al.*²²⁸ have approaches that work if the single site distribution is discrete (the other quoted approaches require an absolutely continuous component for this distribution).

In higher dimensions, the two main approaches to localization are due to Fröhlich and Spencer⁸⁷ (see also von Dreifus and Klein²⁷⁰) and to Aizenman and Molchanov.¹⁰ (See also Aizenman and Graf⁸ and Aizenman *et al.*¹¹) Basically, these authors and the many papers that extend their ideas prove dense point spectrum in regimes where the coupling constant is large or

one is near the edge of the spectrum. It is believed—but not proven—that in suitable regimes when $\nu \geq 3$, there is absolutely continuous spectrum.

For almost periodic models, one can have any kind of spectral type. The almost Mathieu model has been almost entirely analyzed and the spectral type shows a great variety. Recall this is the discrete model with potential

$$V_{\alpha,\lambda,\theta}(n) = \lambda \cos(\pi \alpha n + \theta),$$

where λ, α are fixed parameters and θ runs through Ω . Then

(i) If $\lambda < 2$, there is always (i.e., for any irrational α) lots of a.c. spectrum and it is known for some α and believed for all α that is all there is (see Last,¹⁶⁹ Gesztesy and Simon,⁹² Gordon *et al.*,¹⁰⁰ Jitomirskaya;¹³⁴ the earliest results of this genre are due to Dinaburg and Sinai⁶⁷).

(ii) If $\lambda = 2$ and α is an irrational whose continued fraction integers are unbounded (almost all α have this property), then the spectrum is known to be purely singular continuous for almost all θ (see Gordon *et al.*¹⁰⁰).

(iii) If $\lambda > 2$ and α is an irrational with good Diophantine properties ($|\alpha - p/q| \geq Cq^{-l}$ for some C, l and all $p, q, \in \mathbb{Z}$), then for a.e. θ , the spectrum is dense pure point (Jitomirskaya;¹³⁴ see also Bourgain and Goldstein³⁸).

(iv) If $\lambda > 2$ and α is irrational, there are always lots of θ (a dense G_δ) for which the spectrum is purely singular continuous (Jitomirskaya and Simon¹³⁶). For some α , like those in (iii), the set while a dense G_δ has measure 0. For Liouville α (irrational α 's with $\overline{\lim}(1/q) \ln|\sin \pi \alpha q| = -\infty$), the spectrum is purely singular continuous (Avron and Simon²² using results of Gordon⁹⁹).

In general, for almost periodic models, the spectral type is dependent on the number theoretic properties of the frequencies. Among the general spectral results known for almost periodic models is that the spectrum is everywhere constant on Ω (rather than only almost everywhere constant; Avron and Simon²²) and that the essential support of the a.c. spectrum is everywhere constant (Last and Simon¹⁷⁰). It is known [see (iv)] that $\bar{\sigma}_{pp}$ and σ_{sc} may only be almost everywhere constant and fail to be constant on all of Ω .

VIII. TWO-BODY HAMILTONIANS

Hamiltonians of the form $-\Delta + V$ where $V(x) \rightarrow 0$ at infinity are often referred to as two-body Hamiltonians since the Hamiltonian of two particles with a potential $W(\mathbf{r}_1 - \mathbf{r}_2)$ reduces to $-\Delta + V$ (where V is a multiple of W depending on the masses) after removal of the center of mass. The issues are essentially the same as for one-dimensional decaying potentials as discussed in Sec. V.

With regard to the negative spectrum, again Weyl's criterion easily shows that $\sigma_{ess}(H) = [0, \infty)$ so that H has only discrete spectrum of finite multiplicity in $(-\infty, 0)$ and only 0 can be an accumulation point. Typical is:

Theorem VIII.1: For $\alpha \in \mathbb{Z}^p$, let χ_α be the characteristic function of the unit cube about α . Let $V: \mathbb{R} \rightarrow \mathbb{R}$. Suppose $V \in K_\nu$ and that as $\alpha \rightarrow \infty$, $\|\chi_\alpha V\|_{K_\nu} \rightarrow 0$. Then $\sigma_{ess}(-\Delta + V) = [0, \infty)$.

As for whether $N(V)$, the number of negative bound states (counting multiplicity, i.e., $N(V) = \dim E_{(-\infty, 0)}(H)$), is finite or infinite, there is considerable literature. The earliest bound is due to Birman³² and Schwinger²²⁷ for $\nu = 3$. It states

$$N(V) \leq \frac{1}{(4\pi)^2} \int \frac{|V(x)||V(y)|}{|x-y|^2} dx dy \quad (\nu=3). \tag{VIII.1}$$

Perhaps the most famous bound is that of Cwickel,⁵² Lieb,¹⁷⁷ and Rosenbljum.²²⁰

$$N(V) \leq L_{0,\nu} \int |V(x)|^{\nu/2} dx \quad (\nu \geq 3). \tag{VIII.2}$$

One reason this is of special interest is that for nice V 's as $\lambda \rightarrow \infty$, $N(\lambda V)/\int |\lambda V|^{\nu/2} dx$ converges to a universal constant (see Sec. XIV). In particular, (VIII.1) has the wrong large λ behavior while (VIII.2) has the right such behavior. (Simon²⁴³ had the first bounds with the right large λ behavior for nice enough V 's; he also conjectured (VIII.2).)

As in the one-dimensional case, there are Lieb–Thirring-type bounds on the moments of the negative eigenvalues e_j of $-\Delta + V$

$$\sum_j |e_j|^{\gamma} \leq L_{\gamma,\nu} \int dx |V(x)|^{\gamma+\nu/2} dx$$

for $\gamma > 0$ if $\nu = 2$ and $\gamma \geq 0$ if $\nu \geq 3$. These were proven first in Lieb and Thirring.¹⁸⁵ There has been considerable literature on the best values of $L_{\gamma,\nu}$. In particular, a recent pair of papers of Laptev and Weidl¹⁶⁸ and Hundertmark *et al.*¹²⁴ has obtained a breakthrough in understanding the ν dependence of $L_{\gamma,\nu}$. In particular, they show that for $\gamma \geq \frac{3}{2}$, $L_{\gamma,\nu}$ is given by the quasiclassical value. On the other hand, it is known that $L_{\gamma=0,\nu} > L_{\gamma=0,\nu}^{q.c.}$, the quasiclassical value for all ν (Helffer and Robert^{109,110}).

For a review of the literature on bounds on the number of eigenvalues, especially the subtle two-dimensional case, see Birman and Solomyak.³⁶

The absence of eigenvalues at positive energies is a specialized issue largely independent of the rest of the analysis of positive spectrum. Given the examples of Wigner–von Neumann and related ones of Naboko and Simon discussed in Sec. V, one needs some condition on the falloff or lack of oscillations. Here is a simple result:

Theorem VIII.2: *Let $V(x) = V_1(x) + V_2(x)$ where $|x||V_1(x)| \rightarrow 0$ and $|(x \cdot \nabla)V_2(x)| \rightarrow 0$. Then $-\Delta + V$ has no eigenvalues in $[0, \infty)$.*

Remarks:

(1) The stated theorem requires local regularity (V_1 bounded near infinity and V_2 is C^1), but there are extensions that allow local singularities.

(2) Rellich²¹⁶ proved that if V has compact support, there are no positive energy eigenvalues. Theorem VIII.2 when $V_2 = 0$ is due to Kato¹⁵⁰ and the full result to Agmon² and Simon.²³⁵

(3) See Froese *et al.*⁸⁶ for another result of this genre; we will discuss their result further in Sec. IX.

The methods we will discuss below typically show that $\sigma_{pp} \cap (0, \infty)$ is finite; one can then usually use Theorem VIII.2 to prove that the set is actually empty.

As for positive spectrum, it is intimately related to scattering theory. Given two self-adjoint operators A, B , one says the wave operators exist if

$$\Omega^\pm(A, B) = s\text{-}\lim_{t \rightarrow \mp\infty} e^{itA} e^{-itB} P_{ac}(B)$$

exists where P_{ac} is the projection onto the a.c. subspace for B . We say they are complete if $\text{Ran } \Omega^\pm(A, B) = \text{Ran } P_{ac}(A)$, in which case $\Omega^\pm(A, B)$ are unitary maps of $\text{Ran } P_{ac}(B)$ to $\text{Ran } P_{ac}(A)$ which intertwine A and B . See Reed and Simon,²¹³ Baumgärtel and Wollenberg,²⁵ or Yafaev²⁷⁷ (or many other books) for a discussion of the physics involved.

The development of abstract scattering theory is closely intertwined (pun intended) to its applications to Schrödinger operators. Fundamental work was done by Jauch,¹³³ Cook,⁵¹ Rosenblum,²²¹ Kato,¹⁴⁹ Birman,³³ and Birman and Krein.³⁵

The basic result for positive spectrum for “short-range” potentials is:

Theorem VIII.3: *Let V be such that $(1 + |x|)^{1+\epsilon} V(x) \in L^p + L^\infty(\mathbb{R}^n)$ for $\max(2, \nu/2) < p < \infty$ and let $H = -\Delta + V$ and $H_0 = -\Delta$. Then $\Omega^\pm(H, H_0)$ exist and are complete. Moreover, H has no singular continuous spectrum and any eigenvalues in $(0, \infty)$ are isolated (from other eigenvalues) and of finite multiplicity.*

Remarks:

(1) The first results on absence of singular continuous spectrum depended on eigenfunction expansions and were obtained by Povzner²⁰⁹ (V 's of compact support) and Ikebe¹²⁹ (V 's which

were $O(|x|^{-2-\epsilon})$ at infinity). The earliest results on completeness of wave operators depended on the trace class theory of scattering (of Rosenblum²²¹ and Kato¹⁴⁹) and were obtained by Kuroda.^{166,167} From 1960 to 1972, the decay was successively improved until Agmon³ obtained the $O(|x|^{-1-\epsilon})$ result quoted.

(2) Enss⁷⁷ has a different, quite physical, approach to this result. Enss' work depends in part on an earlier geometric characterization of the continuous subspace of a Schrödinger operator by Ruelle²²² and Amrein and Georgescu.¹⁴ This is sometimes called the RAGE theorem after the initials of the authors.

(3) It is known (e.g., Dollard⁶⁹) that if $V(x) = O(|x|^{-1})$, $\Omega^\pm(H, H_0)$ may not exist.

For long-range behavior decaying slower than $O(|x|^{-1})$, there are results if ∇V decays faster than $O(|x|^{-1-\epsilon})$. Basically, there is only a.c. spectrum at positive energy if $V = V_1 + V_2$ with $V_1 = O(|x|^{-1-\epsilon})$ and $x \cdot \nabla V_2 = O(|x|^{-\epsilon})$. For details, see Lavine,¹⁷² Agmon and Hörmander,⁶ and Hörmander.¹²² These works use modified wave operators as introduced by Dollard.⁶⁹

IX. N-BODY HAMILTONIANS

Let \tilde{H} be the Hamiltonian of N particles in \mathbb{R}^{ν} . Explicitly, \tilde{H} is an operator on $L^2(\mathbb{R}^{\nu N})$ given by $\tilde{H} = \tilde{H}_0 + V$ where

$$\tilde{H}_0 = - \sum_{j=1}^N \frac{1}{2m_j} \Delta_{x_j}$$

with $x = (x_1, \dots, x_N)$ a point in $\mathbb{R}^{\nu N} = \mathbb{R}^{\nu} \times \mathbb{R}^{\nu} \times \dots \times \mathbb{R}^{\nu}$ (N times) and

$$V = \sum_{i < j} V_{ij}(x_i - x_j),$$

with V_{ij} a function in \mathbb{R}^{ν} which decays at infinity. There is a standard way of removing the center of mass and getting an associated Hamiltonian H on $L^2(\mathbb{R}^{\nu(N-1)})$. For a more extensive review of the subject than this brief discussion, see Hunziker and Sigal.¹²⁸

For any partition a of $\{1, \dots, N\}$ into disjoint subsets, one defines $I(a) = \sum_{(i,j) \in a} V_{ij}$ over the pairs (i, j) in distinct clusters and $H(a) = H - I(a)$.

The issues one faces are similar to those in the two-body case but often more subtle. The first thing one needs to establish about N -body systems is where the essential spectrum of H lies. The result involves

$$\Sigma(a) = \inf \text{spec}(H(a)), \tag{IX.1}$$

$$\Sigma = \min_{\#a \geq 2} (\Sigma(a)). \tag{IX.2}$$

Σ is the minimum energy the system can have after it is broken into two pieces moved very far from each other. That makes the following physically attractive:

Theorem IX.1: (HVZ Theorem) *Suppose each V_{ij} viewed as an operator on $L^2(\mathbb{R}^{\nu})$ obeys $V_{ij}(-\Delta_{ij} + 1)^{-1}$ is compact. Then*

$$\sigma_{\text{ess}}(H) = [\Sigma, \infty).$$

Remarks:

(1) The name ‘‘HVZ’’ comes from work of Hunziker,¹²⁶ van Winter,²⁶⁹ and Zhislin²⁸⁰ who first proved it.

(2) The original proofs used resolvent equations; a geometric proof was later found by Enss⁷⁶ and Simon.²⁴⁴

The next issue is whether the discrete spectrum is finite or infinite. A great deal of attention has been paid to atomic or ionic Hamiltonians. Define on $L^2(\mathbb{R}^{3N})$:

$$H_M(N, Z) = \sum_{i=1}^N \left(-\Delta_i - \frac{Z}{|x_i|} \right) + \frac{1}{M} \sum_{i < j} \nabla_i \nabla_j + \sum_{i < j} \frac{1}{|x_i - x_j|},$$

which describes N electrons moving around a nucleus of charge Z and mass M . A basic result states that neutral atoms and positive ions always have an infinite number of bound states:

Theorem IX.2: (Zhislin²⁸⁰) *If $N \leq Z$, $\dim E_{(-\infty, \Sigma)}(H_M(N, Z)) = \infty$ for any M (including $M = \infty$).*

Remarks:

(1) The first result of this genre was Kato¹⁴⁵ who proved the result if $N = Z = 2$ and $M = \infty$ (Helium). He did not properly handle $M < \infty$ because he did not use the right coordinate systems. As shown by Simon,²³⁶ Kato's idea, which involved placing $N - 1$ electrons in the ground states for the $N - 1$ ion and the N th in a hydrogen-like state around the core, can prove Theorem IX.2.

- (2) This result holds even if one adds Fermi statistics (see, e.g., Simon²³⁶).
- (3) If Z is not restricted to be an integer, the proper condition is $N < Z + 1$.

As for negative ions, we have

Theorem IX.3: (Zhislin²⁸¹) $\dim E_{(-\infty, \Sigma)}(H_M(Z + 1, Z)) < \infty$

Remarks:

- (1) This result also has a geometric proof by Sigal²³¹ and Simon.²⁴⁴
- (2) This result may not be true for fermion electrons because the $N - 1$ problem may have a degenerate ground state which allows one with a nonzero dipole moment.
- (3) While it is presumably true that $\dim E_{(-\infty, \Sigma)}(H_M(N, Z)) < \infty$ for all $N \geq Z + 1$, that is not known.

Finally, with regard to bound states of atoms, there is the issue of when $\dim E_{(-\infty, \Sigma)} = 0$. The result is the following:

Theorem IX.4: Let $M = \infty$.

- (a) (Ruskai^{223,224} and Sigal^{231,232}) *For any Z , there is an $N_0(Z)$ so that for $N \geq N_0(Z)$, there is no spectrum in $(-\infty, \Sigma)$. $N_0(Z)$ denotes the smallest N_0 for which this is true.*
- (b) (Lieb *et al.*¹⁸¹) *For fermions, $N_0(Z)/Z \rightarrow 1$ as $Z \rightarrow \infty$.*
- (c) (Benguria and Lieb²⁸) *Without Fermi statistics, $N_0(Z) > 1.2Z$ for Z large.*
- (d) (Lieb¹⁷⁸) $N_0(Z) \leq 2Z$.

Remarks:

- (1) If $N \geq N_0$, then $\inf \text{spec}(H(N, Z)) = \inf \text{spec}(H(N_0, Z)) < \inf \text{spec}(H(N_0 - 1, Z))$.
- (2) Some of these results hold if $M < \infty$.

With short-range potentials, the situation is simple if the bottom of the essential spectrum is two body. Define

$$\Sigma_3 = \min_{\#(a) \geq 3} (\Sigma(a)).$$

Then (see Cycon *et al.*,⁵³ Sec. 3.9)

Theorem IX.5: (Sigal²³¹) *Suppose $\Sigma_3 > \Sigma$, $\nu \geq 3$, and each V_{ij} lies in $L^{\nu/2}(\mathbb{R}^\nu)$. Then $\dim E_{(-\infty, \Sigma)}(H) < \infty$.*

On the other hand, if $\Sigma_3 = \Sigma$, there can be an infinite number of bound states even if the V_{ij} 's have compact support (in x_{ij}). In particular, if $N = 3$, $V_{12} = V_{23} = V_{13} = -c\chi_1$, with χ the characteristic function of a unit ball and c chosen so that $\inf \text{spec}(H) = 0$ but $\inf \text{spec}(H + \varepsilon V) < 0$ for all $\varepsilon > 0$, it is known that $\dim E_{(-\infty, 0)}(H) = \infty$. This is known as the Efimov effect after work of Efimov.^{74,75} For proofs of this phenomenon, see Yafaev²⁷⁶ and Ovchinnikov and Sigal.²⁰¹

In analyzing the spectrum of H on $[\Sigma, \infty)$, a particular class of physically significant energies occurs, the thresholds. For each partition a of $\{1, \dots, N\}$ with $\#a \geq 2$, there is a natural decompo-

sition of $L^2(\mathbb{R}^{\nu(N-1)}) = \mathcal{H}_a \otimes \mathcal{H}^a$ where \mathcal{H}_a are functions of $x_i - x_j$ with i and j in the same cluster of a and \mathcal{H}^a are functions of $R_\alpha - R_\beta$, where R_α is the center of mass of a cluster (see Ref. 128 for an elegant way of doing this kinematics). Under the decomposition $H(a) = H_a \otimes I + I \otimes T^a$. H_a is the internal energy of the cluster and T^a the kinetic energy of the cluster centers of mass. $\mathcal{I}(a)$ is the set of eigenvalues of H_a (with the condition that if $\#(a) = N$, so H_a is 0 on \mathbb{C} , then $\mathcal{I}(a) = \{0\}$). The set of thresholds is defined as

$$\mathcal{I} = \bigcup_a \mathcal{I}(a).$$

Note: An energy in $\mathcal{I}(a)$ is a sum of eigenvalues of individual cluster Hamiltonians. In particular, the statement in the theorems below that the set of thresholds is a closed countable set follows by induction from the other statement that eigenvalues can only accumulate at thresholds.

The three-body problem turns out to have some aspects that make it simpler than the general N -body problem, and Faddeev⁷⁹ and later Enss⁷⁸ (using very different methods) have fairly complete results on spectral and scattering theory for $N = 3$. We will focus here on results that apply for all N .

Historically, the first aspect of the continuous spectrum for general N -body systems controlled was the absence of a singular continuous spectrum. The earliest result required analyticity of the potentials but included atoms:

Theorem IX.6: (Balslev and Combes²³) *Suppose each $V_{ij}(x) = f_{ij}(x_i - x_j)$ where f_{ij} is a function on $\mathbb{R}^\nu \setminus \{0\}$ that obeys*

$$A(\theta) = V(e^\theta x)(-\Delta + 1)^{-1}$$

is compact and has an analytic continuation from $\theta \in \mathbb{R}$ to $\{\theta \mid |\operatorname{Im} \theta| < \varepsilon\}$ for some $\varepsilon > 0$. Then $\sigma_{\text{sc}}(H) = \emptyset$.

Moreover,

- (i) *Any eigenvalue of H in $\mathbb{R} \setminus \mathcal{I}$ is of finite multiplicity, and eigenvalues can only accumulate at thresholds.*
- (ii) *The set of eigenvalues union thresholds is a closed countable set.*

Remarks:

- (1) Such potentials are called dilation analytic.
- (2) This result was first proven for two-body systems by Aguilar and Combes.⁷
- (3) See Simon^{237,238} for extensions of this result.

The most general results on absence of singular continuous spectrum depend on the ideas of Mourre.¹⁹⁶

Theorem IX.7: *Suppose $V_{ij}(x) = f_{ij}(x_i - x_j)$ where f_{ij} is a function on \mathbb{R}^ν that obeys (as operators on $L^2(\mathbb{R}^\nu)$)*

- (i) *$f_{ij}(x)(-\Delta + 1)^{-1}$ is compact;*
- (ii) *$(-\Delta + 1)^{-1} x \cdot \nabla f_{ij} (-\Delta + 1)^{-1}$ is compact.*

Then $\sigma_{\text{ess}}(H)$ is empty. Moreover, any eigenvalue in $\mathbb{R} \setminus \mathcal{I}$ is discrete, eigenvalues can only accumulate at thresholds, and the set of eigenvalues and thresholds is a closed countable set.

Remarks:

(1) This theorem was proven for $N = 3$ by Mourre.¹⁹⁶ His methods were extended and elucidated by Perry *et al.*²⁰⁵ who obtained the general N -body result. Substantial simplifications of the proof were found by Froese and Herbst.⁸⁵

(2) Condition (ii) does not require that f_{ij} be smooth because $\nabla f_{ij} = [\nabla, f_{ij}]$ and $\nabla(-\Delta + 1)^{-1}$ is bounded. Basically, (i), (ii) hold if $f_{ij} = f_{ij}^{(1)} + f_{ij}^{(2)}$, where $x f_{ij}^{(1)} (-\Delta + 1)^{-1}$ is compact and $f_{ij}^{(2)}$ is smooth with $(x \cdot \nabla) f_{ij}^{(2)} (-\Delta + 1)^{-1}$ and $f_{ij}^{(2)} (-\Delta + 1)^{-1}$ compact.

(3) Froese and Herbst⁸⁵ have some general results that imply that $\mathcal{I} \cap (0, \infty) = \emptyset$ (see Theorem 4.19 in Cycon *et al.*⁵³).

Finally, there has been extensive study of scattering theory and completeness. For each cluster with $\#(a) \geq 2$, let P_a on \mathcal{H}_a be the projection onto the point spectrum of H_a and let $P(a) = P_a \otimes I$, the projection onto vectors which are bound within the clusters and arbitrary for the centers of mass coordinates. The cluster wave operators are defined by

$$\Omega^\pm(a) = s\text{-}\lim_{t \rightarrow \mp\infty} e^{+itH} e^{-itH(a)} P(a). \tag{IX.3}$$

$\text{Ran}(\Omega^+(a))$ are those states which in the distant past look like bound clusters (corresponding to the partition a) moving freely relative to one another.

The existence of cluster wave operators (IX.3) was proven first by Hack.¹⁰³ It is not hard to see (e.g., Theorem XI.36 in Reed and Simon²¹³) that for $a \neq b$, $\text{Ran } \Omega^+(a)$ is orthogonal to $\text{Ran } \Omega^+(b)$. Asymptotic completeness is the statement that

$$\bigoplus_{\#(a) \geq 2} \text{Ran}(\Omega^+(a)) = \mathcal{H}_{\text{ac}}(H),$$

where $\mathcal{H}_{\text{ac}}(H)$ is the absolutely continuous subspace for H . After fairly general results for $N=3$ (Faddeev⁷⁹ and Enns⁷⁸) and for general N with weak coupling (Iorio and O'Carroll¹³¹) and repulsive potentials (Lavine¹⁷¹), Sigal and Soffer²³³ solved the general result. Their theorem is

Theorem IX.8: (Sigal and Soffer²³³) *If each $V_{ij}(x) = f_{ij}(x_i - x_j)$ where $|(D^\alpha f_{ij}(x))| \leq C(1 + |x|)^{-|\alpha| - \varepsilon - 1}$ for all multiindices with $|\alpha| \leq 2$, then asymptotic completeness holds.*

Extensions and clarifications of this work are due to Graf,¹⁰¹ Hunziker,¹²⁷ and Yafaev.²⁷⁸ Long-range potentials are discussed in Dereziński,⁶⁵ Sigal and Soffer,²³⁴ and Dereziński and Gerard.⁶⁶

X. CONSTANT ELECTRIC AND MAGNETIC FIELDS

Quantum mechanics with a potential and constant electric or magnetic field played a critical role experimentally and theoretically in the earliest days of the subject, and there has been considerable mathematical literature on the spectral properties of these operators. The basic Stark Hamiltonian on $L^2(\mathbb{R}^{\nu})$ is

$$H = -\Delta + Ex_1 + V(x), \tag{X.1}$$

where V is short range. A key role has been played by an explicit formula of Avron and Herbst¹⁸ for the operator when $V=0$, viz.,

$$\exp(-it(-\Delta + x_1)) = \exp(-it^3/3) \exp(-itx_1) \exp(-it\Delta + ip_1 t^2), \tag{X.2}$$

where $p_1 = (1/i)(\partial/\partial x_1)$. Classically in an electric field, a particle has $x_1 = N - ct^2$ as $t \rightarrow \infty$ and (X.2) realizes this with the $p_1 t^2$ term. It means the borderline for short range is $|x|^{-1/2-\varepsilon}$ rather than $|x|^{-1-\varepsilon}$. The result is

Theorem X.1: *Suppose $|V(x)| \leq C(1 + |x|)^{-\varepsilon}(1 + |x_1|)^{-1/2-\varepsilon}$. Then H given by (X.1) has complete wave operators and empty singular continuous spectrum. Eigenvalues are isolated and of finite multiplicity.*

This result and ones similar to it are discussed by Herbst,¹¹⁴ Yajima,²⁷⁹ and Simon.²⁴⁹ Multiparticle completeness in electric fields has been studied by Herbst *et al.*,¹¹⁶ and Adachi and Tamura.¹

There is much literature on both constant and variable magnetic fields but an extensive review of it is beyond the scope of this paper. One can begin looking at the literature by consulting a series by Avron *et al.*¹⁹⁻²¹ and Chapter 6 of Cycon *et al.*⁵³ and references therein. See also Sec. XII.

XI. COULOMB ENERGIES

While much of the mathematical theory of nonrelativistic quantum mechanics has focused on general potentials, nature uses the Coulomb potential and there is considerable literature on binding energies of Coulomb systems, especially as some parameter goes to infinity. Section IX (see Theorem IX.4) already discussed one such result. We will only introduce some seminal themes; consult Lieb¹⁷⁹ for a review of the subject.

The most famous of these results is the stability of matter. In its simplest form, it concerns the Hamiltonian

$$H(N, M; R_1, \dots, R_M) = - \sum_{i=1}^N \Delta_i - \sum_{i, \alpha} \frac{1}{|x_i - R_\alpha|} + \sum_{i < j} \frac{1}{|x_i - x_j|} + \sum_{\alpha < \beta} \frac{1}{|R_\alpha - R_\beta|}$$

of N electrons moving in the field of M infinitely massive protons. Let \mathcal{H}_f be the functions on $L^2(\mathbb{R}^{3N})$ thought of as functions $\psi(x_1, \dots, x_N)$ of N variables in \mathbb{R}^3 which are antisymmetric, that is,

$$\psi(x_{\pi(1)}, \dots, x_{\pi(n)}) = (-1)^\pi \psi(x_1, \dots, x_N)$$

for any permutation π ; that is, \mathcal{H}_f is the wave function with Fermi statistics (we ignore spin which is easily accommodated). Define

$$E(N, M) = \inf_{\substack{\psi \in \mathcal{H}_f \\ R_1, \dots, R_M}} \langle \psi, H(N, M; R_1, \dots, R_M) \psi \rangle.$$

Stability of matter states that

$$E(N, M) \geq -c(N + M). \tag{XI.1}$$

Among other things, this bound is important because it is equivalent to the fact that the radius of a chunk of matter with $N=M$ does not shrink to zero as $N \rightarrow \infty$.

The first proof of (XI.1) was obtained by Dyson and Lenard^{72,73} with a constant C that was many powers of ten too large. Lieb and Thirring¹⁸⁶ found an elegant proof with a constant C that is on the order of magnitude of Rydbergs. The result (XI.1) fails if one does not impose Fermi statistics (see Dyson⁷¹ and Conlon *et al.*⁵⁰). Extensions that involve relativistic kinetic energy, magnetic and/or radiation fields can be found in Conlon,⁴⁹ Lieb *et al.*,¹⁸⁰ and Fefferman *et al.*⁸¹

Another Coulomb energy problem that has been extensively studied is the total binding energy in the limit of large of Z . One defines

$$H(N, Z) = \sum_{i=1}^N \left(-\Delta_i - \frac{Z}{|x_i|} \right) + \sum_{i < j} \frac{1}{|x_i - x_j|}$$

on \mathcal{H}_f and

$$E(N, Z) = \inf_{\psi \in \mathcal{H}_f} \langle \psi, H(N, Z) \psi \rangle$$

and

$$E(Z) = \min_N E(N, Z).$$

One knows that

$$E(Z) = \alpha Z^{7/3} + \beta Z^2 + \gamma Z^{5/3} + o(Z^{5/3}). \tag{XI.2}$$

The α term is given by Thomas–Fermi theory and this leading asymptotics was proven by Lieb and Simon.¹⁸² The β term is called the Scott correction and it was established by a combination of ideas of Hughes¹²³ and Siedentop and Weikard.^{229,230} The full asymptotics (XI.2) was obtained by Fefferman and Seco.⁸² Results for large Z and large magnetic field can be found in Lieb *et al.*^{183,184}

XII. EIGENVALUE PERTURBATION THEORY

Some of Schrödinger’s earliest papers on quantum mechanics concerned eigenvalue perturbation theory. Kato’s book¹⁵³ is a source of detailed information on what we will call regular and asymptotic perturbation theory below. A review of some of the other aspects can be found in Reed and Simon²¹² and Simon.²⁵⁴

If A is self-adjoint and B is A -bounded in the sense of (III.3), and if E_0 is a simple eigenvalue of A , then for β small, $A + \beta B$ has a unique eigenvalue $E(\beta)$ near E_0 and $E(\beta)$ is analytic in β . This is a result of Rellich²¹⁵ and Kato.^{142,143} An example is

$$-\Delta_1 - \Delta_2 - \frac{1}{|x_1|} - \frac{1}{|x_2|} - \frac{1}{Z} \frac{1}{|x_1 - x_2|} \tag{XII.1}$$

about $1/Z=0$ which is equivalent after scaling (of space and energy) to

$$-\Delta_1 - \Delta_2 - \frac{Z}{|x_1|} - \frac{Z}{|x_2|} + \frac{1}{|x_1 - x_2|}.$$

The numerical radius of convergence in $|1/Z|$ is about 1.06 so $H(Z=2)$ and $H(Z=1)$ are both included. Kato¹⁴⁷ developed the theory for form perturbations. Rellich and Kato included degenerate eigenvalues.

Titchmarsh^{265,266} and Kato¹⁴⁶ also developed the theory of asymptotic situations like the anharmonic oscillator

$$-\frac{d^2}{dx^2} + x^2 + \beta x^4, \tag{XII.2}$$

where each eigenvalue $E_n(\beta)$ for $\beta > 0$ has an asymptotic series

$$E_n(\beta) \sim \sum_{n=0}^{\infty} a_n \beta^n$$

even though this series can be divergent (and is for the case (XII.2), as shown by Bender and Wu²⁷). See Herbst and Simon¹¹⁷ for an example where an asymptotic series converges but to the wrong answer! See Simon²⁵² for a study of multiwell problems.

In some cases, including (XII.2), it is known that the divergent perturbation series can be made to give the right eigenvalue with a summability method, either Padé approximation (Loeffel *et al.*¹⁸⁷) or Borel summation (Graffi *et al.*¹⁰²). Borel summability is also known to work for the Zeeman series for hydrogen–hydrogen perturbed by turning on a constant magnetic field; see Avron *et al.*²¹ and Avron *et al.*¹⁷

In certain cases, eigenvalues are perturbed into resonances, the subject of Sec. XIII. For eigenvalues embedded in continuous spectrum under regular perturbations (like (XII.1)), the convergence of the perturbation series for a resonance and its related time-dependent perturbation theory and the Fermi golden rule is discussed in Simon.^{237,238} For Stark Hamiltonians, the basic paper is Herbst.¹¹⁵ Harrell and Simon¹⁰⁷ found the leading resonance asymptotics in this case.

XIII. RESONANCES

Almost everything we have discussed so far has involved a single operator and properties invariant under unitary transformations. The notion of resonances has got to involve additional structure. For example, the operators $-\Delta - |x|^{-1} - Fx = H(F)$ are unitarily equivalent for all $F \neq 0$. But according to the physics lore, there is a resonance with an F -dependent position. We will not emphasize the extra structure, but it is there. We will focus on two definitions of resonances: one suitable for potentials that decay very rapidly (see Zworski^{284,285} for reviews) and the method of complex scaling already discussed in a different context in Sec. IX. (See Reed and Simon²¹² and Simon²⁴⁵ for reviews.)

Let ν be an odd dimension, let V be a bounded potential of compact support on \mathbb{R}^ν , and for $\text{Re } \kappa > 0$, define

$$B(\kappa) = |V|^{1/2} (-\Delta + \kappa^2)^{-1} V^{1/2},$$

where $V^{1/2} = |V|^{1/2} \text{sgn}(V)$. Then $-\kappa^2$ is an eigenvalue of $-\Delta + V$ if and only if -1 is an eigenvalue of $B(\kappa)$. Since ν is odd, $B(\kappa)$ has an analytic continuation as a compact operator-valued function of κ to all of \mathbb{C} (when $\nu=1$, there is a simple pole at $\kappa=0$ but $\kappa B(\kappa)$ is entire). If $\text{Re } \kappa < 0$ and -1 is an eigenvalue of $B(\kappa)$, we say $-\kappa^2$ is a resonance of $-\Delta + V$.

Froese⁸³ has a lovely formula that relates resonances defined by this method to scattering theory. For all κ , $B(\kappa) - B(-\kappa)$ is trace class so $(1 + B(-\kappa))(1 + B(\kappa))^{-1}$ is 1 plus trace class and has a determinant as an operator on $L^2(\mathbb{R}^\nu)$. For k real and $S(k)$, the S matrix on $L^2(S^{\nu-1})$,

$$\det(S(k)) = \det((1 + B(-ik))(1 + B(ik))^{-1}),$$

so resonances are related to poles of the analytic continuation of S .

There has been considerable literature on the number of resonances. Let $N(R)$ be the number of resonances with energy E obeying $|E| < R$. In one dimension, one has a complete result:

Theorem XIII.1: (Zworski²⁸²) *Let $\nu=1$ and suppose $[a, b]$ is the convex hull of the support of V . Then*

$$\lim_{R \rightarrow \infty} R^{-1/2} N(R) = \frac{2}{\pi} |b - a|.$$

Remarks:

(1) The result depends on a theorem of Titchmarsh and Cartwright on the zeros of Fourier transforms of functions of compact support.

(2) Froese⁸³ has obtained some results for cases when a potential decays faster than any exponential but may not have compact support.

In higher dimensions, much less is known. Zworski²⁸³ proved that for V of compact support, $N(R) \leq C(R+1)^{\nu/2}$ (see also Froese⁸⁴). On the other hand for general V 's, it is only known (Sá Barreto and Zworski²²⁵) that $\lim_{R \rightarrow \infty} N(R) = \infty$.

Suppose V is a dilation analytic potential in the sense of Theorem IX.6. Let

$$H(\theta) = -e^{-2\theta} \Delta + V(e^\theta \tau).$$

Because of the analyticity assumption, $H(\theta)$ is analytic in $\{\theta \mid |\text{Im}(\theta)| < \alpha\}$ for some α . Then Aguilar and Combes⁷ found the essential spectrum of $H(\theta)$ for $N=2$ and Balslev and Combes²³ for general N :

Theorem XIII.2: $\sigma_{\text{ess}}(H(\theta)) = \cup_{E \in \mathcal{I}(\theta)} (E + e^{-2\theta} \mathbb{R})$

Remarks:

(1) $\mathcal{I}(\theta)$ is the thresholds of $H(\theta)$ defined analogously to the case $\theta=0$. It is not hard to see that $\sigma_{\text{ess}}(H(\theta))$ and $\mathcal{I}(\theta)$ depend only on $\text{Im } \theta$.

(2) If $\text{Im } \theta > 0$, $\sigma_{\text{ess}}(H(\theta)) \cap \mathbb{R}$ consists precisely of \mathcal{I} . Basically as we increase $\text{Im } \theta$ from 0, the essential spectrum rotates about the thresholds. In doing that, it can uncover resonances.

Resonances defined by this method have been used by quantum chemists for numerical calculations as well as a theoretical tool. Simon^{237,238} used it to study the Fermi golden rule and Harrell and Simon¹⁰⁷ to prove various one-dimensional tunneling estimates.

Avron¹⁶ used these ideas to study large-order perturbation theory for hydrogen in a magnetic field; a rigorous proof of his results was obtained by Helffer and Sjöstrand.¹¹²

Herbst¹¹⁵ has extended the ideas to Hamiltonians with constant electric field. Among his results is the surprising one that if $0 < \text{Im } \theta < \pi/3$, then $-e^{-2\theta}\Delta + e^\theta x$ has empty spectrum!

XIV. THE QUASICLASSICAL LIMIT

There has been considerable literature on the connection between quantum and classical mechanics. Much of it has focused on what happens as $\hbar \rightarrow 0$, but there are other limiting situations where a classical or semiclassical picture is appropriate—for example, the large Z limit of atoms. We will touch on some of the subjects considered, but the literature is vast. Robert²¹⁹ has an excellent review of those results obtained for very smooth potentials using the Fourier integral operator methods pioneered by Hörmander and Maslov. Therefore, I will not try to cover these results here. We note that in Sec. XI, we referenced the Thomas–Fermi limit, which is quasiclassical.

Consider first the $\hbar \downarrow 0$ limit. Let $H_\hbar = -(\hbar^2/2m)\Delta + V$. Kac^{140,141} had the idea that the small \hbar limit of $\exp(-sH_\hbar)$ was the same as the zero time limit in Brownian motion. This allows one to prove under great generality that the quantum partition function $\text{Tr}(\exp(-sH_\hbar))$ approaches a classical partition function as $\hbar \downarrow 0$; see, for example, Theorem 10.1 in Simon.²⁴⁶ The earliest results I know of on this subject are due to Berezin.³¹

Quantum dynamics, $e^{-isH_\hbar/\hbar}\psi_\hbar$, on suitable states ψ_\hbar make an elegant classical limit—one takes ψ_\hbar to be a coherent state which collapses to a single point in phase space as $\hbar \downarrow 0$. Such results were found by Hagedorn^{104–106} (similar methods were developed independently by Ralston²¹⁰).

Since $-\hbar^2\Delta + V = \hbar^2[-\Delta + \hbar^{-2}V]$, the small \hbar limit is the same as the large coupling constant limit for $-\Delta + \lambda V$. In particular, if $N(V) = \dim E_{(-\infty, 0)}(-\Delta + V)$, the quantity discussed in Sec. VIII, one has

Theorem XIV.1: *Let $v \geq 3$ and $V \in L^{v/2}(\mathbb{R}^v)$. Then $\lim_{\lambda \rightarrow \infty} N(\lambda V)/\lambda^{v/2} = (2\pi)^{-v} \tau_v \int_{V \leq 0} (-V(x))^{v/2} d^v x$, where τ_v is the volume of a unit ball in \mathbb{R}^v .*

Remarks:

(1) This theorem is quasiclassical since the right side is $(2\pi)^{-v}$ times the volume of the classical phase space region where $p^2 + V(x) \leq 0$.

(2) The historical thread for this theorem goes back to a celebrated paper of Weyl²⁷⁴ on Dirichlet Laplacians. Theorems like XIV.1 with stronger conditions on V are due to Birman and Borzov,³⁴ Kac,¹⁴¹ Martin,¹⁹¹ and Tamura.²⁶³ See Reed and Simon,²¹² Theorem XIII.80) for the proof under the stated assumptions.

Let $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$ in a fairly regular way (e.g., suppose V is an elliptic polynomial). Then $-\Delta + V$ has discrete spectrum and the asymptotics of the number of eigenvalues $\dim E_{(-\infty, \alpha]}(-\Delta + V)$ as $\alpha \rightarrow \infty$ is determined by phase space. Results of this type go back to Titchmarsh,²⁶⁷ see also Reed and Simon,²¹² Theorem XIII.81). Similarly, if $V(x) \rightarrow 0$ but so slowly that $N(V) = \infty$, for example, $V(x) \sim -|x|^{-\beta}$ with $0 < \beta < 2$, then the divergence of $\dim E_{(-\infty, \alpha]}(-\Delta + V)$ as $\alpha \uparrow 0$ is sometimes given by quasiclassical considerations; see Brownell and Clark,⁴⁰ McLeod,¹⁹⁴ and Reed and Simon,²¹² Theorem XIII.82).

ACKNOWLEDGMENTS

I would like to thank Michael Aizenman, Brian Davies, Percy Deift, Fritz Gesztesy, Dirk Hundertmark, Walter Hunziker, and Rowan Killip for useful input.

This material is based upon work supported by the National Science Foundation under Grant No. DMS-9707661. The Government has certain rights in this material.

REFERENCES

This order of references, which departs from the usual practice of this journal, was permitted on an exceptional basis by the editor for the convenience of the reader.

- ¹T. Adachi and H. Tamura, "Asymptotic completeness for long-range many-particle systems with Stark effect, II," *Commun. Math. Phys.* **174**, 537–559 (1996).
- ²S. Agmon, "Lower bounds for solutions of Schrödinger-type equations in unbounded domains," Proceedings International Conference on Functional Analysis and Related Topics, University of Tokyo Press, Tokyo, 1969.
- ³S. Agmon, "Spectral properties of Schrödinger operators and scattering theory," *Ann. Norm. Sup. Pisa C1 Sci II* **2**, 151–218 (1975).
- ⁴S. Agmon, "Lectures on Exponential Decay of Solutions of Second-Order Elliptic Equations," Princeton University Press, Princeton (1982).
- ⁵S. Agmon, "On positive solutions of elliptic equations with periodic coefficients in R^N , spectral results and extensions to elliptic operators on Riemannian manifolds," in *Differential Equations*, edited by I. Knowles and R. Lewis (North Holland, Amsterdam, 1984).
- ⁶S. Agmon and L. Hörmander, "Asymptotic properties of solutions of differential equations with simple characteristics," *J. Anal. Math.* **30**, 1–38 (1976).
- ⁷J. Aguilar and J. M. Combes, "A class of analytic perturbations for one-body Schrödinger Hamiltonians," *Commun. Math. Phys.* **22**, 269–279 (1971).
- ⁸M. Aizenman and G. M. Graf, "Localization bounds for an electron gas," *J. Phys. A* **31**, 6783–6806 (1998).
- ⁹M. Aizenman and E. Lieb, "On semiclassical bounds for eigenvalues of Schrödinger operators," *Phys. Lett.* **66A**, 427–429 (1978).
- ¹⁰M. Aizenman and S. Molchanov, "Localization at large disorder and at extreme energies: An elementary derivation," *Commun. Math. Phys.* **157**, 245–277 (1993).
- ¹¹M. Aizenman, J. H. Schenker, R. M. Friedrich, and D. Hundertmark "Constructive fractional-moment criteria for localization in random operators," *Physica A* (to be published).
- ¹²M. Aizenman and B. Simon, "Brownian motion and Harnack's inequality for Schrödinger operators," *Commun. Pure Appl. Math.* **35**, 209–271 (1982).
- ¹³W. Allegretto, "Positive solutions and spectral properties of second order elliptic operators," *Pac. J. Math.* **92**, 15–25 (1981).
- ¹⁴W. Amrein and V. Georgescu, "On the characterization of bound states and scattering states in quantum mechanics," *Helv. Phys. Acta* **46**, 635–658 (1973).
- ¹⁵F. V. Atkinson, "On the location of the Weyl circles," *Proc. R. Soc. Edinburgh, Sect. A: Math.* **88A**, 345–356 (1981).
- ¹⁶J. Avron, "Bender-Wu formulas for the Zeeman effect in hydrogen," *Ann. Phys. (N.Y.)* **131**, 73–94 (1981).
- ¹⁷J. Avron *et al.*, "The Bender-Wu formula, $SO(4, 2)$ dynamical group and the Zeeman effect in hydrogen," *Phys. Rev. Lett.* **43**, 691–693 (1979).
- ¹⁸J. Avron and I. Herbst, "Spectral and scattering theory of Schrödinger operators related to the Stark effect," *Commun. Math. Phys.* **52**, 239–254 (1977).
- ¹⁹J. Avron, I. Herbst, and B. Simon, "Schrödinger operators with magnetic fields, I. General interactions," *Duke Math. J.* **45**, 847–883 (1978).
- ²⁰J. Avron, I. Herbst, and B. Simon, "Schrödinger operators with magnetic fields, II. Separation of center of mass in homogeneous magnetic fields," *Ann. Phys. (N.Y.)* **114**, 431–451 (1978).
- ²¹J. Avron, I. Herbst, and B. Simon, "Schrödinger operators with magnetic fields, III. Atoms in homogeneous magnetic field," *Commun. Math. Phys.* **79**, 529–572 (1981).
- ²²J. Avron and B. Simon, "Almost periodic Schrödinger operators, II. The integrated density of states," *Duke Math. J.* **50**, 369–391 (1983).
- ²³E. Balslev and J. M. Combes, "Spectral properties of many-body Schrödinger operators with dilation-analytic interactions," *Commun. Math. Phys.* **22**, 280–294 (1971).
- ²⁴V. Bargmann, "On the number of bound states in a central field of force," *Proc. Natl. Acad. Sci. USA* **38**, 961–966 (1952).
- ²⁵H. Baumgärtel and M. Wollenberg, *Mathematical Scattering Theory* (Birkhäuser, Basel, 1983).
- ²⁶E. D. Belokolos, A. I. Bobenko, V. Z. Enol'skii, A. R. Its, and V. B. Matveev, *Algebro-Geometric Approach to Nonlinear Integrable Equations* (Springer, Berlin, 1994).
- ²⁷C. Bender and T. T. Wu, "Anharmonic oscillator," *Phys. Rev.* **184**, 1231–1260 (1969).
- ²⁸R. Benguria and E. Lieb, "A proof of the stability of highly negative ions in the absence of the Pauli principle," *Phys. Rev. Lett.* **50**, 1771–1774 (1983).
- ²⁹J. Berezanskii, "On expansion according to eigenfunctions of general self-adjoint differential operators," *Dokl. Akad. Nauk SSSR* **108**, 379–382 (1956).
- ³⁰J. Berezanskii, "Expansions in eigenfunctions of self-adjoint operators," *Transl. Math. Mono.* **17**, Am. Math. Soc., Providence, RI (1968).
- ³¹F. A. Berezin, Wick and anti-Wick operator symbols, *Mat. Sb.* **86**, 578–610 (1971) [*Math. USSR Sb.* **15**, 577–606 (1971)].
- ³²M. S. Birman, "On the number of eigenvalues in a quantum scattering problem," *Vest. LSU* **16**, 163–166 (1961).
- ³³M. S. Birman, "Conditions for the existence of wave operators," *Dokl. Akad. Nauk SSSR* **143**, 506–509 (1962).
- ³⁴M. S. Birman and V. V. Borzov, "On the asymptotics of the discrete spectrum of some singular differential operators," *Topics Math. Phys.* **5**, 19–30 (1972).

- ³⁵M. S. Birman and M. G. Krein, "On the theory of wave operators and scattering operators," Dokl. Akad. Nauk SSSR **144**, 475–478 (1962).
- ³⁶M. S. Birman and M. Z. Solomyak, "Estimates for the number of negative eigenvalues of the Schrödinger operator and its generalizations," Advances Soviet Mathematics, edited by M. S. Birman (American Mathematical Society, Providence, RI, 1991), Vol. 7, pp. 1–55.
- ³⁷G. Borg, "Uniqueness theorems in the spectral theory of $y'' + (\lambda - q(x))y = 0$," Proceedings 11th Scandinavian Congress of Mathematicians, Johan Grundt Tanums Forlag, Oslo, 1952, pp. 276–287.
- ³⁸J. Bourgain and M. Goldstein, "On nonperturbative localization with quasiperiodic potential," preprint.
- ³⁹F. Browder, "Eigenfunction expansions for formally self-adjoint partial differential operators, I, II," Proc. Natl. Acad. Sci. USA **42**, 769–771 (1956); **42**, 870–872 (1956).
- ⁴⁰F. H. Brownell and C. W. Clark, "Asymptotic distribution of the eigenvalues of the lower part of the Schrödinger operator spectrum," J. Math. Mech. **10**, 31–70 (1961).
- ⁴¹R. Carmona, "Regularity properties of Schrödinger and Dirichlet semigroups," J. Funct. Anal. **17**, 227–237 (1974).
- ⁴²R. Carmona, "One-dimensional Schrödinger operators with random or deterministic potentials: New spectral types," J. Funct. Anal. **51**, 229–258 (1983).
- ⁴³R. Carmona, A. Klein, and F. Martinelli, "Anderson localization for Bernoulli and other singular potentials," Commun. Math. Phys. **108**, 41–66 (1987).
- ⁴⁴R. Carmona and J. Lacroix, *Spectral Theory of Random Schrödinger Operators* (Birkhäuser, Boston, 1990).
- ⁴⁵K. Chadan and P. C. Sabatier, *Inverse Problems in Quantum Scattering Theory*, 2nd ed., (Springer, New York, 1989).
- ⁴⁶M. Christ and A. Kiselev, "Absolutely continuous spectrum for one-dimensional Schrödinger operators with slowly decaying potentials: Some optimal results," J. Am. Math. Soc. **11**, 771–797 (1998).
- ⁴⁷J. M. Combes and L. Thomas, "Asymptotic behavior of eigenfunctions for multi-particle Schrödinger operators," Commun. Math. Phys. **34**, 251–270 (1973).
- ⁴⁸M. Combes and J. Ginibre, "Spectral and scattering theory for the Schrödinger operator with strongly oscillating potentials," Ann. Inst. Henri Poincaré, Sect. A **A24**, 17–29 (1976).
- ⁴⁹J. Conlon, "The ground state energy of a classical gas," Commun. Math. Phys. **94**, 439–458 (1984).
- ⁵⁰J. Conlon, E. Lieb, and H.-T. Yau, "The $N^{7/5}$ law for charged bosons," Commun. Math. Phys. **116**, 417–448 (1988).
- ⁵¹J. Cook, "Convergence of the Moller wave matrix," J. Math. and Phys. **36**, 82–87 (1957).
- ⁵²M. Cwikel, "Weak type estimates for singular values and the number of bound states of Schrödinger operators," Ann. Math. **106**, 93–102 (1977).
- ⁵³H. Cycon, R. Froese, W. Kirsch, and B. Simon, *Schrödinger Operators With Application to Quantum Mechanics and Global Geometry* (Springer, Berlin, 1987).
- ⁵⁴E. B. Davies, "Properties of Green's functions of some Schrödinger operators," J. London Math. Soc. **7**, 473–491 (1973).
- ⁵⁵E. B. Davies, *Heat Kernels and Spectral Theory* (Cambridge University Press, Cambridge, 1989).
- ⁵⁶E. B. Davies, " L^p spectral theory of higher-order elliptic differential operators," Bull. London Math. Soc. **29**, 513–546 (1997).
- ⁵⁷E. B. Davies and Y. Safarov (eds.), *Spectral Theory and Geometry*, London Math. Soc. Lecture Notes, Vol. 273 (Cambridge University Press, Cambridge, 1999).
- ⁵⁸P. Deift and R. Hempel, "On the existence of eigenvalues of the Schrödinger operator $H - \lambda W$ in a gap of $\sigma(H)$," Commun. Math. Phys. **103**, 461–490 (1986).
- ⁵⁹P. Deift, W. Hunziker, B. Simon, and E. Vock, "Pointwise bounds on eigenfunctions and wave packets in N -body quantum systems, IV," Commun. Math. Phys. **64**, 1–34 (1978).
- ⁶⁰P. Deift and R. Killip, "On the absolutely continuous spectrum of one-dimensional Schrödinger operators with square summable potentials," Commun. Math. Phys. **203**, 341–347 (1999).
- ⁶¹P. Deift and E. Trubowitz, "Inverse scattering on the line," Commun. Pure Appl. Math. **32**, 121–251 (1979).
- ⁶²F. Delyon, "Apparition of purely singular continuous spectrum in a class of random Schrödinger operators," J. Stat. Phys. **40**, 621–630 (1985).
- ⁶³F. Delyon, Y. Levy, and B. Souillard, "Anderson localization for multidimensional systems at large disorder or large energy," Commun. Math. Phys. **100**, 463–470 (1985).
- ⁶⁴F. Delyon, B. Simon, and B. Souillard, "From power pure point to continuous spectrum in disordered systems," Ann. Inst. Henri Poincaré Phys. Theor. **42**, 283–309 (1985).
- ⁶⁵J. Dereziński, "Asymptotic completeness of long-range N -body quantum systems," Ann. Math. **138**, 427–476 (1993).
- ⁶⁶J. Dereziński and C. Gérard, *Scattering Theory of Classical and Quantum N -Particle Systems* (Springer, Berlin, 1997).
- ⁶⁷E. Dinaburg and Ya. Sinai, "The one-dimensional Schrödinger equation with a quasiperiodic potential," Funct. Anal. Appl. **9**, 279–289 (1975).
- ⁶⁸R. K. Dodd, J. C. Eilbeck, J. D. Gibbon, and H. C. Morris, *Solitons and Nonlinear Wave Equations* (Academic, London, 1982).
- ⁶⁹J. Dollard, "Asymptotic convergence and the Coulomb interaction," J. Math. Phys. **5**, 729–738 (1964).
- ⁷⁰B. A. Dubrovin, V. B. Matveev, and S. P. Novikov, "Nonlinear equations of Korteweg-de Vries type, finite-zone linear operators, and Abelian varieties," Russ. Math. Surveys **31:1**, 59–146 (1976).
- ⁷¹F. Dyson, "Ground-state energy of a finite system of charged particles," J. Math. Phys. **8**, 1538–1545 (1967).
- ⁷²F. Dyson and A. Lenard, "Stability of matter, I," J. Math. Phys. **8**, 423–434 (1967).
- ⁷³F. Dyson and A. Lenard, "Stability of matter, II," J. Math. Phys. **9**, 698–711 (1968).
- ⁷⁴V. N. Efimov, "Energy levels arising from resonant two-body forces in a three-body system," Phys. Lett. **B33**, 563–564 (1970).

- ⁷⁵ V. N. Efimov, "Weakly-bound states of three resonantly-interacting particles," *Sov. J. Nucl. Phys.* **12**, 589–595 (1971).
- ⁷⁶ V. Enss, "A note on Hunziker's theorem," *Commun. Math. Phys.* **52**, 233–238 (1977).
- ⁷⁷ V. Enss, "Asymptotic completeness for quantum-mechanical potential scattering, I. Short-range potentials," *Commun. Math. Phys.* **61**, 285–291 (1978).
- ⁷⁸ V. Enss, "Completeness of three-body quantum scattering," in *Dynamics and Processes*, edited by P. Blanchard and L. Streit, Lecture Notes in Math, Vol. 1031 (Springer, Berlin, 1983), pp. 62–88.
- ⁷⁹ L. Faddeev, *Mathematical Aspects of the Three Body Problem in Quantum Scattering Theory* (Steklov Institute, 1963).
- ⁸⁰ W. Faris and R. Lavine, "Commutators and self-adjointness of Hamiltonian operators," *Commun. Math. Phys.* **35**, 39–48 (1974).
- ⁸¹ C. Fefferman, J. Fröhlich, and G. M. Graf, "Stability of ultraviolet-cutoff quantum electrodynamics with non-relativistic matter," *Commun. Math. Phys.* **190**, 309–330 (1997).
- ⁸² C. Fefferman and L. Seco, "On the energy of a large atom," *Bull. Am. Math. Soc.* **23**, 525–530 (1990).
- ⁸³ R. Froese, "Asymptotic distribution of resonances in one dimension," *J. Diff. Eqns.* **137**, 251–272 (1997).
- ⁸⁴ R. Froese, "Upper bounds for the resonance counting function in odd dimension," *Can. J. Math.* **50**, 538–546 (1998).
- ⁸⁵ R. Froese and I. Herbst, "A new proof of the Mourre estimate," *Duke Math. J.* **49**, 1075–1085 (1982).
- ⁸⁶ R. Froese, I. Herbst, M. Hoffmann-Ostenhof, and T. Hoffmann-Ostenhof, " L^2 -exponential lower bounds to solutions of the Schrödinger equation," *Commun. Math. Phys.* **87**, 265–286 (1982/83).
- ⁸⁷ J. Fröhlich and T. Spencer, "Absence of diffusion in the Anderson tight binding model for large disorder or low energy," *Commun. Math. Phys.* **88**, 151–184 (1983).
- ⁸⁸ L. Garding, "Eigenfunction expansions connected with elliptic differential operators," *Tolte Skend. Mat. Lund*, 44–45 (1953).
- ⁸⁹ I. Gel'fand, "Expansion in characteristic functions of an equation with periodic coefficients," *Dokl. Akad. Nauk SSSR* **73**, 1117–1120 (1950).
- ⁹⁰ I. M. Gel'fand and B. M. Levitan, "On the determination of a differential equation from its special function," *Izv. Akad. Nauk SSSR Ser. Mat.* **15**, 309–360 (1951) (Russian); [*Amer. Math. Soc. Transl. Ser. 2* **1**, 253–304 (1955)].
- ⁹¹ F. Gesztesy and B. Simon, "A short proof of Zheludev's theorem," *Trans. Am. Math. Soc.* **335**, 329–340 (1993).
- ⁹² F. Gesztesy and B. Simon, "The xi function," *Acta Math.* **176**, 49–71 (1996).
- ⁹³ F. Gesztesy and B. Simon, "A new approach to inverse spectral theory, II. General real potentials and the connection to the spectral measure," *Ann. Math.* (to be published).
- ⁹⁴ F. Gesztesy and B. Simon, "On local Borg-Marchenko uniqueness results," *Commun. Math. Phys.* (to be published).
- ⁹⁵ D. J. Gilbert, "On subordinacy and analysis of the spectrum of Schrödinger operators with two singular endpoints," *Proc. R. Soc. Edinburgh, Sect. A: Math.* **112**, 213–229 (1989).
- ⁹⁶ D. J. Gilbert, "On subordinacy and spectral multiplicity for a class of singular differential operators," *Proc. R. Soc. Edinburgh, Sect. A: Math.* **128**, 549–584 (1998).
- ⁹⁷ D. J. Gilbert and D. B. Pearson, "On subordinacy and analysis of the spectrum of one-dimensional Schrödinger operators," *J. Math. Anal. Appl.* **128**, 30–56 (1987).
- ⁹⁸ I. Goldsheid, S. Molchanov, and L. Pastur, "A pure point spectrum of the stochastic one-dimensional Schrödinger equation," *Funct. Anal. Appl.* **11**, 1–10 (1977).
- ⁹⁹ A. Gordon, "On the point spectrum of one-dimensional Schrödinger operators," *Usp. Mat. Nauk* **31**, 257–258 (1976).
- ¹⁰⁰ A. Gordon, S. Jitomirskaya, Y. Last, and B. Simon, "Duality and singular continuous spectrum in the almost Mathieu equation," *Acta Math.* **178**, 169–183 (1997).
- ¹⁰¹ G. M. Graf, "Asymptotic completeness for N-body short-range quantum systems: A new proof," *Commun. Math. Phys.* **132**, 73–101 (1990).
- ¹⁰² S. Graffi, V. Grecchi, and B. Simon, "Borel summability: Application to the anharmonic oscillator," *Phys. Lett.* **32B**, 631–634 (1970).
- ¹⁰³ M. N. Hack, "Wave operators in multichannel scattering," *Nuovo Cimento* **13**, 231–236 (1959).
- ¹⁰⁴ G. Hagedorn, "Semiclassical quantum mechanics, I. The $\hbar \rightarrow 0$ limit for coherent states," *Commun. Math. Phys.* **71**, 77–93 (1980).
- ¹⁰⁵ G. Hagedorn, *Semiclassical Quantum Mechanics, II. The Large Mass Asymptotics for Coherent States* (Pitman, Boston, 1981).
- ¹⁰⁶ G. Hagedorn, "Semiclassical quantum mechanics, III. The large order asymptotics and more general states," *Ann. Phys. (N.Y.)* **135**, 58–70 (1981).
- ¹⁰⁷ E. Harrell and B. Simon, "The mathematical theory of resonances which have exponentially small widths," *Duke Math. J.* **47**, 845–902 (1980).
- ¹⁰⁸ B. Helffer, M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, and M. Owen, "Nodal sets for groundstates of Schrödinger operators with zero magnetic field in non-simply connected domains," *Commun. Math. Phys.* **202**, 629–649 (1999).
- ¹⁰⁹ B. Helffer and D. Robert, "Riesz means of bounded states and semi-classical limit connected with a Lieb-Thirring conjecture," *Asymp. Anal.* **3**, 91–103 (1990).
- ¹¹⁰ B. Helffer and D. Robert, "Riesz means of bounded states and semi-classical limit connected with a Lieb-Thirring conjecture, II," *Ann. Inst. Henri Poincaré Phys. Theor.* **53**, 139–147 (1990).
- ¹¹¹ B. Helffer and J. Sjöstrand, "Multiple wells in the semiclassical limit, I," *Commun. Partial Diff. Eqns.* **9**, 337–408 (1984).
- ¹¹² B. Helffer and J. Sjöstrand, "Résonances en limite semi-classique," *Mém. Soc. Math. France (N.S.)* No. **24**, 24–25 (1986).
- ¹¹³ R. Hempel and J. Voigt, "The spectrum of a Schrödinger operator in $L^p(\mathbb{R}^n)$ is p -independent," *Commun. Math. Phys.* **104**, 243–250 (1986).

- ¹¹⁴I. Herbst, "Unitary equivalences of Stark-Hamiltonians," *Math. Z.* **155**, 55–70 (1977).
- ¹¹⁵I. Herbst, "Dilation analyticity in constant electric field, I. The two-body problem," *Commun. Math. Phys.* **64**, 279–298 (1979).
- ¹¹⁶I. Herbst, J. Møller, and E. Skibsted, "Asymptotic completeness for N -body Stark Hamiltonians," *Commun. Math. Phys.* **174**, 509–535 (1996).
- ¹¹⁷I. Herbst and B. Simon, "Some remarkable examples in eigenvalue perturbation theory," *Phys. Lett.* **78B**, 304–306 (1978).
- ¹¹⁸I. Herbst and S. Sloan, "Perturbations of translation invariant positivity preserving semigroups in $L^2(R)$," *Trans. Am. Math. Soc.* **236**, 325–360 (1978).
- ¹¹⁹H. Hess, R. Schrader, and D. Uhlenbrock, "Domination of semigroups and generalization of Kato's inequality," *Duke Math. J.* **44**, 893–904 (1977).
- ¹²⁰M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, and N. Nadirashvili, "Interior Hölder estimates for solutions of Schrödinger equations and the regularity of nodal sets," *Commun. Partial Diff. Eqns.* **20**, 1241–1273 (1995).
- ¹²¹M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, and H. Stremmitzer, "Local properties of Coulombic wave functions," *Commun. Math. Phys.* **163**, 185–215 (1994).
- ¹²²L. Hörmander, *The Analysis of Linear Partial Differential Operators, IV. Fourier Integral Operators* (Springer, Berlin, 1994).
- ¹²³W. Hughes, "An atomic energy lower bound that agrees with Scott's correction," *Adv. Math.* **79**, 213–270 (1990).
- ¹²⁴D. Hundertmark, A. Laptev, and T. Weidl, "New bounds on the Lieb-Thirring constants," *Invent. Math.* (to be published).
- ¹²⁵D. Hundertmark, E. Lieb, and L. Thomas, "A sharp bound for an eigenvalue moment of the one-dimensional Schrödinger operator," *Adv. Theor. Math. Phys.* **2**, 719–731 (1998).
- ¹²⁶W. Hunziker, "On the spectra of Schrödinger multiparticle Hamiltonians," *Helv. Phys. Acta* **39**, 451–462 (1966).
- ¹²⁷W. Hunziker, "Mathematical theory of N -body quantum systems," *Helv. Phys. Acta* **71**, 26–43 (1998).
- ¹²⁸W. Hunziker and I. M. Sigal, "The quantum N -body problem," preprint.
- ¹²⁹T. Ikebe, "Eigenfunction expansions associated with the Schrödinger operators and their application to scattering theory," *Arch. Ration. Mech. Anal.* **5**, 1–34 (1960).
- ¹³⁰T. Ikebe and T. Kato, "Uniqueness of the self-adjoint extension of singular elliptic differential operators," *Arch. Ration. Mech. Anal.* **9**, 77–92 (1962).
- ¹³¹R. Iorio and M. O'Carroll, "Asymptotic completeness for the multi-particle Schrödinger Hamiltonians with weak potentials," *Commun. Math. Phys.* **27**, 137–145 (1972).
- ¹³²A. R. Its and V. B. Matveev, "Schrödinger operators with finite-gap spectrum and N -soliton solutions of the Korteweg-de Vries equation," *Theor. Math. Phys.* **23**, 343–355 (1975).
- ¹³³J. M. Jauch, "Theory of the scattering operator," *Helv. Phys. Acta* **31**, 127–158 (1958).
- ¹³⁴S. Jitomirskaya, "Metal-insulator transition for the almost Mathieu operator," *Ann. Math.* (to be published).
- ¹³⁵S. Jitomirskaya and Y. Last, "Dimensional Hausdorff properties of singular continuous spectra," *Phys. Rev. Lett.* **76**, 1765–1769 (1996).
- ¹³⁶S. Jitomirskaya and B. Simon, "Operators with singular continuous spectrum, III. Almost periodic Schrödinger operators," *Commun. Math. Phys.* **165**, 201–205 (1994).
- ¹³⁷G. Kac, "Expansion in characteristic functions of self-adjoint operators," *Dokl. Akad. Nauk SSSR* **119**, 19–22 (1958).
- ¹³⁸I. S. Kac, "On the multiplicity of the spectrum of a second-order differential operator," *Sov. Math. Dokl.* **3**, 1035–1039 (1962).
- ¹³⁹I. S. Kac, "Spectral multiplicity of a second-order differential operator and expansion in eigenfunction," *Izv. Akad. Nauk SSSR, Ser. Mat.* **27**, 1081–1112 (1963).
- ¹⁴⁰M. Kac, "On some connections between probability theory and differential equations," *Proceedings 2nd Berkeley Symposium on Mathematical Statistics and Probability*, University of California Press, Berkeley and Los Angeles, 1951, pp. 189–215.
- ¹⁴¹M. Kac, "On the asymptotic number of bound states for certain attractive potentials" in *Topics in Functional Analysis*, edited by I. Gohberg and M. Kac (Academic, New York, 1978), pp. 159–167.
- ¹⁴²T. Kato, "On the convergence of the perturbation method, I," *Prog. Theor. Phys.* **4**, 514–523 (1949).
- ¹⁴³T. Kato, "On the convergence of the perturbation method, II," *Prog. Theor. Phys.* **5**, 95–101 (1950); **5**, 207–212 (1950).
- ¹⁴⁴T. Kato, "Fundamental properties of Hamiltonian operators of Schrödinger type," *Trans. Am. Math. Soc.* **70**, 195–211 (1951).
- ¹⁴⁵T. Kato, "On the existence of solutions of the helium wave equation," *Trans. Am. Math. Soc.* **70**, 212–218 (1951).
- ¹⁴⁶T. Kato, "On the convergence of the perturbation method," *J. Fac. Sci., Univ. Tokyo, Sect. I* **6**, 145–226 (1951).
- ¹⁴⁷T. Kato, "Perturbation theory of semi-bounded operators," *Math. Ann.* **125**, 435–447 (1953).
- ¹⁴⁸T. Kato, "On the eigenfunctions of many particle systems in quantum mechanics," *Commun. Pure Appl. Math.* **10**, 151–171 (1957).
- ¹⁴⁹T. Kato, "Perturbation of continuous spectra by trace class operators," *Proc. Jpn. Acad.* **33**, 260–264 (1957).
- ¹⁵⁰T. Kato, "Growth properties of solutions of the reduced wave equation with variable coefficients," *Commun. Pure Appl. Math.* **12**, 403–425 (1959).
- ¹⁵¹T. Kato, "Schrödinger operators with singular potentials," *Isr. J. Math.* **13**, 135–148 (1972).
- ¹⁵²T. Kato, "Remarks on Schrödinger operators with vector potentials," *Int. Equ. Operator Theory* **1**, 103–113 (1978).
- ¹⁵³T. Kato, *Perturbation Theory for Linear Operators*, 2nd ed. (Springer, Berlin, 1980).
- ¹⁵⁴C. E. Kenig, *Harmonic Analysis Techniques for Second Order Elliptic Boundary Value Problems* (American Mathematical Society, Providence, RI, 1994).

- ¹⁵⁵R. Killip, "Perturbations of one-dimensional Schrödinger operators preserving the a.c. spectrum," preprint.
- ¹⁵⁶A. Kiselev, "Absolutely continuous spectrum of one-dimensional Schrödinger operators and Jacobi matrices with slowly decreasing potentials," *Commun. Math. Phys.* **179**, 377–400 (1996).
- ¹⁵⁷A. Kiselev, "Stability of the absolutely continuous spectrum of the Schrödinger equation under slowly decaying perturbations and a.e. convergence of integral operators," *Duke Math. J.* **94**, 619–646 (1998).
- ¹⁵⁸A. Kiselev, Y. Last, and B. Simon, "Modified Prüfer and EFGP transforms and the spectral analysis of one-dimensional Schrödinger operators," *Commun. Math. Phys.* **194**, 1–45 (1998).
- ¹⁵⁹S. Kotani, "Lyapunov exponents and spectra for one-dimensional random Schrödinger operators," *Contemporary Mathematics* (American Mathematical Society, Providence, RI, 1986), Vol. 50.
- ¹⁶⁰S. Kotani and N. Ushiroya, "One-dimensional Schrödinger operators with random decaying potentials," *Commun. Math. Phys.* **115**, 247–266 (1988).
- ¹⁶¹V. Kovalenko and Yu. Semenov, "Some problems on expansions in generalized eigenfunctions of the Schrödinger operator with strongly unique potentials," *Rus. Math. Surveys* **33**, 119–157 (1978).
- ¹⁶²M. G. Krein, "Solution of the inverse Sturm-Liouville problem," *Dokl. Akad. Nauk SSSR* **76**, 21–24 (1951).
- ¹⁶³M. G. Krein, "Determination of the density of a nonhomogeneous symmetric cord by its frequency spectrum," *Dokl. Akad. Nauk SSSR* **76**, 345–348 (1951).
- ¹⁶⁴M. G. Krein, "On determination of the potential of a particle from its S -function," *Dokl. Akad. Nauk SSSR* **105**, 433–436 (1955).
- ¹⁶⁵H. Kunz and B. Souillard, "Sur le spectre des operateurs aux differences finies aleatoires," *Commun. Math. Phys.* **78**, 201–246 (1980).
- ¹⁶⁶S. Kuroda, "Perturbations of continuous spectra by unbounded operators, I," *J. Math. Soc. Jpn.* **11**, 247–262 (1959).
- ¹⁶⁷S. Kuroda, "Perturbations of continuous spectra by unbounded operators, II," *J. Math. Soc. Jpn.* **12**, 243–257 (1960).
- ¹⁶⁸A. Laptev and T. Weidl, "Sharp Lieb-Thirring inequalities in high dimensions," *Acta Math* **184**, 87–111 (2000).
- ¹⁶⁹Y. Last, "A relation between a.c. spectrum of ergodic Jacobi matrices and the spectra of periodic approximants," *Commun. Math. Phys.* **151**, 183–192 (1993).
- ¹⁷⁰Y. Last and B. Simon, "Eigenfunctions, transfer matrices, and absolutely continuous spectrum of one-dimensional Schrödinger operators," *Invent. Math.* **135**, 329–367 (1999).
- ¹⁷¹R. Lavine, "Absolute continuity of Hamiltonian operators with repulsive potentials," *Proc. Am. Math. Soc.* **22**, 55–60 (1969).
- ¹⁷²R. Lavine, "Absolute continuity of positive spectrum for Schrödinger operators with long range potentials," *J. Funct. Anal.* **12**, 30–54 (1973).
- ¹⁷³H. Leinfelder and C. Simader, "Schrödinger operators with singular magnetic vector potentials," *Math. Z.* **176**, 1–19 (1981).
- ¹⁷⁴B. Ya. Levin, "Fourier- and Laplace-type transformations by means of solutions of a second-order differential equation," *Dokl. Akad. Nauk SSSR* **106**, 187–190 (1956).
- ¹⁷⁵B. M. Levitan, *Theory of Generalized Translation Operators*, 2nd ed. (Nauka, Moscow, 1973).
- ¹⁷⁶B. M. Levitan, *Inverse Sturm-Liouville Problems* (VNU Science, Utrecht, 1987).
- ¹⁷⁷E. Lieb, "Bounds on the eigenvalues of the Laplace and Schrödinger operators," *Bull. Am. Math. Soc.* **82**, 751–753 (1976).
- ¹⁷⁸E. Lieb, "Bound on the maximum negative ionization of atoms and molecules," *Phys. Rev. A* **29**, 3018–3028 (1984).
- ¹⁷⁹E. Lieb, "The stability of matter: From atoms to stars," *Bull. Am. Math. Soc.* **22**, 1–49 (1990).
- ¹⁸⁰E. Lieb, M. Loss, and J. P. Solovej, "Stability of matter in magnetic fields," *Phys. Rev. Lett.* **75**, 985–989 (1995).
- ¹⁸¹E. Lieb, I. M. Sigal, B. Simon, and W. Thirring, "Asymptotic neutrality of large Z ions," *Phys. Rev. Lett.* **52**, 994–996 (1984).
- ¹⁸²E. Lieb and B. Simon, "The Thomas-Fermi theory of atoms, molecules, and solids," *Adv. Math.* **23**, 22–116 (1977).
- ¹⁸³E. Lieb, J. P. Solovej, and J. Yngvason, "Asymptotics of heavy atoms in high magnetic fields, I. Lowest Landau band regions," *Commun. Pure Appl. Math.* **47**, 513–591 (1994).
- ¹⁸⁴E. Lieb, J. P. Solovej, and J. Yngvason, "Asymptotics of heavy atoms in high magnetic fields, II. Semiclassical regions," *Commun. Math. Phys.* **161**, 77–124 (1994).
- ¹⁸⁵E. Lieb and W. Thirring, "Bound for the kinetic energy of fermions which proves the stability of matter," *Phys. Rev. Lett.* **35**, 687–689 (1975).
- ¹⁸⁶E. Lieb and W. Thirring, "Inequalities for the moments of the eigenvalues of the Schrödinger Hamiltonian and their relation to Sobolev inequalities," in *Studies in Mathematical Physics: Essays in Honor of Valentine Bargmann*, edited by E. Lieb, B. Simon, and A. Wightman (Princeton University Press, Princeton, 1976), pp. 269–303.
- ¹⁸⁷J. Loeffel, A. Martin, B. Simon, and A. Wightman, "Padé approximants and the anharmonic oscillator," *Phys. Lett.* **30B**, 656–658 (1969).
- ¹⁸⁸V. A. Marchenko, "Certain problems in the theory of second-order differential operators," *Dokl. Akad. Nauk SSSR* **72**, 457–460 (1950).
- ¹⁸⁹V. A. Marchenko, "Some questions in the theory of one-dimensional linear differential operators of the second order, I," *Am. Math. Soc. Transl.* **101**, 1–104 (1973).
- ¹⁹⁰V. A. Marchenkov, *Sturm-Liouville Operators and Applications* (Birkhäuser, Basel, 1986).
- ¹⁹¹A. Martin, "Bound states in the strong coupling limit," *Helv. Phys. Acta* **45**, 140–148 (1972).
- ¹⁹²H. McKean and E. Trubowitz, "Hill's operator and hypoelliptic function theory in the presence of infinitely many branch points," *Commun. Pure Appl. Math.* **29**, 143–226 (1976).
- ¹⁹³H. McKean and P. van Moerbeke, "The spectrum of Hill's equation," *Invent. Math.* **30**, 217–274 (1975).
- ¹⁹⁴J. B. McLeod, "The distribution of eigenvalues for the hydrogen atom and similar cases," *Proc. London Math. Soc.* **11**, 139–158 (1961).

- ¹⁹⁵ A. Melin, "Operator methods for inverse scattering on the real line," *Commun. Partial Diff. Eqns.* **10**, 677–766 (1985).
- ¹⁹⁶ E. Mourre, "Absence of singular continuous spectrum for certain selfadjoint operators," *Commun. Math. Phys.* **78**, 391–408 (1981).
- ¹⁹⁷ S. N. Naboko, "On the dense point spectra of Schrödinger and Dirac operators," *Theor. Math. Phys.* **68**, 18–28 (1986).
- ¹⁹⁸ E. Nelson (unpublished).
- ¹⁹⁹ S. Novikov, S. V. Manakov, L. P. Pitaevskii, and V. E. Zakharov, *Theory of Solitons* (Consultants Bureau, New York, 1984).
- ²⁰⁰ A. O'Connor, "Exponential decay of bound state wave functions," *Commun. Math. Phys.* **32**, 319–340 (1973).
- ²⁰¹ Yu. Ovchinnikov and I. M. Sigal, "Number of bound states of three-body systems and Efimov's effect," *Ann. Phys. (N.Y.)* **123**, 274–295 (1979).
- ²⁰² L. Pastur, "Spectral properties of disordered systems in the one-body approximation," *Commun. Math. Phys.* **75**, 179–196 (1980).
- ²⁰³ L. Pastur and A. Figotin, *Spectra of Random and Almost-Periodic Schrödinger Operators* (Springer, Berlin, 1992).
- ²⁰⁴ D. B. Pearson, "Singular continuous measures in scattering theory," *Commun. Math. Phys.* **60**, 13–36 (1978).
- ²⁰⁵ P. Perry, I. M. Sigal, and B. Simon, "Spectral analysis of multiparticle Schrödinger operators," *Ann. Math.* **114**, 519–567 (1981).
- ²⁰⁶ J. Piepenbrink, "Nonoscillatory elliptic equations," *J. Diff. Eqns.* **15**, 541–550 (1974).
- ²⁰⁷ Y. Pinchover, "On positive solutions of second-order elliptic equations, stability results and classification," *Duke Math. J.* **57**, 955–980 (1988).
- ²⁰⁸ A. Ya. Povzner, "On differential equations of Sturm-Liouville type on a half-axis," *Mat. Sb.* **23**, 3–52 (1948) (Russian); [*Am. Math. Soc. Transl.* **4**, 24–101 (1962)].
- ²⁰⁹ A. Ya. Povzner, "On the expansion of arbitrary functions in terms of the eigenfunctions of the operator $-\Delta u + cu$," *Mat. Sb.* **43**, 109–156 (1953), (Russian); [*Am. Math. Soc. Transl.* **60**, 268–302 (1967)].
- ²¹⁰ J. Ralston, "On the construction of quasimodes associated with stable periodic orbits," *Commun. Math. Phys.* **51**, 219–242 (1976), **67**, 91 (1979) (Erratum).
- ²¹¹ M. Reed and B. Simon, *Methods of Modern Mathematical Physics, II. Fourier Analysis, Self-Adjointness* (Academic, New York, 1975).
- ²¹² M. Reed and B. Simon, *Methods of Modern Mathematical Physics, IV. Analysis of Operators* (Academic, New York, 1978).
- ²¹³ M. Reed and B. Simon, *Methods of Modern Mathematical Physics, III. Scattering Theory* (Academic, New York, 1979).
- ²¹⁴ M. Reed and B. Simon, *Methods of Modern Mathematical Physics, I. Functional Analysis*, rev. ed. (Academic, New York, 1980).
- ²¹⁵ F. Rellich, "Störungstheorie der Spektralzerlegung I-V," *Math. Ann.* **113**, 600–619 (1937); **113**, 677–685 (1937); **116**, 555–570 (1939); **117**, 356–382 (1940); **118**, 462–484 (1942).
- ²¹⁶ F. Rellich, "Über das asymptotische Verhalten der Lösungen von $\Delta u + \lambda u = 0$ in unendlichen Gebieten," *Jber. Deutsch. Math. Verein* **53**, 57–65 (1943).
- ²¹⁷ C. Remling, "A probabilistic approach to one-dimensional Schrödinger operators with sparse potentials," *Commun. Math. Phys.* **185**, 313–323 (1997).
- ²¹⁸ C. Remling, "The absolutely continuous spectrum of one-dimensional Schrödinger operators with decaying potentials," *Commun. Math. Phys.* **193**, 151–170 (1998).
- ²¹⁹ D. Robert, "Semi-classical approximation in quantum mechanics. A survey of old and recent mathematical results," *Helv. Phys. Acta* **71**, 44–116 (1998).
- ²²⁰ G. V. Rosenbljum, "The distribution of the discrete spectrum for singular differential operators," *Dokl. Akad. Nauk SSSR* **202**, 1012–1015 (1972) (Russian); [*Sov. Math. Dokl.* **13**, 245–249 (1972)].
- ²²¹ M. Rosenblum, "Perturbations of continuous spectrum and unitary equivalence," *Pac. J. Math.* **7**, 997–1010 (1957).
- ²²² D. Ruelle, "A remark on bound states in potential scattering theory," *Nuovo Cimento A* **61A**, 655–662 (1969).
- ²²³ M. B. Ruskai, "Absence of discrete spectrum in highly negative ions," *Commun. Math. Phys.* **82**, 457–469 (1982).
- ²²⁴ M. B. Ruskai, "Absence of discrete spectrum in highly negative ions, II," *Commun. Math. Phys.* **85**, 325–327 (1982).
- ²²⁵ A. Sá Barreto and M. Zworski, "Existence of resonances in potential scattering," *Commun. Pure Appl. Math.* **49**, 1271–1280 (1996).
- ²²⁶ M. Schechter, *Spectra of Partial Differential Operators* (North Holland, Amsterdam, 1971).
- ²²⁷ J. Schwinger, "On the bound states of a given potential," *Proc. Natl. Acad. Sci. USA* **47**, 122–129 (1961).
- ²²⁸ C. Shubin, R. Vakilian, and T. Wolff, "Some harmonic analysis questions suggested by Anderson-Bernoulli models," *Geom. Funct. Anal.* **8**, 932–964 (1998).
- ²²⁹ H. Siedentop and R. Weikard, "On the leading energy correction for the statistical model of the atom: Interacting case," *Commun. Math. Phys.* **112**, 471–490 (1987).
- ²³⁰ H. Siedentop and R. Weikard, "On the leading correction of the Thomas-Fermi model: Lower bound," *Invent. Math.* **97**, 159–193 (1989).
- ²³¹ I. M. Sigal, "Geometric methods in the quantum many-body problem. Nonexistence of very negative ions," *Commun. Math. Phys.* **85**, 309–324 (1982).
- ²³² I. M. Sigal, "How many electrons can a nucleus bind?," *Ann. Phys. (N.Y.)* **157**, 307–320 (1984).
- ²³³ I. M. Sigal and A. Soffer, "The N -particle scattering problem: Asymptotic completeness for short-range systems," *Ann. Math.* **126**, 35–108 (1987).
- ²³⁴ I. M. Sigal and A. Soffer, "Asymptotic completeness of N -particle long-range scattering," *J. Am. Math. Soc.* **7**, 307–334 (1994).

- ²³⁵B. Simon, "On positive eigenvalues of one-body Schrödinger operators," *Commun. Pure Appl. Math.* **22**, 531–538 (1969).
- ²³⁶B. Simon, "On the infinitude or finiteness of the number of bound states of an N -body quantum system, I," *Helv. Phys. Acta* **43**, 607–630 (1970).
- ²³⁷B. Simon, "Quadratic form techniques and the Balslev-Combes theorem," *Commun. Math. Phys.* **27**, 1–9 (1972).
- ²³⁸B. Simon, "Resonances in N -body quantum systems with dilation analytic potentials and the foundations of time-dependent perturbation theory," *Ann. Math.* **97**, 247–274 (1973).
- ²³⁹B. Simon, "Essential self-adjointness of Schrödinger operators with positive potentials," *Math. Ann.* **201**, 211–220 (1973).
- ²⁴⁰B. Simon, "On the number of bound states of two-body Schrödinger operators—A review," in *Studies in Mathematical Physics: Essays in Honor of Valentine Bargmann*, edited by E. Lieb, B. Simon, and A. Wightman (Princeton University Press, Princeton, 1976), pp. 305–326.
- ²⁴¹B. Simon, "Universal diamagnetism of spinless Bose systems," *Phys. Rev. Lett.* **36**, 1083–1084 (1976).
- ²⁴²B. Simon, "The bound state of weakly coupled Schrödinger operators in one and two dimensions," *Ann. Phys. (N.Y.)* **97**, 279–288 (1976).
- ²⁴³B. Simon, "Analysis with weak trace ideals and the number of bound states of Schrödinger operators," *Trans. Am. Math. Soc.* **224**, 367–380 (1976).
- ²⁴⁴B. Simon, "Geometric methods in multiparticle quantum systems," *Commun. Math. Phys.* **55**, 259–274 (1977).
- ²⁴⁵B. Simon, "Complex scaling: A rigorous overview," *Int. J. Quantum Chem.* **14**, 529–542 (1978).
- ²⁴⁶B. Simon, *Functional Integration and Quantum Physics* (Academic, New York, 1979).
- ²⁴⁷B. Simon, "Kato's inequality and the comparison of semigroups," *J. Funct. Anal.* **32**, 97–101 (1979).
- ²⁴⁸B. Simon, "Maximal and minimal Schrödinger forms," *J. Oper. Th.* **1**, 37–47 (1979).
- ²⁴⁹B. Simon, "Phase space analysis of simple scattering systems. Extensions of some work of Enss," *Duke Math. J.* **46**, 119–168 (1979).
- ²⁵⁰B. Simon, "Schrödinger semigroups," *Bull. Am. Math. Soc.* **7**, 447–526 (1982).
- ²⁵¹B. Simon, "Some Jacobi matrices with decaying potential and dense point spectrum," *Commun. Math. Phys.* **87**, 253–258 (1982).
- ²⁵²B. Simon, "Semiclassical analysis of low lying eigenvalues, I. Non-degenerate minima: Asymptotic expansions," *Ann. Inst. Henri Poincaré Phys. Theor.* **38**, 295–307 (1983).
- ²⁵³B. Simon, "Semiclassical analysis of low lying eigenvalues, II. Tunneling," *Ann. Math.* **120**, 89–118 (1984).
- ²⁵⁴B. Simon, "Fifty years of eigenvalue perturbation theory," *Bull. Am. Math. Soc.* **24**, 303–319 (1991).
- ²⁵⁵B. Simon, "Operators with singular continuous spectrum, I. General operators," *Ann. Math.* **141**, 131–145 (1995).
- ²⁵⁶B. Simon, "Bounded eigenfunctions and absolutely continuous spectra for one-dimensional Schrödinger operators," *Proc. Am. Math. Soc.* **124**, 3361–3369 (1996).
- ²⁵⁷B. Simon, "Some Schrödinger operators with dense point spectrum," *Proc. Am. Math. Soc.* **125**, 203–208 (1997).
- ²⁵⁸B. Simon, "A new approach to inverse spectral theory, I. Fundamental formalism," *Ann. Math.* **150**, 1–29 (1999).
- ²⁵⁹B. Simon, "A Feynman-Kac formula for unbounded semigroups," *Proceedings International Conference on Infinite Dimensional (Stochastic) Analysis and Quantum Physics, Leipzig, 1999* (to be published).
- ²⁶⁰B. Simon, "Schrödinger Operators in the Twenty-First Century," *Mathematical Physics 2000* (Imperial College Press, London, to be published).
- ²⁶¹B. Simon and T. Wolff, "Singular continuous spectrum under rank one perturbations and localization for random Hamiltonians," *Commun. Pure Appl. Math.* **39**, 75–90 (1986).
- ²⁶²F. Stummel, "Singuläre elliptische Differentialoperatoren in Hilbertschen Räumen," *Math. Ann.* **132**, 150–176 (1956).
- ²⁶³H. Tamura, "The asymptotic eigenvalue distribution for nonsmooth elliptic operators," *Proc. Jpn. Acad.* **50**, 19–22 (1974).
- ²⁶⁴L. Thomas, "Time dependent approach to scattering from impurities in a crystal," *Commun. Math. Phys.* **33**, 335–343 (1973).
- ²⁶⁵E. C. Titchmarsh, "Some theorems on perturbation theory, I," *Proc. R. Soc. London, Ser. A* **200**, 34–46 (1949).
- ²⁶⁶E. C. Titchmarsh, "Some theorems on perturbation theory, II," *Proc. R. Soc. London, Ser. A* **201**, 473–479 (1950).
- ²⁶⁷E. C. Titchmarsh, *Eigenfunction Expansions*, Part I, II, (Oxford University Press, London), Part I, 2nd ed. (1962); Part II (1958).
- ²⁶⁸E. Trubowitz, "The inverse problem for periodic potentials," *Comm. Pure Appl. Math.* **30**, 321–337 (1977).
- ²⁶⁹C. van Winter, "Theory of finite systems of particles I," *Mat. Fys. Skr. Danske Vid. Selsk.* **1**, 1–60 (1964).
- ²⁷⁰H. von Dreifus and A. Klein, "A new proof of localization in the Anderson tight binding model," *Commun. Math. Phys.* **124**, 285–299 (1989).
- ²⁷¹J. von Neumann, *Mathematical Foundations of Quantum Mechanics* (Springer, Berlin, 1932).
- ²⁷²J. von Neumann and E. Wigner, "Über merkwürdige diskrete Eigenwerte," *Z. Phys.* **30**, 465–467 (1929).
- ²⁷³J. Weidmann, "Zur Spektraltheorie von Sturm-Liouville-Operatoren," *Math. Z.* **98**, 268–302 (1967).
- ²⁷⁴H. Weyl, "Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen," *Math. Ann.* **71**, 441–469 (1911).
- ²⁷⁵H. Weyl, *The Theory of Groups and Quantum Mechanics* (Dover, New York, 1931).
- ²⁷⁶D. R. Yafaev, "On the theory of the discrete spectrum of the three-particle Schrödinger operator," *Math. USSR Sb.* **23**, 535–559 (1974).
- ²⁷⁷D. R. Yafaev, *Mathematical Scattering Theory* (American Mathematical Society, Providence, RI, 1992).
- ²⁷⁸D. R. Yafaev, "Radiation conditions and scattering theory for N -particle Hamiltonians," *Commun. Math. Phys.* **154**, 523–554 (1993).

- ²⁷⁹K. Yajima, "Spectral and scattering theory for Schrödinger operators with Stark-effect," J. Fac. Sci., Univ. Tokyo, Sect. 1 **26**, 377–390 (1979).
- ²⁸⁰G. M. Zhislin, "Discussion of the spectrum of Schrödinger operator for systems of many particles," Tr. Mosk. Mat. Obs. **9**, 81–128 (1960).
- ²⁸¹G. M. Zhislin, "On the finiteness of the discrete spectrum of the energy operator of negative atomic and molecular ions," Teor. Mat. Fiz. **7**, 571–578 (1971).
- ²⁸²M. Zworski, "Distribution of poles for scattering on the real line," J. Funct. Anal. **73**, 277–296 (1987).
- ²⁸³M. Zworski, "Sharp polynomial bounds on the number of scattering poles," Duke Math. J. **59**, 311–323 (1989).
- ²⁸⁴M. Zworski, "Counting scattering poles," in *Spectral and Scattering Theory*, edited by M. Ikawa (Marcel Dekker, New York, 1994), pp. 301–331.
- ²⁸⁵M. Zworski, "Resonances in physics and geometry," Notices Am. Math. Soc. **46**, 319–328 (1999).