

A Family of Rhodium Complexes with Selective Toxicity towards Mismatch Repair-Deficient Cancers

Kelsey M. Boyle, and Jacqueline K. Barton

J. Am. Chem. Soc., **Just Accepted Manuscript** • Publication Date (Web): 05 Apr 2018

Downloaded from <http://pubs.acs.org> on April 5, 2018

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

A Family of Rhodium Complexes with Selective Toxicity towards Mismatch Repair-Deficient Cancers

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Kelsey M. Boyle and Jacqueline K. Barton*
Division of Chemistry and Chemical Engineering
California Institute of Technology
Pasadena CA, 91125

* to whom correspondence should be addressed at jkbarton@caltech.edu

ABSTRACT

Rhodium metalloinsertors are a unique set of metal complexes that bind specifically to DNA base pair mismatches *in vitro* and kill mismatch repair (MMR)-deficient cells at lower concentrations than their MMR-proficient counterparts. A family of metalloinsertors containing rhodium-oxygen ligand coordination, termed “**Rh–O**” metalloinsertors, has been prepared and shown to have a significant increase in both overall potency and selectivity towards MMR-deficient cells regardless of structural changes in the ancillary ligands. Here we describe DNA-binding and cellular studies with the second generation of **Rh–O** metalloinsertors in which an ancillary ligand is varied in both steric bulk and lipophilicity. These complexes, of the form $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$, all include the O-containing PPO ligand (PPO = 2-(pyridine-2-yl)propan-2-ol) and the aromatic inserting ligand chrysi (5,6-chrysene quinone diimine) but differ in the identity of their ancillary ligand L, where L is a phenanthroline or bipyridyl derivative. The **Rh–O** metalloinsertors in this family all show micromolar binding affinities for a 29-mer DNA hairpin containing a single CC mismatch. The complexes display comparable lipophilic tendencies and pK_a values of 8.1-9.1 for dissociation of an imine proton on the chrysi ligand. In cellular proliferation and cytotoxicity assays with MMR-deficient cells (HCT116O) and MMR-proficient cells (HCT116N), the complexes containing the phenanthroline-derived ligands show highly selective cytotoxic preference for the MMR-deficient cells at nanomolar concentrations. Using mass spectral analyses, it is shown that the complexes are taken into cells through a passive mechanism and exhibit low accumulation in mitochondria, an off-target organelle that, when targeted by parent metalloinsertors, can lead to non-selective cytotoxicity. Overall, these **Rh–O** metalloinsertors have distinct and improved behavior compared to previous generations of parent metalloinsertors, making them ideal candidates for further therapeutic assessment.

INTRODUCTION

Over the past 70 years, DNA and its associated metabolic processes have proven to be fruitful targets for the design of new therapeutic agents.¹ Many of the most common FDA-approved chemotherapeutics work by binding DNA, such as the DNA-crosslinking agent cisplatin and the DNA-intercalating agent doxorubicin.²⁻⁵ Despite the prevalence of these drugs in the clinic, there are many drawbacks to their design and mechanisms of action. In many cases, the drugs target a generic DNA structure that is common to both healthy and cancerous cells. The incidental targeting of healthy tissue can result in dramatic and often dose-limiting side effects, such as emesis and nephrotoxicity.⁶ To circumvent these off-target effects, it is essential to identify new therapeutic targets that are almost exclusively found within cancerous tissues and cells.

In our research, we focus on one such target: DNA base pair mismatches. Mismatches occur regularly in cells due to polymerase errors or interaction with exogenous compounds.⁷ In healthy cells, these errors are corrected by the mismatch repair (MMR) machinery of the cell. However, in many solid tumors or tumors of Lynch syndrome patients, mutations in MMR proteins severely down-regulate or completely inactivate repair.^{8,9} As a result, these cancers contain a relative abundance of DNA base pair mismatches compared to healthy cells, making mismatches a potential biomarker for selective cancer therapy.

Mismatched base pairs have been targeted through the design of metal complexes, called rhodium metalloinsertors, which selectively and non-covalently bind these lesions.¹⁰ Rhodium metalloinsertors contain a sterically expansive aromatic chrysi (5,6-chrysenequinone diimine) ligand that is capable of π -stacking with DNA bases. Due to steric bulk, however, the chrysi ligand is unable to easily intercalate into well-matched DNA, and instead primarily interacts with

1
2
3 DNA at thermodynamically destabilized sites, such as mismatches or abasic sites.¹¹ The ability
4 of a prototypical metalloinsertor, $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{3+}$ (bpy = 2,2'-bipyridine), to selectively bind
5 DNA mismatches has been verified using both *in vitro* binding assays and crystallographic
6 studies.^{12–15} Crystallographic and NMR studies show that this complex binds DNA mismatches
7 *via* metalloinsertion, a non-covalent binding mode in which the complex inserts into DNA at the
8 mismatched site from the minor groove, ejects the mismatched DNA bases, and π -stacks with the
9 flanking well-matched base pairs.¹⁴ This mismatch-targeting ability has also been seen in human
10 cell culture experiments, with metalloinsertors exhibiting enhanced cytotoxicity in MMR-
11 deficient cell lines relative to their MMR-proficient counterparts.^{15,16} This result is in stark
12 contrast to most DNA-targeting therapeutics, such as the aforementioned cisplatin and
13 doxorubicin, which are selective towards MMR-*proficient* cell lines over MMR-deficient cell
14 lines, leading to the development of resistance in MMR-deficient tumors following
15 treatment.^{17,18}

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Several generations of metalloinsertors have been synthesized since $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{3+}$,
which has led to the recent discovery of a potent and selective family of rhodium
metalloinsertors containing a pyridyl-alcohol ligand and unique Rh–O ligand coordination
(Figure 1).¹⁹ This Rh–O ligand coordination is structurally distinct from earlier generations of
parent metalloinsertors, which contained solely Rh–N coordination.²⁰ Furthermore, these **Rh–O**
metalloinsertors were found to have improved potency and selectivity towards MMR-deficient
cancer cells over MMR-proficient cancer cells. Surprisingly, this high potency and cell
selectivity was seen across a variety of metalloinsertors containing O-coordinated ligands that
differed significantly in size and structure (spanning methyl, pyridyl, phenyl, and hexyl

functionalization), suggesting the biological activities of **Rh–O** metalloinsertors are not perturbed by ligand substitution off of the O-containing site.

Here, a family of rhodium metalloinsertors was designed and synthesized as variations of the **Rh–O** metalloinsertor $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$ (phen = 1,10-phenanthroline). These complexes, of the form $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$, all include the O-containing PPO ligand but differ in the identity of their ancillary ligand, L, where L = bpy, HDPA (2,2'-dipyridylamine), 4,7-DMP (4,7-dimethyl-1,10-phenanthroline), 5,6-DMP (5,6-dimethyl-1,10-phenanthroline), and DIP (4,7-diphenyl-1,10-phenanthroline) (Figure 2). The ancillary ligand substitution alters the steric bulk and lipophilicity of these complexes, which can ultimately affect DNA-binding properties and biological activity.^{20,21} Each complex described, even the most lipophilic and sterically bulky, shows biological selectivity towards MMR-deficient cell lines, further demonstrating that the Rh–O ligand framework is amenable to a wide array of functionalization. To better understand the trends in biological activity of these complexes, each metalloinsertor was examined for binding affinity to mismatched DNA, pK_a, lipophilicity, whole cell uptake, and subcellular localization into the nucleus and mitochondria. The results indicate that minimizing uptake of the complexes into the mitochondria may be a key factor in ensuring high biological selectivity and support that these **Rh–O** complexes exhibit distinct differences in metalloinsertor-DNA binding and cell activation compared to parent metalloinsertors.

EXPERIMENTAL METHODS

Materials. Commercially available chemicals were used as received. All reagents and Sephadex ion-exchange resin were obtained from Sigma-Aldrich with the following exceptions. RhCl₃ was purchased from Pressure Chemical, Inc. Dowex ion-exchange beads were purchased

1
2
3 from Acros Organics. Analytical standards for Rb and transition metals were purchased from
4 Analytical West and Ultra Scientific, respectively. MTT and ELISA assay kits were obtained
5 from Roche. Pierce BCA assay kit and NP40 were purchased from Thermo Scientific. Sep-pak
6 C18 solid-phase extraction (SPE) cartridges were purchased from Waters Chemical Co. Cell
7 culture media and supplements were purchased from Invitrogen. Tissue culture flasks and plates
8 were obtained from Corning. ^{32}P labeled ATP was purchased from Perkin Elmer. UreaGel
9 supplies were purchased from National Diagnostics. Microbiospin columns were purchased from
10 BioRad.

11
12
13
14
15
16
17
18
19
20
21
22 *Synthesis and Characterization of Metal Complexes.* $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]\text{Cl}_2$ and
23 $[\text{Rh}(\text{bpy})_2(\text{chrysi})]\text{Cl}_3$ were synthesized following the published protocols.^{19,22} New metal
24 complexes were synthesized in a similar manner to published procedures.^{19,20,23} A description of
25 the general synthetic procedures are below. Complete synthetic details for each complex,
26 including specific amounts (masses, volumes, and ratios) as well as slight deviations from the
27 synthetic scheme below, can be found in the SI.

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Synthesis of $[\text{Rh}(\text{L})\text{Cl}_4][\text{K or H}_3\text{O}]$ complexes. For L = bpy, 4,7-DMP, 5,6-DMP, and
DIP: $\text{RhCl}_3 \cdot 3\text{H}_2\text{O}$ (1 equiv.) and KCl (1 equiv.) were refluxed in methanol for 2 h at 98 °C.
Ligand (L, 1 equiv.) was added in a minimum volume of methanol and refluxed for 4 h, during
which the deep red solution turned to golden-brown precipitate. The solution was filtered over a
medium fritted filter, rinsed with methanol, and dried under vacuum to produce $[\text{Rh}(\text{L})\text{Cl}_4]\text{K}$.
Crude yield: 84% (bpy), 86% (4,7-DMP), 91% (5,6-DMP), 95% (DIP). For L = HDPA:
 $\text{RhCl}_3 \cdot 3\text{H}_2\text{O}$ (1 equiv.) was refluxed in concentrated HCl (38% w/v) for 3 h at 98 °C. Ligand (L,
2 equiv.) was added in a minimum volume of HCl, followed immediately by boiling water. The
solution was refluxed for 16 h, then cooled to 4 °C. The golden precipitate was filtered over a

1
2
3 Buchner funnel and dried under vacuum to produce $[\text{Rh}(\text{L})\text{Cl}_4][\text{H}_3\text{O}]$. Crude yield: 100%
4
5 (HDPA).

6
7
8 *Synthesis of $[\text{Rh}(\text{L})(\text{NH}_3)_4][\text{OTf}]_3$.* $[\text{Rh}(\text{L})\text{Cl}_4][\text{K or H}_3\text{O}]$ (1 equiv.) was added to an
9
10 oven-dried 25 mL Schlenk flask and degassed under argon. Neat triflic acid (HOTf, 10 g, excess)
11
12 was added to the flask under positive argon pressure, producing a deep red solution. The flask
13
14 was purged to remove newly formed HCl gas and stirred for 16 h. The solution was then added
15
16 dropwise to cold, stirring ether at $-78\text{ }^\circ\text{C}$ to produce a yellow-brown precipitate. The precipitate
17
18 was filtered over a medium fritted filter and rinsed with additional cold ether. The product,
19
20 $[\text{Rh}(\text{L})(\text{OTf})_4][\text{K or H}_3\text{O}]$, was combined with NH_4OH (28% w/v) and stirred at $40\text{ }^\circ\text{C}$ for 1 h.
21
22 The solvent was removed under vacuum and the product was suspended in minimal ethanol,
23
24 precipitated with ether, filtered over a medium fritted filter, and dried further under vacuum to
25
26 produce $[\text{Rh}(\text{L})(\text{NH}_3)_4][\text{OTf}]_3$. Crude yields of 42% (bpy), 10% (HDPA), 15% (4,7-DMP), 77%
27
28 (5,6-DMP), 72% (DIP).
29
30
31

32
33 *Synthesis of $[\text{Rh}(\text{L})(\text{chrysi})(\text{NH}_3)_2][\text{OTf}]_3$.* $[\text{Rh}(\text{L})(\text{NH}_3)_4][\text{OTf}]_3$ (1 equiv.) was
34
35 combined with 5,6-chryseno-quinone (1 equiv.) and a mixture of acetonitrile, water, and NaOH,
36
37 and stirred for 1-12 h at ambient temperature. The solution changed from bright orange (the color
38
39 of free ligand) to red-brown (for L = bpy, HDPA, 5,6-DMP, and DIP) or green-brown (for
40
41 L=4,7-DMP) with no precipitate. The reaction was quenched with HCl, producing a deep red
42
43 solution, and the solvent was removed under vacuum. The products from L=bpy, HDPA, and
44
45 DIP were purified using a C18 SepPak, pre-equilibrated with 0.1% TFA (aq, TFA =
46
47 trifluoroacetic acid) and eluted with 1:3 MeCN:0.1% TFA (aq). The products from L=4,7-DMP
48
49 and 5,6-DMP were purified by HPLC using a reverse phase C18 column with gradient elution
50
51 from 15:85 MeCN:0.1% TFA (aq) to 95:5 MeCN:0.1% TFA (aq) over 30 min. Products were in
52
53
54
55
56
57
58
59
60

1
2
3 the form of $[\text{Rh}(\text{L})(\text{chrysi})(\text{NH}_3)_2][\text{TFA}]_3$. Crude yields of 33% (bpy), 51% (HDPA), 46% (4,7-
4
5 DMP), 62% (5,6-DMP), 100% (DIP).
6

7
8 *Synthesis of $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]\text{Cl}_2$.* $[\text{Rh}(\text{L})(\text{chrysi})(\text{NH}_3)_2][\text{TFA}]_3$ (1 equiv.) was
9
10 combined with PPO in a mixture of ethanol and water and refluxed 16 h (for L = bpy, 4,7-DMP,
11
12 5,6-DMP, and DIP) or 7 days (for L=HDPA). The solvent was removed under vacuum and the
13
14 product was purified by HPLC using the method described above for L = bpy, HDPA, 4,7-DMP,
15
16 and DIP. For L = 5,6-DMP, an isocratic method of 30:70 MeCN:0.1% TFA (aq) was used. For L
17
18 = bpy, HDPA, and 4,7-DMP, the purified product was converted to the chloride salt using
19
20 Sephadex QAE resin charged with MgCl_2 . For L = 5,6-DMP and DIP, the purified product was
21
22 converted to the chloride salt using Dowex 1x2 500-100 mesh ion exchange resin. Purified yields
23
24 of 30% (bpy), 10% (HDPA), 10% (4,7-DMP), 23% (5,6-DMP), 33% (DIP).
25
26
27

28
29 *Characterization of $[\text{Rh}(\text{bpy})(\text{chrysi})(\text{PPO})](\text{TFA})_2$.* LCQ-MS (cation): m/z calc. 650.1
30
31 ($\text{M}-1\text{H}^+$), 325.6 (M^{2+}); obs. 650.0, 325.8. UV-Vis (H_2O): 259nm ($59,800 \text{ M}^{-1} \text{ cm}^{-1}$), 287nm
32
33 ($43,100 \text{ M}^{-1} \text{ cm}^{-1}$), 298nm ($37,100 \text{ M}^{-1} \text{ cm}^{-1}$), 312nm ($32,000 \text{ M}^{-1} \text{ cm}^{-1}$), 435nm ($10,000 \text{ M}^{-1} \text{ cm}^{-1}$)
34
35 ^1H NMR (500 MHz, Acetonitrile- d_3) δ 13.44 (br s, 1.2H), 11.89 (br s, 2H), 9.45 (d, $J = 5.6$
36
37 Hz, 1H), 9.36 (d, $J = 5.7$ Hz, 0.6H), 8.80 (d, $J = 8.0$, 1.4 Hz, 1H), 8.71 (d, $J = 5.3$ Hz, 0.6H), 8.62
38
39 (d, $J = 8.2$ Hz, 0.6H), 8.60-8.54 (m, 2.6H), 8.43-8.26 (m, 8H), 8.26-8.21 (m, 1H), 8.14 (d, $J =$
40
41 8.2, 1.5 Hz, 0.6H), 8.06-7.89 (m, 4.8H), 7.85-7.78 (m, 1,6H), 7.77-7.68 (m, 3.2H), 7.68-7.61 (m,
42
43 2.2H), 7.60-7.52 (m, 2.6H), 7.31 (d, 0.6H), 7.29-7.21 (m, 2.6H), 1.91 (s, 3H), 1.87 (s, 1.8H), 1.58
44
45 (s, 4.8H), purified as a 1:0.6 mixture of diastereomers.
46
47
48

49
50 *Characterization of $[\text{Rh}(\text{HDPA})(\text{chrysi})(\text{PPO})](\text{TFA})_2$.* LCQ-MS (cation): m/z calc. 665.2
51
52 ($\text{M}-1\text{H}^+$), 333.1 (M^{2+}); obs. 665.3, 333.3. UV-Vis (H_2O): 259nm ($60,400 \text{ M}^{-1} \text{ cm}^{-1}$), 283nm
53
54 ($45,900 \text{ M}^{-1} \text{ cm}^{-1}$), 326nm ($18,600 \text{ M}^{-1} \text{ cm}^{-1}$), 440nm ($8,500 \text{ M}^{-1} \text{ cm}^{-1}$). ^1H NMR (500 MHz,
55
56
57
58
59
60

1
2
3 Acetonitrile- d_3) δ 12.49 (br s, 1H), 12.04 (br s, 1H), 8.72 (dd, $J = 8.0, 1.3$ Hz, 1H), 8.50 (d, $J =$
4 6.0 Hz, 1H), 8.38-8.31 (m, 3H), 8.31-8.23 (m, 2H), 8.20-8.13 (m, 2H), 8.08-8.00 (m, 2H), 7.98
5
6 (td, $J = 8.6, 1.6$ Hz, 1H), 7.94-7.81 (m, 4H), 7.69 (m, 3H), 7.51 (ddd, $J = 7.6, 6.0, 1.4$ Hz, 1H),
7
8 7.23 (ddd, $J = 7.4, 6.1, 1.3$ Hz, 1H), 7.17 (ddd, $J = 7.4, 6.2, 1.4$ Hz, 1H), 1.78 (s, 3H), 1.56 (s,
9
10 3H), purified as a single diastereomer.
11
12
13

14
15 *Characterization of [Rh(4,7-DMP)(chrysi)(PPO)](TFA)₂*. LCQ-MS (cation): m/z calc.
16
17 702.2 (M-1H⁺), 351.6 (M²⁺); obs. 702.3, 351.8. UV-Vis (H₂O): 269nm (106,800 M⁻¹ cm⁻¹),
18
19 440nm (11,400 M⁻¹ cm⁻¹). ¹H NMR (500 MHz, Acetonitrile- d_3) δ 13.31 (br s, 0.8H), 11.75 (br s,
20
21 2H), 9.50 (d, $J = 5.4$ Hz, 1H), 9.42 (d, $J = 5.4$ Hz, 0.4H), 8.86 (dd, $J = 5.5, 0.9$ Hz, 1H), 8.83 (dd,
22
23 $J = 8.0, 1.3$ Hz, 1H), 8.73 (d, $J = 5.4$ Hz, 0.4H), 8.47 (d, $J = 2.5$ Hz, 0.4H), 8.46-8.35 (m, 4.2H),
24
25 8.34 (d, $J = 8.2$ Hz, 0.4H), 8.27 (d, $J = 8.8$ Hz, 1H), 8.21-8.17 (m, 1.4H), 8.16 (d, $J = 8.1$ Hz,
26
27 0.4H), 8.08 (dd, $J = 5.4, 1.0$ Hz, 1H), 8.04 (d, $J = 5.4$ Hz, 0.4H), 8.00 (dd, $J = 7.5, 1.7$ Hz, 1H),
28
29 7.97-7.92 (m, 2.4H), 7.84 (m, 1.8H), 7.77 (m, 1.4H), 7.61-7.51 (m, 5.2H), 7.19-7.15 (m, 0.4H),
30
31 7.10-7.03 (m, 2.8H), 3.05 (s, 3H), 3.04 (s, 1.2H), 3.02 (s, 1.2H), 2.99 (s, 3H), 1.95 (s, 3H), 1.92
32
33 (s, 1.2H), 1.62 (s, 3H), 1.61 (s, 1.2H), purified as a 1:0.4 mixture of diastereomers.
34
35
36
37

38
39 *Characterization of [Rh(5,6-DMP)(chrysi)(PPO)](TFA)₂*. LCQ-MS (cation): m/z calc.
40
41 702.2 (M-1H⁺), 351.6 (M²⁺); obs. 702.3, 351.8. UV-Vis (H₂O): 267nm (80,600 M⁻¹ cm⁻¹), 280nm
42
43 (81,700 M⁻¹ cm⁻¹), 438nm (10,500 M⁻¹ cm⁻¹). ¹H NMR (500 MHz, Acetonitrile- d_3) δ 13.40 (br s,
44
45 0.3H), 11.77 (br s, 1H), 9.68 (d, $J = 5.2$ Hz, 1H), 9.59 (d, $J = 5.1$ Hz, 0.3H), 9.06-8.97 (m, 3.9H),
46
47 8.84-8.89 (m, 1.3H), 8.43-8.37 (m, 2.6H), 8.34 (d, $J = 8.2$ Hz, 0.3H), 8.29-8.14 (m, 5.2H), 8.02-
48
49 7.97 (m, 2.3H), 7.96-7.89 (m, 2.6H), 7.83-7.73 (m, 1.6H), 7.57 (td, $J = 7.4, 1.4$ Hz, 2H), 7.55-
50
51 7.50 (m, 1H), 7.17 (d, $J = 5.7$ Hz, 0.3H), 7.10-7.02 (m, 2.3H), 2.91 (s, 0.9H), 2.90 (s, 0.9H), 2.89
52
53
54
55
56
57
58
59
60

(s, 3H), 2.87 (s, 3H), 1.93 (s, 3H), 1.90 (s, 0.9H), 1.58 (s, 3.9H), purified as a 1:0.3 mixture of diastereomers.

Characterization of [Rh(DIP)(chrysi)(PPO)]Cl₂. LCQ-MS (cation): m/z calc. 826.2 (M-1H⁺); obs. 826.3. UV-Vis (H₂O): 267nm (103,000 M⁻¹ cm⁻¹). ¹H NMR (500 MHz, Methanol-*d*₄) δ 9.74 (dd, J = 5.5, 0.9 Hz, 1H), 9.70 (dd, J = 5.5, 0.8 Hz, 0.5H), 8.89 (m, 1.5H), 8.76 (m, 1.5H), 8.58-8.46 (m, 4.5H), 8.40-8.28 (m, 6H), 8.14-7.98 (m, 4.5H), 7.81-7.59 (m, 15H), 7.56-7.49 (m, 1.5H), 7.41-7.33 (m, 6H), 7.34-7.23 (m, 3H), 2.07 (s, 3H), 2.02 (s, 1.5H), 1.70 (s, 1.5H), 1.69 (s, 3H), purified as a 1:0.5 mixture of diastereomers.

Enantiomeric Separation of [Rh(phen)(chrysi)(PPO)]Cl₂. Purified [Rh(phen)(chrysi)(PPO)][TFA]₂ was dissolved in 1:1 ethanol:water and HPLC purified on an Astec CYCLOBOND chiral column using an isocratic elution method of 40:60 ACN:0.1 M KPF₆ (aq) over 37 min. The column was periodically rinsed with 40:60 MeCN:H₂O to remove KPF₆ buildup. Separated enantiomers were collected and exchanged to the chloride salt using Sephadex QAE resin pre-equilibrated with MgCl₂. The enantiomeric nature of the collected fractions was verified using circular dichroism (CD) as follows: 200 μM solutions of Δ- and Λ-[Rh(phen)(chrysi)(PPO)]Cl₂ were made in aqueous solution and their CD spectra recorded in 1 nm increments on an Aviv 62DS spectropolarimeter under a N₂ atmosphere at ambient temperature. The spectra were recorded a second time 30 d later to assess decomposition or racemization of the sample, and none was observed.

Determination of Extinction Coefficients. Aqueous solutions of each [Rh(L)(chrysi)(PPO)]Cl₂ complex were made and a UV-Visible spectrum was recorded for each. The solutions were diluted 50x, 100x, 500x, and 1000x in 2% HNO₃. The dilutions were analyzed for Rh content *via* ICP-MS (inductively coupled plasma mass spectrometry) and the

1
2
3 concentration was determined by comparison to a standard curve. Extinction coefficients were
4
5 determined from the UV-Visible absorbance measurement of the initial solution and the Rh
6
7 concentration of the dilutions following Beer's law ($A = \epsilon lc$). L = DIP was observed to
8
9 significantly adsorb onto plastics, therefore PTFE (polytetrafluoroethylene) and PFA
10
11 (perfluoroalkoxy alkane) coated materials were used in the workup and analysis of its extinction
12
13 coefficient.
14
15

16
17 *Partition Coefficient Determination.* One-octanol and 10 mM Tris-HCl, pH 7.4 were pre-
18
19 equilibrated with each other by vigorously shaking the phases together. A solution of each
20
21 metalloinsertor was made in octanol and the UV-Visible spectrum of the solution recorded. Each
22
23 solution was combined with an equal volume of aqueous buffer and shaken using a foam insert
24
25 on a Vortex-Genie 2 running at maximum speed for 16 h. The samples were centrifuged to
26
27 separate the aqueous and octanol phases and a UV-Visible spectrum of each octanol fraction was
28
29 recorded. The baseline value obtained at 800 nm was used to normalize the spectra to a common
30
31 zero point. The absorbance of the ~260 nm peak in the final spectrum was compared to the initial
32
33 spectrum to determine the partition coefficient following the literature.²⁴ The partition
34
35 coefficients from three experiments were measured for each $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]\text{Cl}_2$ complex
36
37 and averaged to give the partition coefficient.
38
39
40
41

42
43 *pK_a Determination of Metalloinsertors.* A ~25 μM solution of each metalloinsertor was
44
45 made in 100 mM NaCl. The pH of the sample was adjusted to 4.5 using HCl (10 mM). NaOH
46
47 (10 mM) was titrated into the solution, with stirring. The pH and UV-Visible spectrum were
48
49 recorded after each base addition, up to a pH of 10.5. A back titration to pH 6 was performed to
50
51 check for decomposition, and none was observed. Spectra were corrected for baseline and
52
53 volume changes. The absorbance of the ~430 nm peak was plotted against pH and fit to a
54
55
56
57
58
59
60

1
2
3 sigmoidal curve in OriginPro v8.5, and the pK_a was determined as the inflection point of the
4
5 curve. Three pK_a titrations were performed for each $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]\text{Cl}_2$ complex and
6
7 averaged to give an average pK_a value.
8
9

10 *Binding Constant Determination.* A DNA hairpin (5'-
11 GGCAGGXATGGCTTTTTGCCATYCCTGCC-3', where **XY**=CG or CC for a well-matched or
12
13 mismatched hairpin, respectively) was radiolabeled with γ - ^{32}P ATP and prepared following the
14
15 literature.^{10,19,22} Full details of DNA preparation and purification can be found in the SI. A 4 μM
16
17 solution of the photocleaving metalloinsertor $[\text{Rh}(\text{bpy})_2(\text{chrysi})]\text{Cl}_3$ and solutions containing 0-
18
19 400 μM of a competing metalloinsertor, $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]\text{Cl}_2$ (which does not photocleave
20
21 DNA), were made in MilliQ water. Five μL of the $[\text{Rh}(\text{bpy})_2(\text{chrysi})]\text{Cl}_3$ solution, 5 μL of the
22
23 competing metalloinsertor, and 10 μL of the hairpin DNA were combined to create a solution
24
25 containing 1 μM $[\text{Rh}(\text{bpy})_2(\text{chrysi})]\text{Cl}_3$, 0-100 μM competing metalloinsertor, and 1 μM DNA.
26
27 The samples were irradiated with an Oriel 1000 W Hg/Xe solar simulator (340-440 nm) for 20
28
29 min. After irradiation, solvent was removed from the samples and the samples were counted on a
30
31 scintillation counter to determine the necessary exposure time (with 300,000 cpm needing a 1
32
33 hour exposure) and they were suspended in a denaturing formamide loading dye. Samples were
34
35 electrophoresed on a 20% denaturing polyacrylamide urea gel.
36
37
38
39
40
41

42 A phosphor screen was exposed to the polyacrylamide gel and imaged using a Typhoon
43
44 FLA 9000 biomolecular imager. The ratio of photocleaved to uncleaved DNA was quantified
45
46 using ImageQuant TL software. The ratio was plotted against the concentration of
47
48 $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]\text{Cl}_2$ and fit to a sigmoidal curve in OriginPro v8.5 to determine the
49
50 inflection point of the fit. The binding affinity of the competing metalloinsertor was calculated in
51
52 Mathematica 9.0 by solving simultaneous equilibria involving DNA, $[\text{Rh}(\text{bpy})_2(\text{chrysi})]\text{Cl}_3$, and
53
54
55
56
57
58
59
60

1
2
3 [Rh(L)(chrysi)(PPO)]Cl₂. Three photocleavage titrations were performed for each
4
5 [Rh(L)(chrysi)(PPO)]Cl₂ complex and averaged to give the binding affinity.
6

7
8 *Melting Temperature Analysis.* Melting temperature analysis was performed on a
9
10 Beckman DU 7400 spectrophotometer equipped with a T_m Analysis Accessory. The short
11
12 oligomer, 5'-CGGACTCCG-3' (underline denotes mismatch), was purchased from IDT DNA
13
14 and purified by HPLC. Samples containing 11 μM ssDNA (ultimately 5.5 μM dsDNA and
15
16 mismatches) and 6 μM of [Rh(phen)(chrysi)(PPO)]Cl₂, [Rh(bpy)₂(chrysi)]Cl₃ or no metal
17
18 complex were prepared in phosphate buffer (5 mM phosphate, 50 mM NaCl, pH 7.0). Samples
19
20 were heated at a rate of 0.5 °C/min and absorbance was measured at 260 nm every 0.5 °C
21
22 between 10 °C and 50 °C. Data from three experiments was combined and fit to a sigmoidal
23
24 curve in OriginPro v8.5 and the T_m was taken as the inflection point of the curve.
25
26

27
28 *Cell Culture.* HCT116N and HCT116O cells were grown in RPMI (Roswell Park
29
30 Memorial Institute) 1640 media supplemented with 10% FBS (fetal bovine serum), 2 mM L-
31
32 glutamine, 0.1 mM non-essential amino acids, 1 mM sodium pyruvate, 100 units/mL penicillin
33
34 and streptomycin, and 100 μg/mL Geneticin (G418). The cells were incubated in tissue culture
35
36 flasks or plates at 37 °C in a 5% CO₂ atmosphere. All cell studies were performed with the
37
38 chloride salt of each metalloinsertor.
39
40

41
42 *Cell Proliferation ELISA.* Cell proliferation ELISA (enzyme-linked immunosorbent
43
44 assay) was performed following the manufacturers instructions. Briefly, 2×10³ HCT116N or
45
46 HCT116O cells in 100 μL media were plated into each well of a 96-well plate. The cells were
47
48 allowed to adhere for 24 h before the addition of 100 μL of media containing various
49
50 concentrations of rhodium metalloinsertor. The plates were incubated for an additional 48 h
51
52 before the rhodium-containing media was replaced with fresh media, with which the cells were
53
54
55
56
57
58
59
60

1
2
3 allowed to grow for the remainder of a 72 h period. Cells were then treated with an excess of the
4 unnatural nucleic acid, BrdU (bromodeoxyuridine), for 24 h during which time it could be
5
6 incorporated into newly synthesized DNA. Cells were then fixed, labeled with a BrdU antibody,
7
8 and quantified using a colorimetric substrate solution and stop solution. Absorbance was
9
10 measured at 450 nm (background subtracted at 690 nm). Decrease in cellular proliferation was
11
12 determined for each metalloinsertor concentration through comparison to untreated cells.
13
14
15
16
17 Outliers were removed using a modified Thompson Tau test. An additional variation of this
18
19 assay was performed in which the cells were treated with rhodium metalloinsertor for 24 h, then
20
21 directly treated with BrdU in fresh media.
22

23
24 *MTT Cytotoxicity Assay.* Cell proliferation MTT (MTT = 2-(4,5-dimethylthiazol-2-yl)-
25
26 2,5-diphenyltetrazolium bromide) assays were performed following the manufacturers
27
28 instructions. Briefly, 5×10^4 HCT116N or HCT116O cells in 100 μ L media were plated into each
29
30 well of a 96-well plate. Various concentrations of a rhodium metalloinsertor were added to each
31
32 well. The cells were allowed to incubate for 72 h before treatment with MTT for 4 h, during
33
34 which time MTT could be converted into formazan by metabolically active cells. The formazan
35
36 crystals were solubilized and quantified by absorbance at 570 nm (background subtracted at 690
37
38 nm). Viability was determined for each metalloinsertor concentration through comparison to
39
40 untreated cells. Outliers were removed using a modified Thompson Tau test. An additional
41
42 variation of this assay was performed in which the cells were allowed to adhere to the 96-well
43
44 plate overnight before treated with rhodium metalloinsertor for 24 h, followed by MTT
45
46
47 treatment.
48
49

50
51 *Uptake and Localization Experiments.* Whole-cell uptake, mitochondrial localization, and
52
53 nuclear localization of metalloinsertors were determined following published methods.²⁵ Prior to
54
55
56
57
58
59
60

1
2
3 whole-cell, mitochondrial, and nuclear rhodium determination, 24-hour ELISA and MTT assays
4
5 were performed to determine a metalloinsertor concentration that would not result in significant
6
7 cell death by MTT but showed some anti-proliferative effect by ELISA. The concentrations used
8
9 in the uptake and localization studies of the [Rh(L)(chrysi)(PPO)]Cl₂ family were 0.2 μM for
10
11 L=DIP, 0.5 μM for L=phen, bpy, HDPA, 4,7-DMP, and 5,6-DMP, and 10 μM for
12
13 [Rh(bpy)₂(chrysi)]Cl₃, which was included as a control.
14
15

16
17 *Assay for Whole-Cell Rhodium Concentration.* Whole-cell uptake experiments were
18
19 performed following published protocols.²⁰ Briefly, 1×10⁶ HCT116N or HCT116O cells were
20
21 plated into 6-well tissue culture treated plates and allowed to adhere for 24 h. Media was
22
23 aspirated from the cells and fresh media containing a metalloinsertor was added to each well.
24
25 Cells were allowed to incubate for an additional 0.5-24 h with the Rh-containing media. After
26
27 incubation, media was aspirated and the cells were rinsed with PBS (phosphate buffered saline,
28
29 pH 7.4) to remove surface rhodium. Cells were lysed directly in the well using 1 mL of 1% SDS
30
31 solution. These samples were transferred to microcentrifuge tubes and sonicated for 10 s at 20%
32
33 amplitude on a Qsonica Ultrasonic sonicator. Cell lysate was combined with an equal volume
34
35 2% HNO₃. This solution was analyzed for Rh content on an Agilent 8800 Triple Quadrupole
36
37 ICP-MS and the concentration of Rh in each sample was determined by comparison to a standard
38
39 curve (ranging from 1-100 ppb Rh) and normalized using the protein content of each sample.
40
41 The protein content of each sample was determined using a Pierce BCA assay, following the
42
43 manufacturer's instructions.
44
45
46
47
48

49
50 *Assay for Mitochondrial Rhodium Concentration.* Mitochondrial uptake experiments
51
52 were performed following published protocols.^{20,26} Briefly, 1.5×10⁷ HCT116N and HCT116O
53
54 cells were plated in T75 tissue culture treated flasks. The cells were allowed to adhere for 24 h,
55
56
57
58
59
60

1
2
3 after which media was aspirated from each flask and restored with 20 mL media containing a
4 rhodium metalloinsertor. The cells were allowed to grow in the presence of Rh-containing media
5 for 24 h, then harvested using 0.05% trypsin over 5 minutes. Cells were pelleted by
6 centrifugation at 1200 rpm for 5 min. The pellet was rinsed and suspended in PBS, then pelleted
7 again and the PBS removed. The cell pellet was suspended in 500 μ L mitochondrial extraction
8 buffer (200 mM mannitol, 68 mM sucrose, 50 mM PIPES, 50 mM KCl, 5 mM EGTA, 2 mM
9 $MgCl_2$, 1 mM DTT added just before use, and protease inhibitors added just before use) and
10 incubated on ice for 20 min. Each sample was homogenized by 35 passes thorough a 21-gauge
11 needle and syringe. The resultant solution was centrifuged for 5 min at 750 rpm. The supernatant
12 of each sample was transferred to a 1.5 mL microcentrifuge tube and centrifuged for 10 min at
13 14,000 g. The supernatant was decanted and the resulting pellet was the mitochondrial fraction.
14 SDS (800 μ L of a 1% solution) was added to the pellet and sonicated for 10 s at 40% amplitude
15 on a Qsonica Ultrasonic sonicator. Mitochondrial lysate was combined with an equal volume of
16 2% nitric acid. This solution was analyzed for Rh content on an Agilent 8800 Triple Quadrupole
17 ICP-MS and the concentration of Rh in each sample was determined by comparison to a standard
18 curve (ranging from 1-100 ppb Rh) and normalized using the protein content of each sample.
19 The protein content of each sample was determined using a Pierce BCA assay, following the
20 manufacturer's instructions.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 *Assay for Nuclear Rhodium Concentration.* Nuclear uptake experiments were performed
45 following published protocols.²⁰ Briefly, 1×10^7 HCT116N and HCT116O cells were plated in
46 T75 tissue culture treated flasks. The cells were allowed to adhere for 24 h before the media was
47 aspirated and restored with 20 mL media containing a rhodium metalloinsertor. The cells were
48 allowed to grow in the presence of Rh-containing media for 24 h, then harvested using 0.05%
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 trypsin over 5 minutes. Cells were pelleted by centrifugation at 1200 rpm for 5 min. The pellet
4 was rinsed and suspended in PBS, then pelleted and the PBS removed. Each cell pellet was
5
6 was rinsed and suspended in PBS, then pelleted and the PBS removed. Each cell pellet was
7
8 suspended in 1 mL hypotonic buffer (20 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM MgCl₂),
9
10 transferred to a microcentrifuge tube, and incubated on ice for 15 min. NP-40 (50 μL of a 10%
11
12 solution) was added to each sample, vortexed for 10 s at the highest setting, and centrifuged at
13
14 3000 g for 10 min. The supernatant was decanted and the resulting pellet was the nuclear
15
16 fraction. SDS (800 μL of a 1% solution) was added to the pellet and then sonicated for 10 s at
17
18 40% amplitude on a Qsonica Ultrasonic sonicator. Nuclear lysate was combined with an equal
19
20 volume of 2% HNO₃. This solution was analyzed for Rh content on an Agilent 8800 Triple
21
22 Quadrupole ICP-MS and the concentration of Rh in each sample was determined by comparison
23
24 to a standard curve (ranging from 1-100 ppb Rh) and normalized using the protein content of
25
26 each sample. The protein content of each sample was determined using a Pierce BCA assay,
27
28 following the manufacturer's instructions.
29
30
31

32
33 *Assay for Uptake of Metalloinsertors.* Mechanism of uptake experiments were adapted
34
35 from published protocols.²⁷ RbCl and [Ru(DIP)(dppz)]Cl₂ were used as positive and negative
36
37 controls, respectively. Briefly, 1×10⁶ HCT116N or HCT116O cells were plated into 6-well tissue
38
39 culture treated plates and allowed to adhere for 24 h. Metabolic inhibitors (5 μM oligomycin in
40
41 ethanol and 50 mM 2-deoxy-D-glucose) or control solutions (5 mM glucose and ethanol) were
42
43 added to the cell culture media and samples were incubated for 1 h. Media was removed by
44
45 aspiration and each well was washed with PBS. Media (3 mL) containing the **Rh-O**
46
47 metalloinsertor [Rh(phen)(chrysi)(PPO)]Cl₂ (0.5 μM), the parent metalloinsertor
48
49 [Rh(bpy)₂(chrysi)]Cl₃ (10 μM), [Ru(DIP)(dppz)]Cl₂ (2 μM), or RbCl (25 μM) was then added to
50
51 each well and incubated for 1 h. Media was aspirated and cells were rinsed with PBS to remove
52
53
54
55
56
57
58
59
60

1
2
3 surface rhodium, ruthenium, or rubidium. Cells were lysed directly in the well using 1 mL of 1%
4
5 SDS solution. Samples were transferred to microcentrifuge tubes and sonicated for 10 s at 20%
6
7 amplitude on a Qsonica Ultrasonic sonicator. Cell lysate was combined with an equal volume of
8
9 2% HNO₃ and analyzed for Rh, Ru, and Rb content on an Agilent 8800 Triple Quadrupole ICP-
10
11 MS, and the concentration of Rh, Ru, or Rb in each sample was determined by comparison to a
12
13 standard curve (ranging from 1-100 ppb) and normalized using the protein content of each
14
15 sample. The protein content of each sample was determined using a Pierce BCA assay, following
16
17 the manufacturer's instructions.
18
19
20
21
22
23

24 RESULTS

25
26 *Establishing the Enantiomeric Activity of [Rh(phen)(chrysi)(PPO)]²⁺.* Enantiomeric
27
28 separation was performed for the complex [Rh(phen)(chrysi)(PPO)]²⁺ to establish the interaction
29
30 of its Δ- and Λ-enantiomers with DNA *in vitro* and in MMR-deficient or -proficient cells in
31
32 culture. The Δ- and Λ- enantiomers of [Rh(phen)(chrysi)(PPO)]²⁺ were isolated with >90% and
33
34 >95% enantiomeric excess, respectively (Supplementary Figure S1). Circular dichroism
35
36 experiments confirmed the enantiomeric nature of the isolated complexes, and no racemization
37
38 was observed at ambient temperature over 1 month (Supplementary Figure S1). Competition
39
40 titrations between [Rh(phen)(chrysi)(PPO)]²⁺ and the photocleaving metalloinsertor
41
42 [Rh(bpy)₂(chrysi)]³⁺ in the presence of ³²P-radiolabeled DNA containing a CC mismatch
43
44 revealed both enantiomers are capable of binding mismatched DNA base pairs with similar
45
46 affinity (10⁶ M⁻¹, Table 1).¹⁰ Furthermore, both enantiomers were found to have selective
47
48 cytotoxic effects towards MMR-deficient cells over MMR-proficient cells in MTT experiments
49
50 (Supplementary Figure S2). These studies confirm that both enantiomers of the PPO-containing
51
52
53
54
55
56
57
58
59
60

1
2
3 metalloinsertor, $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$, exhibit binding properties towards mismatched
4 DNA that are consistent with a previous generation of **Rh-O** metalloinsertors. These **Rh-O**
5 complexes show no enantiomeric preference in binding DNA, unlike parent metalloinsertors,
6 which show a high enantiomeric preference for the Δ -isomer in binding DNA.^{15,19}
7
8
9
10
11

12 *Binding of Metalloinsertors to a Single Base Pair Mismatch.* The binding affinities of
13 $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertors to DNA containing a single CC mismatch were
14 determined. The $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ complexes do not photocleave DNA upon irradiation,
15 so their binding affinities were assayed *via* a competition titration with $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{3+}$, a
16 complex known to photocleave DNA selectively upon mismatch binding and irradiation.²² A CC
17 mismatch was used as it is highly destabilized relative to other mismatches and therefore
18 undergoes significant photocleavage in the presence of $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{3+}$. A constant
19 concentration of $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{3+}$ and varying concentrations of the competing
20 $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertor were incubated with a DNA hairpin containing a single
21 CC mismatch, irradiated, and the DNA photocleavage products were separated on a denaturing
22 gel. The ratio of photocleaved DNA to intact DNA was plotted against the log of the rhodium
23 concentration and fit to a sigmoidal curve (Figure S3). The inflection point of the sigmoidal fit
24 was used to determine the binding affinity of the competing $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$
25 metalloinsertor by solving simultaneous equilibria equations using the known binding affinity of
26 $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{3+}$. The binding affinities of these complexes are shown in Table 1. All
27 complexes were tested as racemic mixtures and exhibit binding affinities in the range of 2.4 to
28 $7.2 \times 10^6 \text{ M}^{-1}$ (Table 1). Despite differences in ligand steric bulk, all **Rh-O** metalloinsertors
29 tested have binding affinities within one order of magnitude of each other, and thus bind DNA
30 with comparable affinity.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Binding was assessed further via melting temperature analysis. A short, palindromic
4 DNA sequence containing a central CC mismatch was incubated in the presence of the parent
5 metalloinsertor, $[\text{Rh}(\text{bpy})_2(\text{chrysi})]\text{Cl}_3$, or the **Rh-O** metalloinsertor,
6 $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]\text{Cl}_2$. The chosen DNA sequence has a low T_m and therefore exists as
7 ssDNA at room temperature.¹³ In the presence of metalloinsertor, however, the DNA anneals and
8 the melting temperature increases dramatically to 44.9 ± 0.6 and 41.3 ± 0.5 °C for
9 $[\text{Rh}(\text{bpy})_2(\text{chrysi})]\text{Cl}_3$ and $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]\text{Cl}_2$, respectively (Supplementary Figure
10 S3B). These results are in good agreement with the results of the DNA binding assay describe
11 above and corroborate the result that parent and **Rh-O** metalloinsertors have comparable
12 binding affinities to mismatches in DNA, with $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]\text{Cl}_2$ stabilizing DNA to a
13 slightly lesser extent than $[\text{Rh}(\text{bpy})_2(\text{chrysi})]\text{Cl}_3$.

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28 *pK_a Determination of Metalloinsertors.* The pK_a values of $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$
29 metalloinsertors were assessed *via* spectroscopic pH titrations (Table 1, Supplementary Figures
30 S4-S8). The absorbance of a 435-440 nm peak, which corresponds to a charge transfer located on
31 the chrysi ligand, was plotted against the pH of the solution for each complex.²⁸ Data were fit to
32 a sigmoidal curve and the inflection point was taken as the pK_a of the complex, specifically of
33 the imine proton on the chrysi ligand. All **Rh-O** metalloinsertors exhibited pK_a values in the
34 range of 8.1 to 9.1, which are above physiological pH (Table 1), indicating that the chrysi ligands
35 of these complexes remain protonated in cell culture media or within cells. It has been shown
36 previously that fully protonated chrysi ligands, which are seen with **Rh-O** metalloinsertors,
37 buckle in contrast to the deprotonated chrysi ligands of the parent metalloinsertors, which are
38 completely flat and thus easy to stack with the DNA base pairs once inserted.¹⁹
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *Partition Coefficient and Lipophilicity of Metalloinsertors.* The [Rh(L)(chrysi)(PPO)]Cl₂
4 family of metalloinsertors was designed to vary in lipophilicity, and the partition coefficients of
5 each [Rh(L)(chrysi)(PPO)]²⁺ metalloinsertor were determined between aqueous buffer (10 mM
6 Tris-HCl, pH 7.4) and 1-octanol according to literature methods.²⁴ Absorbance measurements at
7 the ~260 nm peak were made in the 1-octanol phase before and after equilibration with the
8 aqueous phase. These absorbance values were compared to determine the partition coefficient,
9 log P (Table 1, Supplementary Figures S9-S12). The log P values followed the expected trend
10 with the least bulky complexes ([Rh(bpy)(chrysi)(PPO)]²⁺ and [Rh(HDPA)(chrysi)(PPO)]²⁺)
11 having the lowest log P values and the bulkiest complex ([Rh(DIP)(chrysi)(PPO)]²⁺) having the
12 greatest log P value. Surprisingly, despite their cationic nature, under these conditions the
13 [Rh(L)(chrysi)(PPO)]²⁺ metalloinsertors are all lipophilic and have partition coefficients favoring
14 octanol over water, ranging from 0.68 to >2.0.

15
16
17 *Cytotoxic and Anti-Proliferative Effects in MMR-Deficient and -Proficient Cells.* The
18 ability of metalloinsertors to selectively kill or impair growth of MMR-deficient cells is a critical
19 factor in their potential value as chemotherapeutic agents.^{19,29} In this structure-activity
20 relationship study, we used ELISA and MTT assays to determine the effect of ligand substitution
21 on biological activity in MMR-deficient and -proficient cells. The ELISA was used to determine
22 the inhibitory effects on DNA synthesis and the MTT assay was performed to establish levels of
23 cytotoxicity. For the ELISA, each metalloinsertor was incubated with HCT116N (MMR-
24 proficient) or HCT116O (MMR-deficient) cells at various concentrations before treatment with
25 the unnatural nucleic acid BrdU. Colorimetric antibody treatment allowed the relative BrdU
26 incorporation into DNA to be quantified, and cellular proliferation was then determined as the
27 ratio of BrdU incorporation between metalloinsertor-treated cells and untreated control cells. The
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 results of the 48-hour metalloinsertor treatment are shown in Figure 3, and the results of a 24-
4
5 hour treatment are shown in Supplemental Figure S15. All $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$
6
7 metalloinsertors exhibit anti-proliferative activity with selectivity towards the MMR-deficient
8
9 cell line. The maximum proliferation difference (referred to as selectivity) between the cell lines
10
11 and the concentration at which this selectivity occurs (referred to as potency) are as follows: $77 \pm$
12
13 10% at 400 nM for $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$, $78 \pm 18\%$ at $2 \mu\text{M}$ for $[\text{Rh}(\text{bpy})(\text{chrysi})(\text{PPO})]^{2+}$,
14
15 $47 \pm 10\%$ at $25 \mu\text{M}$ for $[\text{Rh}(\text{HDPA})(\text{chrysi})(\text{PPO})]^{2+}$, $66 \pm 6\%$ at 400 nM for $[\text{Rh}(4,7-$
16
17 $\text{DMP})(\text{chrysi})(\text{PPO})]^{2+}$, $67 \pm 5\%$ at 400 nM for $[\text{Rh}(5,6\text{-DMP})(\text{chrysi})(\text{PPO})]^{2+}$, and $70 \pm 23\%$ at
18
19 160 nM for $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$.
20
21
22
23

24 For the MTT assay, each metalloinsertor was incubated with HCT116N (MMR-
25
26 proficient) or HCT116O (MMR-deficient) cells at various concentrations before the addition of
27
28 MTT, which can be converted into formazan by mitochondrial reductase activity in a functioning
29
30 cell. Colorimetric measurements of formazan allow the relative viability to be quantified, and
31
32 cellular viability is then determined as the ratio of formazan produced between metalloinsertor-
33
34 treated cells and untreated control cells. The results of the 72-hour treatment are shown in Figure
35
36 4 and the results of the 24-hour treatment are shown in Supplemental Figure S16. All
37
38 $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertors exhibit cytotoxic activity with selectivity towards the
39
40 MMR-deficient cell line. The maximum proliferation difference between the cell lines and the
41
42 concentration at which this difference occurs are as follows: $52 \pm 5\%$ at 300 nM for
43
44 $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$, $30 \pm 7\%$ at $2 \mu\text{M}$ for $[\text{Rh}(\text{bpy})(\text{chrysi})(\text{PPO})]^{2+}$, $13 \pm 11\%$ at $32 \mu\text{M}$
45
46 for $[\text{Rh}(\text{HDPA})(\text{chrysi})(\text{PPO})]^{2+}$, $46 \pm 8\%$ at 600 nM for $[\text{Rh}(4,7\text{-DMP})(\text{chrysi})(\text{PPO})]^{2+}$, $49 \pm$
47
48 3% at 600 nM for $[\text{Rh}(5,6\text{-DMP})(\text{chrysi})(\text{PPO})]^{2+}$, and $39 \pm 6\%$ at 640 nM for
49
50 $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$.
51
52
53
54
55
56
57
58
59
60

1
2
3 *Whole-Cell Uptake, Mechanism of Uptake, and Organelle Localization.* To better
4 understand the range of biological activities of these complexes, cellular uptake and mechanism
5 of uptake were examined *via* ICP-MS based assays. 24-hour ELISA and MTT assays were
6 performed to determine a suitable concentration for uptake and localization studies (which were
7 performed over a 24-hour timescale). To minimize cell death in this assay, a factor which can
8 complicate data interpretation, suitable dosing was determined to be at a concentration at which
9 there was noticeable anti-proliferative effects in the HCT116O cells *via* ELISA but no significant
10 cytotoxicity *via* MTT assay. Whole cell uptake studies were performed with each
11 [Rh(L)(chrysi)(PPO)]²⁺ complex at 0.5 μM with the exception of [Rh(DIP)(chrysi)(PPO)]²⁺,
12 which was performed at 0.2 μM due to its high cytotoxicity at 0.5 μM. For whole cell uptake
13 studies, cells were incubated with metalloinsertors for 24 h before they were lysed and analyzed
14 for rhodium content *via* ICP-MS, with rhodium concentrations normalized to the protein content
15 of each sample. The whole cell uptakes of each metalloinsertor in HCT116O cells are shown in
16 Figure 5 (results in HCT116N cells are similar and shown in Supplementary Figure S17).
17 Overall, all [Rh(L)(chrysi)(PPO)]²⁺ complexes exhibit uptake into cells at concentrations within
18 one order of magnitude of each other. The uptake of these complexes correlates generally with
19 their lipophilicity values, with the least lipophilic complexes ([Rh(HDPA)(chrysi)(PPO)]²⁺ and
20 [Rh(bpy)(chrysi)(PPO)]²⁺) having the poorest uptake and the most lipophilic complex
21 ([Rh(DIP)(chrysi)(PPO)]²⁺) having the highest uptake. Lipophilicity has long been correlated
22 with an increase in cellular uptake and a resultant increase in drug potency.^{30,31}

23
24 In addition to examining whole cell uptake of the [Rh(L)(chrysi)(PPO)]²⁺
25 metalloinsertors, the uptake over time and the mechanism of uptake were also examined. In the
26 former experiment, cells were incubated with a metalloinsertor for 0.5, 1, 3, 6, 9, or 24 h before
27

1
2
3 being lysed and analyzed for rhodium content by ICP-MS. The whole-cell uptake over time of
4 these metalloinsertors in HCT116O cells is shown in Figure 5 (results in HCT116N cells are
5 similar and shown in Supplementary Figure S17). The complexes appear to show significant
6 increases in uptake over the first 3-6 h of incubation with cells, followed by plateau with no
7 evidence of significant efflux during a 24-hour period. These results are consistent with previous
8 studies on metalloinsertors.²⁰

9
10
11
12
13
14
15
16
17 A metabolic inhibition assay was performed to better understand the mechanism of
18 cellular uptake of $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertors. HCT116N and HCT116O cells were
19 pre-treated with the metabolic inhibitors oligomycin A, an inhibitor of oxidative
20 phosphorylation, and 2-deoxy-D-glucose, an inhibitor of glycolysis.²⁷ Metabolic inhibition
21 depletes cellular ATP (adenosine triphosphate), so any compound that is taken into the cell *via*
22 an active, ATP-dependent mechanism should have reduced uptake in metabolically depleted
23 cells. Conversely, complexes taken into the cell *via* a passive mechanism, such as passive
24 diffusion, are not affected by metabolic inhibition and therefore the drug should accumulate in
25 inhibited and uninhibited cells at similar concentrations. $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$ and the
26 parent metalloinsertor, $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{3+}$, were studied to determine if the mechanism of
27 metalloinsertor uptake was ATP-dependent. The compounds RbCl and $[\text{Ru}(\text{dppz})(\text{DIP})_2]^{2+}$ were
28 included as positive and negative controls, respectively. The rubidium ion of RbCl is transported
29 into the cell by Na,K-ATPase, an ATP-dependent ion pump, while $[\text{Ru}(\text{dppz})(\text{DIP})_2]^{2+}$ has
30 previously been shown to enter the cell *via* passive diffusion.^{27, 32} Cells were treated with each
31 compound for 1 h before they were lysed and analyzed by ICP-MS for metal content. As
32 rubidium, ruthenium, and rhodium are not naturally present in cells or cell culture reagents, all
33 three elements can be analyzed as low-background analytes by ICP-MS. The results of each
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 compound in HCT116O cells are shown in Figure 6 (results in HCT116N cells are similar and
4 shown in Supplementary Figure S18). As expected, RbCl showed a significant decrease in
5 uptake when pre-treated with metabolic inhibitors and $[\text{Ru}(\text{dppz})(\text{DIP})_2]^{2+}$ was unaffected by
6 inhibitor pre-treatment. Similar to $[\text{Ru}(\text{dppz})(\text{DIP})_2]^{2+}$, $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$ and
7 $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{2+}$ were also unaffected by inhibitor pre-treatment, suggesting these complexes
8 are also taken into the cell via an ATP-independent mechanism, such as passive diffusion. Since
9 these complexes are all lipophilic and cationic, passive diffusion is a reasonable uptake
10 mechanism, with the negative membrane potential driving diffusion and relatively high
11 lipophilicity facilitating the process as the molecules can more readily partition into the cellular
12 membranes.³³

13
14
15
16
17
18
19
20
21
22
23
24
25
26 Subcellular localization into the nucleus (the on-target organelle) and mitochondria (a
27 major off-target organelle) were also examined by an ICP-MS assay. Localization studies were
28 performed with each $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertor at 0.5 μM with the exception of
29 $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$, which was performed at 0.2 μM . For localization studies, cells were
30 incubated with metalloinsertors for 24 h before they were lysed and analyzed for rhodium
31 content via ICP-MS, with rhodium concentrations normalized to the protein content of each
32 sample. The whole cell uptakes of each metalloinsertor in HCT116O cells are shown in Figure 7
33 (results in HCT116N cells are similar and shown in Supplementary Figure S19). Overall, all
34 $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ complexes have comparable nuclear uptakes and mitochondrial uptakes
35 to one another with the exception of $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$, which has nuclear and
36 mitochondrial uptakes that are 2-3 times higher than other complexes despite being dosed at a
37 lower concentration. All complexes appear to enter the nucleus at high enough concentrations to
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 bind DNA mismatches, with a significant enrichment in nuclear concentration over the
4
5 extracellular concentration of rhodium (Supplementary Table S1).
6
7
8
9

10 **DISCUSSION**

11
12 Early generations of rhodium metalloinsertors, which exclusively contain Rh–N ligand
13
14 coordination, are a richly studied family of metal complexes that can selectively bind to DNA
15
16 base pair mismatches and lead to selective cell death in MMR-deficient cells. Across multiple
17
18 studies, these metalloinsertors were determined to have several characteristic and consistent
19
20 behaviors. Through *in vitro* experiments, we have observed that only the Δ - enantiomer of these
21
22 Rh–N coordinated complexes is capable of binding mismatches in B-form DNA.³⁴ In cellular
23
24 studies, these metalloinsertors have been observed to selectively kill cells in concentration ranges
25
26 of 5–40 μM .^{20,21} In one structure-activity relationship study, the steric bulk of the ancillary
27
28 ligands on a metalloinsertor was seen to influence DNA binding properties and, ultimately, alter
29
30 cellular selectivity.²¹ In another structure-activity relationship study, the lipophilicity of the
31
32 ancillary ligands on a metalloinsertor was seen to dramatically influence its subcellular
33
34 localization within a cell and, again, alter cellular selectivity.²⁰
35
36
37
38
39

40 While the above trends seem to ring true across parent metalloinsertors containing
41
42 exclusively Rh–N ligand coordination, the recent emergence of a new family of metalloinsertors
43
44 that contain Rh–O ligand coordination has challenged many of these characteristics and
45
46 behaviors.¹⁹ For instance, both enantiomers of **Rh–O** metalloinsertors are capable of binding
47
48 DNA mismatches *in vitro*, and are furthermore capable of inducing selective cellular toxicity at
49
50 *nanomolar* concentrations. Additionally, changes in lipophilicity and steric bulk of the O-
51
52 containing ligand seemed to have little, if any, effect on DNA binding affinity and cellular
53
54
55
56
57
58
59
60

1
2
3 selectivity. This remarkable shift in metalloinsertor activity revealed that these **Rh–O** complexes
4
5 have distinct *in vitro* characteristics and biological properties from their parent metalloinsertor
6
7 complexes. As such, a new family of **Rh–O** metalloinsertors has been synthesized,
8
9 characterized, and investigated for biological activity. In contrast to the first generation of **Rh–O**
10
11 metalloinsertors in which the O-containing ligand was varied, in this new family an ancillary
12
13 ligand was varied and the O-containing ligand was kept constant. This family is of the form
14
15 $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$, where L = bpy, phen, HDPA, 4,7-DMP, 5,6-DMP, and DIP. This ligand
16
17 variation influences many features of the metalloinsertor, including steric bulk and lipophilicity,
18
19 both of which have previously been seen to affect DNA binding and cellular activity of the
20
21 parent metalloinsertors.^{20,21} In studying this family of complexes, we aimed to test the unique
22
23 biological activity of metalloinsertors containing the **Rh–O** ligand framework and begin to
24
25 understand the high potency and improved selectivity exhibited by these metalloinsertors over
26
27 parent metalloinsertors and other DNA-binding complexes.
28
29
30
31
32

33 *Robustness of Biological Activity of the Rh–O Ligand Framework.* A primary aim of this
34
35 structure-activity relationship study was to determine if altering the ancillary ligand of **Rh–O**
36
37 metalloinsertors would significantly affect the biological activity of these complexes. Biological
38
39 activity was assessed through both ELISA and MTT assays in two cell lines, HCT116N and
40
41 HCT116O. These cells are derived from the same colorectal carcinoma cell line but differ
42
43 primarily in that HCT116N cells are MMR-proficient whereas HCT116O cells are MMR-
44
45 deficient.³⁵ For this reason, HCT116O cells have a higher relative abundance of DNA
46
47 mismatches over HCT116N cells and therefore should be more sensitive to mismatch-targeting
48
49 metalloinsertors.³⁶
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Indeed, all complexes prepared showed highly selective anti-proliferative or cytotoxic
4 effects toward the MMR-deficient cells over the MMR-proficient cells in both ELISA (Figure 3)
5 and MTT assays (Figure 4), with the exception of $[\text{Rh}(\text{HDPA})(\text{chrysi})(\text{PPO})]^{2+}$, which only
6 shows activity in the ELISA. While selectivity was seen for all complexes, the effective
7 concentrations varied by two orders of magnitude across the family. For instance,
8 $[\text{Rh}(\text{HDPA})(\text{chrysi})(\text{PPO})]^{2+}$ has very low potency and little selectivity compared to other **Rh-O**
9 metalloinsertors. Although it does appear to interfere selectively with DNA synthesis *via* ELISA,
10 this biological interaction does not appear significant enough to produce cytotoxic effects in the
11 MTT assay, even at high drug concentrations (Figure 4). HDPA is the only ligand containing a
12 labile proton and the only ligand that forms a 6-ring chelate with the metal, and it seems possible
13 that these structural features ultimately influence the biological activity of the
14 $[\text{Rh}(\text{HDPA})(\text{chrysi})(\text{PPO})]^{2+}$. It is possible that the 6-member chelate could cause structural
15 aberrations and the proton on HDPA could cause hydrogen-bonding interactions that ultimately
16 alter DNA-binding or DNA-processing by proteins, which could cause a decrease in toxicity.
17 $[\text{Rh}(\text{bpy})(\text{chrysi})(\text{PPO})]^{2+}$ has the second lowest potency of this new family, though remarkably
18 this complex still shows higher potency than the parent metalloinsertors containing only Rh-N
19 coordination.²⁰ The phenanthroline-derived metalloinsertors, $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$,
20 $[\text{Rh}(4,7\text{-DMP})(\text{chrysi})(\text{PPO})]^{2+}$, and $[\text{Rh}(5,6\text{-DMP})(\text{chrysi})(\text{PPO})]^{2+}$, all show comparable
21 nanomolar potencies and selectivities in the ELISA and MTT assays.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 Perhaps the most surprising biological activity is seen with $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$.
48 Historically, metalloinsertors containing the bulky DIP ligand have shown no selectivity for the
49 MMR-deficient cell line.²¹ This lack of selectivity was attributed to substantially lower mismatch
50 binding affinities (10^4 M^{-1} for $[\text{Rh}(\text{DIP})_2(\text{chrysi})]^{3+}$) owing to ancillary bulk, as well as off-target
51
52
53
54
55
56
57
58
59
60

1
2
3 localization into the mitochondria, a property that is common with lipophilic cations.^{20,37}
4
5 [Rh(DIP)(chrysi)(PPO)]²⁺, however, *does* exhibit selective cytotoxicity towards MMR-deficient
6
7 cells over proficient cells in both the ELISA and MTT assays. In fact, [Rh(DIP)(chrysi)(PPO)]²⁺
8
9 displays a similar selectivity and ~2-fold higher potency than [Rh(phen)(chrysi)(PPO)]²⁺ when
10
11 measured by ELISA (Figure 3).
12
13

14
15 Overall, these results confirm that **Rh–O** metalloinsertor biological selectivity is
16
17 minimally influenced by substitution at the ancillary ligand.¹⁹ Thus far, all of the **Rh–O**
18
19 metalloinsertors, derivatized at the O-containing ligand or ancillary ligand, have exhibited
20
21 selectivity in ELISA and/or MTT assays, regardless of steric bulk or lipophilicity, factors that
22
23 had heavily influenced (and sometimes abolished) the selectivity of parent metalloinsertors. It is
24
25 noteworthy that this selectivity profile, wherein the **Rh–O** metalloinsertors selectively kill
26
27 MMR-deficient cells, is shared with the parent complexes and is in stark contrast to what is seen
28
29 with all other DNA-targeting therapeutics, which preferentially kill MMR-proficient cells.^{17,18}
30
31 Although parent and **Rh–O** metalloinsertors share this unique selectivity profile and have similar
32
33 *in vitro* binding properties, suggesting they should interact with DNA in a similar way, the
34
35 **Rh–O** metalloinsertors are dramatically more potent than the parent metalloinsertors, with
36
37 nearly all **Rh–O** complexes (with the sole exception being [Rh(HDPA)(chrysi)(PPO)]²⁺) having
38
39 greater cytotoxicity in MMR-deficient cells than *any* of the parent metalloinsertors. It stands to
40
41 reason, then, that the high potency and selectivity of these **Rh–O** complexes does not reflect a
42
43 difference in DNA binding affinity from the parent complexes, but rather it must instead reflect a
44
45 difference in structure associated with the DNA-metalloinsertor lesion. That is, if the frequency
46
47 of DNA binding is comparable between the **Rh–O** and parent metalloinsertors, the lesion formed
48
49 by **Rh–O** metalloinsertors must activate a cellular response at lower concentrations.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *Uptake Characteristics.* Although the $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ family shows consistent
4 activity towards MMR-deficient cells, the selectivities and potencies of these complexes vary
5 significantly across the family from 160 nM to 25 μM . It was initially hypothesized that these
6 differences in biological activity could be due to differences in cellular uptake. In particular, it
7 seemed possible that the least potent complexes, $[\text{Rh}(\text{HDPA})(\text{chrysi})(\text{PPO})]^{2+}$ (which has almost
8 no cytotoxic properties at 40 μM) and $[\text{Rh}(\text{bpy})(\text{chrysi})(\text{PPO})]^{2+}$ (which has nearly 10-fold lower
9 potency than $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$), could be less effective due to low uptake. Similarly, it
10 was proposed that increased uptake could be responsible for the high potency of
11 $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$. Indeed, it does seem possible that uptake may explain some of the
12 observed potency trends: despite being dosed at 0.2 μM , $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$ exhibits
13 similar uptake to $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$, which was dosed at 0.5 μM . The finding suggests
14 that $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$ may induce biological effects at half the concentration of
15 $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$ as a result of complexes exhibiting similar uptakes at these
16 concentrations. However, uptake alone appears insufficient to explain the potencies of other
17 complexes. For instance, $[\text{Rh}(\text{HDPA})(\text{chrysi})(\text{PPO})]^{2+}$ and $[\text{Rh}(\text{bpy})(\text{chrysi})(\text{PPO})]^{2+}$ have
18 comparably low uptake into the cell despite a >10-fold difference in activity.

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Organelle-specific uptake is also worthy of consideration when examining the activity of
these complexes. Studies on previous generations of parent metalloinsertors bearing solely Rh-N
ligand coordination showed that off-target mitochondrial uptake is strongly influenced by ligand
lipophilicity, with the most lipophilic parent metalloinsertors having high mitochondrial uptake
and low selectivity for MMR-deficient cells.^{20,38} Surprisingly, *all Rh-O* metalloinsertors studied
here are more lipophilic than *any* of the parent metalloinsertors described above, yet all **Rh-O**
complexes exhibit selective cytotoxicity towards MMR-deficient cells, making their selectivity

1
2
3 patterns distinct from trends followed by the parent metalloinsertors. To better understand this
4
5 marked change in trends, on-target nuclear localization and off-target mitochondrial localization
6
7 experiments were performed to assess the biological activity of $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$
8
9 complexes, particularly DIP, which shows selectivity despite its very high lipophilicity.
10
11

12 As indicated, all $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertors enter the nuclei to a similar
13
14 extent and at high enough concentrations to bind DNA mismatches (Figure 7 and Supplementary
15
16 Table S1). Similarly, all $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertors enter the mitochondria to a
17
18 comparable extent. Although nuclear and mitochondrial uptake cannot be compared directly
19
20 (since each is normalized to the total protein in the organelle), the localization patterns of **Rh-O**
21
22 versus parent metalloinsertors can be compared (See Supplementary Figure S20). This
23
24 comparison shows that, unlike their Rh-N coordinated predecessors, **Rh-O** metalloinsertor
25
26 localization into the mitochondria is not significantly influenced by lipophilicity. In fact, despite
27
28 being lipophilic, **Rh-O** complexes exhibit uptake profiles that are comparable to *hydrophilic*
29
30 parent metalloinsertors (which have low mitochondrial uptake) and are distinct from *lipophilic*
31
32 parent metalloinsertors (which have high mitochondrial uptake). This trend in localization is
33
34 consistent with the biological activity we observed; similar to the hydrophilic parent
35
36 metalloinsertors, **Rh-O** complexes are highly selective and show little off-target cytotoxicity.
37
38 Overall, these data indicate that **Rh-O** metalloinsertors are able to maintain their high selectivity
39
40 and potency because the ligand substitutions do not strongly influence their subcellular
41
42 localization. Since these complexes exhibit low mitochondrial uptake, off-target mitochondria-
43
44 induced toxicity does not overwhelm the biological response, and the selective nuclear- and
45
46 mismatch-mediated response can prevail.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 It is also interesting to note that both MMR-proficient HCT116N cells and MMR-
4 deficient HCT116O cells had comparable levels of uptake and similar localization profiles,
5 showing that metalloinsertors enter HCT116N and HCT116O cells at the same rate, through the
6 same passive mechanism, and to the same extent (Figure 5 and 6 and Supplementary Figures S17
7 and S18). These details support the idea that the biological selectivity seen in these cells is not a
8 feature of different cellular uptake or elimination properties. Furthermore, the nuclear uptake into
9 the MMR-deficient and proficient cells are comparable (Figure 7 and Supplementary Figure
10 S19). Therefore, with similar concentrations of metalloinsertors entering the nuclei and similar
11 mismatch binding affinities, any DNA-mediated cytotoxicity must result from a difference in
12 how the drugs interact with the DNA. Rationally, this difference must depend upon an increased
13 mismatch targeting in MMR-deficient cells, where DNA base pair mismatches are more
14 abundant.³⁶

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31 *Source of Potency for the Rh-O Metalloinsertors.* Although MMR-deficient cells have a
32 relative abundance of mismatches compared to MMR-proficient cells, the total number of
33 mismatches formed during each cellular replication is ultimately small due to the high fidelity
34 and proofreading abilities of polymerases. It is clear, therefore, that the lesion formed by parent
35 metalloinsertors must be significantly potent such that even a small number of metalloinsertor-
36 DNA lesions can result in selective cell death. Moreover, despite their similar mismatch binding
37 affinities, the **Rh-O** metalloinsertors are even more potent than parent metalloinsertors, and
38 therefore these **Rh-O** metalloinsertors must produce a unique lesion structure at the mismatched
39 site that can activate a response at even lower concentrations (and therefore fewer
40 metalloinsertor-DNA lesions) than parent metalloinsertors.

1
2
3 Does the increase in potency depend upon a difference in how these **Rh–O**
4 metalloinsertors bind to DNA within the cell?¹⁹ As discussed above, both the Δ - and Λ -
5 enantiomers of $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$ can bind to DNA mismatches *in vitro* and selectively
6 kill MMR-deficient cells in culture. This behavior is distinct from parent metalloinsertors, for
7 which only the Δ -enantiomer can bind mismatches and produce biological effects.¹⁵ The ability
8 of both enantiomers of **Rh–O** metalloinsertors to bind mismatched DNA suggests the binding
9 interaction must be fundamentally distinct from that of the parent metalloinsertors; these new
10 **Rh–O** metalloinsertors must bind DNA in a way that can accommodate the Λ -enantiomer.
11
12
13
14
15
16
17
18
19
20

21 Furthermore, some evidence suggests that even the DNA-binding ability of the Δ -
22 enantiomer may be altered in these **Rh–O** metalloinsertors. Previously, it was observed that
23 bulky parent metalloinsertors, such as $[\text{Rh}(\text{DIP})_2(\text{chrysi})]^{3+}$, exhibited poor binding affinities (10^4
24 M^{-1}) and could not easily be modeled to fit into a mismatched DNA lesion due to significant
25 steric clashing between the DIP ligands and the DNA backbone.²¹ In contrast, significant
26 differences in ancillary ligand steric bulk have minimal effect on the binding affinities of **Rh–O**
27 metalloinsertors, which all bind to DNA with micromolar affinity. Even the most sterically bulky
28 complex, $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$, has a relatively high affinity for mismatched DNA (10^6M^{-1})
29 despite containing the bulky DIP ligand. It therefore seems that the inclusion of the DIP ligand
30 is not sufficient to preclude DNA binding, and perhaps this dramatic increase in binding affinity
31 of a DIP-containing metalloinsertor may indicate that a new binding interaction exists that can
32 accommodate the steric bulk of these **Rh–O** metalloinsertors.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 Another consideration is the conformation of the chrysi ligand of these new **Rh–O**
50 metalloinsertors. All **Rh–O** complexes have chrysi imine pK_a values above physiological pH,
51 indicating that they remain protonated in the intracellular environment. This protonation results
52
53
54
55
56
57
58
59
60

1
2
3 in steric clashing between the imine proton and an aromatic proton in the chrysi system and, as a
4
5 result, the chrysi ligand becomes buckled relative to the rhodium center to relieve the steric
6
7 strain.¹⁹ This is in stark contrast to parent metalloinsertors which deprotonate at cellular pH and
8
9 therefore do not exhibit steric clashing between the imine and aromatic protons. As a result, the
10
11 chrysi ligand lays planar in these parent metalloinsertors. Distortion of the chrysi ligand, the
12
13 ligand that interacts most intimately with the DNA, likely disrupts the overall metalloinsertor-
14
15 DNA binding interaction, further suggesting there is likely a difference in how **Rh-O** and parent
16
17 metalloinsertors bind to DNA.¹⁹

18
19
20
21
22 Lastly, the **Rh-O** complexes reported here are lipophilic ($\log P > 0$), whereas comparable
23
24 parent metalloinsertors are hydrophilic ($\log P < 0$).³⁸ This change in lipophilicity could alter the
25
26 way **Rh-O** complexes interact with the hydrophobic bases of DNA or even DNA-processing
27
28 proteins that may be responsible for recognizing the DNA-metalloinsertor lesion. Overall, these
29
30 results suggest that the **Rh-O** metalloinsertors bind to DNA differently than parent
31
32 metalloinsertors. While these complexes still appear to undergo metalloinsertion, as evidenced
33
34 by their ability to bind mismatched DNA with high affinity, it is unclear how their binding might
35
36 be distinct from parent metalloinsertors. It seems possible that a subtle difference in the extent or
37
38 orientation of mismatched base ejection or in the unwinding of the DNA helix by the
39
40 metalloinsertor could ultimately result in a difference in how that lesion is recognized or
41
42 processed within the cell, which could lead to overall cellular response and increased potency.
43
44 Crystallographic studies of **Rh-O** metalloinsertors with DNA are currently underway to
45
46 investigate the potential difference between parent and **Rh-O** metalloinsertor binding.

47
48
49
50
51
52 *Implications for Future Metalloinsertor Design.* The $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ family of
53
54 metalloinsertors described herein display biological selectivity and potency that are maintained
55
56
57
58
59
60

1
2
3 across various ligand frameworks varying in size and lipophilicity. When compared with other
4
5 **Rh–O** metalloinsertors in which the PPO-type ligand is varied, metalloinsertors containing the
6
7 **Rh–O** motif are consistent in their biological selectivity (and, to a large extent, potency) for
8
9 MMR-deficient cells regardless of significant alterations to their ancillary ligands. It has
10
11 previously been shown that the metalloinsertors with DIP ligands and PPO-type ligands cannot
12
13 be easily modeled into a mismatched DNA lesion due to steric clashes with the DNA
14
15 structure.^{19,21} Despite steric bulk, **Rh–O** metalloinsertors have comparable binding affinities to
16
17 parent metalloinsertors and significantly improved biological activity. Furthermore, these
18
19 complexes show little enantioselectivity; both isomers bind DNA and show high potency, further
20
21 supporting that their metalloinsertion binding interaction markedly differs from parent
22
23 metalloinsertors. Taken together, these observations show that the **Rh–O** metalloinsertor
24
25 framework has great potential for the design of new therapeutics and for the attachment of new
26
27 payloads, while maintaining biological selectivity.^{39–43} The consistently high potency and cell
28
29 selectivity of these complexes is unique and provides the basis for new generations of
30
31 metalloinsertors and metalloinsertor conjugates.
32
33
34
35
36
37
38
39

40 **Associated Content**

41
42 Some experimental methods, representative binding affinity determination, pH titrations,
43
44 enantiomeric characterization, further DNA binding characterization, and cytotoxicity of the two
45
46 enantiomers as well as cellular uptake data. This material is available free of charge via the
47
48 Internet at <http://pubs.acs.org>.
49
50

51 **Author Information**

52
53 Corresponding Author: jkbarton@caltech.edu
54
55
56
57
58
59
60

1
2
3 Notes: The authors declare no competing financial interest.
4
5
6

7
8 **Acknowledgements**
9

10 Financial support for this work from the NIH (GM33309) and AMGEN is gratefully
11 acknowledged. We thank the Department of Defense for supporting K.M.B. through the National
12 Defense Science & Engineering Graduate Fellowship (NDSEG) Program. We also thank Paige
13 Gannon for assisting in uptake and localization studies of the complexes. This project benefitted
14 from the use of instrumentation made available by the Caltech Environmental Analysis Center
15 and we gratefully acknowledge guidance from Dr. Nathan Dalleska. We also thank the Moore
16 Foundation Center for the Chemistry of Cellular Signaling. The Caltech Center for Catalysis and
17 Chemical Synthesis and Dr. Scott Virgil are gratefully acknowledged for assistance with
18 enantiomeric separation.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

- (1) Cheung-Ong, K.; Giaever, G.; Nislow, C. *Chem. Biol.* **2013**, *20*, 648–659.
- (2) Wang, D.; Lippard, S. J. *Nat. Rev. Drug Discov.* **2005**, *4*, 307–320.
- (3) Kelland, L. *Nat. Rev. Cancer* **2007**, *7*, 573–584.
- (4) Nitiss, J. L. *Nat. Rev. Cancer* **2009**, *9*, 338–350.
- (5) *DNA-targeting Molecules as Therapeutic Agents*, 2nd ed.; Waring, M. J., Ed.; Chemical Biology; The Royal Society of Chemistry, 2018.
- (6) Shoshan, M. C.; Linder, S. *Expert Opin. Drug Metab. Toxicol.* **2008**, *4*, 273–280.
- (7) Tham, K. C.; Kanaar, R.; Lebbink, J. H. G. *DNA Repair (Amst.)*. **2016**, *38*, 75–83.
- (8) Arzimanoglou, I. I.; Gilbert, F.; Barber, H. R. K. *Cancer* **1998**, *82*, 1808–1820.
- (9) Pino, M. S.; Mino-Kenudson, M.; Wildemore, B. M.; Ganguly, A.; Batten, J.; Sperduti, I.; Iafrate, A. J.; Chung, D. C. *J. Mol. Diagnostics* **2009**, *11*, 238–247.
- (10) (a) Jackson, B. A.; Barton, J. K. *J. Am. Chem. Soc.* **1997**, *119*, 12986–12987. (b) Jackson, B. A.; Barton, *Biochemistry* **2000**, *39*, 6176–6182
- (11) Zeglis, B. M.; Boland, J. A.; Barton, J. K. *J. Am. Chem. Soc.* **2008**, *130*, 7530–7531.
- (12) Jackson, B. A.; Alekseyev, V. Y.; Barton, J. K. *Biochemistry* **1999**, *38*, 4655–4662.
- (13) Cordier, C.; Pierre, V. C.; Barton, J. K. *J. Am. Chem. Soc.* **2007**, *129*, 12287–12295.
- (14) (a) Pierre, V. C.; Kaiser, J. T.; Barton, J. K. *Proc. Natl. Acad. Sci.* **2007**, *104*, 429–434; (b) Zeglis, B.M.; Pierre, V.C.; Kaiser, J.T.; Barton, J.K. *Biochem.* **2009**, *48*, 4247–4253.
- (15) Hart, J. R.; Glebov, O.; Ernst, R. J.; Kirsch, I. R.; Barton, J. K. *Proc. Natl. Acad. Sci.* **2006**, *103*, 15359–15363.
- (16) Bailis, J. M.; Gordon, M. L.; Gurgel, J. L.; Komor, A. C.; Barton, J. K.; Kirsch, I. R. *PLoS One* **2013**, *8*, e78726.
- (17) Fink, D.; Nebel, S.; Aebi, S.; Zheng, H.; Cenm, B.; Nehmã, A.; Christen, D.; Howell, S. B. *Cancer Res.* **1996**, *56*, 4881–4886.
- (18) Fink, D.; Nebel, S.; Norris, P. S.; Aebi, S.; Kim, H. K.; Haas, M.; Howell, S. B. *Br. J. Cancer* **1998**, *77*, 703–708.
- (19) Komor, A. C.; Barton, J. K. *J. Am. Chem. Soc.* **2014**, *136*, 14160–14172.
- (20) Komor, A. C.; Schneider, C. J.; Weidmann, A. G.; Barton, J. K. *J. Am. Chem. Soc.* **2012**, *134*, 19223–19233.
- (21) Ernst, R. J.; Song, H.; Barton, J. K. *J. Am. Chem. Soc.* **2009**, *131*, 2359–2366.
- (22) Zeglis, B. M.; Barton, J. K. *Nat. Protoc.* **2007**, *2*, 357–371.
- (23) Lee, J. R.; Lung, J. W. *J. Chinese Chem. Soc.* **2003**, *50*, 227–232.
- (24) Sangster, J. *J. Phys. Chem. Ref. Data* **1989**, *18*, 1111–1229.
- (25) Komor, A. C.; Schneider, C. J.; Weidmann, A. G.; Barton, J. K. *J. Am. Chem. Soc.* **2012**, *134*, 19223–19233.
- (26) Ahmad, K. A.; Iskandar, K. B.; Hirpara, J. L.; Clement, M. V.; Pervaiz, S. *Cancer Res.* **2004**, *64*, 7867–7878.
- (27) Puckett, C. A.; Barton, J. K. *Biochemistry* **2008**, *47*, 11711–11716.
- (28) Sitlani, A.; Long, E. C.; Pyle, A. M.; Barton, J. K. *J. Am. Chem. Soc.* **1992**, *114*, 2303–2312.
- (29) Bailis, J. M.; Weidmann, A. G.; Mariano, N. F.; Barton, J. K. *Proc. Natl. Acad. Sci.* **2017**, *114*, 6948–6953.
- (30) Reithofer, M. R.; Bytzeck, A. K.; Valiahdi, S. M.; Kowol, C. R.; Groessler, M.; Hartinger, C. G.; Jakupec, M. A.; Galanski, M.; Keppler, B. K. *J. Inorg. Biochem.* **2011**, *105*, 46–51.
- (31) Waring, M. J.; Johnstone, C. *Bioorganic Med. Chem. Lett.* **2007**, *17*, 1759–1764.

- 1
2
3 (32) Landero Figueroa, J. A.; Stiner, C. A.; Radzyukevich, T. L.; Heiny, J. A. *Sci. Rep.* **2016**,
4 6, 20551.
5 (33) Liu, X.; Testa, B.; Fahr, A. *Pharm. Res.* **2011**, 28, 962–977.
6 (34) Hart, J. R.; Glebov, O.; Ernst, R. J.; Kirsch, I. R.; Barton, J. K. *Proc. Natl. Acad. Sci.*
7 **2006**, 103, 15359–15363.
8 (35) Koi, M.; Umar, A.; Chauhan, D. P.; Cherian, S. P.; Carethers, J. M.; Kunkel, T. A.;
9 Boland, C. R. *Cancer Res.* **1994**, 54, 4308–4312.
10 (36) Glaab, W. E.; Tindall, K. R. *Carcinogenesis* **1997**, 18, 1–8.
11 (37) Modica-Napolitano, J. S.; Aprille, J. R. *Adv. Drug Deliv. Rev.* **2001**, 49, 63–70.
12 (38) Weidmann, A. G.; Komor, A. C.; Barton, J. K. *Philosophical Trans. R. Soc.* **2013**, 371,
13 20120117.
14 (39) Weidmann, A. G.; Barton, J. K. *Inorg. Chem.* **2015**, 54, 9626–9636.
15 (40) Petitjean, A.; Barton, J. K. *J. Am. Chem. Soc.* **2004**, 126, 14728–14729.
16 (41) Chari, R. V. J. *Acc. Chem. Res.* **2008**, 41, 98–107.
17 (42) Zeglis, B. M.; Barton, J. K. *J. Am. Chem. Soc.* **2006**, 128, 5654–5655.
18 (43) Nano, A.; Boynton, A. N.; Barton, J. K. *J. Am. Chem. Soc.* **2017**, 139, 17301–17304.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: General structure of the newest generation of **Rh-O** metalloinsertors (left), which show improved potency and selectivity over parent metalloinsertors that have exclusively Rh-N coordination (right). R has been varied between methyl, phenyl, pyridyl, and hexyl groups and N—N has been varied between several bpy, phen, and HDPA derivatives.




Figure 2. Chemical structures of [Rh(L)(chrysi)(PPO)]²⁺ family of rhodium metalloinsertors, with the PPO ligand shown in red and the changing ancillary ligand shown in blue.

Table 1: Binding affinity, pK_a, and Log P values for each metalloinsertor

Metalloinsertor	Binding Constant ($\times 10^6 \text{ M}^{-1}$) ^a	pK _a (2+ to 1+)	Log P
Δ -[Rh(phen)(chrysi)(PPO)] ²⁺	6.6	–	–
Λ -[Rh(phen)(chrysi)(PPO)] ²⁺	9.2	–	–
<i>rac</i> -[Rh(phen)(chrysi)(PPO)] ²⁺	5.5 ^b	8.3 \pm 0.3 ^a	1.4 \pm 0.1
<i>rac</i> -[Rh(bpy)(chrysi)(PPO)] ²⁺	7.2	8.9 \pm 0.1	0.68 \pm 0.07
<i>rac</i> -[Rh(HDPA)(chrysi)(PPO)] ²⁺	3.0	9.1 \pm 0.1	0.69 \pm 0.08
<i>rac</i> -[Rh(4,7-DMP)(chrysi)(PPO)] ²⁺	1.5	9.1 \pm 0.1	1.1 \pm 0.1
<i>rac</i> -Rh[(5,6-DMP)(chrysi)(PPO)] ²⁺	2.3	9.0 \pm 0.3	0.71 \pm 0.01
<i>rac</i> -[Rh(DIP)(chrysi)(PPO)] ²⁺	1.6	8.1 \pm 0.1	> 2.0 ^c

^a binding affinities measured using the DNA hairpin 5'-GGCAGGCATGGCTTTTGGCCATCCTGCC-3' (underline denotes mismatch) in 100 mM NaCl, 20 mM NaPi, pH 7.1 buffer. Competition titrations were performed against the photocleaving metalloinsertor [Rh(bpy)₂(chrysi)]Cl₃.

^b Values from reference 19

^c The change in absorbance in the [Rh(DIP)(chrysi)(PPO)]²⁺-containing 1-octanol phase before and after equilibration with the aqueous phase was too small to accurately and consistently measure.




Figure 3: Cellular proliferation ELISA for the $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertors in MMR-deficient (HCT1160, red circles) and MMR-proficient (HCT116N, blue squares) cells. Cells were incubated with various concentrations of metalloinsertor for 48 h before treatment with BrdU. Cell proliferation is shown as %BrdU incorporated into DNA compared to untreated control cells. Error is shown as the standard deviation of 5 replicates.




Figure 4: Cellular viability MTT assay for the $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertors in MMR-deficient (HCT1160, red circles) and MMR-proficient (HCT116N, blue squares) cells. Cells were incubated with various concentrations of metalloinsertor for 72 h before treatment with MTT. Cell proliferation is shown as % viability from MTT metabolism, compared to untreated control cells. Error is shown as the standard deviation of 5 replicates.




Figure 5: Whole-cell rhodium uptake assays. (*left*) Rhodium accumulation in HCT1160 was measured by ICP-MS analysis after a 24 hour incubation with $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ metalloinsertors (where **L** = **phen**, **bpy**, **HDPA**, **4,7-DMP**, **5,6-DMP**, or **DIP**). (*right*) Rhodium accumulation over time in HCT1160 cells was measured by ICP-MS for three metalloinsertors, $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$ (**phen**), $[\text{Rh}(\text{DIP})(\text{chrysi})(\text{PPO})]^{2+}$ (**DIP**), and the parent metalloinsertor $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{3+}$ (**Rh-BC**). Rhodium content was normalized to protein content determined by BCA assay, and the results of four independent trials were averaged with error shown as the standard deviation.




Figure 6: Uptake assay for rhodium metallosinsertors and controls. $[\text{Rh}(\text{phen})(\text{chrysi})(\text{PPO})]^{2+}$ (**phen**), $[\text{Rh}(\text{bpy})_2(\text{chrysi})]^{3+}$ (**Rh-BC**), $[\text{Ru}(\text{DIP})_2(\text{chrysi})]^{2+}$ (**Ru-DIP**), and **RbCl** accumulation in HCT1160 cells was measured by ICP-MS analysis after treatment with or without metabolic inhibitors (oligomycin and 2-deoxy-D-glucose). Rhodium, ruthenium, and rubidium contents were normalized to protein content determined by BCA assay. Each experiment was performed in triplicate and averaged, with error shown as the standard deviation. Each experiment was repeated three times with similar outcomes (not shown). RbCl* indicates that Rb concentrations for RbCl have been lowered by a factor of 500 in this graphic.




Figure 7: ICP-MS assay for nuclear and mitochondrial uptake of rhodium metalloinsertors. Rhodium accumulation in HCT1160 cells was measured by ICP-MS analysis after a 24 hour incubation with $[\text{Rh}(\text{L})(\text{chrysi})(\text{PPO})]^{2+}$ (where **L** = **phen**, **bpy**, **HDPA**, **4,7-DMP**, **5,6-DMP** or **DIP**). Rhodium content was normalized to protein content of each organelle fraction determined by BCA assay. The results of 4 independent studies were averaged with error shown as the standard deviation.

Table of Contents Image:

