

Almost Periodic Hill's Equation and the Rings of Saturn

Joseph E. Avron^(a) and Barry Simon^(b)

Department of Mathematics, California Institute of Technology, Pasadena, California 91125

(Received 30 January 1981)

Incommensurate perturbations of classical orbits lead to an almost periodic Hill's operator whose spectrum, we argue, is a Cantor set, but one with large Lebesgue measure. Applied to the rings of Saturn, this implies that the complex groove structure in the rings approximates a Cantor set. We also emphasize the possible relevance of the sun in producing "side gaps" which magnify the apparent gap size.

PACS numbers: 95.10.Ce, 03.65.Db, 96.30.Mh

One of the striking discoveries of Voyager I is that the structure of the rings of Saturn is exceedingly complex. Prior to this information, one of the more popular explanations of the structure of the rings was as an effect of resonances with the moons of Saturn.¹ Our goal here is to examine whether a resonance picture can possibly provide a structure as complicated as that observed. We will study the question within a linear stability analysis which will lead to the spectral analysis of an almost periodic Hill's equation. Our conclusion is that the structure of such a spectrum is sufficiently rich so that it *may* account for the main features of the ring structure. More definitive conclusions will have to await, among other things, further study of this spectral analysis which we hope this note will stimulate.

Linear stability is an approximation and as such does not have all the deep structure of the fully nonlinear equations of classical mechanics, as born out by the Kolmogorov-Arnold-Moser (KAM) theory.² However, it has the advantages of being conceptually simple, relatively easy to estimate orders of magnitude and the right qualitative features. Indeed, one can regard certain aspects of our analysis as a kind of "poor man's KAM." Moreover, since the basic equation turns out to be a time-independent Schrodinger equation (see below), one can use intuition³ from that subject in this analysis even though the analogy is on the level of the mathematics and not on the basic physics.

The main features of the theory of the rings of Saturn which emerges are as follows:

(1) The largest individual gaps are on the order of 20 km⁴ and such large gaps only occur near the position of a classical resonance, e.g., at the Cassini division.

(2) In an idealized model where one considers point particles running for an infinite time, with no collisions, the rings have the structure of a Cantor set (see below), i.e., the gaps are dense.

(3) Large scale features like the Cassini division are aggregates of a large number of gaps separated by thin ringlets, something consistent with the Voyager observations. We will argue that a long period can conspire with a moon resonance to produce resonance "side gaps" and therefore an aggregate of gaps. One candidate⁵ for this long period term is the sun.

Let us recall Hill's basic analysis of linear stability. Consider a closed orbit, $\vec{v}(t)$, in the x - y plane of Newton's equation $\ddot{\vec{v}} = \vec{F}[\vec{v}(t)]$. If we replace $\vec{v}(t)$ by $\vec{v}(t) + \hat{z}w(t)$ and keep first order terms in w we find $Hw = 0$ with

$$H = -(d^2/dt^2) + V(t), \quad (1)$$

where $V(t) \equiv -\partial_z F_z[\vec{v}(t)]$. If H has zero in its (continuous) spectrum w stays bounded; otherwise, there is a solution which grows exponentially. If the original orbit is not in a plane or if w is not perpendicular to that plane, H is replaced by a vector-valued Hill's operator.

To describe our picture of the spectrum of the relevant Hill's operator, we first describe certain Cantor sets.

A Cantor set is a subset of the real line which is closed, has no isolated points, and which is nowhere dense, i.e., whose complement is dense. The original set constructed by Cantor had the additional feature of zero-Lebesgue measure. In contrast we shall concentrate on Cantor sets with large Lebesgue measure. An example is as follows: Pick a sequence of odd (positive) integers $n_j = 2l_j + 1$ and expand $x = l_0 + \sum_{j=1}^{\infty} a_j / (n_1 \cdots n_j)$, $a_j = 0, \dots, n_j - 1$. The set of points with no $a_j = l_j$ is a Cantor set and its Lebesgue measure in $[l_0, l_0 + 1]$ is $\prod_{j=1}^{\infty} (1 - n_j^{-1})$, which will be close to 1 if $\sum n_j^{-1}$ is small.

We shall argue that the spectrum of (1) with $V(t)$ quasiperiodic is a Cantor set with a large Lebesgue measure. By quasiperiodic we mean $V(t) = f(\omega_1 t, \dots, \omega_n t)$, $f(t_1 + m_1, \dots, t_n + m_n) = f(t_1, \dots, t_n)$, m_i integers and ω_i incommensurate. The

evidence we have for this claim is the following⁶:

(1) For weak coupling, i.e., $V(t) = gf(t)$, g small, take $f(t) = \sum \vec{f}(\vec{m}) e^{2\pi i \vec{m} \cdot \vec{t}}$ with $\vec{f}(\vec{m}) \neq 0$, $|\vec{m}| \geq m_0$, and $\sum |\vec{f}(\vec{m})| < \infty$. First order perturbation theory⁷ gives gaps of order $g|\vec{f}(\vec{m})| \neq 0$ centered at energies $E_{\vec{m}} = (\pi \vec{m} \cdot \vec{\omega})^2$ whose total measure is small. The $E_{\vec{m}}$ are dense in $[0, \infty)$.

(2) There is a different class of almost periodic potentials (limit-periodic⁸) for which Cantor spectrum with large Lebesgue measure has been proven rigorously by Moser⁹ and independently by Avron and Simon.¹⁰

(3) Azbel¹¹ analyzed yet another class of potentials in the strong coupling limit. He argues that the spectrum has all the scales of the continued fraction expansion of the periods, something compatible with a Cantor set.

(4) A lattice version of Eq. (1) (a difference equation) has been analyzed numerically by Hofstadter¹² in another context and suggests a Cantor set for the spectrum.

Since Cantor sets have no scale, what appears to be a wide band on one scale is, in fact, several bands separated by narrow gaps on a finer scale. The aggregation of small individual gaps and bands to form apparent gaps and bands on a larger scale plays an important role in the sequel. To see how this may come about consider the spectrum of the Schrodinger operator

$$(-d^2/dt^2) + \lambda \cos(2\pi t) + \mu \cos(2\pi t/n), \quad (2)$$

with λ and μ fairly small, and n a large number so that μn is not too small. We take n an integer so we can analyze (2) in terms of Floquet theory although we assume the qualitative features we find are true even if n is nonintegral. Consider the gaps near energy 1. If $\mu = 0$, there is one gap of size λ . If $\lambda = 0$, there are many gaps at energy approximately $[(n-k)/n]^2$ (k an integer near 0) but the asymptotics of the Mathieu equation¹³ suggests that the size is $O[(\mu e^2)^n]$ which is very small. If both μ and λ are nonzero, the lowest order perturbation theory contribution to the gap which wants to open at energy $[(n-k)/n]^2$ is, for fixed k , $O[\lambda(\mu n)^k]$, the n 's coming from small energy denominators in perturbation theory. This argument suggests that when λ and μ are small but $\mu n \sim 1$, (2) has, in addition to its "main" gap of size λ near energy 1, many other "side gaps."

Now consider a test particle moving in a Keplerian orbit, $v_0(t)$, in the equatorial plane of Saturn in an idealized situation where we initially ignore the moons and the sun. As is well known all these orbits are stable. Now study the effect

of the moons with small inclinations.¹⁴ One should apply the linear stability equation to an actual orbit, $u(t)$, near $v_0(t)$, but this orbit is obviously very difficult to compute exactly and indeed it may not even exist in situations where linear stability predicts unstable orbits. We therefore make an approximation: Let $\vec{F}_0(r)$ and $\vec{F}_1(r)$ denote the forces on a ring particle due to Saturn and the moons, respectively, so $\vec{v}_0 = \vec{F}_0(\vec{v}_0)$. Since \vec{F}_1 is small an approximate solution is $\vec{v} \sim \vec{v}_0 + \vec{u}$ with

$$\ddot{\vec{u}} = (\vec{u} \cdot \nabla) \cdot \vec{F}_0[\vec{v}_0(t)] + \vec{F}_1[\vec{v}_0(t)]. \quad (3)$$

$\nabla \vec{F}_0$ has eigenvalues $[(2\pi/T_r)^2, (2\pi/T_r)^2, 0]$ with T_r the period of the Keplerian orbit and $\vec{F}_1 = \sum \vec{q}_i$, \vec{q}_i the disturbance function

$$\vec{q}_i = \nabla \{ Gm_i [|\vec{v}_0(t) - \vec{r}_i(t)|^{-3} - |\vec{r}_i(t)|^{-3} \vec{v}_0(t) \cdot \vec{r}_i(t)] \}. \quad (4)$$

m_i are the masses and \vec{r}_i the distance from Saturn of the satellite i . $\vec{u}(t)$ can be calculated by classical perturbation methods.¹⁵

If we apply linear stability to $v_0 + \mu$, we find for the virtual perturbation w

$$\ddot{w} = (\nabla_{\vec{r}} F_0)[v_0(t)]w + \nabla_{\vec{r}} F_1[u_0(t)]w + (\nabla_{\vec{r}} u \cdot \nabla F_0)[u_0(t)]w, \quad (5)$$

where we keep the first variation of the perturbation F_1 and the first and second variation of F_0 . Since $\nabla_{\vec{r}} F_0 = (2\pi/T_r)^2$, the period, and hence the radius, correspond to the energy in a Schrodinger equation. The situation here is somewhat complicated by the fact that the potential $V(t)$ is a (slowly varying) function of the period. In units of $(2\pi/T_r)^2$, the moon i contributes to the ∇F_1 term in (5) an amplitude of the order $(m_i/m_\Omega) |r_i/(r-r_i)|^3$. This is of order 10^{-7} for five moons of Saturn. The contribution to the $(\nabla \nabla F_0)u$ term has amplitude of order $(m_i/m_\Omega) |r_i/(r-r_i)|^3 \{n^2[(\Omega_i/\omega_r) - 1]^2\}^{-1}$, where n is the driving harmonic. This is again typically of order 10^{-7} unless ω_r is near resonance and the curly brackets are small. Thus the second term dominates near resonances. A simple perturbation argument for the Hill's equation gives an estimate of the width of an individual gap due to the moon Mimas alone at the Cassini division to be of order of 20 km, which agrees with numerical solutions of the three-body problem.¹ Consider now the effects of Mimas and the sun together. The sun has a long period of $\sim 10^4$ d. The amplitude in $(\vec{u} \cdot \nabla) F$ due to the sun is of order Ω_\odot/ω_r .¹⁶ This gives a situation like Eq. (2) with μn of order unity providing

a mechanism for producing side bands. This resolves a serious difficulty that the theory of satellite resonance of Mimas alone had, namely, that Mimas' gap is too small, by orders of magnitude, to account for the Cassini division. In the full problem, apparent gaps can be many orders of magnitude larger than the individual gaps.

Estimating the size of individual gaps (grooves) inside the large apparent rings is more delicate (as is the shift of the center of the Cassini division). In fact, low orders of perturbation theory give unphysically small gaps in this region. We would like to believe that this structure is related to the interplay of the various periods so that small denominators make low orders of perturbation theory suspect. If true, the structure of the rings away from moon resonances requires an understanding of the almost periodic Schrodinger equation beyond what we currently have.

In summary, we feel that a resonance picture can describe some of the observed qualitative features of the rings of Saturn although it remains to be seen whether further study of the almost periodic Hill operator will verify the intuition based on perturbation theory and whether, even if viable, nature chooses this mechanism. We emphasize that linear stability is a single particle theory and is difficult to reconcile if there are many collisions. A competing theory to the satellite resonance of Goldreich and Tremaine¹⁷ posits the gross ring structure as a collective phenomenon.

We thank J. Moser and R. Johnson for telling us of their work before publication, A. Dekel, L. Yaffe, and particularly P. Goldreich for useful discussions and criticism, I. Mikolic-Torreira for writing a computer program for us, and F. Dyson and P. Lax for encouragement. We would also like to thank the S. Fairchild Visiting Scholar Program and the California Institute of Technology Mathematics Department for their hospitality. This research is supported in part by the National Science Foundation under Grant No. MCS-78-01885.

^(a)On leave from Physics Department, Princeton University, Princeton, N. J. 08544.

^(b)On leave from Physics and Mathematics Departments, Princeton University, N. J. 08544. Sherman Fairchild Visiting Scholar at California Institute of Technology, 1980-81.

¹J. B. Pollack, *Space Sci. Rev.* **18**, 3 (1975); F. A. Franklin *et al.*, *Icarus* **12**, 338 (1970), and **15**, 80 (1971).

²See the discussion by V. Arnold, in *Classical Mechanics* (Springer, New York, 1978).

³We emphasize that we are not saying that the rings of Saturn have anything to do with quantum mechanics; only that an equation is the same so quantum mechanical intuition is useful.

⁴This is consistent with the number obtained by Franklin *et al.*, Ref. 1, by numerically integrating the restricted three-body problem.

⁵We examined two other candidates; precession of the quadrupole moment of Saturn due to Titan and precession of this moment due to the sun. The former has too small an amplitude because the angle of precession is exceedingly small (about 10^{-5} rad) and this angle enters quadratically; the latter has such a large period (about 10^{10} d) that it operates on a scale where a single particle theory must be inadequate.

⁶We emphasize that there is earlier important work on almost periodic Schrodinger operators; in addition to that of J. Moser (to be published), M. Ya Azbel [*Phys. Rev. Lett.* **43**, 1954 (1979); see also *Zh. Eksp. Teor. Fiz.* **44**, 980 (1963) [*Sov. Phys. JETP* **17**, 665 (1963)], and **46**, 929 (1964) [**19**, 634 (1964)]] and D. R. Hofstadter [*Phys. Rev. B* **14**, 2239 (1976)]; we mention B. A. Dubrovin *et al.*, *Russian Math. Surveys* **31**, 59 (1976); E. Dynaburg and Ja. G. Sinai, *Functional. Anal. Appl.* **9**, 8 (1975); S. Aubry, *Ann. Israel Phys. Soc.* **3**, 133 (1980); R. Johnson, to be published; J. Bellisard and D. Testard, to be published.

⁷Because of small divisors, perturbation theory diverges but it should be a useful guide.

⁸A typical limit periodic potential is $\sum a_n \cos(\kappa/2^n)$ with $\sum |a_n| < \infty$.

⁹See Moser, Ref. 6. While our work (J. Avron and B. Simon, to be published) is independent of Moser's, his certainly precedes ours by some months.

¹⁰See Avron and Simon, Ref. 9.

¹¹See Azbel, Ref. 6.

¹²See Hofstadter, Ref. 6.

¹³E. Harrell, to be published; see also J. Avron and B. Simon, to be published.

¹⁴Iapetus and Phoebe have large inclinations but their effect is much smaller than the five main moons.

¹⁵W. M. Smart, *Celestial Mechanics* (Longmans, Green, New York, 1953); D. Brower and G. M. Clemence, *Methods of Celestial Mechanics* (Academic, New York, 1961).

¹⁶Although the disturbance function is of order $(\Omega_{\odot}/\omega_r)^2$, it is known from Delaunay lunar theory that there are corrections to the orbit of order $\gamma(\Omega_{\odot}/\omega_r)$ with period $2\Omega_{\odot}$ where $\gamma = \sin I$, with I the inclination. $\gamma \sim 0.45$ in the present case. Consequently, one has to consider the full matrix valued Hill's operator.

¹⁷P. Goldreich and S. Tremaine, *Icarus* **34**, 227-239, 240-253 (1978).