

Appearances of regeneration from naked masses of protoplasm have sometimes been noted, but never in examples in which it had been established that there were no small, fully formed cells from which the growth might have arisen. It is, of course, possible that cultural conditions will be found under which masses of naked protoplasm will revert to cellular growth, such as has been reported by Nečas<sup>4</sup> in *Saccharomyces cerevisiae* and by Pease<sup>11</sup> in *Proteus vulgaris*.

<sup>1</sup> C. Weibull, *Symposium Soc. Gen. Microbiol.*, 6, 111, 1956.

<sup>2</sup> J. Lederberg, these PROCEEDINGS, 42, 574, 1956; K. McQuillen, *Biochim. et Biophys. Acta*, 27, 410, 1958.

<sup>3</sup> K. McQuillen, *Symposium Soc. Gen. Microbiol.*, 6, 127, 1956; S. Spiegelman, in *The Chemical Basis of Heredity*, ed. W. D. McElroy and B. Glass (Baltimore: Johns Hopkins Press, 1957), pp. 232-267; J. Spizizen, these PROCEEDINGS, 43, 694, 1957; D. Fraser, H. R. Mahler, A. L. Shug, and C. A. Thomas, Jr., these PROCEEDINGS, 43, 939, 1957; E. Chargaff, H. M. Schulman, and H. S. Shapiro, *Nature*, 180, 151, 1957.

<sup>4</sup> O. Nečas, *Biol. Zentr.*, 75, 268, 1956; A. A. Eddy and D. H. Williamson, *Nature*, 179, 1252, 1957.

<sup>5</sup> Mary R. Emerson, in manuscript. Our mutant strain, Em 11200, has been shown to be allelic to a number of others of independent origin: B 135 and M 16 obtained from Dr. D. D. Perkins, Stanford University; "wooly," from Dr. H. L. K. Whitehouse, Cambridge University; and "ginger" and an unnamed mutant from Dr. J. R. S. Fincham, University of Leicester. From published descriptions it is likely that these are also allelic to the "cut" mutant of Dr. H. Kuwana (*Cytologia*, 18, 235, 1953).

<sup>6</sup> Mary B. Mitchell and H. K. Mitchell, these PROCEEDINGS, 38, 442, 1952.

<sup>7</sup> Mary B. Mitchell, H. K. Mitchell, and A. Tissieres, these PROCEEDINGS, 39, 606, 1953.

<sup>8</sup> G. W. Beadle and E. L. Tatum, these PROCEEDINGS, 27, 499, 1941.

<sup>9</sup> The hemicellulase preparation was obtained from the Nutritional Biochemical Corporation, Cleveland 28, Ohio, who inform us that it is a concentrate of microbiological origin, containing cellulase, gumase, and maltase in addition to hemicellulase. A single test in our laboratory indicated strong chitinase activity.

<sup>10</sup> J. Lederberg and Jacqueline St. Clair, *J. Bacteriol.*, 75, 143, 1958.

<sup>11</sup> Phyllis Pease, *J. Gen. Microbiol.*, 17, 64, 1957.

## THE REPLICATION OF DNA IN *ESCHERICHIA COLI*\*

BY MATTHEW MESELSON AND FRANKLIN W. STAHL

GATES AND CRELLIN LABORATORIES OF CHEMISTRY,† AND NORMAN W. CHURCH LABORATORY OF CHEMICAL BIOLOGY, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CALIFORNIA

Communicated by Max Delbrück, May 14, 1958

*Introduction.*—Studies of bacterial transformation and bacteriophage infection<sup>1-5</sup> strongly indicate that deoxyribonucleic acid (DNA) can carry and transmit hereditary information and can direct its own replication. Hypotheses for the mechanism of DNA replication differ in the predictions they make concerning the distribution among progeny molecules of atoms derived from parental molecules.<sup>6</sup>

Radioisotopic labels have been employed in experiments bearing on the distribution of parental atoms among progeny molecules in several organisms.<sup>6-9</sup> We anticipated that a label which imparts to the DNA molecule an increased density might permit an analysis of this distribution by sedimentation techniques. To this end, a method was developed for the detection of small density differences among




FIG. 1.—Ultraviolet absorption photographs showing successive stages in the banding of DNA from *E. coli*. An aliquot of bacterial lysate containing approximately  $10^8$  lysed cells was centrifuged at 31,410 rpm in a CsCl solution as described in the text. Distance from the axis of rotation increases toward the right. The number beside each photograph gives the time elapsed after reaching 31,410 rpm.

macromolecules.<sup>10</sup> By use of this method, we have observed the distribution of the heavy nitrogen isotope  $N^{15}$  among molecules of DNA following the transfer of a uniformly  $N^{15}$ -labeled, exponentially growing bacterial population to a growth medium containing the ordinary nitrogen isotope  $N^{14}$ .

*Density-Gradient Centrifugation.*—A small amount of DNA in a concentrated solution of cesium chloride is centrifuged until equilibrium is closely approached.

The opposing processes of sedimentation and diffusion have then produced a stable concentration gradient of the cesium chloride. The concentration and pressure gradients result in a continuous increase of density along the direction of centrifugal force. The macromolecules of DNA present in this density gradient are driven by the centrifugal field into the region where the solution density is equal to their own buoyant density.<sup>11</sup> This concentrating tendency is opposed by diffusion, with the result that at equilibrium a single species of DNA is distributed over a band whose width is inversely related to the molecular weight of that species (Fig. 1).

If several different density species of DNA are present, each will form a band at the position where the density of the CsCl solution is equal to the buoyant density of that species. In this way DNA labeled with heavy nitrogen ( $N^{15}$ ) may be




FIG. 2—*a*: The resolution of  $N^{14}$  DNA from  $N^{15}$  DNA by density-gradient centrifugation. A mixture of  $N^{14}$  and  $N^{15}$  bacterial lysates, each containing about  $10^8$  lysed cells, was centrifuged in CsCl solution as described in the text. The photograph was taken after 24 hours of centrifugation at 44,770 rpm. *b*: A microdensitometer tracing showing the DNA distribution in the region of the two bands of Fig. 2*a*. The separation between the peaks corresponds to a difference in buoyant density of  $0.014 \text{ gm. cm.}^{-3}$

resolved from unlabeled DNA. Figure 2 shows the two bands formed as a result of centrifuging a mixture of approximately equal amounts of  $N^{14}$  and  $N^{15}$  *Escherichia coli* DNA.

In this paper reference will be made to the apparent molecular weight of DNA samples determined by means of density-gradient centrifugation. A discussion has been given<sup>10</sup> of the considerations upon which such determinations are based, as well as of several possible sources of error.<sup>12</sup>

*Experimental.*—*Escherichia coli* B was grown at  $36^\circ \text{C}$ . with aeration in a glucose salts medium containing ammonium chloride as the sole nitrogen source.<sup>13</sup> The growth of the bacterial population was followed by microscopic cell counts and by colony assays (Fig. 3).

Bacteria uniformly labeled with  $N^{15}$  were prepared by growing washed cells for

14 generations (to a titer of  $2 \times 10^8$ /ml) in medium containing  $100 \mu\text{g}/\text{ml}$  of  $\text{N}^{15}\text{H}_4\text{Cl}$  of 96.5 per cent isotopic purity. An abrupt change to  $\text{N}^{14}$  medium was then accomplished by adding to the growing culture a tenfold excess of  $\text{N}^{14}\text{H}_4\text{Cl}$ , along with ribosides of adenine and uracil in experiment 1 and ribosides of adenine, guanine, uracil, and cytosine in experiment 2, to give a concentration of  $10 \mu\text{g}/\text{ml}$  of each riboside. During subsequent growth the bacterial titer was kept between




FIG. 3.—Growth of bacterial populations first in  $\text{N}^{15}$  and then in  $\text{N}^{14}$  medium. The values on the ordinates give the actual titers of the cultures up to the time of addition of  $\text{N}^{14}$ . Thereafter, during the period when samples were being withdrawn for density-gradient centrifugation, the actual titer was kept between  $1$  and  $2 \times 10^8$  by additions of fresh medium. The values on the ordinates during this later period have been corrected for the withdrawals and additions. During the period of sampling for density-gradient centrifugation, the generation time was  $0.81$  hours in Experiment 1 and  $0.85$  hours in Experiment 2.

$1$  and  $2 \times 10^8$ /ml by appropriate additions of fresh  $\text{N}^{14}$  medium containing ribosides.

Samples containing about  $4 \times 10^9$  bacteria were withdrawn from the culture just before the addition of  $\text{N}^{14}$  and afterward at intervals for several generations. Each sample was immediately chilled and centrifuged in the cold for 5 minutes at  $1,800 \times g$ . After resuspension in  $0.40$  ml. of a cold solution  $0.01 M$  in  $\text{NaCl}$  and  $0.01 M$  in ethylenediaminetetra-acetate (EDTA) at pH 6, the cells were lysed by the addition of  $0.10$  ml. of 15 per cent sodium dodecyl sulfate and stored in the cold.




FIG. 4—*a*: Ultraviolet absorption photographs showing DNA bands resulting from density-gradient centrifugation of lysates of bacteria sampled at various times after the addition of an excess of  $N^{14}$  substrates to a growing  $N^{15}$ -labeled culture. Each photograph was taken after 20 hours of centrifugation at 44,770 rpm under the conditions described in the text. The density of the CsCl solution increases to the right. Regions of equal density occupy the same horizontal position on each photograph. The time of sampling is measured from the time of the addition of  $N^{14}$  in units of the generation time. The generation times for Experiments 1 and 2 were estimated from the measurements of bacterial growth presented in Fig. 3. *b*: Microdensitometer tracings of the DNA bands shown in the adjacent photographs. The microdensitometer pen displacement above the base line is directly proportional to the concentration of DNA. The degree of labeling of a species of DNA corresponds to the relative position of its band between the bands of fully labeled and unlabeled DNA shown in the lowermost frame, which serves as a density reference. A test of the conclusion that the DNA in the band of intermediate density is just half-labeled is provided by the frame showing the mixture of generations 0 and 1.9. When allowance is made for the relative amounts of DNA in the three peaks, the peak of intermediate density is found to be centered at  $50 \pm 2$  per cent of the distance between the  $N^{14}$  and  $N^{15}$  peaks.

For density-gradient centrifugation, 0.010 ml. of the dodecyl sulfate lysate was added to 0.70 ml. of CsCl solution buffered at pH 8.5 with 0.01 *M* tris(hydroxymethyl)aminomethane. The density of the resulting solution was 1.71 gm. cm.<sup>-3</sup> This was centrifuged at 140,000 × *g*. (44,770 rpm) in a Spinco model E ultracentrifuge at 25° for 20 hours, at which time the DNA had essentially attained sedimentation equilibrium. Bands of DNA were then found in the region of density 1.71 gm. cm.<sup>-3</sup>, well isolated from all other macromolecular components of the bacterial lysate. Ultraviolet absorption photographs taken during the course of each centrifugation were scanned with a recording microdensitometer (Fig. 4).

The buoyant density of a DNA molecule may be expected to vary directly with the fraction of N<sup>15</sup> label it contains. The density gradient is constant in the region between fully labeled and unlabeled DNA bands. Therefore, the degree of labeling of a partially labeled species of DNA may be determined directly from the relative position of its band between the band of fully labeled DNA and the band of unlabeled DNA. The error in this procedure for the determination of the degree of labeling is estimated to be about 2 per cent.

*Results.*—Figure 4 shows the results of density-gradient centrifugation of lysates of bacteria sampled at various times after the addition of an excess of N<sup>14</sup>-containing substrates to a growing N<sup>15</sup>-labeled culture.

It may be seen in Figure 4 that, until one generation time has elapsed, half-labeled molecules accumulate, while fully labeled DNA is depleted. One generation time after the addition of N<sup>14</sup>, these half-labeled or "hybrid" molecules alone are observed. Subsequently, only half-labeled DNA and completely unlabeled DNA are found. When two generation times have elapsed after the addition of N<sup>14</sup>, half-labeled and unlabeled DNA are present in equal amounts.

*Discussion.*—These results permit the following conclusions to be drawn regarding DNA replication under the conditions of the present experiment.

1. *The nitrogen of a DNA molecule is divided equally between two subunits which remain intact through many generations.*

The observation that parental nitrogen is found only in half-labeled molecules at all times after the passage of one generation time demonstrates the existence in each DNA molecule of two subunits containing equal amounts of nitrogen. The finding that at the second generation half-labeled and unlabeled molecules are found in equal amounts shows that the number of surviving parental subunits is twice the number of parent molecules initially present. That is, the subunits are conserved.

2. *Following replication, each daughter molecule has received one parental subunit.*

The finding that all DNA molecules are half-labeled one generation time after the addition of N<sup>14</sup> shows that each daughter molecule receives one parental subunit.<sup>14</sup> If the parental subunits had segregated in any other way among the daughter molecules, there would have been found at the first generation some fully labeled and some unlabeled DNA molecules, representing those daughters which received two or no parental subunits, respectively.

3. *The replicative act results in a molecular doubling.*

This statement is a corollary of conclusions 1 and 2 above, according to which each parent molecule passes on two subunits to progeny molecules and each progeny

molecule receives just one parental subunit. It follows that each single molecular reproductive act results in a doubling of the number of molecules entering into that act.

The above conclusions are represented schematically in Figure 5.

*The Watson-Crick Model.*—A molecular structure for DNA has been proposed by Watson and Crick.<sup>15</sup> It has undergone preliminary refinement<sup>16</sup> without alteration of its main features and is supported by physical and chemical studies.<sup>17</sup> The structure consists of two polynucleotide chains wound helically about a common axis. The nitrogen base (adenine, guanine, thymine, or cytosine) at each level




FIG. 5.—Schematic representation of the conclusions drawn in the text from the data presented in Fig. 4. The nitrogen of each DNA molecule is divided equally between two subunits. Following duplication, each daughter molecule receives one of these. The subunits are conserved through successive duplications.

on one chain is hydrogen-bonded to the base at the same level on the other chain. Structural requirements allow the occurrence of only the hydrogen-bonded base pairs adenine-thymine and guanine-cytosine, resulting in a detailed complementarity between the two chains. This suggested to Watson and Crick<sup>18</sup> a definite and structurally plausible hypothesis for the duplication of the DNA molecule. According to this idea, the two chains separate, exposing the hydrogen-bonding sites of the bases. Then, in accord with the base-pairing restrictions, each chain serves as a template for the synthesis of its complement. Accordingly, each daughter molecule contains one of the parental chains paired with a newly synthesized chain (Fig. 6).

The results of the present experiment are in exact accord with the expectations of the Watson-Crick model for DNA duplication. However, it must be emphasized that it has not been shown that the molecular subunits found in the present experiment are single polynucleotide chains or even that the DNA molecules studied here correspond to single DNA molecules possessing the structure proposed by Watson and Crick. However, some information has been obtained about the molecules and their subunits; it is summarized below.




FIG. 6.—Illustration of the mechanism of DNA duplication proposed by Watson and Crick. Each daughter molecule contains one of the parental chains (*black*) paired with one new chain (*white*). Upon continued duplication, the two original parent chains remain intact, so that there will always be found two molecules each with one parental chain.

The DNA molecules derived from *E. coli* by detergent-induced lysis have a buoyant density in CsCl of 1.71 gm. cm.<sup>-3</sup>, in the region of densities found for T2 and T4 bacteriophage DNA, and for purified calf-thymus and salmon-sperm DNA. A highly viscous and elastic solution of N<sup>14</sup> DNA was prepared from a dodecyl sulfate lysate of *E. coli* by the method of Simmons<sup>19</sup> followed by deproteinization with chloroform. Further purification was accomplished by two cycles of preparative density-gradient centrifugation in CsCl solution. This purified bacterial DNA was found to have the same buoyant density and apparent molecular weight,  $7 \times 10^6$ , as the DNA of the whole bacterial lysates (Figs. 7, 8).

*Heat Denaturation.*—It has been found that DNA from *E. coli* differs importantly from purified salmon-sperm DNA in its behavior upon heat denaturation.

Exposure to elevated temperatures is known to bring about an abrupt collapse of the relatively rigid and extended native DNA molecule and to make available for acid-base titration a large fraction of the functional groups presumed to be blocked by hydrogen-bond formation in the native structure.<sup>19, 20, 21, 22</sup> Rice and Doty<sup>22</sup> have reported that this collapse is not accompanied by a reduction in molecular weight as determined from light-scattering. These findings are corroborated by density-gradient centrifugation of salmon-sperm DNA.<sup>23</sup> When this material is




FIG. 7.—Microdensitometer tracing of an ultraviolet absorption photograph showing the optical density in the region of a band of  $N^{14}$  *E. coli* DNA at equilibrium. About 2  $\mu$ g. of DNA purified as described in the text was centrifuged at 31,410 rpm at 25° in 7.75 molal CsCl at pH 8.4. The density gradient is essentially constant over the region of the band and is 0.057 gm./cm.<sup>4</sup>. The position of the maximum indicates a buoyant density of 1.71 gm. cm.<sup>-3</sup>. In this tracing the optical density above the base line is directly proportional to the concentration of DNA in the rotating centrifuge cell. The concentration of DNA at the maximum is about 50  $\mu$ g./ml.




FIG. 8.—The square of the width of the band of Fig. 7 plotted against the logarithm of the relative concentration of DNA. The divisions along the abscissa set off intervals of 1 mm.<sup>2</sup>. In the absence of density heterogeneity, the slope at any point of such a plot is directly proportional to the weight average molecular weight of the DNA located at the corresponding position in the band. Linearity of this plot indicates monodispersity of the banded DNA. The value of the slope corresponds to an apparent molecular weight for the Cs·DNA salt of  $9.4 \times 10^6$ , corresponding to a molecular weight of  $7.1 \times 10^6$  for the sodium salt.

kept at 100° for 30 minutes either under the conditions employed by Rice and Doty or in the CsCl centrifuging medium, there results a density increase of 0.014 gm. cm.<sup>-3</sup> with no change in apparent molecular weight. The same results are obtained if the salmon-sperm DNA is pre-treated at pH 6 with EDTA and sodium dodecyl sulfate. Along with the density increase, heating brings about a sharp reduction in the time required for band formation in the CsCl gradient. In the absence of an increase in molecular weight, the decrease in banding time must be ascribed<sup>10</sup> to an increase in the diffusion coefficient, indicating an extensive collapse of the native structure.




FIG. 9.—The dissociation of the subunits of *E. coli* DNA upon heat denaturation. Each smooth curve connects points obtained by microdensitometry of an ultraviolet absorption photograph taken after 20 hours of centrifugation in CsCl solution at 44,770 rpm. The baseline density has been removed by subtraction. A: A mixture of heated and unheated  $N^{15}$  bacterial lysates. Heated lysate alone gives one band in the position indicated. Unheated lysate was added to this experiment for comparison. Heating has brought about a density increase of  $0.016 \text{ gm. cm.}^{-3}$  and a reduction of about half in the apparent molecular weight of the DNA. B: Heated lysate of  $N^{15}$  bacteria grown for one generation in  $N^{14}$  growth medium. Before heat denaturation, the hybrid DNA contained in this lysate forms only one band, as may be seen in Fig. 4. C: A mixture of heated  $N^{14}$  and heated  $N^{15}$  bacterial lysates. The density difference is  $0.015 \text{ gm. cm.}^{-3}$

The decrease in banding time and a density increase close to that found upon heating salmon-sperm DNA are observed (Fig. 9, A) when a bacterial lysate containing uniformly labeled  $N^{15}$  or  $N^{14}$  *E. coli* DNA is kept at  $100^\circ \text{ C.}$  for 30 minutes in the CsCl centrifuging medium; but the apparent molecular weight of

the heated bacterial DNA is reduced to approximately half that of the unheated material.

Half-labeled DNA contained in a detergent lysate of  $N^{15}$  *E. coli* cells grown for one generation in  $N^{14}$  medium was heated at  $100^{\circ}$  C. for 30 minutes in the CsCl centrifuging medium. This treatment results in the loss of the original half-labeled material and in the appearance in equal amounts of two new density species, each with approximately half the initial apparent molecular weight (Fig. 9, B). The density difference between the two species is  $0.015$  gm. cm.<sup>-3</sup>, close to the increment produced by the  $N^{15}$  labeling of the unheated DNA.

This behavior suggests that heating the hybrid molecule brings about the dissociation of the  $N^{15}$ -containing subunit from the  $N^{14}$  subunit. This possibility was tested by a density-gradient examination of a mixture of heated  $N^{15}$  DNA and heated  $N^{14}$  DNA (Fig. 9, C). The close resemblance between the products of heating hybrid DNA (Fig. 9 B) and the mixture of products obtained from heating  $N^{14}$  and  $N^{15}$  DNA separately (Fig. 9, C) leads to the conclusion that the two molecular subunits have indeed dissociated upon heating. Since the apparent molecular weight of the subunits so obtained is found to be close to half that of the intact molecule, it may be further concluded that the subunits of the DNA molecule which are conserved at duplication are single, continuous structures. The scheme for DNA duplication proposed by Delbrück<sup>24</sup> is thereby ruled out.

To recapitulate, both salmon-sperm and *E. coli* DNA heated under similar conditions collapse and undergo a similar density increase, but the salmon DNA retains its initial molecular weight, while the bacterial DNA dissociates into the two subunits which are conserved during duplication. These findings allow two different interpretations. On the one hand, if we assume that salmon DNA contains subunits analogous to those found in *E. coli* DNA, then we must suppose that the subunits of salmon DNA are bound together more tightly than those of the bacterial DNA. On the other hand, if we assume that the molecules of salmon DNA do not contain these subunits, then we must concede that the bacterial DNA molecule is a more complex structure than is the molecule of salmon DNA. The latter interpretation challenges the sufficiency of the Watson-Crick DNA model to explain the observed distribution of parental nitrogen atoms among progeny molecules.

*Conclusion.*—The structure for DNA proposed by Watson and Crick brought forth a number of proposals as to how such a molecule might replicate. These proposals<sup>6</sup> make specific predictions concerning the distribution of parental atoms among progeny molecules. The results presented here give a detailed answer to the question of this distribution and simultaneously direct our attention to other problems whose solution must be the next step in progress toward a complete understanding of the molecular basis of DNA duplication. What are the molecular structures of the subunits of *E. coli* DNA which are passed on intact to each daughter molecule? What is the relationship of these subunits to each other in a DNA molecule? What is the mechanism of the synthesis and dissociation of the subunits in vivo?

*Summary.*—By means of density-gradient centrifugation, we have observed the distribution of  $N^{15}$  among molecules of bacterial DNA following the transfer of a uniformly  $N^{15}$ -substituted exponentially growing *E. coli* population to  $N^{14}$  medium.

We find that the nitrogen of a DNA molecule is divided equally between two physically continuous subunits; that, following duplication, each daughter molecule receives one of these; and that the subunits are conserved through many duplications.

\* Aided by grants from the National Foundation for Infantile Paralysis and the National Institutes of Health.

† Contribution No. 2344.

<sup>1</sup> R. D. Hotchkiss, in *The Nucleic Acids*, ed. E. Chargaff and J. N. Davidson (New York: Academic Press, 1955), p. 435; and in *Enzymes: Units of Biological Structure and Function*, ed. O. H. Gaebler (New York: Academic Press, 1956), p. 119.

<sup>2</sup> S. H. Goodgal and R. M. Herriott, in *The Chemical Basis of Heredity*, ed. W. D. McElroy and B. Glass (Baltimore: Johns Hopkins Press, 1957), p. 336.

<sup>3</sup> S. Zamenhof, in *The Chemical Basis of Heredity*, ed. W. D. McElroy and B. Glass (Baltimore: Johns Hopkins Press, 1957), p. 351.

<sup>4</sup> A. D. Hershey and M. Chase, *J. Gen. Physiol.*, **36**, 39, 1952.

<sup>5</sup> A. D. Hershey, *Virology*, **1**, 108, 1955; **4**, 237, 1957.

<sup>6</sup> M. Delbrück and G. S. Stent, in *The Chemical Basis of Heredity*, ed. W. D. McElroy and B. Glass (Baltimore: Johns Hopkins Press, 1957), p. 699.

<sup>7</sup> C. Levinthal, these PROCEEDINGS, **42**, 394, 1956.

<sup>8</sup> J. H. Taylor, P. S. Woods, and W. L. Hughes, these PROCEEDINGS, **43**, 122, 1957.

<sup>9</sup> R. B. Painter, F. Forro, Jr., and W. L. Hughes, *Nature*, **181**, 328, 1958.

<sup>10</sup> M. S. Meselson, F. W. Stahl, and J. Vinograd, these PROCEEDINGS, **43**, 581, 1957.

<sup>11</sup> The buoyant density of a molecule is the density of the solution at the position in the centrifuge cell where the sum of the forces acting on the molecule is zero.

<sup>12</sup> Our attention has been called by Professor H. K. Schachman to a source of error in apparent molecular weights determined by density-gradient centrifugation which was not discussed by Meselson, Stahl, and Vinograd. In evaluating the dependence of the free energy of the DNA component upon the concentration of CsCl, the effect of solvation was neglected. It can be shown that solvation may introduce an error into the apparent molecular weight if either CsCl or water is bound preferentially. A method for estimating the error due to such selective solvation will be presented elsewhere.

<sup>13</sup> In addition to NH<sub>4</sub>Cl, this medium consists of 0.049 M Na<sub>2</sub>HPO<sub>4</sub>, 0.022 M KH<sub>2</sub>PO<sub>4</sub>, 0.05 M NaCl, 0.01 M glucose, 10<sup>-3</sup> M MgSO<sub>4</sub>, and 3 × 10<sup>-6</sup> M FeCl<sub>3</sub>.

<sup>14</sup> This result also shows that the generation time is very nearly the same for all DNA molecules in the population. This raises the questions of whether in any one nucleus all DNA molecules are controlled by the same clock and, if so, whether this clock regulates nuclear and cellular division as well.

<sup>15</sup> F. H. C. Crick and J. D. Watson, *Proc. Roy. Soc. London, A*, **223**, 80, 1954.

<sup>16</sup> R. Langridge, W. E. Seeds, H. R. Wilson, C. W. Hooper, M. H. F. Wilkins, and L. D. Hamilton, *J. Biophys. and Biochem. Cytol.*, **3**, 767, 1957.

<sup>17</sup> For reviews see D. O. Jordan, in *The Nucleic Acids*, ed. E. Chargaff and J. D. Davidson (New York: Academic Press, 1955), **1**, 447; and F. H. C. Crick, in *The Chemical Basis of Heredity*, ed. W. D. McElroy and B. Glass (Baltimore: Johns Hopkins Press, 1957), p. 532.

<sup>18</sup> J. D. Watson and F. H. C. Crick, *Nature*, **171**, 964, 1953.

<sup>19</sup> C. E. Hall and M. Litt, *J. Biophys. and Biochem. Cytol.*, **4**, 1, 1958.

<sup>20</sup> R. Thomas, *Biochim. et Biophys. Acta*, **14**, 231, 1954.

<sup>21</sup> P. D. Lawley, *Biochim. et Biophys. Acta*, **21**, 481, 1956.

<sup>22</sup> S. A. Rice and P. Doty, *J. Am. Chem. Soc.*, **79**, 3937, 1957.

<sup>23</sup> Kindly supplied by Dr. Michael Litt. The preparation of this DNA is described by Hall and Litt (*J. Biophys. and Biochem. Cytol.*, **4**, 1, 1958).

<sup>24</sup> M. Delbrück, these PROCEEDINGS, **40**, 783, 1955.