

KROTZ 12577-12578

Terms & Conditions

Electronic Supporting Information files are available without a subscription to ACS Web Editions. The American Chemical Society holds a copyright ownership interest in any copyrightable Supporting Information. Files available from the ACS website may be downloaded for personal use only. Users are not otherwise permitted to reproduce, republish, redistribute, or sell any Supporting Information from the ACS website, either in whole or in part, in either machine-readable form or any other form without permission from the American Chemical Society. For permission to reproduce, republish and redistribute this material, requesters must process their own requests via the RightsLink permission system. Information about how to use the RightsLink permission system can be found at <http://pubs.acs.org/page/copyright/permissions.html>.



J-12578-1m1

Supplementary Material

**Assembly of DNA Recognition Elements on an Octahedral Rhodium Intercalator:
Predictive Recognition of 5'-TGCA-3' by Δ -[Rh[(R,R)-Me₂trien]phi]³⁺**

Achim H. Krotz, Brian P. Hudson, and Jacqueline K. Barton*
Division of Chemistry and Chemical Engineering
California Institute of Technology
Pasadena, CA 91125

Figure 1S. Autoradiogram of an 8% denaturing polyacrylamide gel after photocleavage of the 5'-[³²P]-labeled EcoRV/EcoO109I fragment of SV40 DNA: lanes 1 and 2, Maxam-Gilbert A+G and C+T reactions, respectively; lane 3, untreated fragment; lane 4, fragment irradiated in the absence of rhodium complex; lanes 5-10, fragment irradiated in the presence of Δ -[Rh(en)₂phi]³⁺, Δ -R, Δ -S, Λ -[Rh(en)₂phi]³⁺, Λ -S, and Λ -R, respectively. Part of the sequence containing preferred cleavage sites is shown to the right. Strong sites of cleavage for Δ -R correspond to sequences containing ≥ 3 bases within the sequence 5'-TGCA-3'. Irradiations were conducted using a HeCd laser at 325 nm for 20 min. Samples contained 100 nM rhodium and 1 μ M base pairs in 5 mM Tris, 50 mM NaCl, pH 7.0.

