

Supplementary Materials for

Stepwise phosphorylation of leukotriene B₄ receptor 1 defines cellular responses to leukotriene B₄

Yoshimitsu Nakanishi, Modong Tan, Takako Ichiki, Asuka Inoue, Jun-ichi Yoshihara, Naoto Maekawa, Itsuki Takenoshita, Keisuke Yanagida, Shinya Yamahira, Satoshi Yamaguchi, Junken Aoki, Teruyuki Nagamune, Takehiko Yokomizo, Takao Shimizu, Motonao Nakamura*

*Corresponding author. Email: moto-nakamura@dls.ous.ac.jp

Published 21 August 2018, *Sci. Signal.* **11**, eaao5390 (2018)
DOI: 10.1126/scisignal.aao5390

This PDF file includes:

- Fig. S1. Properties of N-glycosylation-deficient HA-BLT1/0N.
- Fig. S2. Conserved Ser and Thr residues in the cytoplasmic domains of human, mouse, rat, guinea pig, and zebrafish BLT1.
- Fig. S3. Confirmation of residues essential for HA-BLT1/0N phosphorylation.
- Fig. S4. Phosphorylation of mouse BLT1.
- Fig. S5. Phosphorylation at LTB₄-induced and basal sites through G_i.
- Fig. S6. LTB₄ dose dependency of intracellular [Ca²⁺] increase.
- Fig. S7. Effect of blockage of the [Ca²⁺] increase on BLT1 phosphorylation.
- Fig. S8. Effect of phosphorylation deficiency on the functions of wild-type BLT1.
- Fig. S9. Effect of helix 8 disruption on BLT1 phosphorylation.
- Fig. S10. Importance of phosphorylation for the ligand sensitivity of BLT1.
- Fig. S11. Effect of phosphorylation on β-arrestin binding to BLT1.
- Table S1. Primer sequences used to generate mutant BLT1s.




Fig. S1. Properties of N-glycosylation-deficient HA-BLT1/0N.

HeLa cells were transfected with the indicated receptors. **(A)** Membrane fractions were treated with PNGase-F and subjected to SDS-PAGE. Receptors were detected by immunoblotting (IB) using an antibody recognizing the HA tag. **(B)** Cells were stained with the HA antibody followed by PE-conjugated secondary antibody, and HA-positive cells were sorted. **(C)** Intracellular Ca^{2+} mobilizations elicited by LTB_4 were evaluated in HeLa cells expressing the indicated receptors. All data are representative of at least three independent experiments.




Fig. S3. Confirmation of residues essential for HA-BLT1/0N phosphorylation.

The human BLT1 mutants deficient in the basal (Δ b-phos), LTB₄-induced (Δ i-phos), or both basal and LTB₄-induced (Δ phos) phosphorylations were constructed as indicated in Fig. 2D. (A) HeLa cells expressing the indicated receptors were stimulated with 100 nM LTB₄ for 40 min. Membrane fractions were subjected to Phos-tag SDS-PAGE and SDS-PAGE. Receptors were detected by immunoblotting (IB) for HA tag. (B) Similar experiments as in (A) were carried out in CHO-K1, Ba/F3, and RBL-2H3 cells. All data are representative of at least three independent experiments.




Fig. S4. Phosphorylation of mouse BLT1.

(A) The indicated mutants of HA-tagged mouse BLT1 were constructed to determine the phosphorylated residues. Ser and Thr residues conserved between human and mouse BLT1 are shown in blue. (B) HeLa cells expressing these mutants were stimulated with or without 100 nM LTB₄ for 20 min. The membrane fractions were subjected to Phos-tag SDS-PAGE and immunoblotting for HA. All data are representative of at least three independent experiments.




Fig. S5. Phosphorylation at LTB₄-induced and basal sites through G_i.

(A) HeLa cells expressing the indicated HA-tagged receptors were stained with the antibody recognizing HA followed by the PE-conjugated secondary antibody, and HA-positive cells were sorted. (B) Schematic model of the signaling pathways operating through GPCRs activated by chemoattractants. Although the stimulation of G_i is a prerequisite, activation of both the ERK1/2 and PI3K-AKT pathways is required for chemotaxis, whereas activation of only the PI3K-AKT pathway is necessary for degranulation. (C) HeLa cells expressing HA-BLT1/0N were stimulated as indicated. For PTX treatment, cells were preincubated with 100 ng/ml PTX overnight. Membrane fractions of these cells were subjected to Phos-tag SDS-PAGE and immunoblotting (IB) for HA. All data are representative of at least three independent experiments.




Fig. S6. LTB₄ dose dependency of intracellular [Ca²⁺]_i increase.

Intracellular Ca²⁺ mobilizations elicited by LTB₄ were evaluated in CHO-K1 cells expressing the indicated receptors. For EGTA treatment, the medium was replaced with Ca²⁺-free medium containing 1 mM EGTA prior to LTB₄ application. Data are means ± SEM, *n* = 3 experiments. **P* < 0.05, ***P* < 0.01 by two-way ANOVA followed by Tukey's *post hoc* test. All data are representative of at least three independent experiments.




Fig. S7. Effect of blockage of the $[Ca^{2+}]$ increase on BLT1 phosphorylation.

HeLa cells expressing HA-BLT1/0N were stimulated with 100 nM LTB₄ for 20 min. For EGTA treatment, the medium was replaced with Ca²⁺-free medium containing 1 mM EGTA prior to 100 nM LTB₄ application. For SKF96365 treatment, cells were preincubated with 100 μM SKF96365 for 10 min before LTB₄ application. Membrane fractions were subjected to Phos-tag SDS-PAGE and immunoblotting for HA. All data are representative of three independent experiments.




Fig. S8. Effect of phosphorylation deficiency on the functions of wild-type BLT1.

(A) HeLa cells expressing HA-BLT1/WT or HA-BLT1/WT Δ phos were stained for HA, and HA-positive cells were sorted. (B) Biotinylated HA-BLT1/WT or HA-BLT1/WT Δ phos on the surface of HeLa cells was stimulated with 10 nM LTB₄ for the indicated time. The biotinylated BLT1 molecules were collected by Streptavidin pull-down. Receptors were detected by immunoblotting for HA. (C) LTB₄-induced intracellular Ca²⁺ mobilizations were examined in CHO-K1 cells expressing HA-BLT1/WT or HA-BLT1/WT Δ phos. (D) Chemotaxis and chemokinesis elicited by 10 nM and 100 nM LTB₄ were evaluated in CHO-K1 cells expressing HA-BLT1/WT or HA-BLT1/WT Δ phos. Data are means \pm SEM, $n = 5$ experiments. NS, not significant; ** $P < 0.01$ by two-way ANOVA followed by Tukey's *post hoc* test. (E) Release of β -hexosaminidase in RBL-2H3 cells expressing the indicated receptors was quantified. Column 1, vehicle; column 2, 10 nM LTB₄ for 40 min; column 3, 10 nM LTB₄ for 20 min then addition of 100 nM LTB₄ for 20 min; column 4, A23187 for 20 min. Data are means \pm SEM, $n = 3$ experiments. NS, not significant; *** $P < 0.001$ by one-way ANOVA followed by Tukey's *post hoc* test. All data are representative of at least three independent experiments.




Fig. S9. Effect of helix 8 disruption on BLT1 phosphorylation.

(A) Two HA-BLT1/ON mutants were constructed to examine the importance of helix 8 in the phosphorylation of BLT1 (25). Putative transmembrane domain-7 and helix 8 are labeled as TM7 and Helix-8, respectively. The 5 basal phosphorylation and 2 LTB₄-induced phosphorylation sites are boxed. (B) HeLa cells expressing the indicated receptors were treated with 100 nM LTB₄ for 40 min. Membrane fractions were subjected to Phos-tag SDS-PAGE. Receptors were detected by Western blotting for HA. All data are representative of at least three independent experiments.




Fig. S10. Importance of phosphorylation for the ligand sensitivity of BLT1.

(A) HEK293T cells expressing SRE-NanoLuc, PGK-firefly luciferase and the indicated receptors were treated with 10 nM or 100 nM LTB₄. The ratios of NanoLuc to firefly luciferase activity are shown. Data are means ± SEM, $n = 3$ experiments. NS, not significant; $*P < 0.05$, $***P < 0.001$ by one-way ANOVA followed by Tukey's *post hoc* test. (B) HEK293T cells expressing AP-TGFα, Gα_{q/11} and the indicated receptors were stimulated under various concentrations of LTB₄, and the AP-TGFα release into the medium was determined. Data are means ± SEM, $n = 3$ experiments. $**P < 0.01$, $***P < 0.001$ by two-way ANOVA followed by Tukey's *post hoc* test.




Fig. S11. Effect of phosphorylation on β -arrestin binding to BLT1.

LTB₄-induced β -arrestin recruitment to BLT1 was assessed by using a NanoBiT system. Luminescent signals in HEK293 cells expressing LgBiT- β arr1 or LgBiT- β arr2 together with BLT1/0N or BLT1/0N Δ Phos were measured before and after stimulation with LTB₄. β -arrestin recruitment was expressed as fold-change luminescence signals. Data are means \pm SEM. $n = 4$ experiments. ** $P < 0.01$, *** $P < 0.001$ by Student's t -test.

Table S1. Primer sequences used to generate mutant BLT1s.

| Mutant | Forward primer | Reverse primer |
|-----------------------|--|---|
| 1N(N2A) | gactacgccgccaactacatcttctg | cagaagatgtagtggcggcgtagtc |
| 1N(N164A) | ccctggaaaacggccatgagcctgtgc | gcacaggctcatggccgtttccagg |
| iL-1 | cagaagcgcgctgtcgtgccctg | cagggcagcgcagcgcgcttctg |
| iL-2A | ctagaccgcgactggcggtg | caccgccagtgcgcggtctag |
| iL-2B | ctttgtggcccagaagctacgcgccaaggcg | cgccttggcgcgtagcttctgggccacaag |
| iL-3A | gctgtggtggccgctacgcggacataggg | ccctatgtccgcgtagggccaccacagc |
| iL-3B | ctfccgcgcgcccgcgcggccgc | gcgccggcgcggcgccggcgccggaag |
| CT-1 | ctgctgcgcgcccggcggtgggc | gcccacgcccgcgcgcagcag |
| CT-2 | ctggagggcgcggcgccgagggctcc | ggacgcctcggcggccgcgcctccag |
| CT-3 | ccgagggcggcccgcgcgcccgggg | cccgcggcgcggcgccgctcgg |
| CT-4 | gggggcgccctgggccagccgctagg | cctagcggcctggcccaggcgccccc |
| CT-5 | cccggccctgccgagccctcactgcc | ggcagtgagggcctcggcagggccggg |
| 0N/S313 | ccgagggcgtccgcgcgcccgggg | cccgcggcgcggcgccgacgctcgg |
| 0N/S314 | ccgagggcggccagcgcgcgc | gcgcgcgctggccgctcgg |
| 0N/T315 | ccgagggcggcccacgcgcccgggg | cccgcggcgcgctggcgccgctcgg |
| 0N/S320 | gggggcagcctgggccagccgctagg | cctagcggcctggcccaggctgcccc |
| 0N/T324 | gggggcgccctgggccagaccgctagg | cctagcggcttggcccaggcgccccc |
| Δ b-phos | ctgggccagccgctaggagc | gctcctagcggcctggcccag |
| 0N/S310 | ctggagggcgcgggctccgagggctcc | ggacgcctcggagcccgcgcctccag |
| 0N/T308 | ctggagggcacggcgccgagggctcc | ggacgcctcggcggccgctcctccag |
| Δ i-phos | ctggagggcgcgggctccgagggctcc | ggacgcctcggagcccgcgcctccag |
| Δ phos | ctggagggcgcggcgccgagggcgcc | ggccgctcggcggcccgcgcctccag |
| pMK-HA | gtgccagactacccatggccaactacatcttct | gagggcggatccta |
| BLT1/0N | ggcgtagtctggcactg | taggatccgccctctc |
| mBLT | atggctgcagccaactacatctcctgcagc | tcacttcgaagactcaggaatggtgg |
| mBLT/0N | cagtaaaatggaacgccaggactctgatc | gatcagagtctggcgttccatttactg |
| mBLT/ Δ b-phos | ctggagggcactggctcggaggtggccgcccgc cgccgccccggcgctctgtgtccaggccccgaag gac | gtccttcggggcctggaccagagcgcccccgcg ggggcggcgccaccctccgagccagtgcct ccag |

| | | |
|-----------------------|---|--|
| mBLT/ Δ i-phos | ctggagggcgctggcgcggaggtgtccagacc cgccgcgggggactctggtccagaccccgaag gac | gtccttcggggcttgaccagagtgccccgcg gcgggtgctggacacctccgcgccagcgcctc cag |
| mBLT/ Δ phos | ctggagggcgctggcgcggaggtggccgcgcg cgccgcgggggctctggtccagggcccgaag gac | gtccttcggggccttgaccagagcccccgcg gcgggcggcggccacctccgcgccagcgcct ccag |
| mS313 | ggctcggaggtgtccgcccccgcggggg | ccccgcggcggcggcgggacacctccgagcc |
| mS314 | ggctcggaggtggccagcggcccccgcggggg | ccccgcggcggcggcgtggccacctccgagcc |
| mT315 | ggctcggaggtggccgccaccgcggggg | ccccgcggcgggtggcggccacctccgagcc |
| mS320 | cgccgcgggggactctggtccaggcc | ggcctggaccagagtgccccgcggcg |
| mT324 | cgctctggtccagaccccgaaggac | gtccttcggggcttgaccagagcg |
| mT308 | ggagggcactggcgcggaggtgtccagacc | ggtgctggacacctccgcgccagtgcctcc |
| mS310 | ctggagggcgctggctcggaggtgtcc | ggacacctccgagccagcgcctccag |