

Demosponge steroid biomarker 26-methylstigmastane provides evidence for Neoproterozoic animals

J. Alex Zumberge¹, Gordon D. Love^{1*}, Paco Cárdenas², Erik A. Sperling³, Sunithi Gunasekera², Megan Rohrssen⁴, Emmanuelle Grosjean⁵, John P. Grotzinger⁶ and Roger E. Summons⁷

Sterane biomarkers preserved in ancient sedimentary rocks hold promise for tracking the diversification and ecological expansion of eukaryotes. The earliest proposed animal biomarkers from demosponges (Demospongiae) are recorded in a sequence around 100 Myr long of Neoproterozoic–Cambrian marine sedimentary strata from the Huqf Supergroup, South Oman Salt Basin. This C₃₀ sterane biomarker, informally known as 24-isopropylcholestane (24-ipc), possesses the same carbon skeleton as sterols found in some modern-day demosponges. However, this evidence is controversial because 24-ipc is not exclusive to demosponges since 24-ipc sterols are found in trace amounts in some pelagophyte algae. Here, we report a new fossil sterane biomarker that co-occurs with 24-ipc in a suite of late Neoproterozoic–Cambrian sedimentary rocks and oils, which possesses a rare hydrocarbon skeleton that is uniquely found within extant demosponge taxa. This sterane is informally designated as 26-methylstigmastane (26-mes), reflecting the very unusual methylation at the terminus of the steroid side chain. It is the first animal-specific sterane marker detected in the geological record that can be unambiguously linked to precursor sterols only reported from extant demosponges. These new findings strongly suggest that demosponges, and hence multicellular animals, were prominent in some late Neoproterozoic marine environments at least extending back to the Cryogenian period.

The transition from unicellular protists to multicellular animals constitutes one of the most intriguing and enigmatic events in the evolutionary history of life, largely due to the absence of unambiguous physical fossils for the earliest fauna. The Neoproterozoic rise of eukaryotes¹, including demosponges², in marine environments can be discerned from lipid biomarker records preserved in ancient sedimentary rocks that have experienced a mild thermal history. Molecular phylogenies commonly show that sponges (Porifera) are the sister group of other animals³ and molecular evidence for Neoproterozoic animal life was first proposed based on the occurrence of unusual C₃₀ demosponge-derived steranes informally known as 24-isopropylcholestane (24-ipc) steranes in sedimentary rocks and oils of that age^{2,4}. These steranes are the hydrocarbon remains of 24-isopropylcholesterols and structurally related sterols⁵. The record of 24-ipc steranes commences in Cryogenian-aged sediments in South Oman (around 717–635 million years ago (Ma)^{2,6}) and then occurs continuously through the Ediacaran–Cambrian formations of the Huqf Supergroup of the South Oman Salt Basin. Notably, these steroids also occur as covalently bound constituents fixed within the immobile kerogen phase of the same rocks, which is an important confirmation that these are not younger contaminant compounds that migrated into the rocks².

Demosponges are the only known extant taxon that can biosynthesize 24-ipc precursors as their major sterols. High relative absolute abundances of 24-ipc steranes have now been reported in many other late Neoproterozoic–early Cambrian rocks and oils^{2,4,7–9}. These 24-ipc occurrences—if interpreted correctly—reflect an early

presence of Porifera and provide a conservative minimum time estimate for the origin of animal multicellularity and the sponge body plan. Others have hypothesized that the 24-ipc steranes could be derived from unicellular animal ancestors or have an algal origin¹⁰ since the parent sterols have been reported in trace amounts in some extant pelagophyte algae². The claim that poribacterial sponge symbionts from the candidate phylum *Poribacteria* can make 24-ipc steroids¹¹ has since been shown to be erroneous due to a genome assembly error^{12,13}.

Currently, two chromatographically resolvable series of ancient C₃₀ steranes are known: 24-*n*-propylcholestane (24-npc) and 24-ipc. Demosponges are the most plausible Neoproterozoic–Cambrian source of 24-npc as well as 24-ipc because both are produced by extant demosponges². Foraminifera are another possible source of 24-npc¹⁴. Pelagophyte algae probably account for the 24-npc steranes that are found in Devonian and younger marine sediments and their derived oils¹³. Various recent findings support a pre-Ediacaran origin of animals and sponges, and arguably reinforce the validity of the 24-ipc biomarker record, including: (1) steroid assays and genomic analyses of extant taxa¹³, which suggest that sponges were the most likely Neoproterozoic source biota for 24-ipc steranes; and (2) nuclear and mitochondrial gene molecular clock studies, which consistently support a pre-Ediacaran origin of animals and Neoproterozoic demosponges^{15–18}. The discovery of other sponge biomarkers to augment the 24-ipc sterane record would greatly strengthen evidence for the presence of animals before the appearance of the Ediacara fauna, since the efficacy of the standalone

¹Department of Earth Sciences, University of California, Riverside, Riverside, CA, USA. ²Department of Medicinal Chemistry, Uppsala University, Uppsala, Sweden. ³Department of Geological Sciences, Stanford University, Stanford, CA, USA. ⁴Department of Earth and Atmospheric Sciences, Central Michigan University, Mount Pleasant, MI, USA. ⁵Geoscience Australia, Canberra, Australian Capital Territory, Australia. ⁶Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA. ⁷Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA, USA. *e-mail: glove@ucr.edu

24-ipc sterane record for tracking early demosponges has been contested^{10,12}.

Results and discussion

Here, we report the presence of a new C_{30} sterane designated 26-methylstigmastane (26-mes) in a suite of Neoproterozoic–Cambrian rocks and oils (Fig. 1 and Supplementary Tables 1–3). Furthermore, we attribute this biomarker to demosponges since these are the only known organisms among extant taxa to produce sterols with the same carbon skeleton (Supplementary Information). The abundance of 26-mes sterane biomarkers is of comparable magnitude to 24-ipc and 24-npc (Fig. 1), although the relative proportions of the three main C_{30} sterane compounds can vary from sample to sample (Fig. 1 and Supplementary Tables 1 and 2). Summed C_{30} steranes are typically 1–4% of the total C_{27-30} sterane signal in South Oman rocks, although higher contents >5% can also be found. Our analyses confirm the presence of 26-mes along with 24-ipc steranes in the Neoproterozoic–Cambrian rock extracts and kerogen pyrolysates from South Oman reported previously^{2,8}, as well in Ediacaran–Cambrian-sourced oils from Eastern Siberia⁹ and India⁷, for which representative samples are shown in Fig. 1 (see also Supplementary Table 3). When 26-mes is detectable in Cryogenian to Cambrian age rocks and oils, it is found alongside both the 24-ipc and 24-npc sterane compounds. These three different sterane series constitute only a small subset of all the structural possibilities for C_{30} sterane compounds, which are feasible from adding three additional carbons to a cholestane side chain, and they correspond with three of the most commonly occurring sterane skeletons for C_{30} sterols found in extant demosponges (Fig. 1 and Supplementary Tables 4 and 5). In contrast, 26-mes abundance is typically lower or absent for the small suite of Phanerozoic oils and rocks analysed thus far but can be detected, along with 24-ipc and 24-npc, in some samples but not in the procedural blanks (Supplementary Table 3).

To unequivocally confirm the assignment of the newly identified ancient sterane series as 26-mes, we compared the C_{30} sterane distributions of Neoproterozoic rocks and oils with sterane products derived from sterols of modern sponges comprising demosponges, hexactinellids, homoscleromorphs and calcisponges (Supplementary Tables 4 and 5). We applied catalytic hydroxylation (HyPy)—a mild reductive technique employing high-pressure hydrogen—to transform sterols from sponge biomass into steranes with minimal structural and stereochemical disturbance¹⁹. Only three compatible parent C_{30} sterols, with an identical side chain skeleton to 26-mes, are currently known in extant taxa (Fig. 2), and these were the probable precursors to the sedimentary 26-mes described above (Supplementary Information). *Rhabdastrella globostellata* was used as a model sponge species for initial investigations since its sterols have been previously well characterized^{20,21} and it contains stelliferasterol as the major C_{30} sterol constituent, which was verified for our specimens (Supplementary Fig. 1). We generated a simple C_{30} sterane distribution as expected from the HyPy conversion of the *R. globostellata* sterols, dominated by 26-mes stereoisomers (Supplementary Fig. 2), and these products were used as a sterane standard to unequivocally test for presence or absence of the 26-mes biomarker in modern and ancient samples (Fig. 1). The identification of the fossil 26-mes sterane series was verified by observing co-elution of the $5\alpha,14\alpha,17\alpha(H)$ -20R stereoisomer with the same isomer produced from *R. globostellata* and other extant demosponges (Fig. 1). This co-elution was further confirmed using two different gas chromatography–mass spectrometry (GC-MS) techniques in two different laboratories using different gas chromatography column stationary phases (see Methods).

To better constrain the taxonomic distribution of 26-mes, we supplemented literature reports with targeted analyses of extant




Fig. 1 | MRM-GC-MS ion chromatograms of C_{30} sterane distributions

(414 → 217 Da ion transitions). **a**, Results from Neoproterozoic–Cambrian rock bitumens and oils. **b**, HyPy products from the cells of three modern demosponges (see Supplementary Table 4 for taxonomic assignments). Ancient samples, having undergone protracted burial and alteration, exhibit a more complex distribution of diastereoisomers compared with modern sponge biomass. Four regular sterane diastereoisomers can be found in ancient samples of oil window-maturity ($\alpha\alpha\alpha S$, $\alpha\beta\beta R$, $\alpha\beta\beta S$ and $\alpha\alpha\alpha R$), while two diastereoisomers ($\beta\alpha\alpha R$ and $\alpha\alpha\alpha R$) result from laboratory hydrogenation of individual Δ^5 -sterols in modern sponge biomass. The signal peak for the $\alpha\alpha\alpha S$ geoisomer of 26-mes often co-elutes with other C_{30} steranes, so this isomer peak is usually obscured in chromatograms. Direct correlation with modern sponges uses the $\alpha\alpha\alpha R$ isomer, as shown by the dashed lines. The $\alpha\beta\beta$ isomers show expected enhancement of the signal in 414 → 218 Da ion chromatograms relative to $\alpha\alpha\alpha$ stereoisomers (not shown here). Examples from the Proterozoic rock record show three distinct sterane series co-occurring (24-npc, 24-ipc and 26-mes). The rock from Ghadir Manquill Formation, South Oman, was deposited during the Cryogenian period (probably around 660–635 Ma) and is the oldest example in the rock record known with 24-ipc and 26-mes co-occurring. The 26-mes sterane biomarker was detected in significant amounts in the South Oman rock extracts and kerogen pyrolysates reported previously² (Supplementary Tables 1 and 2). The oil from the Usol'ye Formation, Eastern Siberia, is probably Ediacaran to Early Cambrian in source age⁹, as is the Baghewala-1 oil from India⁷. An 'o' symbol represents 24-npc, a plus 24-ipc and an asterisk 26-mes. Thymo, thymosiosterane (24,26,26'-trimethylcholestane).




Fig. 2 | Chemical structures of stelliferasterol, isostelliferasterol and strongylosterol. These are the three known natural sterol precursors of the 26-mes sterane biomarker^{20,21,23,39,40}. They are found only in certain demosponges, but not detected in other groups of eukaryotes. Note: (1) the methyl-substituent at the terminal position of the sterol side chain, which remains preserved at C-26 in 26-mes; and (2) the unusual double bond positions in the side chains of stelliferasterol and strongylosterol. The biological configuration is 20R for all three sterols, 24R for strongylosterol and stelliferasterol, and 25S for isostelliferasterol. Three stereogenic carbon atoms exist in the side chain of 26-mes (chirality at C-20, C-24 and C-25), but only C-20 stereoisomers give separate compound peaks, producing up to four regular stereoisomers of 26-mes ($\alpha\alpha\alpha S$, $\alpha\beta\beta R$, $\alpha\beta\beta S$ and $\alpha\alpha\alpha R$) in ancient rocks and oils, as has also been found for other sterane compounds.

sponges using HyPy to directly convert sterols into steranes (Supplementary Table 5). Species of *Rhabdastrella* and *Geodia* both produced appreciable amounts of 26-mes after the reductive conversion of sterols to steranes via HyPy treatment (1–9% of total C_{27-30} steranes; Supplementary Table 5). Apart from *Geodia hentscheli*, which only makes conventional sterols for which side chain alkylation is restricted to the C-24 position, 26-mes was the predominant C_{30} sterane product in our *Rhabdastrella* and *Geodia* specimens. Molecular phylogenetic results indicate that these species are closely related within Geodiidae (order Tetractinellida). Additionally, we also detected trace amounts of 26-mes steranes along with 24-ipc and 24-npc in four species of *Aplysina* and *Verongula* (order Verongiida) and one species of *Cymbaxinella* (order Agelasida) (Supplementary Table 5). No 26-mes precursors were detected in a *Jaspis* species, where stelliferasterol and isostelliferasterol were supposedly originally discovered^{20,21}. This is consistent with the belief that the original Great Barrier Reef '*Jaspis stellifera*' specimens were misidentified and were in fact *R. globostellata*²². Our HyPy results for three specimens of *R. globostellata* confirmed that 26-mes sterol precursors were present, as well as in two other *Rhabdastrella* species. Other than the Geodiidae, another known major source of 26-mes steroids is *Petrosia (Strongylophora) cf. durissima*²³ (order Haplosclerida), which can synthesize strongylosterol (Fig. 2) as its dominant single sterol. As not all demosponges make 26-mes, Geodiidae and *P. (S.) cf. durissima* may have retained the ancestral capacity to make terminally methylated C_{30} steroids as major membrane lipids, which has been lost in other demosponge groups.

Our new findings of 26-mes production in *Geodia*, *Rhabdastrella*, *Aplysina* (aspiculate), *Verongula* (aspiculate) and *Cymbaxinella* species suggest that a wider range of demosponge groups might possibly make 26-mes, as well as other terminally methylated steroids, but have not yet been identified. These demosponge species and others can make various unusual C_{29} and C_{30} sterols with terminal methylation in the side chain (Supplementary Information). For example, *Thymosiopsis conglomerans* (order Chondrillida) makes a distinctive C_{30} sterol²⁴, yielding a different sterane skeleton that

has not been detected in the ancient record (Fig. 1). The finding of this extra terminal carbon atom in a variety of sterols from diverse extant demosponges suggests that the capability for 26-methylated sterol side chains probably has a deep origin within the clade. In the case of both 24-ipc² and 26-mes, the ability to make these sterols is phylogenetically widespread within demosponges. Notably, the known extant demosponge species that contain 26-mes as the dominant hydrocarbon core of their C_{30} steroids are different from those that make 24-ipc as major steroids. Specifically, the demosponge family Halichondriidae (*Ciocalypta* (previously *Collocalypta*), *Halichondria* and *Epipolasis*) and the genus *Topsentia* make 24-ipc among their most abundant sterols¹² while 24-ipc constitutes >99% of sterols in *Cymbastela coralliophila* (*Pseudoaxynissa* species in the original publication) from the family Axinellidae⁵.

From a comprehensive database of steroid assays performed on extant organisms from decades of lipid research, alongside our targeted assays here, 26-mes precursor sterols are found only in certain demosponges^{20,21,23} but—to our knowledge—have never been reported from any other group of eukaryotes (Supplementary Information). This evidence of absence applies to diverse groups of algae^{19,25,26}, hexactinellid sponges (ref. ²⁷ and this study) calcisponges (refs ^{2,28} and this study) homoscleromorphs (this study) and unicellular animal outgroups¹³. Indeed, only steroids possessing conventional side chains (with methyl, ethyl or propyl groups or a hydrogen substituent at C-24) have been reported for other sponge classes, heterotrophic protists and these unicellular animal outgroups^{13,14,27,28}, but not steroids with unconventional side chains of any variety. Thus, the finding of 26-mes, together with the 24-ipc steranes in Neoproterozoic rocks and oils, is most parsimoniously explained by an origin from demosponges living in marine settings.

In terms of possible older occurrences of sponge biomarkers, robust evidence for steranes has been reported in some 800–700 Myr Neoproterozoic rocks^{29,30} from the Chuar Group (USA) and Visingsö Group (Sweden). These rocks contain an unusual C_{28} sterane, 26-methylcholestane, informally designated as cryostane (Fig. 3), which has been proposed as a possible ancient sponge or unicellular stem metazoan marker²⁹. Cryostane is also characterized by the unusual terminal side chain methylation at C-26, making it a structural analogue of 26-mes and adding credence to the case for cryostane being a plausible ancient sponge biomarker. However, plausible precursor sterols for cryostane have not yet been found in any extant organisms, despite the discovery of a wide variety of other unconventional steroid structures in modern sponges. The Chuar and Visingsö Group rocks, like all pre-Sturtian-aged samples reported so far, are devoid of 24-ipc, 24-npc and 26-mes steranes. Thus, cryostane cannot currently be applied as a robust animal biomarker until more is known about its biological origins and whether the biosynthetic capacity to make unconventional 26-methylated steroids (Fig. 4) is restricted to demosponges or otherwise. The origins of cryostane are intriguing and a bridging of the cryostane and 26-mes/24-ipc records may signify a continuity of sponge markers persisting through the two Neoproterozoic glaciation events (Fig. 3), but this requires further investigation.

Conclusions

The co-occurrence of 24-ipc and 26-mes steranes constitutes the earliest robust biomarker evidence for Neoproterozoic animals, first detected in the Cryogenian period before the deposition of the Marinoan cap carbonate (>635 Ma), but following the Sturtian glaciation (beginning at <717 Ma and terminating around 660 Ma)^{2,6}. This suggests that demosponges first achieved ecological prominence in Neoproterozoic marine paleoenvironments, at least between 660 and 635 Ma, which is consistent with recent molecular clock predictions for their first appearance^{15–18}. This view from molecular clocks and biomarkers remains to be reconciled




Fig. 3 | A revised Neoproterozoic–Cambrian timeline showing co-occurrences of 26-mes and 24-ipc sterane biomarkers. The South Oman record commences in the Cryogenian period (>635 Ma) after the Sturtian glaciation (terminating at around 660 Ma⁶) and continues throughout the Ediacaran period into the Early Cambrian for Huqf Supergroup rocks (Supplementary Tables 1–2). Other Ediacaran oils also contain the C₃₀ steranes series (Supplementary Table 3), but some Ediacaran rocks are devoid of the C₃₀ sterane series, although they contain predominantly algal steranes with a C₂₉ dominance⁴⁰. The distribution and abundance patterns of the C₃₀ sterane 26-mes have yet to be fully established for the Phanerozoic rock record; however, it can be detected in some Phanerozoic rocks and oils (see Supplementary Table 3). Cryostane is a potentially older biomarker for sponges or unicellular protists, and it has been detected in pre-Sturtian rocks in the 800–717 Myr age range²⁹ but not in older samples. Cryostane is a C₂₈ sterane analogue of 26-mes, but corresponding sterol precursors for cryostane have never been reported from any extant taxa despite the identification of 26-mes demosponge sterols many decades ago^{20,21,23,38,39}. ‘M’ signifies the Marinoan glaciation; cryostane temporal range is represented by red shading although possible occurrences in younger rocks and oils require further investigation; 26-mes and 24-ipc range is represented by the black bar.

with the fossil spicule record, which suggests a later (Cambrian) origin³¹. Future sampling of modern taxa may reveal other sources of 26-mes sterols (see Supplementary Information), but multiple possibilities for taphonomic mega-bias of early sponge body fossils have been identified (see ref. ³² and the references therein), perhaps related to sparse biomineralization and/or silica dissolution and reprecipitation in low-oxygen marine conditions³³, despite overall higher Proterozoic oceanic silica levels. The records could also be reconciled if demosponge spicules evolved convergently in the Cambrian or if aspiculate demosponges were dominant producers of Neoproterozoic 26-mes and 24-ipc.

All available current data indicate that 26-mes steranes are made by diverse species of modern demosponges, and apparently not by any other sponge class (Hexactinellida, Homoscleromorpha or Calcarea) or other extant eukaryote, implying that Neoproterozoic total-group demosponges were the most probable source biota for these biomarkers. As demosponges are derived within Porifera, these data consequently predict the presence of sponges at this time irrespective of whether sponges³ or ctenophores³⁴ are the sister group of all other animals. Thus, this new Neoproterozoic steroid biomarker evidence for demosponges provides a conservative minimum time estimate for the origin of animal multicellularity and the sponge body plan involving feeding with a water canal system.

Methods

Catalytic HyPy of sponge biomass. Continuous-flow HyPy experiments were performed on 30–150 mg of catalyst-loaded sponge biomass at the University of California, Riverside (UCR) as described previously^{2,19}. Freeze-dried sponge biomass was initially impregnated with an aqueous methanol solution of ammonium dioxodithiomolybdate ((NH₄)₂MoO₂S₂) to give a nominal loading of 3–10 wt% catalyst. Ammonium dioxodithiomolybdate reductively decomposes in situ under HyPy conditions above 250 °C to form a catalytically active molybdenum sulfide (MoS₂) phase.

The catalyst-loaded samples were heated in a stainless-steel (316 grade) reactor tube from ambient temperature to 250 °C at 100 °C min⁻¹, then to 460 °C at 8 °C min⁻¹. A hydrogen sweep gas flow rate of 6 dm³ min⁻¹, measured at ambient temperature and pressure, through the reactor bed ensured that the residence times of volatiles generated were of the order of only a few seconds. Products were




Fig. 4 | Chemical structures for a selection of conventional, unconventional and unsaturated sterols found in modern eukaryotic taxa.

Chemical structures numbered 1–15 (top) represent possible variations of the part of the structure labelled ‘R’ in structures A–C (bottom). Conventional sterols show side chain alkylation of the C₂₇ cholestane core skeleton restricted to the C-24 position, while all examples of unconventional sterols shown here have extended side chains arising from terminal methylation at C-26 and/or C-27. Note that the tetracyclic core and/or side chain of sterols may or may not contain an alkene bond, and when alkene bonds are present these sterols are known as unsaturated compounds. In this scheme, cholesterol is compound B1, stelliferasterol is B14 and stronglysterol is B15. The dominant ancient fossil form is sterane (A), with alcohol and alkene bonds removed by chemical reduction.

collected on a silica gel trap cooled with dry ice and recovered for subsequent fractionation using silica gel adsorption chromatography.

HyPy products (hydropyrolysates) of sponge biomass were separated by silica gel adsorption chromatography into aliphatic (alkane + alkene), aromatic and polar (or nitrogen, sulfur or oxygen) compounds by elution with *n*-hexane, *n*-hexane:dichloromethane (DCM) (1:1 v/v) and DCM:methanol (3:1 v/v), respectively. For hydropyrolysates, solvent-extracted activated copper turnings were added to concentrated solutions of aliphatic hydrocarbon fractions to remove all traces of elemental sulfur, which is formed from disproportionation of the catalyst during HyPy. Aliphatic fractions were further purified to a saturated hydrocarbon fraction by the removal of any unsaturated products (alkenes) via silver nitrate-impregnated silica gel adsorption chromatography and elution with *n*-hexane.

Lipid biomarker analysis of ancient rocks/oils. Detailed methods for the extraction and analysis of sedimentary rocks and oils at UCR were described previously^{35–37} and data are shown in Supplementary Table 3. Rock pieces were first trimmed with a water-cooled rock saw to remove outer weathered surfaces (of at least a few mm thickness) and expose a solid inner portion. They were then sonicated in a sequence of ultrapure water, methanol, DCM and hexane before a final rinse with DCM before powdering and bitumen extraction. Rock fragments were powdering in a zirconia ceramic puck mill in a SPEX 8515 shatterbox,

and cleaned between samples by powdering two batches of fired sand (850 °C overnight) and rinsing with the above series of solvents. Typically, 5 g of crushed rock was extracted in a CEM Microwave Accelerated Reaction System at 100 °C in a DCM:methanol (9:1 v/v) mixture for 15 min. Full laboratory procedural blanks with combusted sand were performed in parallel with each batch of rocks to ensure that any background signals were negligible compared with biomarker analyte abundances found in the rocks (typically by at least three orders of magnitude). Saturated hydrocarbon and aromatic fractions for rock bitumens and oils were obtained by silica gel column chromatography; the saturate fractions were eluted with hexane and the aromatic fractions with DCM:hexane (1:1 v/v).

The procedures for ancient biomarker analyses from sedimentary rocks from the Huqf Supergroup of the South Oman Salt Basin performed at the Massachusetts Institute of Technology (results reported in Supplementary Tables 1 and 2) were similar to those described above for UCR protocols, including multiple reaction monitoring (MRM)-GC-MS methods (see below), which have been described in detail previously^{2,8}. Analytical errors for absolute yields of individual hopanes and steranes are estimated at $\pm 30\%$. Average uncertainties in hopane and sterane biomarker ratios are $\pm 8\%$, as calculated from multiple analyses of a saturated hydrocarbon fraction from Australian Geological Survey Organisation (AGSO) and GeoMark Research standard oils ($n = 30$). Full procedural blanks with combusted sand were run in parallel with each batch of samples to quantify any low background signal. Supplementary Tables 1 and 2 contain original data from Love et al.², but now also with 26-mes sterane data added. The yields and ratios verify that significant abundances of 26-mes were detected in all of these samples at a similar order of magnitude abundance to those of 24-ipc steranes.

Sterane analysis using MRM-GC-MS. Saturated hydrocarbon fractions from ancient rocks and oils, as well as from modern sponge HyPy pyrolysates, were analysed by MRM-GC-MS conducted at UCR on a Waters Autospec Premier mass spectrometer equipped with an Agilent 7890A gas chromatograph and DB-1MS coated capillary column (60 m \times 0.25 mm, 0.25 μ m film) using helium as the carrier gas. Typically, 1 μ l of a hydrocarbon fraction dissolved in hexane was injected onto the gas chromatography column in splitless injection mode. The gas chromatography temperature programme consisted of an initial hold at 60 °C for 2 min, heating to 150 °C at 10 °C min⁻¹ followed by heating to 320 °C at 3 °C min⁻¹, and a final hold for 22 min. Analyses were performed via splitless injection in electron impact mode, with an ionization energy of 70 eV and an accelerating voltage of 8 kV. MRM transitions for C₂₇–C₃₅ hopanes, C₃₁–C₃₆ methylhopanes, C₂₁–C₂₂ and C₂₆–C₃₀ steranes, C₃₀ methylsteranes and C₁₉–C₂₆ tricyclic terpanes were monitored. Procedural blanks with pre-combusted sand yielded less than 0.1 ng of individual hopane and sterane isomers per gram of combusted sand³⁷. Polycyclic biomarker alkanes (tricyclic terpanes, hopanes, steranes and so on) were quantified by the addition of a deuterated C₂₉ sterane standard (d₄- $\alpha\alpha$ -24-ethylcholestane (20R)) to saturated hydrocarbon fractions and comparison of the relative peak areas. In MRM analyses, this standard compound was detected using 404 \rightarrow 221 Da ion transition. Cross-talk of the non-sterane signal in 414 \rightarrow 217 Da ion chromatograms from C₃₀ and C₃₁ hopanes was <0.2% of the 412 \rightarrow 191 Da hopane signal (mainly 17 α ,21 β (H)-hopane, which is resolvable from C₃₀ steranes) and <1% of the 426 \rightarrow 191 Da signal, respectively³⁶.

Peak identifications of sponge steranes were confirmed by comparing the retention times with an AGSO oil saturated hydrocarbon standard and Neoproterozoic oils from Eastern Siberia⁴⁹ and India⁷, which were reported previously to contain significant quantities of 24-ipc and which we have now demonstrated contain significant quantities of 26-mes (Supplementary Table 3). Polycyclic biomarkers were quantified assuming equal-mass spectral response factors between analytes and the d₄-C₂₉- $\alpha\alpha$ -24-ethylcholestane (20R) internal standard. Analytical errors for absolute yields of individual hopanes and steranes are estimated at $\pm 30\%$. Average uncertainties in hopane and sterane biomarker ratios are $\pm 8\%$, as calculated from multiple analyses of a saturated hydrocarbon fraction prepared from AGSO and GeoMark Research standard oils ($n = 30$ MRM analyses).

Sterane analysis by gas chromatography triple quadrupole mass spectrometry (GC-QQQ-MS). To confirm the presence of the new 26-mes peak and investigate the retention time of the analyte peaks compared with other C₃₀ steranes (24-npc and 24-ipc), the saturated hydrocarbon fractions from sponge HyPy products and oils from Eastern Siberia and India (Supplementary Table 3) were run on a different instrument employing a different gas chromatography column from that used in the MRM-GC-MS instrument at UCR. GC-QQQ-MS was performed at GeoMark Research (Houston, Texas) on an Agilent 7000A Triple Quad interfaced with an Agilent 7890A gas chromatograph equipped with a J&W Scientific capillary column (DB-5MS+ DG; 60 m \times 0.25 mm i.d., 0.25 μ m film thickness, 10 m guard column). Using helium as the carrier gas, the flow was programmed from 1.2 to 3.2 ml min⁻¹. The gas chromatography oven was programmed from 40 °C (2 min) to 325 °C (25.75 min) at 4 °C min⁻¹. Saturated hydrocarbon fractions were spiked with a mixture of seven internal standards (Chiron Routine Biomarker Internal Standard Cocktail 1). Samples were concentrated without being taken to dryness and were injected in cold splitless mode at 45 °C with the injector temperature ramped at 700 °C min⁻¹ to 300 °C. The mass spectrometry source was operated in electron ionization mode at 300 °C with ionization energy at -70 eV. A number

of molecular ion to fragment transitions were monitored throughout the run; the dwell time was adjusted as needed to produce 3.5 cycles s⁻¹. Exact chromatographic co-elution (with identical retention time in the C₃₀ sterane analytical window) of the $\alpha\alpha$ R diastereoisomer of 26-mes sterane in our ancient oils with the equivalent peak from the modern sponges was demonstrated in 414 \rightarrow 217 Da ion transitions (the parent molecular mass to daughter fragment ion transition for regular (4-desmethyl) C₃₀ sterane compounds).

Extraction and analysis of sterols in modern sponge cells. A total of 18 modern sponge samples were acquired for solvent extraction to monitor their free sterol contents (as trimethylsilyl ethers). Sponge specimens were supplied by co-authors P.C. and E.A.S. and their colleagues, including J. Vacelet, U. Hentschel, K. Peterson, T. Molinski, T. Pérez, H. Tore Rapp, A. Plotkin, J.-S. Hong, Y. M. Huang, S. Rohde, S. Nichols, B. Calcinaï, J. V. Lopez, G. Gatti, B. Ciperling, J.-P. Fonseca, L. Magro, F. Azzini and A. G. Collins, and the Bedford Institute of Oceanography (Dartmouth, Canada). Sponge biomass arrived immersed in ethanol or freeze-dried. Combined ethanol washings for each sample were filtered to remove suspended particulates, concentrated into a small volume and then transferred to a pre-weighed glass vial and blown down carefully under dry N₂ gas. Freeze-dried sponge biomass was extracted via sonication for 25 min in DCM:methanol (3:1 v/v) to recover the total lipid extract.

Total lipid extracts were separated into three fractions, based on polarity, by silica gel absorption chromatography. Approximately 1–5 mg of total lipid extract was adsorbed on the top of a 10 cm silica gel pipette column and then sequentially eluted with 1.5 column volumes of *n*-hexane (fraction 1), 2 column volumes of DCM (fraction 2) and 3 column volumes of DCM:methanol (7:3 v/v) (fraction 3). The alcohol products, including sterols, typically eluted in fraction 2, and approximately 20–50 μ g of this fraction was derivatized with 10 μ l of bis(trimethylsilyl)trifluoroacetamide in 10 μ l of pyridine at 70 °C for 30 min.

Alcohol fractions were then analysed by GC-MS as trimethylsilyl ethers within 36 h of derivatization in full scan mode at UCR using GC-MS on an Agilent 7890A gas chromatography system coupled to an Agilent 5975C inert Mass Selective Detector mass spectrometer. Sample solutions were volatilized via programmed-temperature vaporization injection onto a DB-1MS capillary column (60 m \times 0.32 mm, 0.25 μ m film thickness) and helium was used as the carrier gas. The oven temperature programme used for gas chromatography for the derivatized alcohol fraction consisted of an initial temperature hold at 60 °C for 2 min, followed by an increase to 150 °C at 20 °C min⁻¹, then a subsequent increase to 325 °C at 2 °C min⁻¹ and a hold for 20 min. Data was analysed using ChemStation G10701CA (Version C) software (Agilent Technologies). C₃₀ sterol identifications for the three 26-mes precursors (stelliferasterol (B13), isostelliferasterol (B14) and strongylosterol (B15); see also Fig. 2) in certain *Rhabdastrella* and *Geodia* sponge species were identified from published mass spectral features and relative retention times^{30,21,23,38,39}. Stelliferasterol was the dominant C₃₀ sterol in *R. globostellata* (PC922; Supplementary Figs. 1–2), while strongylosterol, stelliferasterol and isostelliferasterol were found in *Geodia parva* (GpII).

Reporting Summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

The authors declare that the data supporting the findings of this study are available within the paper and its Supplementary Information files.

Received: 22 July 2018; Accepted: 22 August 2018;
Published online: 15 October 2018

References

- Brocks, J. J. et al. The rise of algae in Cryogenian oceans and the emergence of animals. *Nature* **548**, 578–581 (2017).
- Love, G. D. et al. Fossil steroids record the appearance of Demospongiae during the Cryogenian period. *Nature* **457**, 718–721 (2009).
- Simion, P. et al. Large and consistent phylogenomic dataset supports sponges as the sister group to all other animals. *Curr. Biol.* **27**, 958–968 (2017).
- McCaffrey, M. A. et al. Paleoenvironmental implications of novel C₃₀ steranes in Precambrian to Cenozoic age petroleum and bitumen. *Geochim. Cosmochim. Acta* **58**, 529–532 (1994).
- Hofheinz, W. & Oesterhelt, G. 24-isopropylcholesterol and 22-dehydro-24-isopropylcholesterol, novel sterols from a sponge. *Helv. Chim. Acta* **62**, 1307–1309 (1979).
- Rooney, A. D., Strauss, J. V., Brandon, A. D. & Macdonald, F. A. A Cryogenian chronology: two long-lasting synchronous Neoproterozoic glaciations. *Geology* **43**, 459–462 (2015).
- Peters, K. E. et al. Recognition of an Infracambrian source rock based on biomarkers in the Baghewala-1 oil, India. *AAPG Bull.* **79**, 1481–1494 (1995).
- Grosjean, E. et al. Origin of petroleum in the Neoproterozoic–Cambrian South Oman Salt Basin. *Org. Geochem.* **40**, 87–110 (2009).

9. Kelly, A. E., Love, G. D., Zumberge, J. E. & Summons, R. E. Hydrocarbon biomarkers of Neoproterozoic to Lower Cambrian oils from Eastern Siberia. *Org. Geochem.* **42**, 640–654 (2011).
10. Antcliffe, J. B. Questioning the evidence of organic compounds called sponge biomarkers. *Palaeontology* **56**, 917–925 (2013).
11. Siegl, A. et al. Single-cell genomics reveals the lifestyle of Poribacteria, a candidate phylum symbiotically associated with marine sponges. *ISME J.* **5**, 61–70 (2011).
12. Love, G. D. & Summons, R. E. The record of Cryogenian sponges. A response to Antcliffe. *Palaeontology* **58**, 1131–1136 (2015).
13. Gold, D. A. et al. Sterol and genomic analyses validate the sponge biomarker hypothesis. *Proc. Natl Acad. Sci. USA* **113**, 2684–2689 (2016).
14. Grabenstatter, J. et al. Identification of 24-*n*-proplidenecholesterol in a member of the Foraminifera. *Org. Geochem.* **63**, 145–151 (2013).
15. Erwin, D. H. et al. The Cambrian conundrum: early divergence and later ecological success in the early history of animals. *Science* **334**, 1091–1097 (2011).
16. dos Reis, M. et al. Uncertainty in the timing of origin of animals and the limits of precision in molecular timescales. *Curr. Biol.* **25**, 2939–2950 (2015).
17. Dohrmann, M. & Wörheide, G. Dating early animal evolution using phylogenomic data. *Sci. Rep.* **7**, 3599 (2017).
18. Schuster, A. et al. Divergence times in demosponges (Porifera): first insights from new mitogenomes and the inclusion of fossils in a birth–death clock model. *BMC Evol. Biol.* **18**, 114 (2018).
19. Love, G. D. et al. An optimised catalytic hydrolysis method for the rapid screening of microbial cultures for lipid biomarkers. *Org. Geochem.* **36**, 63–82 (2005).
20. Theobald, N. & Djerassi, C. Determination of the absolute configuration of stelleriferasterol and strongylosterol—two marine sterols with ‘extended’ side chains. *Tetrahedron Lett.* **45**, 4369–4372 (1978).
21. Theobald, N., Wells, R. J. & Djerassi, C. Minor and trace sterols in marine invertebrates. 8. Isolation, structure elucidation, and partial synthesis of two novel sterols—stelleriferasterol and isostelleriferasterol. *J. Am. Chem. Soc.* **100**, 7677–7684 (1978).
22. Kennedy, J. A. Resolving the ‘*Jaspis stellifera*’ complex. *Mem. Queensl. Mus.* **45**, 453–476 (2000).
23. Bortolotto, M., Braekman, J. C., Daloz, D. & Tursch, D. Chemical studies of marine invertebrates. XXXXVI. Strongylosterol, a novel C-30 sterol from the sponge *Strongylophora durissima* Dendy. *Bull. Soc. Chim. Belg.* **87**, 539–543 (1978).
24. Vacelet, J. et al. Morphological, chemical and biochemical characterization of a new species of sponge without skeleton (Porifera, Demospongiae) from the Mediterranean Sea. *Zoosystema* **22**, 313–326 (2000).
25. Volkman, J. K. Sterols in microorganisms. *Appl. Microbiol. Biotechnol.* **60**, 495–506 (2003).
26. Kodner, R. B., Pearson, A., Summons, R. E. & Knoll, A. H. Sterols in red and green algae: quantification, phylogeny, and relevance for the interpretation of geologic steranes. *Geobiology* **6**, 411–420 (2008).
27. Blumenberg, M., Thiel, V., Pape, T. & Michaelis, W. The steroids of hexactinellid sponges. *Naturwissenschaften* **89**, 415–419 (2002).
28. Hagemann, A., Voigt, O., Wörheide, G. & Thiel, V. The sterols of calcareous sponges (Calcarea, Porifera). *Chem. Phys. Lipids* **156**, 26–32 (2008).
29. Brocks, J. J. et al. Early sponges and toxic protists: possible sources of cryostane, an age diagnostic biomarker antedating Sturtian Snowball Earth. *Geobiology* **14**, 129–149 (2015).
30. Adam, P., Schaeffer, P. & Brocks, J. J. Synthesis of 26-methylcholestanane and identification of cryostane in mid-Neoproterozoic sediments. *Org. Geochem.* **115**, 246–249 (2018).
31. Botting, J. P. & Muir, L. A. Early sponge evolution: a review and phylogenetic framework. *Palaeoworld* **27**, 1–29 (2018).
32. Muscente, A. D., Marc Michel, F., Dale, J. G. & Xiao, S. Assessing the veracity of Precambrian ‘sponge’ fossils using in situ nanoscale analytical techniques. *Precam. Res.* **263**, 142–156 (2015).
33. Berelson, W. M. et al. Anaerobic diagenesis of silica and carbon in continental margin sediments: discrete zones of TCO₂ production. *Geochim. Cosmochim. Acta* **69**, 4611–4629 (2005).
34. Whelan, N. V. et al. Ctenophore relationships and their placement as the sister group to all other animals. *Nat. Ecol. Evol.* **1**, 1737–1746 (2017).
35. Rohrsen, M., Love, G. D., Fischer, W., Finnegan, S. & Fike, D. A. Lipid biomarkers record fundamental changes in the microbial community structure of tropical seas during the Late Ordovician Hirnantian glaciation. *Geology* **41**, 127–130 (2013).
36. Rohrsen, M., Gill, B. C. & Love, G. D. Scarcity of the C₃₀ sterane biomarker, 24-*n*-propylcholestanane, in Lower Paleozoic marine paleoenvironments. *Org. Geochem.* **80**, 1–7 (2015).
37. Haddad, E. E. et al. Lipid biomarker stratigraphic records through the Late Devonian Frasnian/Famennian boundary: comparison of high- and low-latitude epicontinental marine settings. *Org. Geochem.* **98**, 38–53 (2016).
38. Stoilov, I. L., Thompson, J. E., Cho, J. H. & Djerassi, C. Biosynthetic studies of marine lipids. 9. Stereochemical aspects and hydrogen migrations in the biosynthesis of the triply alkylated side chain of the sponge sterol strongylosterol. *J. Am. Chem. Soc.* **108**, 8235–8241 (1986).
39. Cho, J. H., Thompson, J. E., Stoilov, I. L. & Djerassi, C. Biosynthetic studies of marine lipids. 14. 24(28)-dehydroaplysterol and other sponge sterols from *Jaspis stellifera*. *J. Org. Chem.* **53**, 3466–3469 (1988).
40. Pehr, K. et al. Ediacara biota thrived in oligotrophic and bacterially dominated marine environments across Baltica. *Nat. Commun.* **9**, 1807 (2018).

Acknowledgements

Funding support for this work came from the NASA Astrobiology Institute teams Alternative Earths (NNA15BB03A) and Foundations of Complex Life (NNA13AA90A), NASA Exobiology program (grant number 80NSSC18K1085), NSF Frontiers in Earth System Dynamics programme (grant number 1338810), Agouron Institute, and European Union's Horizon 2020 research and innovation programme through the SponGES project (grant agreement number 679849). Many thanks go to the sponge collectors, including T. Pérez, H. Tore Rapp, A. Plotkin, J.-S. Hong, Y. M. Huang, S. Rohde, S. Nichols, J. V. Lopez, B. Calcinai, G. Gatti, B. Ciperling, J. Boavida, J.-P. Fonseca, L. Magro and F. Azzini. The fieldwork on the Mohns and Knipovich Ridges aboard the RV G. O. Sars in 2014 was supported by the Research Council of Norway through the Centre for Geobiology, UiB (contract number 179560). A *Vazella pourtalesii* specimen was provided by E. Kenchington through funding from the Marine Conservation Target fund of the Department of Fisheries and Oceans Canada. Other specimens were provided by E. Kenchington as Canadian lead for the NEREIDA (NAFO Potential Vulnerable Marine Ecosystems-Impacts of Deep-sea Fisheries) project led by Spain and Canada. We thank K. Ubhayasekera (Department of Chemistry, Uppsala University) for GC-MS analyses, and S. Rajendran and T. Aljazar (Department of Medicinal Chemistry, Uppsala University) for help with the isolation of sponge sterols. We thank Petroleum Development Oman for Neoproterozoic–Cambrian rock samples from the South Oman Salt Basin for ancient biomarker analysis. We are grateful to D. Rocher for GC-QQQ-MS analysis and GeoMark Research (Houston, TX) for providing oil samples from Eastern Siberia.

Author contributions

J.A.Z. and G.D.L. planned the investigation and wrote the manuscript with input from P.C. and E.A.S. J.A.Z. processed and interpreted the lipid biomarker data with help from G.D.L., E.G., R.E.S. and M.R. P.C., E.A.S., R.E.S., S.G., M.R., E.G. and J.P.G. provided comments on drafts of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41559-018-0676-2>.

Reprints and permissions information is available at www.nature.com/reprints.

Correspondence and requests for materials should be addressed to G.D.L.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see [Authors & Referees](#) and the [Editorial Policy Checklist](#).

Statistical parameters

When statistical analyses are reported, confirm that the following items are present in the relevant location (e.g. figure legend, table legend, main text, or Methods section).

n/a | Confirmed

- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- An indication of whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistics including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated
- Clearly defined error bars
State explicitly what error bars represent (e.g. SD, SE, CI)

Our web collection on [statistics for biologists](#) may be useful.

Software and code

Policy information about [availability of computer code](#)

Data collection

No such software was used

Data analysis

No such software was used

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors/reviewers upon request. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

The authors declare that the data supporting the findings of this study are available within the paper and its supplementary information files.

Field-specific reporting

Please select the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/authors/policies/ReportingSummary-flat.pdf](https://www.nature.com/authors/policies/ReportingSummary-flat.pdf)

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	We analyzed and quantified the lipid biomarker content of a large suite of ancient sedimentary rocks and oils, specifically targeting ancient steroid biomarker compounds sourced by demosponges of Neoproterozoic-Cambrian age.
Research sample	We targeted ancient sedimentary rocks of low thermal maturity which allowed excellent preservation of lipid biomarker constituents
Sampling strategy	64 ancient sedimentary rocks samples of Cryogenian-Cambrian age were analyzed to provide stratigraphic coverage of all formations from the Huqf Supergroup, South Oman Salt Basin. Ediacaran-Cambrian oils from Siberia and India, as well as some Phanerozoic source rocks, were also analyzed for comparison.
Data collection	Lipid biomarker analysis was performed at UC Riverside and MIT
Timing and spatial scale	Sedimentary rock and oils were obtained from research projects between 2005 and 2017, mostly supplied by oil companies from sub-surface drilling of wells. The rock and oils used are grouped by geological age and provenance.
Data exclusions	No exclusions, not applicable.
Reproducibility	Our reproducibility for molecular biomarker ratios and yields was assessed using oil hydrocarbon standards ran with each batch of analyses.
Randomization	The rocks and oil samples are grouped by geological age and provenance
Blinding	Not applicable
Did the study involve field work?	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No

Reporting for specific materials, systems and methods

Materials & experimental systems

n/a	Involvement in the study
<input type="checkbox"/>	<input checked="" type="checkbox"/> Unique biological materials
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input type="checkbox"/>	<input checked="" type="checkbox"/> Palaeontology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Human research participants

Methods

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging

Unique biological materials

Policy information about [availability of materials](#)

Obtaining unique materials Modern sponge samples are available from the collections of two of the co-authors, Sperling and Cardenas.

Palaeontology

Specimen provenance Sedimentary rock and oil samples were supplied by Petroleum Development Oman from wells drilled. PDO have approved publication of the data.

Specimen deposition

Rock and oil samples are held in labs of Love at UC Riverside and Summons at MIT

Dating methods

Not applicable

Tick this box to confirm that the raw and calibrated dates are available in the paper or in Supplementary Information.