

AD646 090

CIT - ELECTRON TUBE & MICROWAVE
LABORATORY REPORT

PULSE STIMULATED RADIATION FROM A PLASMA
INVOLVING THE SECOND HARMONIC OF THE GYROFREQUENCY

Wilhelm H. Kegel

Technical Report No. 34

October 31, 1966

CALIFORNIA INSTITUTE OF TECHNOLOGY

PASADENA, CALIFORNIA

CALIFORNIA
INSTITUTE OF
OCT 16 1975
TECHNOLOGY

PULSE STIMULATED RADIATION FROM A PLASMA
INVOLVING THE SECOND HARMONIC OF THE GYROFREQUENCY

Wilhelm H. Kegel

Technical Report No. 34

October 31, 1966

California Institute of Technology
Pasadena, California

A Report on Research Supported by

Office of Naval Research

Electronics Division

Contract Nonr 220(50)

October 1966

PULSE STIMULATED RADIATION FROM A PLASMA INVOLVING THE SECOND HARMONIC OF THE GYROFREQUENCY*

W. H. KEGEL**

California Institute of Technology, Pasadena, California

Received 6 October 1966

If a plasma is excited by two short pulses at the gyrofrequency and its second harmonic, respectively, with a time separation τ , a radiation maximum is predicted to occur at the gyrofrequency a time τ after the second pulse.

We consider a magneto-plasma which is excited by two short microwave pulses, which have a time separation τ . The first pulse is assumed to be at the gyrofrequency, the second at its second harmonic. It is shown that a radiation peak at the gyrofrequency occurs a time τ after the second pulse. This radiation peak is due to the non-linear driving force during the second pulse. This is in contrast to the usual echo, where both pulses are at the gyrofrequency [1-5].

The plasma is considered in the single-particle approach. We account for field inhomogeneities by attributing different gyrofrequencies to different particles. In this model all electrons have at the end of the first pulse (fig. 1a) the same velocity V and the same phase.

$V = e E_1^0 t_1 / 2m$ is given by the strength of the first pulse, E_1^0 being the amplitude of the electric field and t_1 the duration of the pulse. A time τ after the first pulse ($\Delta\omega_c \tau \gg 1$), i.e. at the onset of the second pulse, the electrons are distributed on a circle in the two dimensional rotating velocity space [1] (fig. 1b) due to the differences in gyrofrequency. The second pulse (at $2\omega_c$) induces a quadrupole moment (fig. 1c). If $v(\tau)$ are the coordinates of a given particle in the rotating velocity space at the onset of the second pulse (fig. 1b) and v^* its coordinates at the end of the pulse (fig. 1c), we have in lowest order [6]:

$$v_x^* = v_x(\tau) e^S; \quad v_y^* = v_y(\tau) e^{-S} \quad (1)$$

with

* Research sponsored by the US Office of Naval Research.

** Work performed while the author was on leave from the Institut für Plasmaphysik, Garching bei München, Germany.




Fig. 1. Distribution of particle velocities in a rotating velocity space a) at the end of the first pulse, b) at the onset of the second pulse, c) at the end of the second pulse, d) at a time τ after the second pulse, the first pulse being at ω_c and the second at $2\omega_c$.

$$s = e E_2^0 t_2 / 2cm \quad (2)$$

A time τ after the second pulse, a particle which has a phase angle φ at the onset of the second pulse has moved around the origin by another angle $\varphi \pm 2n\pi$. As a result the ellipse in fig. 1c has been transformed into a circle, the center of which is different from the coordinate origin (fig. 1d). This shows that a macroscopic dipole moment has been formed, i.e. there is a radia-

tion maximum at the gyrofrequency at this time.

For the radiation at the gyrofrequency we have [4,5]:

$$I_c \sim N^2 V^2 \phi_1(t) \quad (3)$$

where N is the number of particles considered and the function $\phi_1(t)$ is given by:

$$\phi_1(t) = \left(\sum_{l=1}^N v_l \right)^{-2} \left| \sum_{l=1}^N v_l \exp[i(\omega_c^l t + \alpha_l)] \right|^2 \quad (4)$$

where α_l is the phase of the l -th particle at $t=0$.

If one determines ϕ_1 , for the radiation peak, i.e. for the time τ after the second pulse [6], one obtains on the assumption $s \ll 1$

$$\phi_1(\tau) \approx s^2 \quad (5)$$

(5) gives a finite amplitude of the radiation peak even for $\tau \rightarrow 0$, in contrast to the results one obtains in the case where both exciting pulses are at ω_c [4,5,7].

The conditions for the occurrence of the *usual* echo (both pulses at ω_c) are of two types: a) General conditions: The electron density must be sufficient for the effect to be observed, the initial temperature must be low enough so that the electrons do not drift out of the observation region, the decay time must be sufficiently long, etc. b) Special conditions which give the essential nonlinearity. For the nonlinearity essentially two different mechanisms have been proposed, namely an energy dependent collision frequency [7], and an energy dependent gyrofrequency [4,5]. In order to demonstrate that in a given experiment a given nonlinearity is the echo forming mechanism, one could try to keep the general conditions fixed

and change only the special conditions until the echo disappears.

In contrast to the usual echo the new effect discussed in this letter (second pulse at $2\omega_c$) depends on the general conditions only, because the nonlinearity is given in this case by the nonlinear driving force during the second pulse, which does not depend on any plasma parameter. So this effect could be used to check independently of the special conditions, whether the general conditions are fulfilled under given experimental conditions.

If other nonlinearities, such as a velocity dependent gyrofrequency or collision cross-section, are taken into account, further radiation maxima may be shown to occur at the gyrofrequency at times 3τ , 5τ , etc. after the second pulse [6].

I should like to acknowledge stimulating discussions with R. W. Gould, K. S. H. Lee and D. Montgomery.

References

1. R. W. Gould, Phys. Letters 19 (1965) 477.
2. G. F. Hermann and R. F. Whitmer, Phys. Rev. 143 (1966) A122.
3. R. M. Hill and D. E. Kaplan, Phys. Rev. Letters 14 (1965) 1062.
4. W. H. Kegel and R. W. Gould, Phys. Letters 19 (1965) 531.
5. W. H. Kegel, Plasma physics (J. Nucl. Energy Part C) in press.
6. W. H. Kegel, Calif. Inst. of Technology, Techn. Report no. 32 Nonr 220 (50), 1966. To be published.
7. D. E. Kaplan and R. M. Hill, Bull. Am. Phys. Soc. 11 (1966) 538.

* * * * *

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
California Institute of Technology Pasadena, California 91109		2b. GROUP	
3. REPORT TITLE			
PULSE STIMULATED RADIATION FROM A PLASMA INVOLVING THE SECOND HARMONIC OF THE GYROFREQUENCY.			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Technical Report (reprint) Oct. 1966			
5. AUTHOR(S) (First name, middle initial, last name)			
Wilhelm H. KEGEL			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
October 31, 1966		2	7
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
Nonr 220(50)		Technical Report No. 34	
b. PROJECT NO.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c.			
d.			
10. DISTRIBUTION STATEMENT			
Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		Office of Naval Research Electronics Division 427 Washington 25, D.C.	
13. ABSTRACT			
If a plasma is excited by two short pulses at the gyrofrequency and its second harmonic, respectively, with a time separation τ , a radiation maximum is predicted to occur at the gyrofrequency at a time τ after the second pulse.			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
plasma radiation cyclotron echo cyclotron harmonic radiation						