

# STRONG REDSHIFT CLUSTERING OF DISTANT GALAXIES<sup>1</sup>

JUDITH G. COHEN<sup>2</sup>, DAVID W. HOGG<sup>3</sup>, MICHAEL A. PAHRE<sup>2</sup> AND ROGER  
BLANDFORD<sup>3</sup>

Received \_\_\_\_\_; accepted \_\_\_\_\_

arXiv:astro-ph/9602120v1 22 Feb 1996

---

<sup>1</sup>Based in large part on observations obtained at the W.M. Keck Observatory, which is operated jointly by the California Institute of Technology and the University of California

<sup>2</sup>Palomar Observatory, Mail Stop 105-24, California Institute of Technology

<sup>3</sup>Theoretical Astrophysics, California Institute of Technology, Mail Stop 130-33, Pasadena, CA 91125

## ABSTRACT

We present initial results from a redshift survey carried out with the Low Resolution Imaging Spectrograph on the 10 m W. M. Keck Telescope of a field 14.6 arc-min<sup>2</sup> in solid angle. In the redshift distribution of the 106 extragalactic objects in this sample we find five strong peaks, with velocity dispersions of  $\sim 500$  km s<sup>-1</sup>. There is evidence for a non-uniform areal density of objects in at least two peaks. These peaks have characteristics (velocity dispersions, density enhancements, spacing, and spatial extent) similar to those of nearby galaxy structures (e.g., walls and clusters), and these are expected in a survey of this kind. We suggest that the prominence of these structures in our survey relative to that in other surveys can be attributed to our  $K$ -selection and dense sampling.

*Subject headings:* Cosmology: observations — Galaxies: redshift and distances — Large-scale structure of Universe

## 1. INTRODUCTION

This is the first in a series of papers describing the results of a deep survey of faint field galaxies in a single field centered at RA(J2000)  $00^h 53^m 23^s.20$ , Dec.  $+12^\circ 33' 57''.5$ . The field is from the HST Medium Deep Survey (Griffiths et al 1994), randomly selected on the basis of high Galactic latitude ( $b = -50^\circ$ ) and low reddening ( $A_V = 0.13$  mag, Burstein & Heiles 1982). Our redshifts were acquired with the Low Resolution Imaging Spectrograph (Oke et al 1995) on the 10 m W. M. Keck Telescope over a rectangular strip  $2 \times 7.3$  arcmin<sup>2</sup> extending north-south, centered on the HST field. Our primary sample comprises all 155 objects with  $K < 20$  mag in the  $2 \times 7.3$  arcmin<sup>2</sup> field. The photometry and the definition of the sample for spectroscopic work is described in Paper II of this series, Pahre et al (1996). We reach higher galaxy surface densities than the  $I$ -selected CFRS survey (LeFèvre et al 1995); our survey complements and extends to fainter objects the  $K$ -selected sample of the Hawaii group (Songaila et al 1994).

Paper III in this series (Cohen et al 1996) provides a detailed description of the redshift survey. Of the 155 objects in the sample, 90 have spectra typical of normal galaxies, three are quasars or broad-lined AGNs and 19 are Galactic stars. Of the remaining objects, 35 were observed, but no redshift could be determined, and eight were not observed at all. The effects of incompleteness in the sample selection and redshift identification are discussed in Papers II and III; both are irrelevant for the present work. The median redshift  $z$  of the 93 extragalactic objects in the main sample is  $z = 0.57$ . An additional 13 galaxy redshifts were determined for objects in this field which were slightly fainter than the  $K < 20$  mag limit or which lie slightly outside the spatial boundaries of the  $K$  selected sample.

## 2. REDSHIFT DISTRIBUTION

The redshift histogram is shown in Figure 1. Of the 106 objects, 40 are in the two strongest peaks, and 64 are in the 5 strongest, i.e. have  $|z - z_p| < 0.020$  for  $z_p = 0.392, 0.429, 0.581, 0.675, \text{ and } 0.766$ . In spite of the fact that there are no clusters apparent in the images of this field, the objects are highly clustered in redshift space. The overdensities in redshift space are at least a factor of 5; and more than 60 percent of the objects in the sample lie in these structures.

### 2.1. Feature significance

We now consider the probability that these apparent features might arise by chance out of a smooth galaxy distribution. This significance calculation is performed not in redshift but in the quantity  $V \equiv c \ln(1 + z)$ . This coordinate is devoid of global meaning but, granted an overall Hubble expansion, corresponds incrementally to local velocity differences. The galaxies are contained in an interval  $4 \times 10^4 < V < 2.4 \times 10^5 \text{ km s}^{-1}$ . As there is no adequate *a priori* understanding of the population from which these galaxies are drawn, the measured data set is used to derive smoothed velocity distributions by the addition of random velocity shifts drawn from a Gaussian distribution with width  $\sigma_V$ . This effectively erases the obvious structure without affecting the overall velocity distribution function. The data set is divided into  $N_b$  uniform velocity bins and the number of galaxies  $n_{0i}$  is counted in the single bin centered on each of the candidate associations. This exercise is repeated using multiple realizations of the smoothed distribution; the mean  $\bar{n}_i$  and standard deviation  $\sigma_{ni}$  are measured for each bin.

A measure of the significance of each feature is the statistic  $X_i = |n_{0i} - \bar{n}_i|/\sigma_{ni}$ . To test the null hypothesis that individual features arise by chance, we also measure the

distribution of  $X_i$  in the smoothed redshift data. Finally, as the appearance of features like these is notoriously sensitive to the binning, we repeat this procedure with different values of  $N_b$  and, for each value of  $N_b$ , a range of shifts of bin “phase,” comparing the maximum values of  $X_i$  measured with the maximal values produced from the smoothed distribution.

To carry this scheme out in practice, a conservatively small smoothing length  $\sigma_V = 2 \times 10^3 \text{ km s}^{-1}$  was adopted,  $N_b$  ranged from 35 to 125 (in integer increments), and for each  $N_b$ , 10 equally-spaced values of bin phase shift from 0 to 1 bin width were allowed. For each value of these parameters, 1000 realizations of the Gaussian smoothed distribution were investigated. The results are given in Table 1: the observed  $z_p$ ; the number of galaxies within the bin coinciding with  $z_p$  (this number changes slightly depending on  $N_b$ ; the value given is that corresponding to the  $N_b$  and phase at which  $X_i$  is the largest; it is not the same as the number assigned to each peak by the  $|z - z_p| < 0.020$  rule); the largest value  $X_{\max}$  of  $X_i$  for that feature inferred from the data seen over the full range in  $N_b$  and phase; and the likelihood of the feature arising by chance, computed as the fraction of realizations of the smoothed distribution in which such a large value of  $X_i$  was found in any bin. The latter quantity shows that the feature at  $z=0.392$  is not significant and the one at  $z=0.766$  is only marginally so. These results are robust to eliminating the few galaxies with redshifts of lower precision, or the 13 not in the  $K$ -selected sample. The estimated significances increase with increasing smoothing length  $\sigma_V$ .

## 2.2. Velocity dispersions

The radial velocity precision of our redshifts is unusually high for a deep redshift survey. We estimate that the uncertainty in  $z$  for those objects with redshifts considered secure and accurate (comprising 80 of the 106 galaxies) is  $\approx 300 \text{ km s}^{-1}$ . The velocity dispersions for the five strongest peaks in the redshift histogram are given in Table 2. The velocity

dispersions of the peaks are only slightly smaller when the objects with low-precision redshifts are excluded. Both because of the biasing effect of including outliers and the spreading of redshifts due to measurement uncertainties, these velocity dispersions should be treated as upper limits.

### 3. ANGULAR DISTRIBUTION

The angular distribution of the entire sample, as well as that of galaxies with  $|z - z_p| < 0.020$  for each of the five peaks, is shown in Figure 2.

To provide a quantitative test of the uniformity of the areal distributions, two-dimensional Kolmogorov-Smirnov tests (Fasano & Franceschini 1987) were applied to the sample. When the areal distribution of the entire sample is compared with a uniform distribution, no evidence for non-uniform distribution is found. For each feature, the areal distribution of objects in the feature (membership defined by maximizing  $X_i$  as described in the previous section) is compared with that of all the objects not in the feature. The 2-d K-S test  $D$  values for the individual peaks are given in Table 3. Two, at  $z=0.392$  and  $z=0.581$ , are significantly non-uniform in areal distribution. These results are not significantly altered if one uses only the 93 objects in the  $K < 20$  mag sample. They are also robust to replacing the distribution of feature non-members with a uniform distribution.

Interestingly, the one feature, at  $z = 0.392$ , judged insignificant in redshift clustering shows significant angular clustering, and the feature, at  $z = 0.766$ , judged marginally significant in redshift clustering is also marginally significant in angular clustering. For this reason we judge both these features at least marginally significant.

### 4. DISCUSSION

#### 4.1. Effects of sample definition decisions

Although there are many other faint object redshift surveys, none have found as much strong redshift clustering as is presented here. We believe that this can be attributed to differences in sample definition. This survey of objects goes to high galaxy number density,  $4 \times 10^4$  per square degree at  $K = 20$ , comparable to the “Hawaii” survey (Songaila et al, 1995) but with greater completeness at the faint end, and deeper than the CFRS (LeFèvre et al 1995) or the  $B$ -selected LDSS-2 survey (Glazebrook et al 1995). Infrared selection is not subject to the same biases towards late-type spirals and irregulars as is found in  $B$ -selected samples. Our redshifts are measured with good precision, allowing good resolution of the peaks and their velocity dispersions.

Finally and most importantly, this sample is not sparse-sampled, i.e., (almost) every object with  $K < 20$  mag in the field is observed, so we do not miss structures of limited angular extent. For very sparsely sampled data, one obtains no statistically valid peaks at all in the redshift distribution, c.f. the CFRS (Lilly et al 1995). As one samples less sparsely, “walls” with substantial velocity dispersions, as in the ESO survey (Bellanger & de Lapparent 1995), appear. Suggestions similar to these regarding the effect of different sampling schemes have been offered by de Lapparent et al (1991) and by Ramella et al (1992) to explain why some redshift surveys did not see structures such as the “Great Wall” while others, e.g., the CFA survey (de Lapparent et al 1986), did.

#### 4.2. Structure morphology

In general, imaging surveys for galaxy structures find clusters and filaments because these appear as angular patches of higher projected number density, but do not find walls because the projected number density is relatively invariant to collapse from

three-dimensional volumes into two-dimensional walls. On the other hand, redshift surveys like this one, in small angular areas, are unlikely to hit compact clusters and filaments, but should pierce any walls. Because the structures in the present sample are seen in the redshift distribution but not in the imaging, there is some evidence that they are wall-like.

The derived velocity dispersions  $\sigma_v$  are small and reminiscent of those of sparse clusters of galaxies. Zabludoff et al (1990) found a median  $\sigma_v = 740 \text{ km s}^{-1}$  for 69 nearby, moderately rich Abell clusters. Our observed  $\sigma_v$  are much smaller than that found for superclusters; for example, Postman, Geller, & Huchra (1988) found  $\sigma_v = 1300 \text{ km s}^{-1}$  for primary members of the seven Abell clusters in the Corona Borealis supercluster, while Small (1996) found  $1800 \text{ km s}^{-1}$  for the entire supercluster. If the cluster Cor Bor itself is excluded, then the value drops to  $< 1000 \text{ km s}^{-1}$  (Postman, Huchra, & Geller 1992; Zucca et al 1993). Small (1996) also observed many individual lines of sight through the supercluster and found  $400 < \sigma_v < 1200 \text{ km s}^{-1}$ ; these lines of sight are comparable in extent to our field, while our derived  $\sigma_v$  are among lowest values measured along individual Cor Bor lines of sight. The velocity dispersion for the Great Wall, however, is  $230 \text{ km s}^{-1}$  measured over  $3^\circ$  cells (Ramella et al 1992), while that for groups in the SSRS2 (DaCosta et al 1994) is  $\sim 200 \text{ km s}^{-1}$ . Our  $\sigma_v$  are thus comparable to Great-Wall-size structures, poor clusters of galaxies, or lines of sight through superclusters. The separations between structures are also comparable: the comoving distances of the five strongest redshift peaks are 915, 981, 1228, 1364, and  $1485h^{-1} \text{ Mpc}$ , while the distance between the Cor Bor supercluster and the one immediately behind it at  $z \approx 0.12$  (Goia et al 1982, Sarazin et al 1982, and Small 1996) is  $130h^{-1} \text{ Mpc}$ . In this regard our redshift distribution is also reminiscent of that found by Broadhurst et al (1990), although the features in ours are not “periodic.”

At  $z = 0.6$ , our field has a size of  $1.7 \times 0.5 h^{-2} \text{ Mpc}^2$ . We take the number density of clusters in the universe to be of order of a few times  $10^{-5} h^3 \text{ Mpc}^{-3}$ , as reported for

Shectman (1985) clusters (Bahcall, 1988) and  $R \geq 1$  Abell galaxy clusters (Postman et al 1996). The fact that we observe 5 clusters along a randomly selected line of sight implies a typical cluster radius of between 2 and  $5 h^{-1}$  Mpc, depending on the exact number density and the world model. These lengths are on the order of, if a bit higher than, locally measured cluster radii, and are consistent with the marginally non-uniform angular distributions we find above. That these structures might be sparse clusters is also consistent with their small measured velocity dispersions.

If we “run the clock backward” on structure formation, it seems unlikely that the huge walls and rich clusters seen locally would completely disperse into a uniform galaxy distribution by redshift of 1. It would be surprising if the high-redshift counterparts of local walls were *not* found in deep redshift surveys. (Rich clusters, on the other hand, only contain a small fraction of all galaxies and are much less likely to be encountered). Many groups have found similar or related structures. Bellanger & de Lapparent (1995), presenting the first results of the ESO-Sculptor Faint Galaxy Redshift Survey for galaxies with  $R < 20.5$  mag over  $0.28 \text{ deg}^2$ , have found what they call “walls” analogous to the Great Wall seen in the local universe. The galaxy distribution they see is strongly clustered in the line of sight, consisting of walls and voids, although their redshift histogram does not show structures as strongly peaked in redshift space as those presented here (for the reasons given above). A structure, interpreted as a normal dense galaxy cluster at high redshift, was found by LeFèvre et al (1994), consisting of a group of 12 galaxies around a quasar at  $z = 0.98$ , with velocity dispersion  $\sigma_v = 955 \text{ km s}^{-1}$  and transverse structure on a characteristic scale  $2 h^{-1}$  Mpc. Of course an observation of transverse structure is not an argument against wall morphology because if a sheet becomes self gravitating it must break up into structures that as wide as the wall is thick (Ostriker & Cowie 1981). Hutchings et al (1995) have detected compact groups of galaxies with a radius of  $\sim 1$  arcmin ( $0.25 h^{-1}$  Mpc) around 14 QSOs with  $z \sim 1.1$ . Ellingson et al (1991) (see also Ellingson & Yee 1994)

find galaxy clusters around QSOs with  $\sigma_v \sim 400\text{--}500 \text{ km s}^{-1}$ . Studies of quasar absorption lines in QSO pairs and in individual objects also suggest the existence of superclusters at  $z \approx 2.5$  (Dinshaw & Impey 1996).

### 4.3. Critical future observations

Given these observations and our interpretation, it is possible to predict the outcome of future observations which are crucial in determining the statistical, morphological, and physical properties of these structures. If they are wall-like, then we expect to see coherence as we extend the survey to adjacent fields. Even if the clumps are sparse clusters, local observations of Cor Bor and the Great Wall suggest that the clumps will group into large two-dimensional sheets. Redshift surveys in adjacent fields are essential to answering these questions of morphology.

There are many redshift surveys undertaken by different groups, and it is important to demonstrate that the differences among these surveys in prominence of structures in the  $z$  distribution is indeed attributable to differences in sample definition. It is imperative that the objects in this survey (and adjacent fields) be reselected with photometric and sparse-sampling criteria matched to other surveys; comparison could then be used to confirm that the structures are common and this field is not “special.” Also, the morphology-density relation (Dressler 1980) suggests that changing to bluer selection bands should reduce the percentage of galaxies lying inside the structures.

Finally, similar samples in widely separated fields will provide us with statistics for these structures, such as typical separations, velocity dispersions, filling factor, etc. It is possible that some of the much-discussed field-to-field variations found in galaxy counts, angular correlation functions, and redshift distributions (e.g., Koo & Kron, 1992) can be

attributed to the field-to-field variations in the walls along the line of sight.

If, as we strongly suspect they will, further observations do substantiate the view that roughly half the old galaxies are localized in walls, then a major challenge will be to determine whether the galaxies in these walls have virialized. At present there appear to be no good observational arguments against this view.

We are grateful to George Djorgovski, Keith Matthews, Gerry Neugebauer, Tom Soifer and Jim Westphal for helpful conversations and to Todd Small and Wal Sargent for permission to use their data on Cor Bor prior to publication. The entire Keck user community owes a huge debt to Bev Oke, Jerry Nelson, Gerry Smith, and many other people who have worked to make the Keck Telescope a reality. We are grateful to the W. M. Keck Foundation, and particularly its president, Howard Keck, for the vision to fund the construction of the W. M. Keck Observatory.

Table 1. Statistical Parameters for Redshift Peaks

$z_p$	No. Galaxies at $N_b(\text{max})$	$X_{max}$	Likelihood
0.392	9	3.4	0.5
0.429	12	6.4	0.005
0.581	20	12.4	0
0.675	8	5.3	0.02
0.766	6	5.0	0.2

Table 2. Velocity Dispersions in Redshift Peaks

$z_p$	$N^a$	$\sigma_v(N)$ (km s <sup>-1</sup> )	$\sigma_v(N - 1)^b$ (km s <sup>-1</sup> )
0.392	9	585	465
0.429	17	685	615
0.581	23	610	410
0.675	8	440	405
0.766	7	975	670

<sup>a</sup>Number of galaxies with  $z$  within 0.02 of  $z_p$ .

<sup>b</sup>Omitting the most discrepant galaxy, velocity dispersion as derived from the remaining  $N - 1$ .

Table 3. Results of 2D K-S Tests for Uniform Spatial Distribution

$z_p$	No. Galaxies at $N_b(\text{max})$	2D K-S Statistic $D$
0.392	9	0.01
0.429	12	0.2
0.581	20	0.01
0.675	8	0.4
0.766	6	0.1

## REFERENCES

- Bahcall, N.A., 1988, ARA&A, 26, 631
- Bellanger, C. & de Lapparent, V., 1995, ApJ, 455, L103
- Broadhurst, T., Ellis, R., Koo, D. & Szalay, A., 1990, Nature, 343, 726
- Burstein, D. & Heiles, C., 1982, AJ, 87, 1165
- Cohen, J.G. et al, (Paper III), 1996, manuscript in preparation
- Da Costa, N. et al, 1994, ApJ, L1
- de Lapparent, V., Geller, M. & Huchra, J.P., 1986, ApJ, 302, L1
- de Lapparent, V., Geller, M. & Huchra, J.P., 1991, ApJ, 369, 273
- Dinshaw, N. & Impey, C.D., 1996, ApJ in press
- Dressler, A. 1980, ApJ, 236, 351
- Ellingson, E., Green, R.F. & Yee, H.K.C., 1991, ApJ, 378, 476
- Ellingson, E. & Yee, H.K.C., 1994, ApJS, 92, 33
- Fasano, G. & Franceschini, A., 1987, MNRAS, 225, 155
- Glazebrook, K., Ellis, R., Colless, M., Broadhurst, T., Allington-Smith, J. & Tanvir, N.,  
1995, MNRAS, 273, 157
- Goia, I.M., Geller, M.J., Huchra, J.P., Maccarao, T., Steiner, J.E. & Stocke, J., 1982, ApJ,  
255, L17
- Griffiths, R.E. et al 1994, ApJ, 437, 67
- Hogg, D.W. et al 1996, manuscript in preparation
- Hutchings, J.B., Crampton, D. & Johnson, A., 1995, AJ, 109, 73
- Koo, D.C. & Kron, R.G., 1992, ARA&A, 30, 613

- LeFèvre, O., Crampton, D., Lilly, S.J., Hammer, F. & Tresse, L., 1994, ApJ, 423, L89
- LeFèvre, O., Crampton, D., Lilly, S.J., Hammer, F. & Tresse, L., 1995, ApJ 455, 60
- Lilly, S.J., LeFèvre, O., Crampton, D., Hammer, F. & Tresse, L., 1995, ApJ, 455, 50
- Oke, J.B., et al, 1995, PASP 107, 3750
- Ostriker, J. & Cowie L.L., 1981, ApJ, 243, L127
- Pahre, M.A. et al, (Paper II), 1996 manuscript in preparation
- Postman, M., Geller, M. & Huchra, 1988, AJ, 95, 267
- Postman, M., Huchra, J.P. & Geller, M.J., 1992, ApJ, 384,404
- Postman M.A., Lubin L.M., Gunn J.E., Oke J.B., Hoessel J.G., Schneider D.P., Christensen J.A., 1996, ApJ, in press
- Ramella, M., Geller, M.J. & Huchra, J.P., 1992, ApJ, 384, 396
- Sarazin, C., Rood, H. & Struble, M., 1982, AA, 108, L7
- Shectman, S., 1985, ApJS, 57. 77
- Small, T.A., 1996, PhD thesis, Caltech
- Songaila, A., Cowie, L.L., Hu, E.M. & Gardner, J.P., 1994, ApJS, 94, 461
- Zabludoff, A.I., Huchra. J.P. & Geller, M.J., 1990, ApJS, 74,1
- Zucca, E., Zamorani, G., Scaramella, R. & Vettolani, G., 1995, ApJ, 407, 470

Fig. 1.— The redshift histogram for the galaxies in our survey. The solid fill denotes galaxies whose redshifts are considered secure, while the open fill denotes galaxies whose redshifts are of lower precision.

Fig. 2.— The distribution of our sample of galaxies projected onto the sky is shown. The first panel shows the entire sample. This is followed by the spatial distribution of the galaxies in each of the 5 strongest peaks in the redshift histogram. The galaxies not included in the  $K < 20$  mag sample are shown as open circles, while those with uncertain  $z$  are shown as open crosses.



