

Related content

New matter effects in terrestrial neutrino oscillation experiments

To cite this article: Kathryn M. Zurek JHEP10(2004)058

View the [article online](#) for updates and enhancements.

- [A road map to solar neutrino fluxes, neutrino oscillation parameters, and tests for new physics](#)
John N. Bahcall and Carlos Peña-Garay
- [Solar neutrinos before and after Neutrino 2004](#)
John N. Bahcall, Maria C. Gonzalez-Garcia and Carlos Peña-Garay
- [Theory of neutrinos: a white paper](#)
R N Mohapatra, S Antusch, K S Babu et al.

Recent citations

- [Distorted neutrino oscillations from time varying cosmic fields](#)
Gordan Krnjaic *et al*
- [Exploring a nonminimal sterile neutrino model involving decay at IceCube](#)
Z. Moss *et al*
- [Probing nonstandard neutrino cosmology with terrestrial neutrino experiments](#)
Akshay Ghalsasi *et al*

RECEIVED: June 9, 2004

REVISED: October 13, 2004

ACCEPTED: October 22, 2004

New matter effects in terrestrial neutrino oscillation experiments

Kathryn M. Zurek

*Institute for Nuclear Theory, University of Washington
Seattle, WA 98195-1550, U.S.A.
E-mail: zurkat@u.washington.edu*

ABSTRACT: Consequences of a new force mediated by a light scalar particle for neutrino oscillation experiments are considered. Such a force could give rise to neutrino masses and mixings whose matter dependence for earth densities is much more significant than the MSW effect. We consider the effects of such a new force on the limits derived from oscillation experiments, and examine how the constraints on neutrino models are altered. Re-analysis of neutrino data, as well as new experiments in which large matter effects are systematically explored, is required to directly probe such physics beyond the standard model.

KEYWORDS: Solar and Atmospheric Neutrinos, Neutrino Physics.

JHEP10(2004)058

Contents

1. Introduction and motivation	1
2. New limits from medium dependence	3
3. Three neutrino models	5
4. Models consistent with LSND	9
4.1 3+1 models	10
4.2 2+2 models	11
5. Summary	11

1. Introduction and motivation

The last ten years have brought great progress with the first direct experimental evidence for the physics beyond the standard model, the discovery of neutrino masses and mixing. In the standard model, the neutrino is a massless particle with matter interactions mediated by W and Z bosons. Wolfenstein [1], Mikheev and Smirnov [2] were the first to show that these standard model interactions give rise to matter dependent masses and mixings from coherent forward scattering in the medium, and that these effects can be significant despite the small cross-sections for such interactions. Subsequently, many authors have considered the medium effects of beyond the standard model neutrino interactions, with an emphasis on flavor changing neutral interactions (for a recent analysis, see [3, 4] and references therein). These medium effects arise from new interactions whose energy scale is above the standard model, and have been shown to be weaker than the standard model interactions. The effects of a new scalar particle coupling only to neutrinos [5], neutrino mass arising from matter-neutrino interactions through such a scalar [6], and the matter effects of a massless string dilaton field coupled to neutrinos [7] have also been considered.

In this paper, we wish to consider matter effects on neutrinos due to interactions with a new, very light and weakly coupled scalar particle, the “acceleron”. In [8], such a new light particle was introduced as the connection between neutrinos and the dark energy, to explain the apparent coincidence that the dark energy scale (10^{-3} eV)⁴ is of about the same scale as the neutrino masses $\delta m_\nu \sim O(10^{-2}$ eV). It was also shown [9] that such new interactions with a light scalar particle could give rise to masses and mixings which are environment dependent. Here we parameterize such a new interaction by Yukawa couplings,

$$V = \lambda_N \phi \bar{N} N + \sum_i \lambda_{\nu_i} \phi \nu_i \nu_i + V_0(\phi), \quad (1.1)$$

where ϕ is the new scalar field, N the nucleon field, and ν_i are neutrino fields, written here in a basis where interaction with the scalar field is diagonal. A light scalar field coupling to neutrinos has been considered before, but without the coupling to matter [5]. In a mean field approximation, such an interaction, would give rise to a neutrino matter potential, which is an effective shift in the neutrino mass,

$$M_i^{\text{eff}} = \frac{\lambda_{\nu_i} \lambda_N}{m_\phi^2} n_N, \quad (1.2)$$

where n_N is the nucleon number density, $m_\phi^2 \equiv V_0''(\bar{\phi})$, and $\bar{\phi}$ is the vev of ϕ , the solution of $V'(\bar{\phi}) = 0$. This minimization equation is equivalent in the mean field approximation to

$$\lambda_N n_N + \sum_i \lambda_{\nu_i} n_{\nu_i} + V_0'(\bar{\phi}) = 0. \quad (1.3)$$

Now if V_0 is a sufficiently flat potential, which is the case if ϕ is a light particle, then $\bar{\phi}$ will vary significantly with n_N (and n_{ν_i} , which is not of interest for the environments considered here). This implies that there are two sources for density dependence in M_i^{eff} : not only the factor of n_N in M_i^{eff} , but also the dependence $m_\phi^2(\bar{\phi}(n_N))$. Therefore, we also note that the form of the density dependence will in general be different than that of the MSW term, which is simply proportional to n_e .

M_i^{eff} in turn modifies the effective neutrino hamiltonian H_{eff} , which in a two neutrino formalism reads,

$$H_{\text{eff}} = \frac{1}{2E} U \begin{pmatrix} (m_1 + M_1^{\text{eff}})^2 & M_3^{\text{eff}}{}^2 \\ M_3^{\text{eff}}{}^2 & (m_2 + M_2^{\text{eff}})^2 \end{pmatrix} U^\dagger + \begin{pmatrix} \sqrt{2}G_f n_e & \\ & \end{pmatrix}, \quad (1.4)$$

where U is the mixing matrix, $\sqrt{2}G_f n_e$ is the usual MSW term, m_i are the vacuum neutrino masses, and M_i^{eff} is some linear combination of those given in eq. (1.2). The result of these new terms in H_{eff} is that the effective mass splittings and mixings (determined from the eigenvalues and eigenvectors of the matrix), and their density dependences, are modified. The modification could be quite significant, as the new terms could yield effects potentially much larger than the standard MSW term. Consistency with gravitational force experiments requires the coupling in the new term $\lambda_N < 10^{-2} m_N/M_{Pl}$ [10]. This small coupling, however, can be compensated by n_N in typical earth densities 3g/cm^3 , and would give rise to a potential of size

$$M_i^{\text{eff}} = 1\text{eV} \left(\frac{\lambda_\nu}{10^{-1}} \right) \left(\frac{\lambda_N}{10^{-21}} \right) \left(\frac{\rho_N}{\rho_N^o} \right) \left(\frac{10^{-6}\text{eV}}{m_\phi} \right)^2, \quad (1.5)$$

where $\rho_N^o = 3\text{g/cm}^3$, so that M_i^{eff}/E could potentially be much larger than the MSW term for typical earth densities, which is of $O(10^{-13}\text{eV})$.

In this paper we will take the point of view that these non-standard matter effects do in fact give rise to masses and mixings that are significantly different in typical earth densities than in the vacuum. We will then consider how our interpretation of terrestrial neutrino oscillation experiments shifts within the framework of these new matter effects.

We wish to stress, however, that while such new matter effects could dominate at earth densities, they would be subdominant to, or of comparable size as, the MSW term near the level crossing point in the sun, hence reproducing the features of an MSW solution to the solar neutrino problem. This is possible because $M^{\text{eff}} \propto n_N$ only if $V_0(\phi)$ is a purely quadratic potential. For a more general form of the potential, $V_0(\phi) \sim \phi^n$, $m_\phi^2 = V_0''(\bar{\phi})$ is density dependent. This implies $M^{\text{eff}} \propto n_N^{1/(n-1)}$, so that the density dependence of the new terms may in fact be quite different from the MSW term. Hence the size and importance of the new terms relative to the MSW term will also be a density dependent quantity.

We also wish to note that, although such a new light particle was originally motivated in connection with the dark energy, non-standard interactions with any new light scalar particle, due to the long range nature of the force enhanced by n_N , could potentially give rise to new significant matter effects. As eq. (1.5) shows, these effects could be much more significant for typical earth densities than those considered from interactions with new heavy particles.

The outline of the paper is as follows. In the first section we make model independent considerations on how the interpretation of neutrino oscillation data is changed when large non-standard interactions are considered. In the second section we apply these new considerations to the standard three neutrino model, and discuss the implications for future experiments. Lastly, we specifically consider the range of possible models consistent with results from LSND.

2. New limits from medium dependence

For simplicity in the analysis, and because we wish to make our considerations independent of a model for how M_i^{eff} depends on density, we consider here only two media: air versus rock, or more generally, any High Density Material (HDM hereafter), which includes any combination of rock, steel, and concrete. We assume that the effects of rock or any HDM on the oscillation parameters are similar, and that they are significantly different from the air parameters. Additionally, because the analysis here is independent of a model for how M_i^{eff} depends on density, no limits from solar neutrinos are derived. To extract the vacuum neutrino masses and mixings from any solar neutrino data requires a model for how H_{eff} and hence $m_\phi^2(\bar{\phi})$ depends on density; whether the masses and mixing are significantly different from the terrestrial values is highly model dependent, and hence will not be considered here. We leave the analysis of solar neutrinos, including the relevant day/night effects, to future work [11]. Lastly, we assume that at the interface of two different media the sudden approximation is valid, so that the evolution of the neutrino phase is continuous. This is valid so long as the Compton wavelength of the neutrino mass in the medium is short compared to the distance traveled in the medium.

With these assumptions, we summarize in table 1 the experimental signals we have to date, their HDM and air pathlengths, and limits derived. There are several modifications to the limits derived when considering large, non-standard in-medium effects. First, as mentioned above, in order to derive limits from the solar neutrino data, we need a model

Signal	Channel	Environment	SI $\Delta m_{\min, \max}^2$ (eV 2)	Medium $\Delta m_{\min, \max}^2$ (eV 2)	ref.
SNO	$\nu_e \rightarrow \nu_e, \nu_\mu, \nu_\tau$	solar-interior	6.5×10^{-5}	8.2×10^{-5}	Unknown
Super-K(solar)	$\nu_e \rightarrow \nu_e, \nu_\mu$	solar-interior	3×10^{-5}	1.9×10^{-4}	Unknown
Super-K(atm)	$\nu_\mu \rightarrow \nu_x$	air/HDM	1.9×10^{-3}	3.0×10^{-3}	1.5×10^{-3}
KamLAND	$\nu_e \rightarrow \nu_x$	HDM	10^{-5}	10^{-4}	10^{-5}
K2K	$\nu_\mu \rightarrow \nu_x$	HDM	1×10^{-3}	4×10^{-3}	1×10^{-3}
LSND	$\nu_\mu \rightarrow \nu_e$	HDM	4×10^{-2}	1.2	4×10^{-2}
					1.2
Null Search	Channel	Environment	SI Δm_{\min}^2 (eV 2)	Medium Δm_{\min}^2 (eV 2)	ref.
KARMEN	$\nu_\mu \rightarrow \nu_e$	$\sim 50\%$ air	5×10^{-2}	0.1	[21]
Bugey	$\nu_e \rightarrow \nu_x$	air	10^{-2}	N/A	[23, 24]
CHOOZ	$\nu_e \rightarrow \nu_x$	$\sim 80 - 90\%$ air	7×10^{-4}	4×10^{-3}	[25, 26]
Palo Verde	$\nu_e \rightarrow \nu_x$	$\sim 95\%$ HDM	2×10^{-3}	2×10^{-3}	[27, 28]
CDHS	$\nu_\mu \rightarrow \nu_x$	Unknown	0.25	Unknown	[29]
NOMAD	$\nu_\mu \rightarrow \nu_\tau$	$\sim 60\%$ HDM	0.7	1.2	[30, 31]
	$\nu_e \rightarrow \nu_\tau$		5.9	9.8	
CHORUS	$\nu_\mu \rightarrow \nu_\tau$	$\sim 60\%$ HDM	0.6	1	[32, 31]
	$\nu_e \rightarrow \nu_\tau$		7.1	11.8	
Future Expmt.	Channel	Environment	SI Δm_{\min}^2 (eV 2)	Medium Δm_{\min}^2 (eV 2)	ref.
MiniBooNE	$\nu_\mu \rightarrow \nu_e$	HDM	2×10^{-2}	2×10^{-2}	[33]
OPERA	$\nu_\mu \rightarrow \nu_\tau$	HDM	10^{-3}	10^{-3}	[34]
MINOS	$\nu_\mu \rightarrow \nu_e, \nu_\mu, \nu_\tau$	HDM	10^{-3}	10^{-3}	[34]

Table 1: Range of allowed mass splittings for standard interactions (SI) versus large, medium dependent interactions. $\Delta m_{\min, \max}^2$ defines the range of values allowed by the signal at 90% C.L. Δm_{\min}^2 is the upper bound on Δm^2 at $\sin^2 2\theta = 1$ at 90% C.L. HDM is High Density Medium.

to determine how $V_0''(\bar{\phi})$, and hence M_i^{eff} , depends on density. Without such a model, we are unable to extract δm^2 and $\sin^2 2\theta$ from the solar neutrino data, as the non-standard dependences of δm^2 and $\sin^2 2\theta$ on solar density are unknown. One might argue that the concordance of the SNO result with the KamLAND reactor neutrino result, two experiments which occur in very different media, indicates that such new matter dependence must be very small. The agreement of the results may indicate that the new couplings considered here are indeed very small and that the standard neutrino picture is in fact correct. However, this need not be the case, as the standard MSW solution may not be highly sensitive to the new effects. For example, the new effects considered here, while leading to significant differences between HDM and air parameters, could saturate above earth densities, leading to only small changes in the MSW solution. In any case, this is a highly model dependent question, and in keeping with the binary HDM/air analysis we conduct here, this is left for future work [11].

Second, the standard analysis of the Super-K data utilizes an up-down ratio of ν_e and ν_μ events [12], one pathlength of which is mostly in air, and the other pathlength mostly in rock, which could potentially have very different mixings and splittings. To find the range of allowed parameters in HDM, one can instead make use of the ratio of upward stopping to upward through-going muon sample [13], with limits as given in the final column of table 1. Notice that these limits are considerably less restrictive than those of the whole set.

Third, for negative searches partially in rock or air, Δm_{\min}^2 is scaled by the fraction of the path in HDM. One can see this by noting that for small $\Delta m^2 L / 4E$, the amplitudes

are simply additive:

$$A(\nu_x \rightarrow \nu_y) = \langle \nu_x | U_n \cdots U_2 U_1 | \nu_y \rangle = A_n(\nu_x \rightarrow \nu_y) + \cdots + A_2(\nu_x \rightarrow \nu_y) + A_1(\nu_x \rightarrow \nu_y), \quad (2.1)$$

where U_n is the evolution matrix in medium n, and $A_n(\nu_x \rightarrow \nu_y)$ is the oscillation amplitude in medium n. To determine the bounds in the medium of interest, we set all other oscillation amplitudes to zero except in the medium of interest, and we find that in medium n

$$P_n(\nu_x \rightarrow \nu_y) \simeq (\Delta m_{\min}^2 L_n / 2E)^2, \quad (2.2)$$

so that Δm_{\min}^2 is simply scaled by the ratio of the pathlength in medium n, L_n , to the total neutrino pathlength. To determine $\sin^2 2\theta_{\min}$ from these null searches, we again take mixings in all other media except the one of interest to zero. Hence, at large Δm^2 ,

$$P(\nu_x \rightarrow \nu_y) = \frac{1}{2} \sin^2 2\theta_{\min}, \quad (2.3)$$

so that $\sin^2 2\theta_{\min}$ is unchanged. This applies, for example, to the CHOOZ reactor experiment, where the upper bound on Δm^2 in HDM is scaled by a factor of 5-10, as the experiment is 80–90% in air, but at large Δm^2 , $\sin^2 2\theta_{\min} = 0.1$ is unchanged. Therefore, the limit on ν_e disappearance in HDM for small Δm^2 is derived, not from CHOOZ, but from the Palo Verde reactor experiment ($\approx 95\%$ in HDM).

Finally, we note that we have no signals, and only constraints from null searches, for the neutrino parameters in air. Therefore, in what remains in this paper we will consider only the range of possible neutrino models admitted by the data in HDM. We will find that the range of possible models and allowed parameters is quite different from those admitted in the standard picture with standard interactions.

3. Three neutrino models

Within a three neutrino framework, the experimental signals given in table 1 are traditionally understood as follows. The parameter space is determined by two mass splittings and three mixing angles. One mixing and mass splitting is determined from the solar neutrino data, which is interpreted as the MSW conversion of $\nu_e \rightarrow \nu_\mu, \nu_\tau$ in the sun, with large mixing and mass splitting $O(10^{-5} \text{ eV}^2)$. The other mass splitting and a second mixing angle is determined from Super-K atmospheric data, which is interpreted as the vacuum oscillation of $\nu_\mu \rightarrow \nu_\tau$, with large mixing and mass splitting $O(10^{-3} \text{ eV}^2)$. The limit on the third mixing angle comes from the CHOOZ experiment $\sin^2 2\theta_{13} < 0.1$. The combination of these limits is illustrated in figure 1a.

The constraints, however, are significantly modified with the new interactions considered here in HDM. This is illustrated in figure 1b. The limits on ν_e disappearance in HDM are derived from KamLAND, with two allowed regions in the range $10^{-5} \text{ eV}^2 < \Delta m^2 < 10^{-4} \text{ eV}^2$, somewhat less restrictive than the limits from SNO $6.5 \times 10^{-5} \text{ eV}^2 < \Delta m^2 < 8.3 \times 10^{-5} \text{ eV}^2$. For ν_μ disappearance, the Super-K upward muon data constrains the parameters in HDM, which is less restrictive in both Δm^2 and $\sin^2 2\theta$ constraints than the




Figure 1: Allowed regions for $\nu_\mu \rightarrow \nu_x$ and $\nu_e \rightarrow \nu_x$ for (a) standard interactions and (b) non-standard medium dependent interactions in HDM (High Density Medium). The curves are parameterizations of the published allowed regions at 90% C.L., with the exception of the KamLAND data, which was published at 95% C.L.

full set utilizing an up-down ratio of events. But although we cite the upward muon data set as the best limits in HDM on ν_μ disappearance, bear in mind that even these limits may not be a rigorous measurement of the HDM parameters, as the analysis contains contamination from a horizon bin whose pathlength is mostly air. For a truly rigorous constraint, a reanalysis of the data would have to be conducted where the horizon bin is removed. The K2K experiment, though less restrictive in $\sin^2 2\theta$, offers the best constraint on Δm^2 in HDM at 90% C.L. The K2K experiment allows additional contours in the range $10^{-2} \text{ eV}^2 \lesssim \Delta m^2 \lesssim 10^{-1} \text{ eV}^2$ at 99% C.L. not shown in the plot.

These modifications in constraints have several implications for the interpretation of the experiments in a three neutrino model. In a conventional analysis CHOOZ limits $\sin^2 2\theta_{13} < 0.1$. With the new interactions considered here, however, the most obvious limit on $\sin^2 2\theta_{13}$ in HDM is derived from Palo Verde. For the allowed region of Super-K parameter space, $\Delta m^2 > 1.5 \times 10^{-3}$, it requires $\sin^2 2\theta_{13} \lesssim 0.5$, a rather unrestrictive limit. In addition, the Super-K experiment has not carried out an analysis on ν_e disappearance mixing angles with only rock pathlengths to restrict $\sin^2 2\theta_{13}$. The limit that we do have on atmospheric ν_e disappearance was derived in the full set utilizing an up-down ratio of events, with both air and rock pathlengths, in order to remove uncertainties as large as 20% in the atmospheric neutrino flux. For ν_e , this up-down asymmetry $A = (U - D)/(U + D)$, where U is the number of upward-going events and D is the number of downward-going events, is measured to be quite small $A_{\nu_e} = -0.009 \pm 0.042 \pm 0.005$ [35], indicating that to high certainty the observed and produced ν_e fluxes are the same.

To derive, however, the allowed mixings in air and rock from A_{ν_e} and A_{ν_μ} would require a new fit with six free parameters,

$$\Delta m_{\text{HDM}}^2, \quad \sin^2 2\theta_{13}^{\text{HDM}}, \quad \sin^2 2\theta_{23}^{\text{HDM}}, \quad \Delta m_{\text{air}}^2, \quad \sin^2 2\theta_{13}^{\text{air}}, \quad \sin^2 2\theta_{23}^{\text{air}}, \quad (3.1)$$

instead of two. As will be shown here, even an uncertainty of 5% in the ratio of observed to produced ν_e fluxes for maximally oscillated neutrinos could potentially allow $\sin^2 2\theta_{13}$ to be quite large. To quantify how large θ_{13} can be consistent with Super-K's lack of observation of ν_e oscillation, we write the ratio of the observed ν_e flux to the ν_e flux produced in the atmosphere:

$$\frac{\Phi_{\nu_e}}{\Phi_{\nu_e}^0} = \frac{\Phi_{\nu_\mu}^0}{\Phi_{\nu_e}^0} P(\nu_\mu \rightarrow \nu_e) + \frac{\Phi_{\nu_e}^0}{\Phi_{\nu_e}^0} P(\nu_e \rightarrow \nu_e). \quad (3.2)$$

We consider the limits from a maximally oscillated neutrino, which gives the tightest constraints on $\sin^2 2\theta_{13}$. Most of such neutrinos have energy ~ 1 GeV and are coming directly from the opposite side of the earth ($\cos \theta \sim -1$). For these neutrinos, the ratio of ν_μ to ν_e flux is [36]

$$\frac{\Phi_{\nu_\mu}^0}{\Phi_{\nu_e}^0} \simeq 2.5 \quad (3.3)$$

and is quite well known (uncertainty $\sim 5\%$). Lower energy neutrinos give a less restrictive bound on θ_{13} because $\Phi_{\nu_\mu}^0/\Phi_{\nu_e}^0$ is smaller at lower energies, while the flux of neutrinos with energies much above 1 GeV drops off rapidly while never becoming fully oscillated. Hence we expect the tightest constraint on θ_{13} to be derived for neutrinos with energy ~ 1 GeV. Substituting this ratio into the relevant oscillation probabilities, we find

$$\frac{\Phi_{\nu_e}}{\Phi_{\nu_e}^0} = 1 + \sin^2 2\theta_{13} (2.5 \sin^2 \theta_{23} - 1) \sin^2 x, \quad (3.4)$$

where we are using the usual conventions θ_{23} for the $\mu - \tau$ mixing and θ_{13} for e-disappearance, and $x = \Delta m_{\text{atm}}^2 L / 4E$. To get a rough estimate of the limits on θ_{13} in HDM from Super-K, we note again that ν_e are not observed to oscillate within some fractional uncertainty ϵ in flux, so we can approximate

$$1 - \epsilon < \frac{\Phi_{\nu_e}}{\Phi_{\nu_e}^0} < 1 + \epsilon. \quad (3.5)$$

For neutrinos which have maximally oscillated, $\sin^2 x = 1$, we find

$$-\epsilon < \sin^2 2\theta_{13} (2.5 \sin^2 \theta_{23} - 1) < \epsilon \quad (3.6)$$

Likewise, we can write for ν_μ disappearance

$$\frac{\Phi_{\nu_\mu}}{\Phi_{\nu_\mu}^0} = P(\nu_\mu \rightarrow \nu_\mu) + \frac{1}{\Phi_{\nu_\mu}^0 / \Phi_{\nu_e}^0} P(\nu_e \rightarrow \nu_\mu), \quad (3.7)$$

so that

$$\frac{\Phi_{\nu_\mu}}{\Phi_{\nu_\mu}^0} = 1 - (2.4 \sin^2 \theta_{23} \cos^2 \theta_{13} - 4 \sin^4 \theta_{23} \cos^4 \theta_{13} + 1.6 \sin^2 \theta_{23} \cos^4 \theta_{13}) \sin^2 x. \quad (3.8)$$




Figure 2: The curves marked $\epsilon = 0.05, 0.1, 0.2$ are the constraints from ν_e disappearance from Super-K given by eq. (3.6) for 5%, 10% and 20% uncertainty in the neutrino flux. The upper curve is the constraint for $\sin^2 2\theta_{atm}$ from eq. (3.9). The shaded regions are the overlap of these constraints which gives the allowed values of $\sin^2 2\theta_{23}$ and $\sin^2 2\theta_{13}$ from Super-K. Dark shaded is for $\epsilon = 0.05$, medium shaded for $\epsilon = 0.1$, and light shaded for $\epsilon = 0.2$.

The Super-K upward muon analysis requires $\sin^2 2\theta_{atm} > 0.8$. Then we have the second constraint

$$0.8 < 2.4 \sin^2 \theta_{23} \cos^2 \theta_{13} - 4 \sin^4 \theta_{23} \cos^4 \theta_{13} + 1.6 \sin^2 \theta_{23} \cos^4 \theta_{13} < 1.0. \quad (3.9)$$

Shown in figure 2 is the allowed regions given by eqs. (3.6), (3.9) for $\epsilon = 0.05, 0.1$ and 0.2. The overlap of the constraints gives the combined allowed region. We can see that while $\sin^2 2\theta_{23}$ is constrained to be quite large, there is much less constraint on $\sin^2 2\theta_{13}$ consistent with the data. In particular, for $\sin^2 2\theta_{23} = 1$, $\sin^2 2\theta_{13}$ could be as large as 0.2 for $\epsilon = 0.05$, or as large as 0.8 for $\epsilon = 0.2$. Such solutions can occur when ν_e depletion is canceled by replenishment through ν_μ oscillation to within an amount ϵ . We conclude that $\sin^2 2\theta_{13}$ could be significantly larger than the value $\sin^2 2\theta_{13} < 0.1$ derived from the standard analysis.

A large $\sin^2 2\theta_{13}$ would also indicate that KamLAND could observe very significant ν_e oscillation to the Super-K mass splitting. This possibility is excluded in the standard three neutrino model through the limits of the CHOOZ experiment, an experiment mostly in air, $\sin^2 2\theta_{13} < 0.1$. As we have shown here, however, the new best limits from Super-K, KamLAND, and Palo Verde cannot exclude this possibility in HDM, and in this case a three neutrino analysis of the KamLAND data would be necessary.

To narrow these allowed regions, a reanalysis of the full Super-K data set would need to be carried out allowing six free parameters on ν_μ and ν_e disappearance, as listed in eq. (3.1). Once this analysis has been carried out, one can determine whether the combination of these parameters with the Palo Verde limits constrain θ_{13} such that a three neutrino analysis of KamLAND is necessary. In addition, in the near future we will have additional signals when the MINOS and OPERA experiments measure Δm_{HDM}^2 , $\sin^2 2\theta_{23}^{HDM}$ through




Figure 3: LSND allowed region along with constraints on ν_e and ν_μ oscillations from null searches (a) Bugey and CDHS with standard interactions and (b) Palo Verde and CDHS with non-standard in-medium interactions. As the fraction of the CDHS beamline in HDM is unknown, the limits for non-standard interactions are at most as restrictive as those shown. The curves are parameterizations of the published allowed regions; in addition irregularities in the Bugey curve for large Δm^2 have been suppressed, as they are irrelevant for the analysis conducted here. Dashed lines indicate limits from null searches, and solid lines allowed regions from positive signals.

direct detection of the *appearance* of $\nu_\mu \rightarrow \nu_\tau$. MINOS will also be able to probe $\sin^2 2\theta_{13}^{\text{HDM}}$ through $\nu_\mu \rightarrow \nu_e$ down to mixing angles $\sin^2 2\theta_{13}^{\text{HDM}} \sim 0.1$. We have shown here that, with these new matter effects, the value of $\sin^2 2\theta_{13}^{\text{HDM}}$ could potentially be much larger than is now thought. Should MINOS find a value of $\sin^2 2\theta_{13}$ in HDM which is larger than allowed by CHOOZ experiment, we have a smoking gun for new matter effects in three neutrino models.

4. Models consistent with LSND

Next we consider how these new matter effects improve the fit of 3+1, 2+2 and three neutrino models to the data with LSND. Shown in figure 3a is the LSND allowed region for $\nu_\mu \rightarrow \nu_e$ oscillations, along with the best constraints for $\nu_e \rightarrow \nu_x$ and $\nu_\mu \rightarrow \nu_x$ oscillations from Bugey and CDHS. The modified constraints in HDM are shown in figure 3b. The Palo Verde experiment, not Bugey, constrains $\sin^2 2\theta_e$ in HDM. The limits from CDHS on ν_μ disappearance in HDM are unknown.¹ In addition, the KARMEN experiment, which is $\sim 50\%$ air no longer constrains LSND. We consider the implications of these new limits in HDM on 3+1 and 2+2 models with LSND, which are disfavored with standard interactions.

¹My thanks to Joe Rothberg and his collaborators at CERN for looking into beam path information. The lapse of time between the experiment and the present, however, has not allowed them to track this information down.

4.1 3+1 models

These medium effects improve the fit of 3+1 models to the KamLAND, Super-K, Bugey, Palo Verde, and LSND results.

In typical 3+1 models, we have three splittings, $O(10^{-5} \text{ eV}^2)$, $O(10^{-3} \text{ eV}^2)$, and the larger LSND splitting $O(0.1 - 1 \text{ eV}^2)$. With only standard interactions, 3+1 models are disfavored [37], and a 3+2 scenario [38] improves the fit somewhat, but only by contriving to set the masses of the sterile neutrino in the region where the constraints from CDHS and Bugey are weakest.

The reason that these models are disfavored is that the combination of the Bugey and CDHS limits on ν_e and ν_μ result in a $\mu - e$ mixing angle which is too small at 90% C.L. to accommodate the LSND result. To quantify this observation, we write the ν_e and ν_μ disappearance angles measured by Bugey and CDHS in terms of the fraction of ν_e or ν_μ in the LSND mass splitting, U_{e4} and $U_{\mu4}$:

$$\sin^2 2\theta_e = 4(1 - |U_{e4}|^2) |U_{e4}|^2 \quad (4.1)$$

$$\sin^2 2\theta_\mu = 4(1 - |U_{\mu4}|^2) |U_{\mu4}|^2. \quad (4.2)$$

Likewise, the LSND $\mu - e$ mixing angle is written in terms of these same quantities

$$\sin^2 2\theta_{\mu e} = 4 |U_{e4}|^2 |U_{\mu4}|^2 = (1 - \cos 2\theta_\mu)(1 - \cos 2\theta_e). \quad (4.3)$$

When the limits from Bugey and CDHS on $\sin^2 2\theta_e$ and $\sin^2 2\theta_\mu$, as shown in figure 3b, are substituted into these expressions, we find that the resulting allowed mixing $\sin^2 2\theta_{\mu e}$ is too small at 90% C.L. to accommodate the LSND result. There is only a small region of LSND parameter space, $0.2 < \Delta m^2 < 0.3$, which, although excluded at 90% C.L. is allowed at 99% C.L. Hence these fits are only marginally allowed within a small region of parameter space.

We immediately see how the situation changes with these new matter effects. Bugey, an experiment mostly in air, can no longer constrain LSND. Instead, the constraint on ν_e disappearance in HDM comes from Palo Verde $\sin^2 2\theta_e < 0.17$, which corresponds to the region of LSND parameter space $\Delta m^2 > 0.1 \text{ eV}^2$. These limits are considerably less restrictive than $\sin^2 2\theta_e < 0.04$ from Bugey, and the result is that the combined limits of Palo Verde and CDHS do not disfavor the LSND result.

The Palo Verde limit implies that if MiniBooNE sees $e - \mu$ oscillations, it should see them with $\Delta m^2 > 0.1 \text{ eV}^2$. Furthermore, if oscillations are detected at MiniBooNE in the region $0.1 \text{ eV}^2 < \Delta m^2 < 0.25 \text{ eV}^2$ (excluded by Bugey in a conventional analysis), we have strong indication for non-standard matter interactions for neutrinos.

An additional prediction of this model is that a reactor experiment, with similar sensitivity as Bugey, but with neutrino pathlengths instead in HDM, would detect ν_e disappearance, provided the CDHS limits are rigorous in rock. The reason is that the combination of CDHS and Bugey limits with standard interactions disfavor at 90% C.L. mixings large enough to accommodate the LSND result. We would expect a signal in the approximate range $0.04 < \sin^2 2\theta < 0.17$, $0.1 \text{ eV}^2 < \Delta m^2 < 1 \text{ eV}^2$. The caveat here is that the CDHS limits be applicable in HDM; or, equivalently, that some other experiment (e.g. Super-K)

limit ν_μ disappearance in HDM, for the mass splittings of interest for LSND, at least as much as CDHS for a given Δm^2 . Currently the material in the neutrino path of CDHS is unknown. If the experiment had been partially in air, such that $\Delta m_{\min}^{2CDHS}/r \gtrsim 0.3$, where r is the fraction of the neutrino pathlength in HDM, and no other experiment places as tight of limits on ν_μ disappearance as the standard CDHS limits, we would have no prediction for a signal from such a reactor experiment.

4.2 2+2 models

Analysis of the SNO, KamLAND, and Super-K data strongly disfavors 2+2 oscillation schemes in the standard interaction picture [37, 39, 40, 41, 42] because 2+2 models with LSND require large mixing with sterile states, which are disfavored by the SNO, KamLAND and Super-K atmospheric data.

To determine the fit of 2+2 models to the data in HDM, a new analysis would need to be carried out. The existing analyses distinguish between active-active versus active-sterile oscillations through the MSW effect, which couples only to active neutrinos. These analyses, however, are only valid if the MSW effect is the dominant matter effect in earth, and hence do not constrain active-sterile mixing in the case that the new non-standard interactions considered here dominate. A new analysis with the new interactions, however, would be highly model dependent, as the form of the potential due to the new interactions must be known. As we have been carrying out only a model independent analysis for such new interactions, we will not perform such an analysis here. They could, however, easily be carried out once the form of the potential is known.

5. Summary

We have examined the effects on neutrino oscillations of a very light scalar mediating a force between neutrinos and ordinary matter, as proposed in [9]. As these new matter effects could be much larger than the MSW effect for typical earth densities, we have considered how these new interactions would significantly modify the constraints on the neutrino parameters in any High Density Medium (HDM). I summarize in table 2 for each parameter the allowed range and experimental signals with standard interactions versus these new in-medium non-standard interactions. In particular, indications for these new matter effects from already planned and running experiments include:

- $\sin^2 2\theta_{13}^{\text{HDM}}$ with a value larger in HDM than the limit given by the CHOOZ experiment;
- Δm_{12}^2 , $\sin^2 2\theta_{12}$ detected by KamLAND in a range not allowed by the SNO results;
- Δm_{23}^2 , $\sin^2 2\theta_{23}$ detected by K2K, OPERA, and MINOS with parameters somewhat modified from the Super-K result;
- Detection of ν_e disappearance in HDM in the region excluded by Bugey, should MiniBooNE confirm the LSND signal.

Any of these signals would be a smoking gun for the new matter effects considered here.

Model	Parameter	SI range		Experiment	HDM range		Experiment
3 neutrino	$\sin^2 2\theta_{13}$	0	0.1	CHOOZ	0	0.5	Super-K, KamLAND K2K, MINOS Reactor expt.
	$\sin^2 2\theta_{12}$	0.76	0.88	SNO	0.4	1	KamLAND
	$\sin^2 2\theta_{23}$	0.92	1	Super-K	0.8	1	Super-K, K2K MINOS, OPERA
	Δm_{12}^2	6.5×10^{-5}	8.2×10^{-5}	SNO	10^{-3}	10^{-5}	KamLAND
	Δm_{23}^2	1.9×10^{-3}	3.0×10^{-3}	Super-K	1×10^{-3}	6×10^{-3}	Super-K, K2K MINOS, OPERA
3+1 neutrino	$\sin 2\theta_{\mu e}$	disfavored		CDHS, Bugey	10^{-3}	10^{-1}	MiniBooNE Reactor expt.
2+2 neutrino	$\sin 2\theta_{\mu s}$	disfavored		Super-K	0	1	MINOS, OPERA
	$\sin 2\theta_{e s}$	disfavored		SNO	0	1	no experiment

Table 2: Parameters of interest for each type of model. Standard interactions (SI) upper and lower limits (90% C.L.) and the experiment which gives that limit (columns three and four). Present HDM limits and experiments which are of interest for measuring these parameters in the near future (columns five and six).

These upcoming and currently running experiments promise to determine the parameters quite accurately in rock. We have, however, no signal for the parameters in air, as well as no information on the form of the density dependence. Without a theory for the density dependence of such a force, we are unable to constrain the parameters from solar neutrinos, and the constraints from short baseline and atmospheric neutrino experiments could be significantly different if the neutrino parameters are sensitively dependent to small variations in density.

Experiments are needed to determine if and how sensitively the neutrino parameters vary from such a new force. In the past it has been assumed that new matter effects were arising from physics above the standard model energy scale, and for typical earth densities were unimportant. It is suggested here that a new attitude be taken in which large non-standard in-medium dependences are searched for in terrestrial neutrino experiments. We may find that terrestrial neutrino oscillations provide the tool to directly explore a wide range of physics beyond the standard model.

Acknowledgments

I am grateful to David Kaplan, Ann Nelson and Neal Weiner for many helpful comments along the way, and for proposing the problem in the first place. Much thanks also to Jeff Wilkes, Richard Gran and Kiyoshi Shiraishi for useful discussions on Super-K, and to Rob Fardon for reading various drafts and making useful suggestions. This work was supported by the US Department of Energy grant DE-FG03-00ER41132.

References

- [1] L. Wolfenstein, *Neutrino oscillations in matter*, *Phys. Rev. D* **17** (1978) 2369.

- [2] S.P. Mikheev and A.Y. Smirnov, *Resonance enhancement of oscillations in matter and solar neutrino spectroscopy*, Sov. J. Nucl. Phys. **42** (1985) 913.
- [3] M.M. Guzzo, P.C. de Holanda and O.L.G. Peres, *Effects of non-standard neutrino interactions on MSW-LMA solution*, Phys. Lett. **B 591** (2004) 1 [[hep-ph/0403134](#)].
- [4] A. Friedland, C. Lunardini and C. Pena-Garay, *Solar neutrinos as probes of neutrino-matter interactions*, Phys. Lett. **B 594** (2004) 347 [[hep-ph/0402266](#)].
- [5] J. Stephenson, G. J., T. Goldman and B.H.J. McKellar, *MSW-like enhancements without matter*, Mod. Phys. Lett. **A 12** (1997) 2391 [[hep-ph/9610317](#)].
- [6] R.F. Sawyer, *Do neutrinos have mass only within matter?*, Phys. Lett. **B 448** (1999) 174 [[hep-ph/9809348](#)].
- [7] R. Horvat, *Medium effects in string-dilaton-induced neutrino oscillations*, Phys. Rev. **D 58** (1998) 125020 [[hep-ph/9802377](#)].
- [8] R. Fardon, A.E. Nelson and N. Weiner, *Dark energy from mass varying neutrinos*, [astro-ph/0309800](#).
- [9] D.B. Kaplan, A.E. Nelson and N. Weiner, *Neutrino oscillations as a probe of dark energy*, Phys. Rev. Lett. **93** (2004) 091801 [[hep-ph/0401099](#)].
- [10] E.G. Adelberger, B.R. Heckel and A.E. Nelson, *Tests of the gravitational inverse-square law*, Ann. Rev. Nucl. Part. Sci. **53** (2003) 77 [[hep-ph/0307284](#)].
- [11] A. Nelson and N. Weiner, ongoing work, 2004.
- [12] SUPER-KAMIOKANDE collaboration, T. Toshito, *Super-Kamiokande atmospheric neutrino results*, [hep-ex/0105023](#).
- [13] SUPER-KAMIOKANDE collaboration, Y. Fukuda et al., *Neutrino-induced upward stopping muons in Super-Kamiokande*, Phys. Lett. **B 467** (1999) 185 [[hep-ex/9908049](#)].
- [14] SNO collaboration, S.N. Ahmed et al., *Measurement of the total active ${}^8\text{B}$ solar neutrino flux at the Sudbury Neutrino Observatory with enhanced neutral current sensitivity*, Phys. Rev. Lett. **92** (2004) 181301 [[nucl-ex/0309004](#)].
- [15] SUPER-KAMIOKANDE collaboration, M.B. Smy et al., *Precise measurement of the solar neutrino day/night and seasonal variation in Super-Kamiokande-I*, Phys. Rev. **D 69** (2004) 011104 [[hep-ex/0309011](#)].
- [16] M. Ishitsuka, *L/E analysis of the atmospheric neutrino data from Super-Kamiokande*, Ph.D. thesis, University of Tokyo, 2004.
- [17] E. Kearns, *Status and future of atmospheric experiments*, talk given at *Weak Interactions and Neutrinos-2003*, Lake Geneva, Wisconsin, U.S.A.
- [18] KAMLAND collaboration, T. Araki et al., *Measurement of neutrino oscillation with KamLAND: evidence of spectral distortion*, [hep-ex/0406035](#).
- [19] K2K collaboration, M.H. Ahn et al., *Indications of neutrino oscillation in a 250 Km long-baseline experiment*, Phys. Rev. Lett. **90** (2003) 041801 [[hep-ex/0212007](#)].
- [20] T. Nakaya, *K2K results*, talk given at *Neutrino 2004*.
- [21] KARMEN collaboration, J. Wolf, *Final LSND and KARMEN-2 neutrino oscillation results*, prepared for International Europhysics Conference on High-Energy Physics (HEP 2001), Budapest, Hungary, 12-18 Jul 2001.

- [22] B. Louis, private communication.
- [23] Y. Declais et al., *Search for neutrino oscillations at 15 meters, 40 meters and 95 meters from a nuclear power reactor at bugey*, *Nucl. Phys. B* **434** (1995) 503.
- [24] M. Abbes et al., *The Bugey-3 neutrino detector*, *Nucl. Instrum. Meth. A* **374** (1996) 164.
- [25] M. Apollonio et al., *Search for neutrino oscillations on a long base-line at the CHOOZ nuclear power station*, *Eur. Phys. J. C* **27** (2003) 331 [[hep-ex/0301017](#)].
- [26] D. Steinberg, private communication.
- [27] F. Boehm et al., *Final results from the Palo Verde neutrino oscillation experiment*, *Phys. Rev. D* **64** (2001) 112001 [[hep-ex/0107009](#)].
- [28] F. Boehm, private communication.
- [29] F. Dydak et al., *A search for muon-neutrino oscillations in the δm^2 range 0.3 eV^2 to 90 eV^2* , *Phys. Lett. B* **134** (1984) 281.
- [30] NOMAD collaboration, P. Astier et al., *Final nomad results on $\nu_\mu \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_{\text{tau}}$ oscillations including a new search for nu/tau appearance using hadronic tau decays*, *Nucl. Phys. B* **611** (2001) 3 [[hep-ex/0106102](#)].
- [31] NOMAD collaboration, J. Altegoer et al., *The NOMAD experiment at the CERN SPS*, *Nucl. Instrum. Meth. A* **404** (1998) 96.
- [32] CHORUS collaboration, E. Eskut et al., *New results from a search for $\nu_\mu \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\tau$ oscillation*, *Phys. Lett. B* **497** (2001) 8.
- [33] MINIBOONE collaboration, G. McGregor, *The MiniBooNE experiment: Status and plans*, *AIP Conf. Proc.* **655** (2003) 58.
- [34] MINOS AND OPERA collaboration, A. Weber, *Status of the LBL oscillation experiments MINOS and OPERA*, prepared for International Europhysics Conference on High Energy Physics (HEP 2001), Budapest, Hungary, 12-18 Jul. 2001.
- [35] A. Kibayashi, *Atmospheric neutrino oscillation analysis using Fluka 3D flux and Super-Kamiokande data*, Ph.D. thesis, University of Hawaii 2002.
- [36] V. Agrawal, T.K. Gaisser, P. Lipari and T. Stanev, *Atmospheric neutrino flux above 1 GeV*, *Phys. Rev. D* **53** (1996) 1314 [[hep-ph/9509423](#)].
- [37] M. Maltoni, T. Schwetz and J.W.F. Valle, *Status of four-neutrino mass schemes: a global and unified approach to current neutrino oscillation data*, *Phys. Rev. D* **65** (2002) 093004 [[hep-ph/0112103](#)].
- [38] M. Sorel, J.M. Conrad and M. Shaevitz, *A combined analysis of short-baseline neutrino experiments in the (3+1) and (3+2) sterile neutrino oscillation hypotheses*, [hep-ph/0305255](#).
- [39] SUPER-KAMIOKANDE collaboration, A. Habig, *Discriminating between $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_{\text{sterile}}$ in atmospheric ν_μ oscillations with the Super-Kamiokande detector*, [hep-ex/0106025](#).
- [40] SNO collaboration, Q.R. Ahmad et al., *Measurement of day and night neutrino energy spectra at sno and constraints on neutrino mixing parameters*, *Phys. Rev. Lett.* **89** (2002) 011302 [[nucl-ex/0204009](#)].

- [41] J.N. Bahcall, M.C. Gonzalez-Garcia and C. Peña-Garay, *If sterile neutrinos exist, how can one determine the total ${}^8\text{B}$ and ${}^7\text{Be}$ solar neutrino fluxes?*, *Phys. Rev.* **D 66** (2002) 035802 [[hep-ph/0204194](#)].
- [42] P.C. de Holanda and A.Y. Smirnov, *Searches for sterile component with solar neutrinos and KamLAND*, [hep-ph/0211264](#).