

Supporting Information for ”Global estimates of the energy transfer from the wind to the ocean, with emphasis on near-inertial oscillations”

M. Mar Flexas,¹ Andrew F. Thompson¹, Hector S. Torres², Patrice Klein², J.

Thomas Farrar³, Hong Zhang^{4,2}, and Dimitris Menemenlis²

Corresponding author: Mar Flexas, Environmental Science and Engineering, California Institute of Technology, 1200 California Blvd., Pasadena, CA 91125 (marf@caltech.edu)

¹California Institute of Technology,
Pasadena, CA 91125

²Jet Propulsion Laboratory, California
Institute of Technology, Pasadena,
California

³Woods Hole Oceanographic Institution,
Woods Hole, Massachusetts.

⁴Joint Institute for Regional Earth
System Science and Engineering, University
of California Los Angeles, Los Angeles,
California, USA.

Contents of this file

1. Figures S1 to S2

Introduction The supporting information included in this document consists of two figures. Figure S1 shows the frequency-wavenumber cospectrum in linear space. Figure S2 shows seasonal maps of surface inertial currents from LLC 1/48 .

Figure S1. Frequency-wavenumber cospectrum of surface kinetic energy fluxes (in W m^{-2}) in three target regions in JFM and JAS months. The three target regions are (a,b) the Kuroshio Extension [31 N–46 N, 156 W–176 W], (c,d) NE Atlantic [38 N–53 N, 23 W–8 E], and (e,f) Southern Ocean, east of Kerguelen Plateau [32 S–47 S, 98 E–113 E]

Figure S2. Seasonal (three-month averaged) maps of surface near-inertial currents (in m s^{-1}) from LLC 1/48 .




Figure S1: Frequency-wavenumber cospectrum of surface kinetic energy fluxes (in W m^{-2}) in three target regions in JFM and JAS months. The three target regions are (a,b) the Kuroshio Extension [31 N–46 N, 156 W–176 W], (c,d) NE Atlantic [38 N–53 N, 23 W–8 E], and (e,f) Southern Ocean, east of Kerguelen Plateau [32 S–47 S, 98 E–113 E].

