

Eigenstate entanglement in the Sachdev-Ye-Kitaev model

Yichen Huang (黄溢辰)*

Institute for Quantum Information and Matter, California Institute of Technology, Pasadena, California 91125, USA

Yingfei Gu (顾颖飞)†

Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA

(Dated: July 10, 2019)

We study the entanglement entropy of eigenstates (including the ground state) of the Sachdev-Ye-Kitaev model. We argue for a volume law, whose coefficient can be calculated analytically from the density of states. The coefficient depends on not only the energy density of the eigenstate but also the subsystem size. Very recent numerical results of Liu, Chen, and Balents confirm our analytical results.

I. INTRODUCTION

Entanglement, a concept of quantum information theory, has been widely used in condensed matter and high energy physics [2, 40, 41, 48] to provide insights beyond those obtained via “conventional” quantities. For ground states of (geometrically) local Hamiltonians, it characterizes quantum criticality [3, 4, 17, 29, 51] and topological order [26, 30, 31]. The scaling of entanglement [10] is quantitatively related to the classical simulability of quantum many-body systems [15, 18, 37, 44, 50].

Besides ground states, it is also important to understand the entanglement of excited eigenstates. Significant progress has been made in this regard [1, 5, 7–9, 14, 19, 20, 22, 23, 33, 36, 43, 52, 53, 55]. For chaotic local Hamiltonians, one expects a volume law: The entanglement entropy of an eigenstate between a subsystem (smaller than half the system size) and its complement scales as the subsystem size with a coefficient depending on the energy density of the eigenstate. Indeed, analytical arguments [7] and numerical simulations [8, 14, 43] strongly suggest that it is to leading order equal to the thermodynamic entropy of the subsystem at the same energy density.

Let us put this result in context. The eigenstate thermalization hypothesis (ETH) states that for expectation values of local observables, a single eigenstate resembles a thermal state with the same energy density [6, 38, 49]. The equivalence of entanglement and thermodynamic entropies is a variant of ETH for the von Neumann entropy.

Over the past few years, the Sachdev-Ye-Kitaev (SYK) model [25, 35, 42] has become a very active research topic in condensed matter and high energy physics. As a quantum mechanical model of $N \gg 1$ Majorana fermions with random all-to-all interactions, it is exactly solvable in the large- N limit, and has an extensive zero-temperature entropy. Entanglement and ETH in the SYK model have recently been studied [12, 16, 21, 27, 47]. In particular, the numerical results of Fu and Sachdev [12] suggest the

breakdown of the equivalence of entanglement and thermodynamic entropies in the SYK model: The ground-state entanglement entropy appears to be slightly less than the maximum ($\frac{1}{2} \ln 2$ per Majorana fermion) and is significantly greater than the zero-temperature entropy.

We argue that due to all-to-all interactions, the equivalence of entanglement and thermodynamic entropies should be modified as follows: The entanglement entropy of an eigenstate (including the ground state) for a subsystem smaller than half the system size is to leading order equal to the thermodynamic entropy of the subsystem at a different energy density, which depends on not only the energy density of the eigenstate but also the subsystem size. This allows us to derive an analytical expression for the scaling of the eigenstate entanglement entropy in the SYK model from the density of states.

The SYK model is maximally chaotic [25, 35] in the sense of the maximum “Lyapunov exponent” [34] for the exponential growth of out-of-time-ordered correlators [24, 28, 39, 45, 46]. Our argument is not related to this dynamical behavior, and should apply to a broad class of chaotic quantum many-body systems.

II. PRELIMINARIES

Definition 1 (entanglement entropy). The entanglement entropy of a bipartite pure state ρ_{AB} is defined as the von Neumann entropy

$$S(\rho_A) = -\text{tr}(\rho_A \ln \rho_A) \quad (1)$$

of the reduced density matrix $\rho_A = \text{tr}_B \rho_{AB}$.

Consider a quantum mechanical system of N Majorana fermions $\chi_1, \chi_2, \dots, \chi_N$, where $N \gg 1$ is an even number. The Hamiltonian of the SYK model is

$$H = \sum_{1 \leq i < j < k < l \leq N} J_{ijkl} \chi_i \chi_j \chi_k \chi_l, \quad \{\chi_i, \chi_j\} = \delta_{ij}, \quad (2)$$

where the coefficients are independent real Gaussian random variables with zero mean $\overline{J_{ijkl}} = 0$ and variance

$$\overline{J_{ijkl}^2} = \frac{3!}{N^3}. \quad (3)$$

* yichen.huang@microsoft.com

† yingfei_gu@g.harvard.edu

We summarize the spectral properties of the SYK model in the thermodynamic limit $N \rightarrow +\infty$:

- The mean energy of H is $\text{tr} H/d = 0$, where $d = 2^{N/2}$ is the dimension of the Hilbert space.
- The distribution of eigenvalues near the mean energy is asymptotically normal with variance $\text{tr}(H^2)/d = \Theta(N)$. Asymptotic normality is a general feature of models with random few-body interactions [11].
- The ground-state energy is $-NE_0 + o(N)$ (extensive), where $E_0 > 0$ is a known constant.
- The density of states at energy NE is [13]

$$D_N(E) \sim e^{\left[\frac{1}{2} \ln 2 - \frac{1}{16} \arcsin^2\left(\frac{E}{E_0}\right)\right]N}. \quad (4)$$

This is an excellent approximation away from the edges of the spectrum, i.e., when $E_0 - |E|$ is not too small.

III. ARGUMENT

We divide the system into two subsystems A and B . Subsystem A consists of M Majorana fermions, where M is even. Assume without loss of generality that $M \leq N/2$. We split the Hamiltonian (2) into three parts:

$$H = H_A + H_\partial + H_B, \quad (5)$$

where $H_{A(B)}$ contains terms acting only on subsystem $A(B)$, and H_∂ consists of cross terms. We observe that

$$H'_A := \left(\frac{N}{M}\right)^{\frac{3}{2}} H_A \quad (6)$$

is the SYK model of M Majorana fermions.

Lemma 1 ([54]). *The thermal state maximizes the von Neumann entropy among all states with the same energy.*

Proof. For completeness, we give a simple proof of this well-known fact. Let

$$\sigma_\beta := \frac{1}{Z_\beta} e^{-\beta H}, \quad Z_\beta := \text{tr} e^{-\beta H} \quad (7)$$

be a thermal state at inverse temperature β . For any density matrix ρ , the relative entropy

$$S(\rho \parallel \sigma_\beta) := \text{tr}(\rho \ln \rho - \rho \ln \sigma_\beta) \geq 0 \quad (8)$$

is nonnegative. If ρ and σ_β have the same energy, then

$$\begin{aligned} S(\rho) &= -\text{tr}(\rho \ln \rho) \leq -\text{tr}(\rho \ln \sigma_\beta) = \beta \text{tr}(\rho H) + \ln Z_\beta \\ &= \beta \text{tr}(\sigma_\beta H) + \ln Z_\beta = -\text{tr}(\sigma_\beta \ln \sigma_\beta) = S(\sigma_\beta). \end{aligned} \quad (9)$$

This completes the proof of Lemma 1. \square

Lemma 2. *To leading order,*

$$S(\rho_A) \leq \ln D_M \left(\frac{\text{tr}(\rho_A H'_A)}{M} \right) \quad (10)$$

for any density matrix ρ_A of subsystem A .

Proof. Consider the SYK model H'_A . The inverse temperature β at which the thermal state $\sigma_\beta := e^{-\beta H'_A} / \text{tr} e^{-\beta H'_A}$ has the same energy as ρ_A is obtained by solving

$$\frac{\int_{-E_0}^{E_0} M \epsilon e^{-\beta M \epsilon} D_M(\epsilon) d\epsilon}{\int_{-E_0}^{E_0} e^{-\beta M \epsilon} D_M(\epsilon) d\epsilon} = \text{tr}(\rho_A H'_A). \quad (11)$$

The saddle-point approximation, which becomes exact in the limit $M \rightarrow +\infty$, gives

$$\beta = \left. \frac{\partial \ln D_M(\epsilon)}{M \partial \epsilon} \right|_{\epsilon = \frac{\text{tr}(\rho_A H'_A)}{M}}. \quad (12)$$

Using Lemma 1,

$$\begin{aligned} S(\rho_A) &\leq S(\sigma_\beta) = \ln \int_{-E_0}^{E_0} e^{\beta(\text{tr}(\rho_A H'_A) - M\epsilon)} D_M(\epsilon) d\epsilon \\ &= \ln D_M \left(\frac{\text{tr}(\rho_A H'_A)}{M} \right). \end{aligned} \quad (13)$$

In the last step, we used the saddle-point approximation and Eq. (12). \square

Theorem 1. *Let $|\psi\rangle$ be an eigenstate of the SYK model (2) with energy NE . To leading order,*

$$S(\rho_A) \leq \ln D_M \left(\left(\frac{M}{N} \right)^{\frac{3}{2}} E \right), \quad (14)$$

where $\rho_A = \text{tr}_B |\psi\rangle\langle\psi|$ is the reduced density matrix.

Proof. To leading order, the quartic Hamiltonian (2) has $N^4/4!$ terms, $M^4/4!$ of which are in H_A . In average,

$$\text{tr}(\rho_A H_A) = \left(\frac{M}{N} \right)^4 NE. \quad (15)$$

Substituting Eq. (6),

$$\text{tr}(\rho_A H'_A) = \left(\frac{M}{N} \right)^{\frac{3}{2}} NE = \left(\frac{M}{N} \right)^{\frac{3}{2}} ME. \quad (16)$$

Hence, Theorem 1 follows from Lemma 2. \square

A slight modification of the proof(s) of Theorem 1 (and Lemma 2) yields

Theorem 2. *For (translationally invariant) local Hamiltonians, the entanglement entropy of an eigenstate between two subsystems is to leading order upper bounded by the thermodynamic entropy of the smaller subsystem at the same energy density.*

Remark. The upper bound in Theorem 2 holds regardless of whether the system is chaotic. Notably, it is not tight in (integrable) free-fermion systems [52]. In chaotic systems, analytical arguments [7] and numerical simulations [8, 14, 43] strongly suggest that the bound is attained.

Since the SYK model is chaotic, one might expect that the upper bound in Theorem 1 is attained. Thus, we have a volume law

$$S(\rho_A) \sim \left[\frac{1}{2} \ln 2 - \frac{1}{16} \arcsin^2 \left(\left(\frac{M}{N} \right)^{\frac{3}{2}} \frac{E}{E_0} \right) \right] M \quad (17)$$

with a coefficient depending on E (the energy density of the eigenstate) and M (the size of the smaller subsystem). This is the leading-order scaling of the eigenstate entanglement entropy, and we are unable to calculate the subleading corrections.

Example 1. Suppose that N is a multiple of 4. For $M = N/2$, the ground-state entanglement entropy scales as

$$\begin{aligned} S(\rho_A) &\sim \left(\frac{1}{2} \ln 2 - \frac{1}{16} \arcsin^2 \frac{1}{2\sqrt{2}} \right) M \\ &\approx (0.34657 - 0.00816)M = 0.33841M. \end{aligned} \quad (18)$$

IV. CONCLUSIONS AND OUTLOOK

In the SYK model, we have argued that the entanglement entropy of an eigenstate with energy NE for a subsystem of size $M \leq N/2$ is to leading order equal to the thermodynamic entropy of the subsystem at energy $(M/N)^{3/2}ME$. Therefore,

- The entanglement entropy of an eigenstate obeys a volume law with the maximum coefficient $\frac{1}{2} \ln 2$

if the subsystem size is a vanishing fraction of the system size. This is because the subsystem is at the mean energy density of the Hamiltonian (6).

- The entanglement entropy of an eigenstate with finite energy density obeys a volume law with a non-maximal coefficient if the subsystem size is a constant fraction of the system size.

In the future, it would be interesting to study the Renyi entanglement entropy of eigenstates of the SYK model. As a generalization of entanglement entropy, the Renyi entanglement entropy reflects the entanglement spectrum (the full spectrum of the reduced density matrix ρ_A). See Refs. [9, 14, 33] for recent results on the Renyi entanglement entropy of eigenstates of chaotic local Hamiltonians.

ACKNOWLEDGMENTS

We would like to thank Xiao-Liang Qi for pointing out Lemma 1. Y.H. acknowledges funding provided by the Institute for Quantum Information and Matter, an NSF Physics Frontiers Center (NSF Grant PHY-1733907). Y.H. is also supported by NSF DMR-1654340. Y.G. is supported by the Gordon and Betty Moore Foundation EPiQS Initiative through Grant GBMF-4306.

Note added.—Very recently, we became aware of related work by Liu *et al.* [32], which studied eigenstate entanglement in the SYK model using different methods. Among other results, the ground-state entanglement entropy was calculated up to $N = 44$ Majorana fermions using exact diagonalization. For $M = N/2$ (the subsystem size is half the system size), the data are well fitted by the expression $0.3375M - 0.666$, which is consistent with our analytical result (18).

-
- [1] B. Bauer and C. Nayak. Area laws in a many-body localized state and its implications for topological order. *Journal of Statistical Mechanics: Theory and Experiment*, 2013(9):P09005, 2013.
 - [2] L. Bombelli, R. K. Koul, J. Lee, and R. D. Sorkin. Quantum source of entropy for black holes. *Physical Review D*, 34(2):373–383, 1986.
 - [3] P. Calabrese and J. Cardy. Entanglement entropy and quantum field theory. *Journal of Statistical Mechanics: Theory and Experiment*, 2004(06):P06002, 2004.
 - [4] P. Calabrese and J. Cardy. Entanglement entropy and conformal field theory. *Journal of Physics A: Mathematical and Theoretical*, 42(50):504005, 2009.
 - [5] L. D’Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol. From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics. *Advances in Physics*, 65(3):239–362, 2016.
 - [6] J. M. Deutsch. Quantum statistical mechanics in a closed system. *Physical Review A*, 43(4):2046–2049, 1991.
 - [7] J. M. Deutsch. Thermodynamic entropy of a many-body energy eigenstate. *New Journal of Physics*, 12(7):075021, 2010.
 - [8] J. M. Deutsch, H. Li, and A. Sharma. Microscopic origin of thermodynamic entropy in isolated systems. *Physical Review E*, 87(4):042135, 2013.
 - [9] A. Dymarsky, N. Lashkari, and H. Liu. Subsystem eigenstate thermalization hypothesis. *Physical Review E*, 97(1):012140, 2018.
 - [10] J. Eisert, M. Cramer, and M. B. Plenio. *Colloquium: Area laws for the entanglement entropy. Reviews of Modern Physics*, 82(1):277–306, 2010.
 - [11] L. Erdős and D. Schröder. Phase transition in the density of states of quantum spin glasses. *Mathematical Physics, Analysis and Geometry*, 17(3-4):441–464, 2014.
 - [12] W. Fu and S. Sachdev. Numerical study of fermion and boson models with infinite-range random interactions. *Physical Review B*, 94(3):035135, 2016.
 - [13] A. M. García-García and J. J. M. Verbaarschot. Analyt-

- ical spectral density of the Sachdev-Ye-Kitaev model at finite N . *Physical Review D*, 96(6):066012, 2017.
- [14] J. R. Garrison and T. Grover. Does a single eigenstate encode the full Hamiltonian? *Physical Review X*, 8(2):021026, 2018.
- [15] S. Gharibian, Y. Huang, Z. Landau, and S. W. Shin. Quantum Hamiltonian complexity. *Foundations and Trends in Theoretical Computer Science*, 10(3):159–282, 2015.
- [16] Y. Gu, A. Lucas, and X.-L. Qi. Spread of entanglement in a Sachdev-Ye-Kitaev chain. *Journal of High Energy Physics*, 2017(9):120, 2017.
- [17] C. Holzhey, F. Larsen, and F. Wilczek. Geometric and renormalized entropy in conformal field theory. *Nuclear Physics B*, 424(3):443–467, 1994.
- [18] Y. Huang. *Classical simulation of quantum many-body systems*. PhD thesis, University of California, Berkeley, 2015.
- [19] Y. Huang. Universal eigenstate entanglement of chaotic local Hamiltonians. *Nuclear Physics B*, 938:594–604, 2019.
- [20] Y. Huang and J. E. Moore. Excited-state entanglement and thermal mutual information in random spin chains. *Physical Review B*, 90(22):220202, 2014.
- [21] N. Hunter-Jones, J. Liu, and Y. Zhou. On thermalization in the SYK and supersymmetric SYK models. *Journal of High Energy Physics*, 2018(2):142, 2018.
- [22] D. A. Huse, R. Nandkishore, V. Oganesyan, A. Pal, and S. L. Sondhi. Localization-protected quantum order. *Physical Review B*, 88(1):014206, 2013.
- [23] J. P. Keating, N. Linden, and H. J. Wells. Spectra and eigenstates of spin chain Hamiltonians. *Communications in Mathematical Physics*, 338(1):81–102, 2015.
- [24] A. Kitaev. Hidden correlations in the Hawking radiation and thermal noise. In *2015 Breakthrough Prize in Fundamental Physics Symposium*, 2014. <https://www.youtube.com/watch?v=0Q9qN8j7EZI>, <http://online.kitp.ucsb.edu/online/entangled15/kitaev/>.
- [25] A. Kitaev. A simple model of quantum holography. In *KITP Program: Entanglement in Strongly-Correlated Quantum Matter*, 2015. <http://online.kitp.ucsb.edu/online/entangled15/kitaev/>.
- [26] A. Kitaev and J. Preskill. Topological entanglement entropy. *Physical Review Letters*, 96(11):110404, 2006.
- [27] I. Kourkoulou and J. Maldacena. Pure states in the SYK model and nearly- AdS_2 gravity. arXiv:1707.02325, 2017.
- [28] A. I. Larkin and Y. N. Ovchinnikov. Quasiclassical method in the theory of superconductivity. *Sov. Phys. JETP*, 28(6):1200–1205, 1969.
- [29] J. I. Latorre, E. Rico, and G. Vidal. Ground state entanglement in quantum spin chains. *Quantum Information and Computation*, 4(1):48–92, 2004.
- [30] M. Levin and X.-G. Wen. Detecting topological order in a ground state wave function. *Physical Review Letters*, 96(11):110405, 2006.
- [31] H. Li and F. D. M. Haldane. Entanglement spectrum as a generalization of entanglement entropy: Identification of topological order in non-abelian fractional quantum Hall effect states. *Physical Review Letters*, 101(1):010504, 2008.
- [32] C. Liu, X. Chen, and L. Balents. Quantum entanglement of the Sachdev-Ye-Kitaev models. *Physical Review B*, 97(24):245126, 2018.
- [33] T.-C. Lu and T. Grover. Renyi entropy of chaotic eigenstates. *Physical Review E*, 99(3):032111, 2019.
- [34] J. Maldacena, S. H. Shenker, and D. Stanford. A bound on chaos. *Journal of High Energy Physics*, 2016(8):106, 2016.
- [35] J. Maldacena and D. Stanford. Remarks on the Sachdev-Ye-Kitaev model. *Physical Review D*, 94(10):106002, 2016.
- [36] Y. O. Nakagawa, M. Watanabe, H. Fujita, and S. Sugiura. Universality in volume-law entanglement of scrambled pure quantum states. *Nature Communications*, 9:1635, 2018.
- [37] T. J. Osborne. Hamiltonian complexity. *Reports on Progress in Physics*, 75(2):022001, 2012.
- [38] M. Rigol, V. Dunjko, and M. Olshanii. Thermalization and its mechanism for generic isolated quantum systems. *Nature*, 452(7189):854–858, 2008.
- [39] D. A. Roberts, D. Stanford, and L. Susskind. Localized shocks. *Journal of High Energy Physics*, 2015(3):51, 2015.
- [40] S. Ryu and T. Takayanagi. Aspects of holographic entanglement entropy. *Journal of High Energy Physics*, 2006(8):45, 2006.
- [41] S. Ryu and T. Takayanagi. Holographic derivation of entanglement entropy from the anti-de Sitter space/conformal field theory correspondence. *Physical Review Letters*, 96(18):181602, 2006.
- [42] S. Sachdev and J. Ye. Gapless spin-fluid ground state in a random quantum Heisenberg magnet. *Physical Review Letters*, 70(21):3339–3342, 1993.
- [43] L. F. Santos, A. Polkovnikov, and M. Rigol. Weak and strong typicality in quantum systems. *Physical Review E*, 86(1):010102, 2012.
- [44] N. Schuch, M. M. Wolf, F. Verstraete, and J. I. Cirac. Entropy scaling and simulability by matrix product states. *Physical Review Letters*, 100(3):030504, 2008.
- [45] S. H. Shenker and D. Stanford. Multiple shocks. *Journal of High Energy Physics*, 2014(12):46, 2014.
- [46] S. H. Shenker and D. Stanford. Multiple shocks in scrambling. *Journal of High Energy Physics*, 2015(5):132, 2015.
- [47] J. Sonner and M. Vielma. Eigenstate thermalization in the Sachdev-Ye-Kitaev model. *Journal of High Energy Physics*, 2017(1):149, 2017. <http://online.kitp.ucsb.edu/online/entangled15/kitaev/>.
- [48] M. Srednicki. Entropy and area. *Physical Review Letters*, 71(5):666–669, 1993.
- [49] M. Srednicki. Chaos and quantum thermalization. *Physical Review E*, 50(2):888–901, 1994.
- [50] F. Verstraete and J. I. Cirac. Matrix product states represent ground states faithfully. *Physical Review B*, 73(9):094423, 2006.
- [51] G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev. Entanglement in quantum critical phenomena. *Physical Review Letters*, 90(22):227902, 2003.
- [52] L. Vidmar, L. Hackl, E. Bianchi, and M. Rigol. Entanglement entropy of eigenstates of quadratic fermionic Hamiltonians. *Physical Review Letters*, 119(2):020601, 2017.
- [53] L. Vidmar and M. Rigol. Entanglement entropy of eigenstates of quantum chaotic Hamiltonians. *Physical Review Letters*, 119(22):220603, 2017.
- [54] A. Wehrl. General properties of entropy. *Reviews of Modern Physics*, 50(2):221–260, 1978.
- [55] Z.-C. Yang, C. Chamon, A. Hamma, and E. R. Mucciolo. Two-component structure in the entanglement spectrum of highly excited states. *Physical Review Letters*, 115(26):267206, 2015.