

anisotropic and nonlinear acoustic effects might also be directly studied.

3) Information regarding the pressure or density modulation of the ground-state and excited-state absorption in the crystal.

The amount of information simultaneously obtained from measurements such as these, concerning both molecular and crystalline properties, is sizable. We hope this process can be naturally extended to other materials as well, providing a unified understanding of their properties.

#### ACKNOWLEDGMENT

Earlier work on the transient grating experiments was supported by the Joint Services Electronics Program at Stanford University and by the National Science Foundation. This work grew out of studies of energy transport in pentacene carried out in collaboration with Prof. M. Fayer and D. Dlott of the Stanford Chemistry Department, and we appreciate the use of molecular crystal samples provided by them.

#### REFERENCES

- [1] D. W. Phillion, D. J. Kuizenga, and A. E. Siegman, "Subnanosecond relaxation time measurements using a transient induced grating method," *Appl. Phys. Lett.*, vol. 27, pp. 85-87, 1975.
- [2] J. R. Salcedo, A. E. Siegman, D. D. Dlott, and M. D. Fayer, "Dynamics of energy transport in molecular crystals: The picosecond transient grating method," *Phys. Rev. Lett.*, vol. 41, pp. 131-133, July 10, 1978; also in *Picosecond Phenomena*, C. V. Shank, E. P. Ippen, and S. L. Shapiro, Eds. Berlin: Springer Series in Chem. Phys., vol. 4, 1978, pp. 240-243.
- [3] D. C. Auth, "New high-power source of coherent microwave phonons," *Appl. Phys. Lett.*, vol. 16, pp. 521-523, June 15, 1970.
- [4] D. W. Phillion, "Evolution of mode-locked pulses and their application to time-resolved spectroscopy," Ph.D. dissertation, Stanford Univ., Stanford, CA, Nov. 1974.
- [5] H. Kogelnik, "Coupled wave theory for thick hologram gratings," *Bell Syst. Tech. J.*, vol. 48, pp. 2909-2947, 1969.
- [6] A. E. Siegman, "Bragg diffraction of a Gaussian beam by a crossed-Gaussian volume grating," *J. Opt. Soc. Amer.*, vol. 67, pp. 545-550, 1977.
- [7] D. R. Dean, "Optically induced diffraction gratings in liquids and solids," Office of Naval Research, Washington, DC, Tech. Rep. TR-73-02.
- [8] M. E. Mack, "Stimulated thermal Rayleigh scattering with picosecond pulses," *Ann. N.Y. Acad. Sci.*, vol. 168, pp. 419-436, 1970.
- [9] R. Herman and M. Gray, "Theoretical prediction of the stimulated thermal Rayleigh scattering in liquids," *Phys. Rev. Lett.*, vol. 19, pp. 824-827, 1967.
- [10] J. M. Donnini, "Effect d'une compression sur les spectres de l'anthracène et du pentacene en solution dans le p-terphenyle a 1.7 K," *J. Chim. Phys.*, II, vol. 11-12, pp. 1542-1545, 1974.
- [11] H. Eichler and H. Stahl, "Time and frequency behavior of sound waves thermally induced by modulated laser pulses," *J. Appl. Phys.*, vol. 44, pp. 3429-3435, Aug. 1973.

## Notes and Lines

### Correction to "Phase Conjugate Optics and Real-Time Holography"

AMNON YARIV

In the above paper,<sup>1</sup> page 656, column 1, Fig. 12 should be replaced with the following figure and caption.




Fig. 12. (a) A holographic exposure and (b) reconstruction.

Manuscript received December 18, 1978.

The author is with the Departments of Electrical Engineering and Applied Physics, California Institute of Technology, Pasadena, CA 91125.

<sup>1</sup>A. Yariv, *IEEE J. Quantum Electron.*, vol. QE-14, pp. 650-660, Sept. 1978.